UC Irvine

UC Irvine Previously Published Works

Title

Ergosterol peroxide from *Rhizoctonia repens*: Composition, conformation, and origin

Permalink

<https://escholarship.org/uc/item/6hq7c2w5>

Journal

Journal of the Chemical Society, Chemical Communications, 0(22)

ISSN

0022-4936

Authors

Arditti, J
Fisch, REMH
Flick, BH

Publication Date

1972-12-01

DOI

10.1039/C39720001217

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Ergosterol Peroxide from *Rhizoctonia repens*: Composition, Conformation, and Origin

By J. ARDITTI,* R. ERNST M. H. FISCH, and B. H. FLICK

(Department of Developmental and Cell Biology, University of California, Irvine, California 92664)

Summary Ergosterol peroxide (ep), synthetic or from fungal extracts, is shown to be a mixture of isomers arising from α - and β -attack by $O_2(^1\Delta_g)$ on ergosterol; the stereochemistry of the peroxide bridge is assigned.

THE origin of ergosterol peroxide (ep), a relatively rare sterol isolated from a number of fungal extracts,^{1a,b} has been questioned. The gummy residue from extractions contains pigments which may function as photosensitizers in the presence of light and air, generating singlet oxygen ($^1\Delta_g$),¹ after which ep is produced by symmetry-allowed

attack² on endogenous ergosterol. The hypothesis was verified for two systems (*Piptoporus betulinus* and *Daedalea quercina*) and it was suggested that ep may be an artifact rather than a natural product.^{1b} We now report new information which supports this contention.

Extraction of fungal mats of *Rhizoctonia repens* M 32 yielded ep (0.03% t.l.c., u.v.) When crude extracts are left, the yield rises, but some ep is always present, even when extraction is followed immediately by analysis.

The stereochemistry of peroxide attachment is known to be 5α , 8α as originally inferred from the rule of rear attack.^{3,4}

However, if an electrophilic reagent has relatively low steric requirements, C-5 approach is not exclusively from the α face.⁵ *A priori*, it seemed reasonable that singlet oxygen ($^1\Delta_g$) would be small enough to permit attack from both faces. Ep was prepared⁶ to give a product which melted sharply and had the properties reported. Signals of the ring B vinyl protons were two superimposed AB patterns (J 9 Hz for both) centred at δ (CDCl₃ vs Me₄Si) 6.33 (large) and 6.38 p.p.m. (small). We attributed these two patterns to the 5α , 8α and 5β , 8β isomers respectively. In spectra run at 220 MHz, the two patterns were partially resolved. Integration showed the minor component to constitute 16% of the mixture.

The n.m.r. data could have had an alternate explanation: *viz.*, conformers of the peroxide bridge, (1) and (2), both derived from α -attack, which interconvert at a rate slow on the n.m.r. time scale. Models suggested (1) and (2) to be of comparable energy as required by this hypothesis. (A 15:85 mixture corresponds to a difference of one kilocalorie per mole in free energy at 25°.)

By the addition of dilute solutions of Eu(dpm)₃,⁷ shifts of as much as 40 Hz in the C-6 and C-7 ep signals could be obtained, and the upfield doublet was always shifted more than the downfield doublet in both AB patterns. Such behaviour is not consistent with a mixture of (1) and (2) but would be expected from (1) and (3) if the proton closest to oxygen is always the downfield one, *i.e.* if the proximate C-O bond is deshielding.⁸ Such is the case for ascaridole, (4),⁹ where the vinyl protons are shifted some 0.2 p.p.m. downfield from the vinyl absorption in the model compound, bicyclo[2,2,2]octene.^{9b} Since the C-6 (more

shifted) proton corresponds to the upfield resonance in the AB pattern, the stereochemistry of the peroxide bridge is as in (1) and (3).

Comparison with "naturally occurring" ep shows that both samples are the same mixture. This finding strongly suggests that the "natural" material also arises from attack of singlet oxygen on ergosterol and hence is an artifact.

We thank Mr. Paul A. Kroon, California Institute of Technology for assistance with the 220 MHz spectra and Dr. J. Nüesch, ETH, Zürich, for a culture of *Rhizoctonia repens* M 32. This work was supported by the Office of Naval Research (US).

(Received, 31st July 1972; Com. 1335.)

¹ (a) *Inter alia*, P. Wieland and V. Prelog, *Helv. Chim. Acta*, 1947, **30**, 1028; (b) H. K. Adam, I. M. Campbell, and N. J. McCorkindale, *Nature*, 1967, **216**, 397.

² A. U. Kahn and D. R. Kearns, *J. Chem. Phys.*, 1968, **48**, 3272; *Advances in Chemistry Series*, 77, American Chemical Society, Washington, D.C., 1968, p. 143; D. R. Kearns, *J. Amer. Chem. Soc.*, 1969, **91**, 6554.

³ L. Fieser, *Experientia*, 1950, **6**, 312.

⁴ The chemical evidence is summarized in L. Fieser and M. Fieser, 'Steroids,' Reinhold, New York, 1959, pp. 244-246.

⁵ *Inter alia*, the reaction of cholest-5-ene with perbenzoic acid demonstrably yields some β -epoxide. H. B. Henbest and T. I. Wrigley, *J. Chem. Soc.*, 1957, 4596.

⁶ A. Windaus and J. Brunken, *Annalen*, 1928, **460**, 225.

⁷ C. C. Hinkley, *J. Amer. Chem. Soc.*, 1969, **91**, 5160.

⁸ J. W. Emsley, J. Feeney, and L. H. Sutcliffe, 'High Resolution Nuclear Magnetic Resonance Spectroscopy,' Pergamon, Oxford, 1965, pp. 136-138.

⁹ (a) Varian n.m.r. catalogue number 276; (b) D. Swern, A. H. Clements, and T. M. Luong, *Anal. Chem.*, 1969, **41**, 412.