UCSF

UC San Francisco Previously Published Works

Title

A randomized trial to assess the effect of a research informational pamphlet on telephone survey completion rates among older Latinos.

Permalink

<https://escholarship.org/uc/item/6kd0p8f3>

Journal

Contemporary clinical trials, 33(4)

ISSN

1551-7144

Authors

Santoyo-Olsson, Jasmine
Phan, Lisa
Stewart, Anita L
et al.

Publication Date

2012-07-01

DOI

10.1016/j.cct.2012.03.004

Peer reviewed

Published in final edited form as:

Contemp Clin Trials. 2012 July ; 33(4): 624–627. doi:10.1016/j.cct.2012.03.004.

A Randomized Trial to Assess the Effect of a Research Informational Pamphlet on Telephone Survey Completion Rates among Older Latinos

Jasmine Santoyo-Olsson, MS^{1,2}, Lisa Phan, BA³, Anita L. Stewart, PhD², Celia Kaplan, DrPH, MA¹, Gina Moreno-John, MD, MPH¹, and Anna M. Nápoles, PhD, MPH¹

¹Medical Effectiveness Research Center for Diverse Populations and the Center for Aging in Diverse Communities, Division of General Internal Medicine, Department of Medicine, University of California San Francisco (UCSF), San Francisco, CA, USA

²Institute for Health and Aging, UCSF, San Francisco, CA, USA

³Marin Community Clinics, San Rafael, CA, USA

Abstract

Purpose of the study—To assess the effects of receiving a research informational pamphlet produced by the federal Office for Human Research Protections on telephone survey participation rates of older Latinos, and illustrate the feasibility of nesting recruitment studies within other funded studies when stand alone funding for recruitment studies is limited.

Design and Methods—Latino patients aged 50 with 1 visit during the preceding year (N=1,314) were sampled from three community clinics and a multi-specialty medical group. Patients were randomly assigned to receive or not receive a pamphlet that contained information on research participation in the initial mailing for the telephone survey study. Survey participation rates were compared between the pamphlet and no pamphlet groups.

Results—In a multivariate model, women (OR=1.4; 95% CI 1.1, 1.8), and those with public insurance (vs. no insurance; OR=1.7; 95% CI 1.1, 2.5) were more likely to participate, while those age 65+ (vs. age 50–54; OR=0.6; 95% CI 0.4, 0.8) were less likely to participate; there was no significant difference by pamphlet group (OR=0.8; 95% CI 0.7, 1.1). Nesting of the randomized trial of the recruitment pamphlet within the funded study required minimal additional resources.

© 2012 Elsevier Inc. All rights reserved.

The corresponding author for this manuscript is: Jasmine Santoyo-Olsson, MS, University of California San Francisco, 3333 California Street, Suite 335, San Francisco, CA 94118-0856, USA, Phone: (415) 514-3355, Fax: (415) 502-8291, jasmine.santoyo-olsson@ucsf.edu.

Lisa Phan, BA, Marin Community Clinics, 3110 Kerner Blvd., San Rafael, CA 94901, USA, Phone: 415-526-8500, lisaph222@yahoo.com

Anita L. Stewart, PhD, University of California San Francisco, 3333 California Street, Suite 340, San Francisco, CA 94118-0646, USA, Phone: 415-502-5207, Fax: 415-502-5208, anita.stewart@ucsf.edu

Celia Kaplan, DrPH, MA, University of California San Francisco, 3333 California Street, Suite 335, San Francisco, CA 94118-0856, USA, Phone: 415-502-5601, Fax: 415-502-8291, celia.kaplan@ucsf.edu

Gina Moreno-John, MD, MPH, University of California San Francisco, 400 Parnassus Avenue, Box 0320, San Francisco, CA 94143-0320, USA, Phone: 415-476-4624, Fax: 415-476-7964, ginam@medicine.ucsf.edu

Anna M. Nápoles, PhD, MPH, University of California San Francisco, 3333 California Street, Suite 335, San Francisco, CA 94118-0856, USA, Phone: 415-476-6290, Fax: 415-502-8291, anapoles@ucsf.edu

Conflicts of Interest: The authors have no financial or other conflicts of interest to disclose.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Implications—Recruitment methods that are more intensive than a pamphlet may be needed to enhance survey participation rates among older Latinos. Nesting recruitment trials within funded studies is a promising and efficient approach for testing recruitment strategies.

Keywords

Latinos; Hispanics; randomized trial; response rate; research participation

1. Introduction

Increasing the participation of ethnic minorities in clinical research is necessary to ensure findings are applicable to these groups. There are few randomized trials comparing the effectiveness of recruitment methods, especially among ethnically diverse groups [1–5]. The recruitment literature consists predominantly of descriptive, observational studies. The federal government has invested in promoting participation in clinical research through the development and distribution of an informational pamphlet by the Office for Human Research Protections (OHRP) titled, “Becoming a Research Volunteer: It’s Your Decision [6].” This pamphlet was designed to address a commonly cited need among diverse populations for more information on research [7,8].

Clearing up misconceptions by providing comprehensible information facilitates an informed decision about research participation. When awareness and opportunity to participate exist [10], minority groups participate in research, including clinical trials, at rates that are comparable to mainstream populations [11]. Distributing information on the nature of research with an invitation to participate in a specific study may be a cost-effective way to enhance participation of underrepresented groups, especially among Latinos who tend to be less familiar with research, but who also tend to be less likely than Whites to refuse to participate once they are better informed [11,12].

An efficient way to build evidence on the efficacy of recruitment methods is to conduct nested studies of these methods within funded studies. The purpose of this study is to: 1) test the effect of the OHRP informational pamphlet on participation in a telephone survey of colorectal cancer (CRC) screening among older Latinos; and 2) demonstrate how health disparities researchers can nest recruitment trials within their funded studies.

2. Material and Methods

2.1 Sample

A randomized trial testing the OHRP brochure’s effect on response rates was nested within a cross-sectional telephone survey study of cultural factors and CRC screening [13]. The sampling frame was derived from electronic databases from four clinics. Inclusion criteria were: 1) self-reported Latino ethnicity; 2) age 50 or older; 3) English- or Spanish-speaking; 4) 1 primary care visit in the preceding year; and 4) no personal history of CRC.

2.2 Materials

The OHRP launched a public campaign promoting research participation, which included distribution of the pamphlet titled, “Becoming a Research Volunteer: It’s Your Decision.” We elected to test the pamphlet’s effect on survey response rates because it offered a low-cost tool that might enhance participation in the parent study. Also, despite the significant investment in the production and distribution of the pamphlet, it had never been formally evaluated. The pamphlet covers what research is, its role in prevention, diagnostic, and treatment advances, and questions for potential participants to ask researchers to enable an

informed decision about participation. It uses simple language, has photos of ethnically diverse individuals, and is available in English and Spanish.

2.3 Procedures

Stratifying by site, we randomized half the sampling frame to receive a standard initial mailing (control group) and half to receive the standard mailing plus the English and Spanish versions of the pamphlet (experimental group). The standard mailing included: 1) a bilingual letter inviting them to a telephone survey about CRC screening; 2) a bilingual, low-literacy study information sheet; and 3) a postage-paid opt-in/out postcard. Participants received \$25 for completing the survey. University of California San Francisco's and a hospital's Institutional Review Boards approved the study.

2.4 Measures

Experimental condition (pamphlet or no pamphlet) was the main predictor. Sociodemographic variables included gender, age, type of health insurance, and clinic site. No other demographic variables were available on the sampling frame.

A dichotomous outcome measure indicated whether or not patients completed the telephone survey. Those receiving the pamphlet were asked four questions: 1) "Did you receive a pamphlet with the letter? The pamphlet was called "Becoming a Research Volunteer" (yes or no); 2) "Did you read the pamphlet?" (yes or no); 3) If "yes" to #2, "In what language(s)?" (Spanish and/or English); and 4) "How much did the pamphlet influence your decision whether or not to participate in the phone survey?" (none, a little, quite a bit, or a lot).

2.5 Statistical Analysis

We compared gender, age, and health insurance characteristics of experimental and control groups using chi-square analysis. We examined the effect of the pamphlet and sociodemographic factors on survey completion using chi-square analyses. Multivariate logistic regression analysis assessed the independent effects of the pamphlet and sociodemographic characteristics on survey completion. Persons who were ineligible, deceased, or for whom we had an incorrect address ($N=88$) were excluded. An additional 36 individuals were dropped due to missing data on health insurance, leaving a final sample of 1,190.

3. Results

Of 1,314 letters mailed to potential participants, 504 completed the survey, for an overall response rate of 38%. The denominator includes patients who were never reached (31%), refused (24%), or were ineligible (7%). Sixty-four percent of participants were women, 30% were age 65 or older, and 44% had no health insurance (Table 1).

There were no differences between the pamphlet and no pamphlet groups on gender ($p=0.62$), age ($p=0.79$), health insurance ($p=0.54$), or clinic site ($p=0.71$). There was no significant difference in survey completion rate between the pamphlet (39%) and no pamphlet group (43%). Latinas were more likely (44%) than Latino men (35%; $p<0.01$) to complete the survey. Latinos age 65 and older were less likely (36%) to complete the survey than those age 55–64 (45%) or 50–54 (42%) ($p<0.05$). Latinos with private health insurance were less likely (35%) than those with no insurance (46%) ($p<0.01$) to complete the survey.

In multivariate analysis, the pamphlet was not associated with survey completion (OR = 0.8; 95% CI 0.7, 1.1). In the total sample, controlling for pamphlet group and other covariates, women (OR=1.4; 95% CI 1.1, 1.8) and those with public insurance (vs. no insurance; OR=1.7; 95% CI 1.1, 2.5) were more likely to complete the survey, while those aged 65+ (vs. aged 50–54; OR=0.6; 95% CI 0.4, 0.8) were less likely to complete the survey (Table A. 1).

Among those receiving the pamphlet, only 47% (n=116) remembered it, substantially less than the proportion that recalled receiving the letter (69%; n=172). Women were more likely (55%) than men (33%) to remember the pamphlet ($p < .001$); those age 65 and older (37%) were less likely than those age 55–64 (45%) or age 50–54 (59%) to remember the pamphlet ($p < .05$). Among those who remembered the pamphlet, women (79%) were more likely than men (47%; $p < 0.01$) to indicate that the pamphlet had influenced their decision to participate quite a bit or a lot (versus a little or not at all).

The nested recruitment study involved minimal additional resources, specifically 21 staff hours (four hours to randomize the sampling frame, three hours for data entry, 10 hours for programming, and four hours to review and synthesize results); no additional postage costs were incurred.

4. Discussion

In this randomized trial of a pamphlet nested within a study of CRC screening, its inclusion in the initial mailing had no effect on response rates among older Latinos. About half of the people to whom the pamphlet was mailed recalled receiving it, mostly women and younger Latinos. Latinas were also more likely than Latino men to indicate that the pamphlet had influenced their decision to participate in the survey. Nesting of the recruitment study within a funded study was possible with minimal additional cost.

Mass mailing of brochures has been shown to be successful in the recruitment of Whites and African Americans [9,14]. Our findings, however, suggest that using pamphlets to recruit Latinos into a telephone survey may be ineffective, although Latina women tended to be more receptive than Latino men. Similarly, two studies that used enhanced mailings (ethnically tailored letters) found that they did not significantly increase recruitment rates among Latinos [1,12,15].

Although direct mailing can be cost-effective for recruiting participants to prevention trials [1,14,16], our findings are consistent with many studies demonstrating that personalized recruitment approaches are necessary to achieve racial/ethnic equity in research participation [4,17,18]. In our study, substantially more people recalled receiving the letter than the pamphlet. The mailing served as an introduction to the follow-up telephone contact in which trained culturally competent interviewers facilitated recruitment by conveying information and addressing low-literacy issues that may be more prevalent among older Latinos [15,19,20].

Similar to other studies, we found that Latinas were more likely to participate than Latino men [2,21]. Gender disparities in labor force participation may explain the higher rates of participation among women [22,23]. Monetary incentives may also be more of an inducement for Latinas who are less likely than Latino men to be working outside the home [12, 24–26].

To eliminate health disparities, much more research is needed on effective strategies for increasing minority enrollment. In the absence of targeted funding, nesting recruitment studies is a useful interim strategy. However, nested studies may only be feasible for low

intensity recruitment methods, such as the one tested in this study. Comparative effectiveness research on more intensive recruitment methods for diverse populations will require dedicated funding and the commitment of researchers.

Acknowledgments

This work was supported by the National Cancer Institute (grant numbers R21CA121176, U01-CA86117); and the Resource Centers for Minority Aging Research program of the National Institute on Aging, the National Institute of Nursing Research, and the National Center on Minority Health and Health Disparities (grant number P30-AG15272). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Cancer Institute, National Institute on Aging, the National Institute of Nursing Research, and the National Center on Minority Health and Health Disparities or the National Institutes of Health.

We are indebted to the staff and patients at Marin Community Clinics, St. Jude Heritage Medical Center, and St. Jude Neighborhood Health Center who made this study possible.

References

1. Kiernan M, Phillips K, Fair JM, King AC. Using direct mail to recruit Hispanic adults into a dietary intervention: an experimental study. *Ann Behav Med.* 2000; 22:89–93. [PubMed: 10892533]
2. Maxwell AE, Bastani R, Glenn BA, Mojica CM, Chang LC. An experimental test of the effect of incentives on recruitment of ethnically diverse colorectal cancer cases and their first-degree relatives into a research study. *Cancer Epidemiol Biomarkers Prev.* 2009; 18:2620–5. [PubMed: 19755646]
3. Napoles-Springer AM, Fongwa MN, Stewart AL, Gildengorin G, Perez-Stable EJ. The effectiveness of an advance notice letter on the recruitment of African Americans and Whites for a mailed patient satisfaction survey. *J Aging Health.* 2004; 16:124S–36S. [PubMed: 15448290]
4. Ramirez AG, Miller AR, Gallion K, San Miguel de Majors S, Chalela P, Garcia Aramburo S. Testing three different cancer genetics registry recruitment methods with Hispanic cancer patients and their family members previously registered in local cancer registries in Texas. *Community Genet.* 2008; 11:215–23. [PubMed: 18417969]
5. Wenzel L, Bowen D, Habbal R, Leighton N, Vu T, Anton-Culver H. Testing targeted approaches to enhance Cancer Genetics Network minority recruitment within Asian populations. *Community Genet.* 2008; 11:234–40. [PubMed: 18417971]
6. Office for Human Research Protections. [Accessed Sept 12, 2011] Becoming a Research Volunteer: It's Your Decision. <http://www.hhs.gov/ohrp/education/brochures/index.html>
7. Levkoff S, Sanchez H. Lessons learned about minority recruitment and retention from the Centers on Minority Aging and Health Promotion. *Gerontologist.* 2003; 43:18–26. [PubMed: 12604742]
8. Napoles-Springer AM, Grumbach K, Alexander M, Moreno-John G, Forte D, Rangel-Lugo M, et al. Clinical research with older African Americans and Latinos: Perspectives from the Community. *Research on Aging.* 2000; 22:668–91.
9. Yancey AK, Ortega AN, Kumanyika SK. Effective recruitment and retention of minority research participants. *Annu Rev Public Health.* 2006; 27:1–28. [PubMed: 16533107]
10. Ford JG, Howerton MW, Lai GY, Gary TL, Bolen S, Gibbons MC, et al. Barriers to recruiting underrepresented populations to cancer clinical trials: a systematic review. *Cancer.* 2008; 112:228–42. [PubMed: 18008363]
11. Wendler D, Kington R, Madans J, Van Wye G, Christ-Schmidt H, Pratt LA, et al. Are racial and ethnic minorities less willing to participate in health research? *PLoS Med.* 2006; 3:e19. [PubMed: 16318411]
12. Napoles-Springer AM, Santoyo J, Stewart AL. Recruiting ethnically diverse general internal medicine patients for a telephone survey on physician-patient communication. *J Gen Intern Med.* 2005; 20:438–43. [PubMed: 15963168]
13. Napoles AM, Santoyo-Olsson J, Farren G, Olmstead J, Cabral R, Ross B, et al. The patient-reported Clinicians' Cultural Sensitivity Survey: a field test among older Latino primary care patients. *Health Expect.* 2011

14. Kennedy BM, Kumanyika S, Ard JD, Reams P, Johnson CA, Karanja N, et al. Overall and minority-focused recruitment strategies in the PREMIER multicenter trial of lifestyle interventions for blood pressure control. *Contemp Clin Trials*. 2010; 31:49–54. [PubMed: 19879377]
15. Calderon JL, Baker RS, Fabrega H, Conde JG, Hays RD, Fleming E, et al. An ethno-medical perspective on research participation: a qualitative pilot study. *MedGenMed*. 2006; 8:23. [PubMed: 16926762]
16. Robinson JL, Fuerch JH, Winiewicz DD, Salvy SJ, Roemmich JN, Epstein LH. Cost effectiveness of recruitment methods in an obesity prevention trial for young children. *Prev Med*. 2007; 44:499–503. [PubMed: 17475318]
17. Paskett ED, Reeves KW, McLaughlin JM, Katz ML, McAlearney AS, Ruffin MT, et al. Recruitment of minority and underserved populations in the United States: the Centers for Population Health and Health Disparities experience. *Contemp Clin Trials*. 2008; 29:847–61. [PubMed: 18721901]
18. Pinsky PF, Ford M, Gamito E, Higgins D, Jenkins V, Lamerato L, et al. Enrollment of racial and ethnic minorities in the Prostate, Lung, Colorectal and Ovarian Cancer Screening Trial. *J Natl Med Assoc*. 2008; 100:291–8. [PubMed: 18390022]
19. Arean PA, Gallagher-Thompson D. Issues and recommendations for the recruitment and retention of older ethnic minority adults into clinical research. *J Consult Clin Psychol*. 1996; 64:875–80. [PubMed: 8916614]
20. Bistricky SL, Mackin RS, Chu JP, Arean PA. Recruitment of African Americans and Asian Americans With Late-Life Depression and Mild Cognitive Impairment. *Am J Geriatr Psychiatry*. 2010
21. Weech-Maldonado R, Elliott MN, Oluwole A, Schiller KC, Hays RD. Survey response style and differential use of CAHPS rating scales by Hispanics. *Med Care*. 2008; 46:963–8. [PubMed: 18725851]
22. United States Department of Labor. Labor force statistics from the Current Population Survey: Women in the labor force, a data book. Vol. 2010. Bureau of Labor Statistics; 2004.
23. Voigt LF, Schwartz SM, Doody DR, Lee SC, Li CI. Feasibility of including cellular telephone numbers in random digit dialing for epidemiologic case-control studies. *Am J Epidemiol*. 2011; 173:118–26. [PubMed: 21071602]
24. Sweeney C, Edwards SL, Baumgartner KB, Herrick JS, Palmer LE, Murtaugh MA, et al. Recruiting Hispanic women for a population-based study: validity of surname search and characteristics of nonparticipants. *Am J Epidemiol*. 2007; 166:1210–9. [PubMed: 17827445]
25. Gonzalez EW, Gardner EM, Murasko D. Recruitment and retention of older adults in influenza immunization study. *J Cult Divers*. 2007; 14:81–7. [PubMed: 19175248]
26. Marcantonio ER, Aneja J, Jones RN, Alsop DC, Fong TG, Crosby GJ, et al. Maximizing clinical research participation in vulnerable older persons: identification of barriers and motivators. *J Am Geriatr Soc*. 2008; 56:1522–7. [PubMed: 18662204]

Table A.1

Correlates of Telephone Survey Completion among Older Latino Primary Care Patients.

	Total N=1,190	No Survey Completed N (%)	Survey Completed N (%)	p-value	Adjusted Odds Ratio for Survey Completion (95% CI) ^a
Experimental Group					
No Pamphlet	620 (52)	353 (57)	267 (43)	0.168	1.0
Pamphlet	570 (48)	347 (61)	223 (39)		0.8 (0.7,1.1)
Gender					
Men	428 (36)	277 (65)	151 (35)	< 0.01	1.0
Women	762 (64)	423 (56)	339 (44)		1.4 (1.1, 1.8)
Age					
50–54	371 (31)	216 (58)	155 (42)		1.0
55–64	459 (39)	254 (55)	205 (45)	0.05	1.1 (0.8, 1.5)
65+	360 (30)	230 (64)	130 (36)		0.6 (0.4, 0.8)
Health insurance					
None	528 (44)	287 (54)	241 (46)		1.0
Public	326 (27)	193 (59)	133 (41)	< 0.01	1.7 (1.1, 2.5)
Private	336 (28)	220 (65)	116 (35)		0.8 (0.5,1.4)

^a Adjusted for health care site.