UC Irvine

UC Irvine Previously Published Works

Title

Effect of water vapor on the destruction of ozone in the stratosphere perturbed by ClX or NO(x) pollutants

Permalink

<https://escholarship.org/uc/item/6md6t850>

Journal

Journal of Geophysical Research, 81(18)

ISSN

0148-0227

Authors

Liu, SC
Donahue, TM
Cicerone, RJ
et al.

Publication Date

1976

DOI

10.1029/JC081i018p03111

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Effect of Water Vapor on the Destruction of Ozone in the Stratosphere Perturbed by ClX or NO_x Pollutants

S. C. LIU, T. M. DONAHUE, R. J. CICERONE, AND W. L. CHAMEIDES

Department of Atmospheric and Oceanic Science, Space Physics Research Laboratory, University of Michigan, Ann Arbor, Michigan 48109

We describe results of a self-consistent one-dimensional coupled flow calculation for O₃, NO_x, HO₂, ClX, H₂O, H₂, CH₄, H₂O₂, and N₂O densities between 10 and 120 km. Our results agree well with observations for the normal mid-latitude atmosphere over this altitude range. We have varied ClX, NO_x, and H₂O independently in our model. We show that the effect of depletion of ozone by ClX is to remove ozone preferentially above 30 km and to lower the altitude of maximum ozone density. This leads to enhanced solar heating of the lower stratosphere and tropopause and suggests the possibility of an increased flux of water into the stratosphere. We show that increasing water vapor in the stratosphere greatly enhances the rate of destruction of O₃ by ClX and also causes an increase in the rate of destruction of O₃ in the NO_x-perturbed atmosphere.

INTRODUCTION

Recently, a number of studies have been published showing the effect on the earth's ozone layer of increasing the concentration of odd chlorine, or ClX (Cl + ClO + HCl) in the stratosphere [Molina and Rowland, 1974; Cicerone et al., 1974; Crutzen, 1974a; Wofsy et al., 1975]. We wish to report here the result of calculations that show the influence of the amount of water vapor in the stratosphere on the effectiveness of a specific level of ClX or NO_x (NO + NO₂ + HNO₃) in ozone removal. Our calculations are for a steady state and cover the range 10–120 km. We employ a one-dimensional coupled flow calculation using Hunten's [1975] eddy diffusion coefficient. Originally, we were motivated to perform this exercise by realizing that in the presence of ClX from chlorocarbons, ozone is preferentially destroyed at high altitudes and the altitude of the peak O₃ concentration tends to be lowered. Thus the heating rate near the tropopause due to absorption by O₃ is thereby increased. This could lead to an increase in the temperature of the tropopause [Ramanathan et al., 1976] and thus decrease the 'cold trap' effect on water vapor [Stanford, 1973]. This in turn would increase the water vapor abundance in the stratosphere. On the other hand, a more uniform reduction of ozone such as by NO_x pollutants might decrease water vapor in the stratosphere. Since HO₂ (H + OH + HO₂) plays such an important role in O₃ chemistry, it seems to us important to assess the effect of changes in water concentration on the natural and the perturbed stratosphere. It is particularly significant in the case of ClX pollution because not only does HO₂ destroy O₃ catalytically in its own right but OH attacks HCl to release Cl and thus reinitiates the ClX catalytic chain reaction.

CALCULATION METHOD AND BOUNDARY CONDITIONS

In our calculation we have self-consistently treated O₃ (O₃ + O), HO₂, NO_x, ClX, H₂O, H₂, CH₄, H₂O₂, and N₂O following the methods developed and described by Liu and Donahue [1974a, b, c, 1975a, 1976]. In the standard (i.e., 'unperturbed') atmosphere we have assumed a mixing ratio of 10⁻³ ppb vol/vol of ClX. In reality there may be about 0.5 ppb odd chlorine due to natural sources in the middle and upper stratosphere as discussed by Cicerone et al. [1975]. Such a background ClX mixing ratio could result from natural CH₃Cl, first detected by Rasmussen (private communication, 1974)

and Lovelock [1975]. In 1975, Rasmussen (private communication, 1975) measured 0.5 ppb CH₃Cl in the troposphere, on average. Our standard model neglects this natural ClX background to emphasize the effects of added ClX. ClX perturbations are modeled by injecting Cl atoms between 25 and 35 km, as is approximately the case for photolysis of CF₂Cl₂ and CFCI₃ [Molina and Rowland, 1974; Cicerone et al., 1974; Crutzen, 1974b; Rowland and Molina, 1975; Wofsy et al., 1975]. Steady injections of this kind lead to a characteristic ClX mixing ratio versus altitude [see, e.g., Cicerone et al., 1974; Wofsy et al., 1975]. The mixing ratio increases with altitude up to about 35 km, where a near-asymptotic value is attained; hereafter in this paper, 'ClX mixing ratio' means this high-altitude asymptote. (Recent measurements of HCl up to 30 km confirm this profile shape [Ackerman, 1976].)

The one-dimensional steady state continuity and diffusive flux equations are solved for the species mentioned above from 10 to 120 km. We use the now standard technique of treating O₃, HO₂, ClX, and NO_x as single species and obtain individual species densities in each family from the photochemical equilibrium conditions. For upper boundary conditions we use the 'flux-to-density' relations described in detail by Liu and Donahue [1974a]. At the lower boundary (10 km) we fix the ratio of the flux to the density of O₃ at the value 0.06 cm s⁻¹. A similar condition, with a ratio of 0.3 cm s⁻¹, is imposed on H₂O₂, NO_x, and ClX. Fixed mixing ratios are set for CH₄, H₂, H₂O, and N₂O at 10 km. In the standard atmosphere their values at 10 km are 1.5, 0.5, 3, and 0.25 ppm, respectively. Photochemical equilibrium is assumed in determining the HO₂ density at the lower boundary. We use the mid-latitude atmosphere (U.S. Standard Atmosphere Supplements, 1966) at equinoctial geometry in computing daily averaged rates of photolysis. The rates at 120 km are listed in Table 1.

The rate constants we used are tabulated in Table 2. We emphasize the importance of the value assumed for the rate constant k_{18} for (R18),

by assuming in these calculations a 'high' value [Hochanadel et al., 1972], 2×10^{-10} cm³ s⁻¹, and a 'low' value [Kaufman, 1975], 2×10^{-11} cm³ s⁻¹, and analyzing in detail the sensitivity of our results to variations of this rate constant. Liu and Donahue [1974c] have shown that Anderson's [1971] measurements of OH densities can be reconciled with this range of values for k_{18} provided certain constraints are placed on a

TABLE 1. Daily Averaged Photodissociation Rates of Relevant Species at 120 km for Equinoctial Conditions at 30° Latitude

Reaction	Rate, s ⁻¹	Reference
O ₂ + <i>hν</i> → O + O	1.26 × 10 ⁻⁶	Hudson <i>et al.</i> [1969] Ditchburn and Young [1961] Metzger and Cook [1964]
O ₃ + <i>hν</i> → O ₂ + O(¹ D)	4.4 × 10 ⁻⁸	Ackerman [1970] DeMore and Raper [1966]
O ₃ + <i>hν</i> → O ₂ + O(³ P)	2.1 × 10 ⁻⁴	Ackerman [1970]
H ₂ O + <i>hν</i> → H + OH	5.0 × 10 ⁻⁶	Thompson <i>et al.</i> [1963] Anderson [1971]
NO ₂ + <i>hν</i> → NO + O	5 × 10 ⁻⁸	Hampson <i>et al.</i> [1973b]
HNO ₃ + <i>hν</i> → OH + NO ₂	8.6 × 10 ⁻⁵	Johnston and Graham [1973]
H ₂ O ₂ + <i>hν</i> → OH + OH	5.3 × 10 ⁻⁵	Schumb <i>et al.</i> [1955]
N ₂ O + <i>hν</i> → N ₂ + O	6.7 × 10 ⁻⁷	Johnston and Selwyn [1975]
H ₂ CO + <i>hν</i> → H ₂ + CO	6 × 10 ⁻⁵	Calvert <i>et al.</i> [1972]
H ₂ CO + <i>hν</i> → HCO + H	4.1 × 10 ⁻⁵	Calvert <i>et al.</i> [1972]
HCl + <i>hν</i> → H + Cl	1.1 × 10 ⁻⁶	Myer and Samson [1970] Inn [1975]
NO ₃ + <i>hν</i> → NO + O ₂	1 × 10 ⁻²	Schott and Davidson [1958]

Solar flux data are taken from Ackerman [1970].

companion reaction destroying HO_x, (R16) between H and HO₂. The lower value of *k*₁₈ usually calls for about 3 times as much HO_x in the stratosphere for a given amount of water. There is, accordingly, less O₃ in models using the low value of

*k*₁₈. The role of (R20), whose rate constant is also poorly known, is discussed by Donahue *et al.* [1976].

Below 10 km we have assumed the amount of ozone present to be 0.54 × 10¹⁸ cm⁻², a quantity taken from the tables of

TABLE 2. Reactions and Rate Constants With References

	Reaction	Rate Constant, cm ³ s ⁻¹ , cm ⁶ s ⁻¹	Reference
R1	Cl + O ₃ → ClO + O ₂	2.4 × 10 ⁻¹¹ exp (-340/ <i>T</i>)	Zahniser <i>et al.</i> [1975]
R2	ClO + O → Cl + O ₂	1.2 × 10 ⁻¹⁰ exp (-250/ <i>T</i>)	Bemand <i>et al.</i> [1973]
R3	ClO + NO → Cl + NO ₂	1.7 × 10 ⁻¹¹	Clyne and Watson [1974]
R4	Cl + CH ₄ → HCl + CH ₃	5.4 × 10 ⁻¹² exp (-1125/ <i>T</i>)	Davis <i>et al.</i> [1975b]*
R5	HCl + OH → Cl + H ₂ O	2 × 10 ⁻¹² exp (-310/ <i>T</i>)	Zahniser <i>et al.</i> [1974]
R6	HCl + O → Cl + OH	1.9 × 10 ⁻¹¹ exp (-3600/ <i>T</i>)	Wong and Belles [1971]
R7	Cl + HO ₂ → HCl + O ₂	2 × 10 ⁻¹¹	estimated
R7a	Cl + H ₂ O ₂ → HCl + HO ₂	6 × 10 ⁻¹⁸	Davis <i>et al.</i> [1975a]
R8	H ₂ + Cl → HCl + H	8 × 10 ⁻¹¹ exp (-2650/ <i>T</i>)	Benson <i>et al.</i> [1969]
R9	H + HCl → Cl + H ₂	1 × 10 ⁻¹¹ exp (-1600/ <i>T</i>)	Benson <i>et al.</i> [1969]
R10	H ₂ O + O(¹ D) → 2OH	3.5 × 10 ⁻¹⁰	Garvin and Hampson [1974]
R11	CH ₄ + O(¹ D) → OH + CH ₃	4 × 10 ⁻¹⁰	Garvin and Hampson [1974]*
R12	CH ₄ + OH → H ₂ O + CH ₃	2.95 × 10 ⁻¹² exp (-1770/ <i>T</i>)	Davis [1973]*
R13	H ₂ + O(¹ D) → OH + H	2.9 × 10 ⁻¹⁰	Garvin and Hampson [1974]
R13a	H ₂ + O → OH + H	3 × 10 ⁻¹⁴ <i>T</i> exp (-4480/ <i>T</i>)	Baulch <i>et al.</i> [1973]
R14	H + O ₃ → OH + O ₂	2.6 × 10 ⁻¹¹	Hampson <i>et al.</i> [1973a]
R15	H + O ₃ + M → HO ₂ + M	6 × 10 ⁻³⁸ exp (290/ <i>T</i>)	Garvin and Hampson [1974]
R16a	H + HO ₂ → H ₂ + O ₂	4.2 × 10 ⁻¹¹ exp (-350/ <i>T</i>)	Garvin and Hampson [1974]
R16b	H + HO ₂ → H ₂ O + O	8.3 × 10 ⁻¹¹ exp (-500/ <i>T</i>)	Lloyd [1973]
R17	OH + O → H + O ₂	4 × 10 ⁻¹¹	Wilson [1973]
R18	OH + HO ₂ → H ₂ O + O ₂	see text	
R19	OH + O ₃ → HO ₂ + O ₂	1.3 × 10 ⁻¹² exp (-950/ <i>T</i>)	Anderson and Kaufman [1973]
R20	HO ₂ + O → OH + O ₂	8 × 10 ⁻¹¹ exp (-500/ <i>T</i>)	Lloyd [1973]
R21	HO ₂ + O ₃ → OH + 2O ₂	1 × 10 ⁻¹⁸ exp (-1250/ <i>T</i>)	Garvin and Hampson [1974]
R22	HO ₂ + HO ₂ → H ₂ O ₂ + O ₂	3 × 10 ⁻¹¹ exp (-500/ <i>T</i>)	Hampson <i>et al.</i> [1973a]
R23	H ₂ O ₂ + OH → HO ₂ + H ₂ O	1.7 × 10 ⁻¹¹ exp (-910/ <i>T</i>)	Hampson <i>et al.</i> [1973a]
R24	HO ₂ + NO → OH + NO ₂	1.1 × 10 ⁻¹¹ exp (-1000/ <i>T</i>)	Davis <i>et al.</i> [1973]
R25	N ₂ O + O(¹ D) → 2NO	1.1 × 10 ⁻¹⁰	Garvin and Hampson [1974]
R25a	N ₂ O + O(¹ D) → N ₂ + O ₂	1.1 × 10 ⁻¹⁰	Garvin and Hampson [1974]
R26	NO + O ₃ → NO ₂ + O ₂	9 × 10 ⁻¹⁸ exp (-1200/ <i>T</i>)	Hampson <i>et al.</i> [1973a]
R27	NO + O + M → NO ₂ + M	7 × 10 ⁻³²	Becker <i>et al.</i> [1973]
R28	NO ₂ + O → NO + O ₂	1.7 × 10 ⁻¹¹ exp (-300/ <i>T</i>)	Baulch <i>et al.</i> [1973]
R29	NO ₂ + OH + M → HNO ₃ + M	1 × 10 ⁻¹¹ / (3.5 × 10 ¹² <i>T</i> ^{2.5} + 2.5 [M])	Anderson <i>et al.</i> [1974]†
R30	HNO ₃ + OH → H ₂ O + NO ₂	0.89 × 10 ⁻¹⁸	Margitan <i>et al.</i> [1975]
R31	NO ₃ + NO → 2NO ₂	8.7 × 10 ⁻¹²	Harker and Johnston [1973]
R32	NO ₂ + O ₃ → NO ₃ + O ₂	1.1 × 10 ⁻¹⁸ exp (-2450/ <i>T</i>)	Garvin and Hampson [1974]
R33	NO ₃ + O + M → NO ₂ + M	1 × 10 ⁻³¹	Hampson <i>et al.</i> [1973b]
R34	O + O ₃ → O ₂ + O ₂	10.0 × 10 ⁻³⁶ exp (510/ <i>T</i>)	Garvin and Hampson [1974]
R35	O + O ₃ → 2O ₂	1.9 × 10 ⁻¹¹ exp (-2300/ <i>T</i>)	Hampson <i>et al.</i> [1973a]
R36	O + O + M → O ₂ + M	2.76 × 10 ⁻³⁴ exp (710/ <i>T</i>)	Campbell and Thrush [1967]

*Methane oxidation chain of Chameides and Walker [1973] is adopted.

† Best fit to the results of Anderson *et al.* [1974] and Tsang [1973].

Hering and Borden [1967] as appropriate to the equinoxes. This quantity is added to the integrated amounts obtained above 10 km in calculating the total column ozone abundance for all of our models.

RESULTS: STANDARD AND CIX-PERTURBED ATMOSPHERE

In Figure 1 we show some of the profiles of species obtained from our calculations for standard conditions and, where appropriate, for high and low values of k_{18} . Figure 2 shows standard ozone profiles for both values of k_{18} . Also shown in Figure 2 is the effect of adding 8 ppb CIX on the ozone densities. The preferential removal of ozone at high altitude is obvious. As a result of this perturbation, the altitude of the ozone maximum is decreased, leading to an increase in the solar and infrared heating rates below the ozone maximum [Ramanathan *et al.*, 1976]. For example, when k_{18} is high and 15 ppb of CIX is present, the ozone maximum is lowered from 24 to 22 km, and the solar heating rate at 10, 15, and 20 km is 13%, 16%, and 19% larger, respectively. Ramanathan *et al.* [1976] have indicated that this kind of change in the ozone distribution might increase the temperature at the tropopause by several degrees. The saturation vapor pressure doubles for a 4°K rise in tropopause temperature and increases by an order of magnitude for a 15°K rise. Thus in the presence of CIX the water vapor concentration at the tropopause would increase [Stanford, 1973]. In the NO_x -perturbed stratosphere, however, the ozone density profile is affected more uniformly [McElroy *et al.*, 1974] and thus could lead to a decrease of water vapor concentration at the tropopause [Ramanathan *et al.*, 1976].

To assess the effect of changing the amount of water vapor in the stratosphere on the ozone distribution, we have repeated the steady state calculations for water vapor mixing ratios at 10 km ranging from 10^{-2} to 30 ppm. When the water vapor mixing ratio at the tropopause is lower than about 1 ppm, the water vapor present in the stratosphere will be produced mostly from methane because oxidation of methane is an important source of water vapor above the tropopause [Hunten and Strobel, 1974; Liu and Donahue, 1974a]. We show in Figure 3 the mixing ratio of hydrogen in the forms of CH_4 , H_2O , and H_2 as a function of altitude for assumed mixing ratios of 0.5 ppb of H_2 , 1.5 ppm of CH_4 at 10 km, and 3 ppm of H_2O in one case and 10^{-2} ppm of H_2O in the other. Plotted is the total hydrogen mixing ratio for each species, i.e., twice the H_2 and H_2O mixing ratios and 4 times the CH_4 mixing ratio. The total mixing ratio ($2[\text{H}_2\text{O}] + 2[\text{H}_2] + 4[\text{CH}_4]$) must be almost constant with altitude, or a very large flux compared to the escape flux of hydrogen would result [Hunten, 1973]. When there is almost no water at 10 km, the 2.8 ppm of H_2O near 50

Fig. 1. Profiles of H_2O , CH_4 , H_2 , and H_2O_2 for high k_{18} (solid line) and low k_{18} (dashed line).

Fig. 2. Ozone profiles for high k_{18} (solid line) and low k_{18} (dashed line) for the standard equinoctial atmosphere at mid-latitudes (solid circles), for this atmosphere with 8 ppb of CIX (crosses), and with H_2O at the tropopause increased by a factor of 10 (open circles).

km is essentially all produced from CH_4 . Above 40 km, H_2 begins to be created as a result of the reaction (R16),

In the case of 3 ppm H_2O at 10 km the H_2 mixing ratio will grow from 0.5 ppm at 10 km to more than 4 ppm at 90 km as a result of conversion of H_2O and CH_4 . But with virtually no H_2O at 10 km the H_2 mixing ratio reaches only a little more than 3 ppm, since the increase comes only from the conversion of methane to H_2O and the subsequent production of odd hydrogen from H_2O .

Figure 2 also shows the effect of increasing the H_2O mixing ratio to 30 ppm on the ozone distribution already modified by the presence of 8 ppb CIX. (We plot profiles for large increases in CIX and H_2O for the sake of obtaining a clear separation of the curves.) Figures 4 and 5 show the percentage reduction in O_3 column abundances as CIX is added to the atmosphere and the H_2O mixing ratio at 10 km is varied. Again, results are shown for two choices of k_{18} .

Before discussing the effects of varying the water vapor source, we call attention to rate constants for the reaction (R1),

Fig. 3. Mixing ratios of $2\text{H}_2\text{O}$, 2H_2 , and 4CH_4 as functions of altitude for two atmosphere models with different water vapor mixing ratios at the tropopause. In the standard model (solid line) the mixing ratios at the tropopause are 6, 6, and 1 ppm for $2\text{H}_2\text{O}$, 4CH_4 , and 2H_2 , respectively. In the other model (dashed line) the corresponding mixing ratios are 2×10^{-3} , 6, and 1 ppm.

Fig. 4. Changes in O_3 column abundance as CIX mixing ratio increases. Results are shown for four mixing ratios of H_2O at the tropopause, 0.01 ppm (open triangles), 3 ppm (solid circles), 9 ppm (crosses), and 30 ppm (open circles), and for high k_{18} (solid line) and low k_{18} (dashed line).

newly measured by Davis *et al.* [1975a] and Zahniser *et al.* [1975], i.e., $2.4 \times 10^{-11} \exp(-340/RT)$, and for the reaction (R4),

where k_4 is $5.4 \times 10^{-12} \exp(-1125/T)$ according to Davis *et al.* [1975b]. In early (pre-1975) studies of stratospheric CIX, k_1 was taken to be $4.3 \times 10^{-11} \exp(-250/T)$ [see, e.g., Stolarski and Cicerone, 1974] based on Clyne and Watson's room temperature rate of 1.85×10^{-11} [Watson, 1974]. Also, k_4 was taken to be $8.8 \times 10^{-11} \exp(-1900/T)$ in the early studies, based on Davis *et al.* [1970]. The use of the new rates k_1 and k_4 in our model has significantly reduced the effect of a given amount of CIX on O_3 compared to that in previous models. In

Fig. 5. Changes in O_3 column abundance as the H_2O mixing ratio at the tropopause is changed. Results are as shown for high k_{18} with 1 ppb of CIX (solid line), low k_{18} with 1 ppb of CIX (dashed line), high k_{18} with 5 ppb of CIX (dash-dotted line), and low k_{18} with 5 ppb of CIX (dotted line).

Fig. 6. Density distribution of methane for the standard atmosphere with high k_{18} (solid line) and low k_{18} (dashed line) and for atmosphere containing 8 ppb of CIX and low k_{18} (dotted line).

this result we agree with Crutzen as quoted by Hammond [1975] and with Wofsy and McElroy [1975].

Our results for the new rates k_1 and k_4 , compared to those for the old k_1 and k_4 , are that 1 ppb of CIX reduces the total amount of O_3 by 0.6% compared to 2.2%, 3 ppb of CIX reduces O_3 by 1.9% compared to 6.4%, and 5 ppb of CIX reduces O_3 by 3.6% compared to 10.4%. These comparisons are made for the larger value of k_{18} . In case the smaller value is more nearly correct the reduction in the integrated O_3 abundance for a given combination of CIX and water vapor concentrations could be much greater, as can be seen in Figures 2, 4, and 5. Thus unless the amount of water vapor in the stratosphere were to change significantly from its present value, the time scale for a specified amount of ozone destruction by CIX could be appreciably greater than first calculated [Cicerone *et al.*, 1974; Crutzen, 1974a; Wofsy *et al.*, 1975]. Even this conclusion is dangerous, however, based as it is on the assumed high value of k_{18} . Donahue *et al.* [1976] discuss this further. The 10% ozone destruction level is reached with less than 6 ppb of CIX if the low value of k_{18} is correct.

Figures 4 and 5 also demonstrate the principal points that we wish to make in this paper. The effects produced by changing H_2O are significant. In particular, when the CIX pollution level is 1 ppb, the percentage ozone reduction goes from 0.6% to 2.9% if the amount of H_2O is tripled (9 ppm at 10 km) and to 9.7% if it is increased by a factor of 10 (for the high value of k_{18}). Again for the low value of k_{18} a much more dramatic effect occurs, i.e., a 6.8% reduction for 1 ppb of CIX and 3 times normal H_2O and a 16.6% reduction for 10 times normal H_2O . Of course if the stratosphere were to dry out for some reason, the changes would occur in the opposite sense.

The reason that water vapor has such a great influence is that it is converted by reaction with $O(^1D)$ into odd hydrogen in the stratosphere. All forms of odd hydrogen destroy ozone catalytically in their own right, but in addition to this direct effect there is a special indirect effect on ozone destruction. The HCl produced by reactions of Cl with CH_4 in reaction (R4) is attached by OH (R5),

thus reinitiating the CIX catalytic chain. Figures 7, 8, 10, and 11 quantitatively demonstrate the nature of the effects just described. Figure 6 shows the methane distribution as a function of altitude for the standard atmosphere with high and low values of k_{18} , where the influence of the enhanced destruction of CH_4 by OH when k_{18} is low is apparent. Much

Fig. 7. Profiles of odd hydrogens and $O(^1D)$ for the standard atmosphere (solid line), with 8 ppb of ClX added and normal H_2O (dashed line) and with 8 ppb of ClX and 3 times H_2O at the tropopause (dash-dotted line). Low value of k_{18} is used here.

more striking, however, is the rapid destruction of methane by Cl in the formation of HCl through reaction (R4) when 8 ppb of ClX is present. At 45 km the methane density has been cut by an order of magnitude compared to the normal value as a result of the presence of ClX. Tropospheric CH_4 is also affected by ClX-induced O_3 losses in the stratosphere, feeding back on the stratosphere [Chameides *et al.*, 1976].

In Figure 7 the densities of HO_x are plotted for the low value of k_{18} in three cases: (1) the standard atmosphere, (2) with 8 ppb of ClX and normal H_2O , and (3) with 8 ppb of ClX and 3 times the normal H_2O . It is interesting that the presence of ClX increases the amount of HO_x considerably below the ozone maximum and decreases it above. The reason is to be found in the preferential destruction of O_3 above the maximum by ClX released from chlorocarbons. This decrease in optical depth enhances $O(^1D)$ production from O_3 at low altitudes, as is demonstrated by the curves showing the $O(^1D)$ densities. Since $O(^1D)$ interacting with H_2O and CH_4 is an important source of HO_x , the result is an increase in HO_x at low altitudes. Conversely, above 40 km, where the atmosphere is optically thin for production of $O(^1D)$, the $O(^1D)$ density is lower because the O_3 density is lower. Thus the HO_x density in the presence of ClX is reduced at higher altitudes. Figure 8 shows the ClX species as they are affected by the addition of water to the stratosphere: the conversion of HCl to Cl by OH can be easily

Fig. 8. Profiles of Cl, ClO, and HCl for atmosphere with 8 ppb of ClX and normal H_2O (solid line) and for atmosphere with 8 ppb of ClX and 3 times H_2O (dashed line). Low value of k_{18} is used here.

Fig. 9. Sinks of O_3 for the standard atmosphere with high k_{18} .

detected below 40 km. Above 50 km, because the H_2O_2 density increases by a larger factor than does the OH density for a given increase in water vapor concentration, Cl is preferentially converted to HCl by the reaction of H_2O_2 with Cl.

In Figures 9 and 10 we show the major sinks of O_3 for the standard atmosphere with high and low values of k_{18} , respectively. These figures show that the odd hydrogen reactions dominate the loss of O_3 above 55 and below 20 km. The Chapman reaction (R35) between O_3 and O dominates only in a narrow range near 50 km, and the NO_x catalytic cycle is in control from 20 to about 40 km, where the ozone density is largest. The effect of increasing the odd hydrogen density by reducing k_{18} is apparent in a comparison of Figures 9 and 10, particularly in the region where the Chapman reaction prevails. This region stretches from 43 to 56 km when k_{18} is large but only from 43 to 47 km when k_{18} is small. Figure 11 in turn, when compared with Figure 10, shows the changes in the dominant sinks of O_3 when 8 ppb of ClX is added. The ClO-O reaction rate in the ClX cycle at 40 km jumps from $3 \times 10^8 \text{ cm}^{-3} \text{ s}^{-1}$ to more than $8 \times 10^8 \text{ cm}^{-3} \text{ s}^{-1}$. The $NO_2 + O$ rate is reduced when ClX becomes dominant because of the decreased amount of O_3 leading to less NO_2 and less O. While the $HO_2 + O_3$ reaction rate is not shown in Figure 11, its dependence upon the addition of ClX and water vapor is similar to that of $OH + O_3$.

Note that for low k_{18} there is a significant amount of H_2O_2 near 25 km as a result of the large concentrations of HO_2

Fig. 10. Same as Figure 9 but with low k_{18} .

Fig. 11. Three important sinks of O_3 for atmosphere with 8 ppb of ClX and normal H_2O (solid line) and for atmosphere with 8 ppb of ClX and 3 times H_2O (dashed line). Low value of k_{18} is used here.

(Figure 1). The reaction (R23)

then becomes an important sink of HO_x and results in about 10% more ozone in the standard atmosphere than would be the case if k_{23} were not included. Even the reaction (R7a)

must be taken into account when k_{18} is small despite the relatively small rate constant recently obtained by Davis *et al.* [1975a]. Without the inclusion of this reaction in the chemical scheme the destruction of ozone would be overestimated by a factor of 1.3 when 1 ppb of ClX is present and by an increasingly greater factor as the amount of ClX increases.

EFFECT OF H_2O ON NO_x -PERTURBED STRATOSPHERE

We have also considered the effects of NO_x and H_2O produced by the 'standard' SST models, at 20 km [Grobecker, 1974]. In Figure 12 we show our results for the globally averaged reduction in O_3 resulting from 100 SST's emitting 9×10^6 NO $cm^{-2} s^{-1}$ between 19 and 21 km. If the stratosphere contains 3 ppm of water vapor, the reduction is 1.22% for the high value of k_{18} and 0.66% for the low value. This means 1.83% and 1% reduction of O_3 in the hemisphere in which the NO_x is dominantly emitted, a result that agrees well with previous calculations (summarized in the publication by the *National Academy of Sciences* [1975]). Note, however, that adding water vapor to the stratosphere causes an enhancement in O_3 reduction, no matter how little is added, whether k_{18} is small or large. With normal or almost normal amounts of H_2O in the stratosphere the reduction in O_3 for a specified artificial source of NO_x turns out to be greater when k_{18} is large and the density of HO_x is small because of the less effective formation of HNO_3 from OH and NO_2 . However, when k_{18} is small, the amount of HO_x required to stay in balance with H_2O increases much more rapidly with the H_2O mixing ratio than when k_{18} is large. For a small increase in the H_2O concentration the removal of O_3 by HO_x becomes more important than the removal by the NO emitted from 100 SST's. This dominance by the HO_x catalytic cycle causes the effect of added H_2O to be much more pronounced for the small value of k_{18} than the large. The same effect is noticeable in Figure 5 for ClX catalysis.

The decrease in ozone density caused by an increase in the

water vapor content of the stratosphere in our model occurs predominantly below about 22 km and above about 45 km. HO_x destruction of O_3 dominates NO_x destruction below 24 km and begins to compete seriously again above 40 km. It is the effect of HO_x below 22 km that causes us to find that increasing the H_2O mixing ratio even by very small amounts causes the ozone column content to decrease, whereas others [McElroy *et al.*, 1974; Crutzen, 1974b] found the opposite. McElroy *et al.* located their lower boundary at 28 km and multiplied the O_3 profile below that level by the change calculated at 28 km [Wofsy, 1974]. This exercise cannot duplicate the large effect of HO_x on O_3 that we find below 24 km. Our disagreement with Crutzen [1974b] may be due to different reaction rate constants. We note that Rao-Vupputuri [1974] also finds that increased H_2O leads to decreased O_3 .

In a sense, our result is disturbing because it suggests that the ozone layer is necessarily unstable against runaway destruction from perturbations (even natural ones that may have occurred in the past) allowing water into the stratosphere. Such a conclusion is not warranted, however, for the nature of the perturbation, the exact nature and location of the change in ozone density, the changes in temperature accompanying the perturbation, and the temperature dependences of all important rate constants must all be considered. Thus a perturbation like the one we describe here resulting from chlorocarbons, which removes the top of the layer of O_3 and causes the temperature to rise below 25 km and fall above 25 km, will cause HO_x to have different effects at high and low altitudes. The rate constants for the steps in the ozone-destroying reactions,

increase with increasing temperature, while the rate constant for the reaction between HO_x and NO_x that produces the sink for NO_x ,

decreases with increasing temperature. Thus as the temperature increases at low altitude, the destruction of ozone from HO_x and from NO_x tends to be augmented, but the opposite is true at high altitude. As a consequence, the ozone will tend to recover and the opacity to grow at high altitude, leading to a decrease in temperature below. Effects such as these certainly must be taken into account in a complete time-dependent theory of the phenomena we are describing in this note.

Fig. 12. Changes in O_3 column abundance in the atmosphere with 100 SST's when the H_2O mixing ratio at the tropopause is changed.

CONCLUSION

This paper draws attention to the very large influence of water vapor in the stratosphere on the catalytic destruction of ozone, particularly by odd chlorine originating from photolysis of chlorocarbon. A change in water vapor concentration, either an increase or a decrease, could conceivably result from changes in the environment affecting the so-called cold trap for water vapor, the production of methane, or both. The destruction of ozone by pollutants is particularly sensitive to the amount of H_2O present in the stratosphere in the case of odd chlorine produced by photolysis of chlorocarbon because of the reconversion of HCl to Cl by OH . The effects considered depend strongly on the value of the rate constant for the reaction of OH with HO_2 producing water vapor, because it determines the amount of odd hydrogen in equilibrium with a given amount of water vapor.

Acknowledgments. This work has greatly benefited from discussion with J. G. Anderson and W. R. Kuhn. The research was supported in part by the Atmospheric Sciences Section, National Science Foundation (NSF) grants DES 74-21598 and DES 74-11478 and by NASA grant NSG-7187. We also acknowledge the National Center for Atmospheric Research (sponsored by NSF) for computer time.

REFERENCES

- Ackerman, M., Ultraviolet solar radiation related to mesospheric processes, *Inst. Aeronaut. Spatiale Belge Brussels, A-77*, 149-159, 1970.
- Ackerman, M., D. Frimout, A. Girard, M. Gottignies, and C. Muller, Stratospheric HCl from infrared spectra, *Geophys. Res. Lett.*, **3**, 81, 1976.
- Anderson, J. G., Rocket measurement of OH in the mesosphere, *J. Geophys. Res.*, **76**, 7820-7824, 1971.
- Anderson, J. G., and F. Kaufman, Kinetics of the reaction $OH(v=0) + O_3 \rightarrow HO_2 + O_2$, *Chem. Phys. Lett.*, **19**, 483-486, 1973.
- Anderson, J. G., J. J. Margitan, and F. Kaufman, Gas phase recombination of OH with NO and NO_2 , *J. Chem. Phys.*, **60**, 3310-3317, 1974.
- Baulch, D. L., D. D. Drysdale, D. G. Horne, and A. E. Lloyd, Evaluated kinetic data for high temperature reactions, in *Homogeneous Gas Phase Reactions of the $H_2-N_2-O_2$ System*, vol. 2, Chemical Rubber Co. Press, Cleveland, Ohio, 1973.
- Becker, K. H., W. Groth, and D. Thran, Mechanism of the air afterglow $NO + O \rightarrow NO_2 + hv$, 14th Combustion Symposium, pp. 353-363, Combust. Inst., Pittsburgh, Pa., 1973.
- Bernand, P. P., M. A. A. Clyne, and R. T. Watson, Reactions of chlorine oxide radicals, 4, Rate constants for the reactions $Cl + OClO$, $O + OClO$, $HO + OClO$, $NO + OClO$, $O + ClO$, *J. Chem. Soc. Faraday Trans., Sect. 1*, **69**, 1356, 1973.
- Benson, S. W., F. R. Cruickshank, and R. Shaw, Iodine monochloride as a thermal source of chlorine atoms: The reaction of chlorine atoms with hydrogen, *Int. J. Chem. Kinet.*, **1**, 29-43, 1969.
- Calvert, J. G., J. A. Kerr, K. L. Demerjian, and R. D. McQuigg, Photolysis of formaldehyde as a hydrogen atom source in the lower atmosphere, *Science*, **175**, 751-752, 1972.
- Campbell, I. M., and B. A. Thrush, The association of oxygen atoms and their combination with nitrogen atoms, *Proc. Roy. Soc., Ser. A*, **296**, 222-231, 1967.
- Chameides, W. L., and J. C. G. Walker, A photochemical theory of tropospheric ozone, *J. Geophys. Res.*, **78**, 8751-8760, 1973.
- Chameides, W. L., S. C. Liu, R. J. Cicerone, and T. M. Donahue, Methane and troposphere-stratosphere feedback, submitted to *J. Geophys. Res.*, 1976.
- Cicerone, R. J., R. S. Stolarski, and S. Walters, Stratospheric ozone destruction by man-made chlorofluoromethanes, *Science*, **185**, 1165-1167, 1974.
- Cicerone, T. J., D. H. Stedman, and R. S. Stolarski, Estimate of late 1974 stratospheric concentration of gaseous chlorine compounds (CIX), *Geophys. Res. Lett.*, **2**, 219-222, 1975.
- Clyne, M. A. A., and R. T. Watson, Kinetic studies of diatomic free radicals using mass spectrometry, 2, Rapid bimolecular reactions involving the $ClO X^* \pi$ radical, *J. Chem. Soc. Faraday Trans., Sect. 1*, **70**, 2250-2259, 1974.
- Crutzen, P. J., Estimates of possible future ozone reductions from continued use of fluoro-chloro-methanes CF_2Cl_2 , $CFCl_3$, *Geophys. Res. Lett.*, **1**, 205-208, 1974a.
- Crutzen, P. J., A review of upper atmosphere photochemistry, *Can. J. Chem.*, **52**, 1569-1581, 1974b.
- Davis, D. D., New rate measurements on the reaction of $O(^3P)$, O_3 , and OH , *Pap 73-501*, Amer. Inst. of Aeronaut. and Astronaut., New York, 1973.
- Davis, D. D., W. Braun, and A. Bass, Reactions of $Cl(^2P_{3/2})$: Absolute rate constants for reaction with H_2 , CH_4 , C_2H_6 , CH_2Cl_2 , C_2Cl_4 , and $C-C_6H_{13}$, *Int. J. Chem. Kinet.*, **2**, 101-114, 1970.
- Davis, D. D., W. W. Wong, S. D. Fischer, R. L. Schiff, J. O. Lephardt, J. E. Prusaczyk, W. A. Payne, L. J. Steif, R. E. Huie, and J. T. Herron, Recent kinetic measurements of the reactions of $O(^3P)$, H , and HO_2 , Proceedings of the Second Conference on the CIAP, DOT-TSC-OST-73-4, U.S. Dep. of Transp., Washington, D. C., 1973.
- Davis, D. D., R. L. Schiff, U. Machado, R. T. Watson, and S. Fischer, Absolute rate constants for elementary reactions of atmospheric importance: Results from the University of Maryland's gas kinetics laboratory, *Rep. 1*, Univ. of Md., College Park, 1975a.
- Davis, D. D., R. L. Schiff, U. Machado, and R. T. Watson, Absolute rate constants for elementary reactions of atmospheric importance: Results from the University of Maryland's gas kinetics laboratory, *Rep. 2*, Univ. of Md., College Park, 1975b.
- DeMore, W. B., and O. F. Raper, Primary processes in ozone photolysis, *J. Chem. Phys.*, **44**, 1780-1783, 1966.
- Ditchburn, R. W., and P. A. Young, The absorption of molecular oxygen between 1850 and 2500 Å, *J. Atmos. Terr. Phys.*, **24**, 127-139, 1962.
- Donahue, T. M., R. J. Cicerone, S. C. Liu, and W. L. Chameides, Effect of odd hydrogen on ozone depletion by chlorine reactions, *Geophys. Res. Lett.*, **3**, 105, 1976.
- Garvin, D., and R. F. Hampson (Eds.), Chemical kinetics data survey 7: Tables of rate and photochemical data for modelling of the stratosphere, *NBSIR 74-430*, 101 pp., Nat. Bur. of Stand., Washington, D. C., 1974.
- Grobecker, A. J., Research program for assessment of stratospheric pollution, *Acta Astron.*, **4**, 179-224, 1974.
- Hammond, A. L., Ozone destruction: Problem's scope grows, its urgency recedes, *Science*, **187**, 1181-1183, 1975.
- Hampson, R. F. (Ed.), W. Braun, R. L. Brown, D. Garvin, J. T. Herron, R. E. Huie, M. J. Kurylo, A. H. Laufer, J. D. McKinley, H. Okabe, M. D. Scheer, W. Tsang, and D. H. Stedman, Survey of photochemical and rate data for twenty-eight reactions of interest in atmospheric chemistry, *J. Phys. Chem. Ref. Data*, **2**, 267-308, 1973a.
- Hampson, R. F. (Ed.), D. Garvin, J. T. Herron, R. E. Huie, M. J. Kurylo, A. H. Laufer, H. Okabe, M. D. Scheer, and W. Tsang, Chemical kinetics survey 6: Photochemical and rate data for twelve gas phase reactions of interest for atmospheric chemistry, *NBSIR 73-207*, 120 pp., Nat. Bur. of Stand., Washington, D. C., 1973b.
- Harker, A. B., and H. S. Johnston, Photolysis of nitrogen dioxide to produce transient O , NO_3 and N_2O_3 , *J. Phys. Chem.*, **77**, 1153-1156, 1973.
- Hering, W. S., and T. R. Borden, *Ozone Sonde Observations Over North America*, vol. 4, *Environ. Res. Pap. 279*, Air Force Cambridge Research Laboratory, Bedford, Mass., 1967.
- Hochanadel, C. J., J. A. Gormley, and P. J. Ogren, Absorption spectrum and reaction kinetics of the HO_2 radical in the gas phase, *J. Chem. Phys.*, **56**, 4426-4432, 1972.
- Hudson, R. D., V. L. Carter, and E. L. Breig, Predissociation in the Schumann-Runge band system of O_2 : Laboratory measurements and atmospheric effects, *J. Geophys. Res.*, **74**, 4079-4086, 1969.
- Hunten, D. M., The escape of light gases from planetary atmospheres, *J. Atmos. Sci.*, **30**, 1481-1494, 1973.
- Hunten, D. M., Vertical transport in atmospheres, in *Atmosphere of Earth and Planets*, edited by B. M. McCormac, pp. 59-72, D. Reidel, Dordrecht, Netherlands, 1975.
- Hunten, D. M., and D. F. Strobel, Production and escape of terrestrial hydrogen, *J. Atmos. Sci.*, **31**, 305-317, 1974.
- Inn, E. C. Y., Absorption coefficient of HCl in the region 1400 to 2200 Å, *J. Atmos. Sci.*, **32**, 2375-2377, 1975.
- Johnston, H., and R. Graham, Gas-phase ultraviolet absorption spectrum of nitric acid vapor, *J. Phys. Chem.*, **77**, 62-63, 1973.
- Johnston, H. S., and G. Selwyn, New cross sections for the absorption of near ultraviolet radiation by nitrous oxide (N_2O), *Geophys. Res. Lett.*, **2**, 549, 1975.

- Kaufman, F., Hydrogen chemistry: Perspective on experiment and theory, in *Atmosphere of Earth and Planets*, edited by B. M. McCormac, pp. 219–232, D. Reidel, Dordrecht, Netherlands, 1975.
- Liu, S. C., and T. M. Donahue, The aeronomy of hydrogen in the atmosphere of earth, *J. Atmos. Sci.*, **31**, 1118–1136, 1974a.
- Liu, S. C., and T. M. Donahue, Mesospheric hydrogen related to exospheric escape mechanisms, *J. Atmos. Sci.*, **31**, 1466–1470, 1974b.
- Liu, S. C., and T. M. Donahue, Realistic model of hydrogen constituents in the lower atmosphere and escape flux from the upper atmosphere, *J. Atmos. Sci.*, **31**, 2238–2242, 1974c.
- Liu, S. C., and T. M. Donahue, The aeronomy of the upper atmosphere of Venus, *Icarus*, **24**, 148–156, 1975a.
- Liu, S. C., and T. M. Donahue, The regulation of hydrogen and oxygen escape from Mars, *Icarus*, in press, 1976.
- Lloyd, A. C., Evaluated and estimated kinetic data for gas phase reactions of the hydroperoxyl radical, *Rep. 10447*, Nat. Bur. of Stand., Washington, D. C., 1973.
- Lovelock, J. E., Natural halocarbons in the air and in the sea, *Nature*, **256**, 193–194, 1975.
- Margitan, J. J., F. Kaufman, and J. G. Anderson, Kinetics of the reaction $\text{OH} + \text{HNO}_3 \rightarrow \text{H}_2\text{O} + \text{NO}_3$, *Int. J. Chem. Kinet., Symp. 1*, 3310–3317, 1975.
- McElroy, M. B., S. C. Wofsy, J. E. Penner, and J. C. McConnell, Atmosphere ozone: Possible impact of stratospheric aviation, *J. Atmos. Sci.*, **31**, 287–303, 1974.
- Metzger, P. H., and G. R. Cook, A reinvestigation of the absorption cross section of molecular oxygen in the 1050–1800 Å region, *J. Quant. Spectros. Radiat. Transfer*, **4**, 107–116, 1964.
- Molina, M. J., and F. S. Rowland, Stratospheric sink for chlorofluoromethanes: Chlorine atom catalyzed destruction of ozone, *Nature*, **249**, 810–812, 1974.
- Myer, J. A., and A. R. Samson, Vacuum-ultraviolet absorption cross section of CO, HCl, and ICN between 1050 and 2100 Å, *J. Chem. Phys.*, **52**, 266–271, 1970.
- National Academy of Sciences, Environmental impact of stratospheric flight, Biological and climatic effects of aircraft emission in the stratosphere, Washington, D. C., 1975.
- Ramanathan, V., L. B. Callis, and R. E. Boughner, Sensitivity of surface temperature and atmospheric temperature to perturbations in the stratospheric concentration of ozone and nitrogen-dioxide, *J. Atmos. Sci.*, in press, 1976.
- Rao-Vupputuri, R. K., A zonally averaged circulation model of the stratosphere incorporating radiative heating and ozone photochemistry in an oxygen-hydrogen-nitrogen atmosphere, Proceedings of the Third Conference on the CIAP, pp. 390–402, Dep. of Transp., Washington, D. C., 1974.
- Rowland, F. S., and M. J. Molina, Chlorofluoromethanes in the environment, *Rev. Geophys. Space Phys.*, **13**, 1–35, 1975.
- Schott, G., and N. Davidson, Shock waves in chemical kinetics: The decomposition of N_2O_5 at high temperatures, *J. Amer. Chem. Soc.*, **80**, 1841–1853, 1958.
- Schumb, W. C., C. N. Satterfield, and R. L. Wentworth, *Hydrogen Peroxide*, pp. 266–291, Reinhold, New York, 1955.
- Stanford, J. L., Possible sink for stratosphere water vapor at the winter antarctic pole, *J. Atmos. Sci.*, **30**, 1431–1436, 1973.
- Stolarski, R. S., and R. J. Cicerone, Stratospheric chlorine: A possible sink for ozone, *Can. J. Chem.*, **52**, 1610–1615, 1974.
- Thompson, B. A., P. Harteck, and R. R. Reeves, Jr., Ultraviolet absorption coefficients of CO_2 , CO, O_3 , H_2O , N_2O , NH_3 , NO, SO_2 , and CH_4 between 1850 and 4000 Å, *J. Geophys. Res.*, **68**, 6431–6436, 1963.
- Tsang, W., Comparison between experimental and calculated rate constant for dissociation and combination reactions involving small polyatomic molecules, *Int. J. Chem. Kinet.*, **5**, 947–963, 1973.
- Watson, R. T., Chemical kinetics data survey 8: Rate constants of ClO_2 of atmospheric interest, *NBSIR 74-516*, Nat. Bur. of Stand., Washington, D. C., 1974.
- Wilson, W. E., Jr., A critical review of the gas-phase reaction kinetics of the hydroxyl radical, *J. Phys. Chem. Ref. Data*, **1**, 535–573, 1973.
- Wofsy, S. C., Atmospheric photochemistry of N-, H-, and Cl-containing radicals, Proceedings of the Third Conference on the CIAP, pp. 359–375, Dep. of Transp., Washington, D. C., 1974.
- Wofsy, S. C., and M. B. McElroy, Stratospheric halogen-ozone chemistry, paper presented at the 169th ACS National Meeting, Amer. Chem. Soc., Philadelphia, Pa., 1975.
- Wofsy, S. C., M. B. McElroy, and N. D. Sze, Freon consumption: Implication for atmospheric ozone, *Science*, **187**, 535–537, 1975.
- Wong, E. L., and F. Belles, Rate measurement for the reaction of hydrogen chloride and deuterium chloride with atomic oxygen, *NASA Tech. Note TN D-6495*, 1971.
- Zahniser, M. S., F. Kaufman, and J. G. Anderson, Kinetics of the reaction of OH with HCl, *Chem. Phys. Lett.*, **27**, 507–510, 1974.
- Zahniser, M. S., F. Kaufman, and J. G. Anderson, Kinetics of $\text{Cl} + \text{O}_3 \rightarrow \text{ClO} + \text{O}_2$, *Chem. Phys. Lett.*, **37**, 226, 1975.

(Received October 14, 1975;
revised February 10, 1976;
accepted February 10, 1976.)