UC Berkeley

UC Berkeley Previously Published Works

Title

Erratum: A finite difference Davidson procedure to sidestep full ab initio hessian calculation: Application to characterization of stationary points and transition state searches (Journal of Chemical Physics (2014) 140 (164115))

Permalink

<https://escholarship.org/uc/item/6mt1z8wj>

Journal

Journal of Chemical Physics, 140(22)

ISSN

0021-9606

Authors

Mallikarjun Sharada, S
Bell, AT
Head-Gordon, M

Publication Date

2014-06-14

DOI

10.1063/1.4883888

Peer reviewed

Erratum: “A finite difference Davidson procedure to sidestep full ab initio hessian calculation: Application to characterization of stationary points and transition state searches” [J. Chem. Phys. 140, 164115 (2014)]

Shaama Mallikarjun Sharada, Alexis T. Bell, and Martin Head-Gordon

Citation: [The Journal of Chemical Physics](#) **140**, 229902 (2014); doi: 10.1063/1.4883888

View online: <https://doi.org/10.1063/1.4883888>

View Table of Contents: <http://aip.scitation.org/toc/jcp/140/22>

Published by the [American Institute of Physics](#)

Articles you may be interested in

[A finite difference Davidson procedure to sidestep full ab initio hessian calculation: Application to characterization of stationary points and transition state searches](#)

The Journal of Chemical Physics **140**, 164115 (2014); 10.1063/1.4871660

[Note: The performance of new density functionals for a recent blind test of non-covalent interactions](#)

The Journal of Chemical Physics **145**, 186101 (2016); 10.1063/1.4967424

[The many-body expansion combined with neural networks](#)

The Journal of Chemical Physics **146**, 014106 (2017); 10.1063/1.4973380

[A time-dependent formulation of multi-reference perturbation theory](#)

The Journal of Chemical Physics **144**, 064102 (2016); 10.1063/1.4941606

[Acceleration of saddle-point searches with machine learning](#)

The Journal of Chemical Physics **145**, 074106 (2016); 10.1063/1.4960708

[MC-PDFT can calculate singlet–triplet splittings of organic diradicals](#)

The Journal of Chemical Physics **148**, 064108 (2018); 10.1063/1.5017132

PHYSICS TODAY

WHITEPAPERS

ADVANCED LIGHT CURE ADHESIVES

Take a closer look at what these environmentally friendly adhesive systems can do

READ NOW

PRESENTED BY
MASTERBOND
ADHESIVES | SEALANTS | COATINGS

Erratum: “A finite difference Davidson procedure to sidestep full *ab initio* hessian calculation: Application to characterization of stationary points and transition state searches” [J. Chem. Phys. **140**, 164115 (2014)]

Shaama Mallikarjun Sharada,¹ Alexis T. Bell,^{1,a)} and Martin Head-Gordon^{2,a)}

¹*Department of Chemical and Biomolecular Engineering, University of California, Berkeley, Berkeley, California 94720, USA*

²*Department of Chemistry, University of California, Berkeley, Berkeley, California 94720, USA*

(Received 4 June 2014; accepted 5 June 2014; published online 12 June 2014)

[<http://dx.doi.org/10.1063/1.4883888>]

Certain key contributions to the field were not presented in the Introduction section of our recent paper.¹ In addition to the variational method described in Ref. 14 and iterative diagonalization methods in Refs. 15–17, we wish to direct the attention of the reader to other significant efforts that have utilized similar principles for various applications. Reiher and Neugebauer developed a mode-tracking algorithm using the Davidson method for calculating only the modes relevant for vibrational analysis.² Similarly, a block-Davidson approach via finite differentiation of the gradient was implemented for calculating hessian-vector products for normal mode analysis.³ Deglmann and Furche developed a method for stationary point characterization with density functional theory where full solution of all coupled-perturbed Kohn-Sham equations, the bottleneck in hessian calculation, could be avoided through iterative calculation of hessian-vector products to determine only the lowest eigenvalues.⁴

We are grateful to Professor Markus Reiher for pointing out the oversight. This work was supported by a grant from Chevron Energy Technology Co.

¹S. Mallikarjun Sharada, A. T. Bell, and M. Head-Gordon, “A finite difference Davidson procedure to sidestep full *ab initio* hessian calculation: Application to characterization of stationary points and transition state searches,” *J. Chem. Phys.* **140**, 164115–164123 (2014).

²M. Reiher and J. Neugebauer, “A mode-selective quantum chemical method for tracking molecular vibrations applied to functionalized carbon nanotubes,” *J. Chem. Phys.* **118**, 1634–1641 (2003).

³A. L. Kaledin, “Gradient-based direct normal-mode analysis,” *J. Chem. Phys.* **122**, 184106–184113 (2005).

⁴P. Deglmann and F. Furche, “Efficient characterization of stationary points on potential energy surfaces,” *J. Chem. Phys.* **117**, 9535–9538 (2002).

^{a)}Authors to whom correspondence should be addressed. Electronic addresses: bell@cchem.berkeley.edu and mhg@cchem.berkeley.edu