

# UC Santa Cruz

## UC Santa Cruz Previously Published Works

### Title

On the resolution of sexual conflict over shared traits.

### Permalink

<https://escholarship.org/uc/item/6n3221bn>

### Journal

Proceedings of the Royal Society B: Biological Sciences, 291(2027)

### Authors

Pennell, Tanya

Mank, Judith

Alonzo, Suzanne

et al.

### Publication Date

2024-07-01

### DOI

10.1098/rspb.2024.0438

Peer reviewed



## Review



**Cite this article:** Pennell TM, Mank JE, Alonzo SH, Hosken DJ. 2024 On the resolution of sexual conflict over shared traits. *Proc. R. Soc. B* **291**: 20240438.

<https://doi.org/10.1098/rspb.2024.0438>

Received: 27 July 2023

Accepted: 5 July 2024

**Subject Category:**

Evolution

**Subject Areas:**

evolution, genetics, ecology

**Keywords:**

sexual conflict, intralocus conflict, resolution, dimorphism, sexual selection

**Author for correspondence:**

David J. Hosken

e-mail: [d.j.hosken@exeter.ac.uk](mailto:d.j.hosken@exeter.ac.uk)

Special Feature: The resolution of evolutionary conflicts within species. Guest edited by J. Arvid Ågren, Göran Arnqvist and Locke Rowe.

# On the resolution of sexual conflict over shared traits

Tanya M. Pennell<sup>1</sup>, Judith E. Mank<sup>2</sup>, Suzanne H. Alonzo<sup>3</sup> and David J. Hosken<sup>1</sup>

<sup>1</sup>Centre for Ecology & Conservation, Faculty of Environment, Science and Economy (ESE), University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

<sup>2</sup>Department of Zoology and Biodiversity Research Centre, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

<sup>3</sup>Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA 95060, USA

TMP, 0000-0003-2186-226X; SHA, 0000-0001-7757-0528; DJH, 0000-0003-2508-3909

Anisogamy, different-sized male and female gametes, sits at the heart of sexual selection and conflict between the sexes. Sperm producers (males) and egg producers (females) of the same species generally share most, if not all, of the same genome, but selection frequently favours different trait values in each sex for traits common to both. The extent to which this conflict might be resolved, and the potential mechanisms by which this can occur, have been widely debated. Here, we summarize recent findings and emphasize that once the sexes evolve, sexual selection is ongoing, and therefore new conflict is always possible. In addition, sexual conflict is largely a multivariate problem, involving trait combinations underpinned by networks of interconnected genes. Although these complexities can hinder conflict resolution, they also provide multiple possible routes to decouple male and female phenotypes and permit sex-specific evolution. Finally, we highlight difficulty in the study of sexual conflict over shared traits and promising directions for future research.

## 1. Introduction

Despite extensive work on the causes and fitness consequences of sexual conflict (differences in the evolutionary interests of the sexes [1–9]), many questions remain, particularly about the quantification and persistence of conflict. There are two general forms of sexual conflict. In one, shared traits that are genetically correlated between the sexes (where male and female relatives have correlated phenotypes because of shared genes) experience sexually antagonistic selection, with different trait values favoured in sperm and egg producers. In the second, fitness differences arise as a result of interactions between the sexes, with different interaction *outcomes* favoured in each. Here, we consider the current state of our understanding of the first type of conflict, with a focus on how it arises and what influences its resolution.

Terminology is key to conceptual frameworks, and the terms intralocus sexual conflict [3,5] and ontogenetic conflict [10] have both been used to describe conflict over shared phenotypes. The first term can be misleading, as it implies single locus effects on single traits, and perhaps biases our thinking towards simplistic resolutions. Many phenotypes are quantitative rather than Mendelian, and intralocus conflict does not fully capture this complexity. Ontogenetic conflict implicates whole developmental pathways and more complicated genetic architecture, which is more biologically realistic [11]. However, it also implies conflict during development, rather than between the sexes. In this review, we use the term ‘sexual conflict over shared traits’ to capture the conflict that arises due to sexually antagonistic selection –selection in opposing directions across the sexes –on traits with a shared genetic architecture. We note that this phrasing is not new (e.g. [12] and see [13]).

Sexual conflict arising from sexually antagonistic selection is as old as the evolution of the sexes. Isogamy without mating types is thought to be the ancestral gametic state [14], and subsequent disruptive selection on gamete size generated anisogamy and hence the sexes [15,16]. Anisogamy describes the universal difference between females (egg producers) and males (sperm producers) [15] and represents the initial form of sexual conflict, with disruptive selection for both small, motile gametes that 'parasitise' large, heavily provisioned gametes. The evolution of anisogamy itself therefore represents the resolution of an initial form of conflict over a shared trait, in this case gamete size, and opened the door to sexual conflict more broadly.

With gamete dimorphism, any gamete size/number trade-off [17] results in a greater abundance of sperm competing for limited eggs. This inevitably leads to sexual selection, and excess sperm generally results in sexual selection acting more strongly on males [4,18,19]. Sexual conflict follows, arising from sexually antagonistic (i.e. divergent sex-specific) selection on traits that have a shared genetic basis between the sexes (e.g. [2]). Thus, with the evolution of anisogamy, both sexual selection and conflict over shared traits are seemingly inevitable, except in idealized situations of strict genetic monogamy with no opportunity for extra-pair matings [20], or in those very rare cases where females and males do not share a genome (e.g. [21]). Sexually antagonistic natural selection also occurs, and although the relative contribution of natural and sexual selection to conflict over shared traits is unknown, sexual selection influences much sexual dimorphism and may play a proportionally larger role in generating conflict (but see [22–24]).

Here, we explore how sexual conflict over shared traits may be weakened or resolved over evolutionary time. At one extreme, strong unresolved conflict occurs when males and females express the same trait values, despite strong sexually antagonistic selection. At the other extreme, complete resolution of conflict over shared traits would mean that both sexes express the phenotypes that maximize the fitness of each sex, and sexually antagonistic selection has ceased. It should also be noted that even when sexual conflict over the shared trait no longer exists, an inherent conflict between the sexes may be ongoing, e.g. females may still do better if males invested more in sperm size to permit the production of smaller more numerous eggs [25].

Although the full resolution of conflict over shared traits may be difficult because males and females in most species share nearly, or even, all of their genome (and sexual selection is ongoing), there are many phenotypic, genetic and genomic lines of evidence that suggest many individual cases of conflict over shared traits can be largely resolved. Most notably, i) phenotypic sexual dimorphism is indicative of at least partial resolution of conflict over shared traits, ii) artificial sex-specific selection can generate dimorphism [26], and iii) selection on traits in one sex need not always produce a correlated response in the other [27]. In addition, recent work has identified a myriad of routes to conflict resolution [11].

In this review, we draw upon the latest research and explore evidence for the resolution of conflict over shared traits and the extent to which genetic independence between the sexes can evolve. We comprehensively review the mechanisms through which resolution may occur, providing an update to earlier articles centred on this topic (e.g. [5,7]). We stress the multivariate nature of conflict, and finally, we suggest future directions of research.


## 2. Towards resolution

### (a) Phenotypic studies of conflict resolution

A trait may be implicated in conflict if there is sexually antagonistic selection acting on the trait, and the trait has an intersexual genetic correlation that deviates from 0 (on a scale of -1 to 1 [2]). Despite the (at times huge) differences between male and female phenotypes [28], the sharing of a genome means traits often have high intersexual genetic correlations, with some >0.8 [29], which may retard responses to sex-specific selection. However, sexually antagonistic selection is expected to break down genetic correlations to permit sex-specific evolution and therefore ameliorate conflict. Consistent with this, weakly negative associations between the degree of sexual dimorphism and the strength of male–female genetic correlations for individual traits have been detected [29–31]. In theory, weaker associations have either allowed more sexual divergence, or antagonistic selection has favoured weaker associations. Note that a negative intersexual genetic correlation for fitness can indicate strong conflict over shared traits, but there may be conflict over traits linked to fitness even when the intersexual genetic correlation for fitness itself is weak or positive [32]. Therefore, assessing selection on individual traits is crucial to our understanding of ongoing conflict.

The relationship between dimorphism and intersexual genetic correlation is complex, and therefore, dimorphism alone cannot predict how freely each sex may respond to selection or the extent to which conflict is resolved. For example, some studies have not recovered an association between intersexual genetic correlations for phenotypic traits and the degree of sexual dimorphism [33,34]. These studies suggest that for many traits (even those that are not obviously dimorphic), sex differences in genetic architecture, heritability, dominance or genetic variance exist [35]. These differences may provide routes to conflict reduction or resolution should these traits come under sexually antagonistic selection and may explain how genetic architecture can be rapidly decoupled under artificial selection. For example, intersexual genetic correlations of floral traits broke down after just five generations of artificial selection [36]. Furthermore, intersexual genetic correlations can be quite high for sexually dimorphic traits (see figure 1 for a hypothetical example). For example, human height, an obviously dimorphic trait, also exhibits a strong intersexual correlation and seems to be subject to sexually antagonistic selection [37]. In this example, trait genetic architecture permits at least partial sexual decoupling of height values, but each sex is not at its optima and conflict persists [37,38].

In addition, intersexual genetic correlations are often described by the term  $r_{mf}$ , which considers genetic covariance between the sexes for a single trait. However, this may not be useful when predicting the evolution of sexual dimorphism and resolution of conflict when multiple traits covary [30,39], which we discuss in more detail below.



**Figure 1.** Sexual dimorphism can exist with high intersexual correlation. In this hypothetical example, mean shared trait values are largely non-overlapping between males (blue) and females (red), panels (a) and (c). However, the intersexual correlation ( $r_{mf}$ ) for a dimorphic trait, often measured by regressing average measurements of female versus male offspring within families, can vary. High  $r_{mf}$  in panel (b) is consistent with broad similarity in genetic architecture between males and females, with additional contribution from the sex determination pathway or sex hormones. Low  $r_{mf}$  in panel (d) is consistent with largely decoupled genetic architecture between the sexes.

### (b) Conflict resolution for correlated phenotypes

Many phenotypes are characterized by genetically correlated suites of traits [40], and genetic covariances are as important as genetic variances in determining phenotypic evolution [41]. For example, measures of genetic dimensionality across a broad taxonomic range show that the amount of additive genetic variation underlying individual traits in univariate analyses is far less than what is available to selection when multiple traits covary [41]. This means that arguments describing individual character evolution in isolation are incomplete because when phenotypes are a web of correlated characters, selection on any trait can ripple through the phenotype to impact other traits in unpredictable ways (e.g. [42]). In addition, it means phenotypes are constrained to some degree by genetic covariances (as well as intersexual genetic covariances), as any individual trait will be linked to other traits. Thus, while considering intersexual genetic correlation for a single trait can reveal constraints, this will not capture the full inter-relatedness of male and female phenotypes [39].

The suite of genetic variances and covariances that underpin a phenotype is described by the genetic variance–covariance matrix,  $\mathbf{G}$ . Furthermore, when considering the sexes and conflict over shared traits, estimates of the genetic variance–covariance matrix for each sex ( $\mathbf{G}_F$  and  $\mathbf{G}_M$ : where the subscripts define female and male matrices), plus the multivariate intersexual covariances for all traits,  $\mathbf{B}$ , are required.  $\mathbf{B}$  is the multivariate analogue of  $r_{mf}$  and has been extensively discussed previously [2,30,39], so we only provide a short summary of how this can affect sexual conflict over shared traits.

Modelling by Lande [2] indicated how intersexual genetic correlations can cause selection on one sex to cause maladaptive changes in the other. However, although the maladaptive phase may only be transitory and not present at equilibrium, evolution towards equilibrium may require many generations, especially for traits with high intersexual correlation [2,29]. Lande's model assumed no evolution of the genetic parameters and both a monomorphic  $\mathbf{G}$  and symmetrical  $\mathbf{B}$ . However, if  $\mathbf{G}$  and  $\mathbf{B}$  are not sexually symmetrical (not the same in both sexes or when transposed), the evolution of sex-specific phenotypes may be easier [39]. For example, if  $\mathbf{G}$  is larger in one sex, then this sex will respond more to the same selection strength, generating sexual dimorphism. Similarly, if  $\mathbf{B}$  is not symmetrical (e.g. when the covariance of trait X in females with trait Y in males is not equal to the covariance of trait Y in females and trait X in males), then one sex will respond more to sexually equivalent selection, and again dimorphism can evolve [39]. Thus, the equivalence of  $\mathbf{G}$  and  $\mathbf{B}$  across the sexes is important to our understanding of the ease with which sexual dimorphism and conflict weakening can evolve.

Simulations that relax some of Lande's [2] assumptions (but assuming low mutation rates and high mutational variance) indicate that the evolution of sexual divergence can be rapid [43]. For example, if  $\mathbf{G}$  and  $\mathbf{B}$  evolve and population size is large ( $n = 4000$ ), sexual dimorphism can evolve rapidly with relatively little change in the intersexual genetic correlation, and with a less intense and fairly short period of (female) maladaptation—between 6000 and 8000 generations [43]. Furthermore, the initial, steepest phase of sexual divergence can occur very quickly—100s of generations. However, with a smaller population

size and/or stronger antagonistic selection, there will be slower sexual divergence and reaching equilibrium can be slow, relying on mutational input rather than standing genetic variation [43].

At present, the evolution of **B** is not well understood empirically or theoretically [30]. However, investigation of within-generation changes to **B** due to sexually antagonistic selection indicates that sexual asymmetry in the strength of selection can reduce male–female genetic covariances [30], and simulations support this inference and indicate **B** can evolve rapidly although perhaps transiently [43]. This is an area that needs additional work. There is, however, more understanding of **G** [40]. Genetic architecture can differ across populations and experimental treatments, with variations in the size and orientation of **G** [40]. Thus, **G** can also evolve rapidly over relatively short timescales (also see [44]).

Importantly, comparison of **G** across the sexes indicates that males and females may never share identical genetic variances and covariances. Estimates from three taxa (house sparrows, mealworms and white campion) show male and female matrices are completely unrelated, meaning selection would cause the sexes to evolve independently [40]. Subsequent work has also shown that while **B** can be asymmetrical and sexual differences can exist in **G**, both can constrain sex-specific evolution ([45]; also see [46]). This is consistent with models showing that conflict resolution may be slow [2] and the evolution of sex differences can generate a conflict load (maladaptation during antagonistic evolution means some individuals do not reproduce: [47]). So, while there can be significant sex differences in **G** and **B**, the sexes can still constrain each other. This is in line with some experimental work showing that generating sexual dimorphism experimentally can take hundreds of generations (e.g. [26]; but see [36]). Examination of candidate sexually antagonistic loci across the whole genome in *Drosophila melanogaster* also reveals long-term constraints on sexual dimorphism, with signatures of conflict persisting for approx. 1 Myr [48].

To summarize, there is a lack of consensus on the speed of resolution, with the precise detail of **G** and **B** likely to be taxon specific (e.g. [49,50]), and their effects also influenced by the strength of sex-specific selection. Meaningful measures of sex-specific selection and a better understanding of the genetic variance–covariance matrix will aid predictions of the evolution of sexual dimorphism. Finally, additional understanding of conflict resolution may be gained through direct assessment of mechanisms (discussed below) that cause long-term changes to both **G** and **B** to allow an uncoupling of male and female phenotypes, but there has been limited research on this to date (e.g. [51]).

### (c) Omics and resolving conflict over shared traits

Simulations indicate that selection should favour across-sex decoupling of traits that are under sexually antagonistic selection [43]. However, for the majority of species, most genes in the genome are present in both sexes, which suggests that the genetic architecture underlying phenotypes is largely shared (figure 2, panels *a–c*). Below, we detail various ways in which decoupling might occur, noting that the highly genetically interconnected (**G**) nature of the phenotype means changes in any trait may have knock-on effects elsewhere in the phenome. Therefore, reducing sexual conflict over one trait could amplify it elsewhere (e.g. [42]). Moreover, although conflict over any one shared phenotype may be resolved, conflict, in general, may be perpetual due to (i) the frequency-dependent nature of sexual selection and (ii) the changing directions and targets of sexual selection, preventing populations from fully reaching equilibrium ([52] and see [53]).


As Fisher [18] noted, sex-specific modifiers (alleles or sex hormones) could alter pleiotropy in sex-specific ways to reduce intersexual genetic correlations. For example, a sex- hormone-dependent mutation that meant an allele was only ever expressed in females would result in sex differences in pleiotropic effects and enable female and male phenotypes to diverge. In addition, any mutation that acts on the initiators of sexual differentiation themselves (e.g. sex hormones), or on their effectiveness in generating differences, would also be favoured by selection [18], and work has demonstrated how testosterone shapes genetic variances (**G**) and covariances (**B**) to facilitate sexual dimorphism [51]. In species with chromosomal sex determination, sex chromosomes trigger cascades leading to sex differentiation, but autosomal genes underpin most subsequent differentiation (e.g. [54]), clearly showing how sex-specific modification can act. Below, we discuss routes to decoupling male and female phenotypes (summarized in table 1) that can involve modifiers. We then discuss sex chromosomes in the context of conflict resolution. Finally, we highlight strategic solutions (sex-ratio adjustment) that may mask the effects of sexually antagonistic alleles. Some of this has been discussed previously (e.g. [5,7,55]), and our goal is to provide an update.

#### (i) Regulatory solutions

Regulatory decoupling (e.g. differential gene expression) can potentially alleviate conflict over shared traits and permit sexually independent evolution. Depending on the dataset, a large proportion of coding genes are expressed differently between the sexes in at least one tissue or cell type [56,57]. Sex-biased expression varies across closely related species and even populations (reviewed in [58]), suggesting it evolves rapidly, as evidenced by rapid responses to artificial selection [59,60]. Other studies have found reduced signatures of conflict associated with genome-wide sex-biased expression patterns [61,62]. Furthermore, negative associations between sex-biased expression and the occurrence of candidate sexually antagonistic loci demonstrate the potential role of regulatory decoupling in conflict resolution [48].

#### (ii) Non-regulatory solutions

Decoupling the sexes does not necessarily require dimorphism in gene regulation itself. Even for genes that are expressed at similar levels in both sexes, genetic variants may affect one sex more, or even predominantly [63], offering a potential route to decouple male and female genetic architecture and conflict reduction. Moreover, even for sexually monomorphic traits, there is evidence that genetic variants have sex-specific effects ([33], figure 2*d*). This could occur through sex-specific alternative



**Figure 2.** Alternative scenarios for genetic architecture and sexual dimorphism. In each panel, small circles outlined in black represent all autosomal or X-linked genes in a developmental pathway underlying a phenotype. Solid lines represent loci that contribute to variation in phenotype (genetic architecture) and the width represents effect sizes. Inner circles represent phenotype distribution in females (red) and males (blue) as well as phenotypic optima (dashed lines). Panel (a): representation of the traditional conceptualization of sexual conflict, where phenotypic optima differ substantially but genetic architectures are largely overlapping. Panel (b): male and female phenotypes can diverge towards phenotypic optima as more of the shared architecture affects only females (red circles and lines) or males (blue circles and lines). Panel (c): non-overlapping phenotypic distributions occur with complete decoupling of male and female architecture of a phenotype. Panel (d): recent work suggests that many monomorphic traits have sex differences in genetic architecture. Panel (e): loci on the Y chromosome (or W in ZW systems) can lead to sex-specific phenotypic distributions even when the remainder of the genetic architecture is shared.

**Table 1.** Separating the sexes. Summaries of the main mechanisms enabling an uncoupling of male and female phenotypes.

main mechanisms to uncouple male- and female-shared phenotypes	description
sex-specific gene regulation	genes are expressed at different levels depending on sex (male or female)
alternative splicing	genes can be spliced differently depending on sex: different combinations of exons are joined to produce functionally different proteins
gene duplication	genes are duplicated and moved to locations where they can be regulated in a sex-specific manner
genomic imprinting	an allele is expressed according to its parent-of-origin, where maternally derived alleles are silenced (via DNA methylation) in males and paternally derived alleles are silenced in females at sexually antagonistic loci
dominance reversal	each allele at a sexually antagonistic locus is partially or completely dominant when expressed in the sex it is favoured and recessive in the sex it harms
sex chromosomes (Y or W chromosome)	the Y (or W) chromosome is a region that is limited to one sex (males or females in ZW systems), which could accumulate male-benefit alleles (or female-benefit alleles in ZW systems)
sex chromosomes (dominance effects)	the X chromosome can facilitate dominance effects, with selection favouring male-benefit alleles that are recessive and female-benefit alleles that are dominant
sex chromosomes (dosage compensation)	the sexes can differ in gene expression and protein production due to differences in sex-chromosome number
sex-ratio adjustment	a mechanism to mask the effects of sexually antagonistic alleles that does not require changes to trait genetic architecture. The sex of offspring is adjusted by the female depending on the assessment of alleles (e.g. male- or female-benefit) carried by an individual (self-assessment) or their mate

splicing—joining of different exon combinations from the same gene to produce functionally different proteins—which occurs post-transcriptionally. Alternative splicing could be important in dimorphism generation because it avoids pleiotropic/functional constraints associated with altering gene expression levels. Although alternative splicing has been implicated in sexual

dimorphism, very little is known about the phenotypic consequences and adaptive significance of sex-specific alternative splicing (reviewed in [64]). However, sex-biased expression and sex-specific alternative splicing are negatively correlated [65,66], suggesting the evolution of one might negate the need for the other. Alternative splicing may also facilitate independent evolution in the sexes [66].

### (iii) Gene duplication

Gene duplication could also facilitate conflict resolution if paralogues adopt sex-specific function ([67,68] and see [69]). In *D. melanogaster* and other taxa, sexual dimorphism may be facilitated by male-biased expression of duplicate genes (reviewed in [69]), which is likely driven by sexual selection in males. In addition, tandem duplicates in *D. melanogaster* apparently resolve conflict over fertility traits [70]. Interestingly, although each autosomal duplicate undergoes sex-biased expression to largely resolve conflict, some antagonistic effects are still present due to residual expression in the 'harmed' sex. The (partial) resolution of this conflict through sex-biased expression and sequence changes occurred reasonably rapidly (approx. 2 Myr) [70].

### (iv) Imprinting

Epigenetic parent-of-origin imprinting, whereby an allele is expressed according to its parent-of-origin, is another potential route to resolving conflict [71]. For example, males (females) that have passed the filter of sex-specific selection are more likely to carry alleles at sexually antagonistic loci that benefit their sex, and so it follows that selection should favour silencing the maternally derived allele in males (or the paternally derived allele in females) [71]. Parent-of-origin effects might also propagate to other linked genes, even those unimprinted [72], altering the expression of multiple genes at once. Imprinting is widespread and occurs in a range of taxa (e.g. [73–75]). However, empirical studies of either sex-dependent imprinting [76–79] or its role in the resolution of conflict over shared traits [80,81] are currently limited.

### (v) Dominance reversal

Dominance reversal, where alleles are partially or completely dominant when expressed in the sex they benefit and recessive otherwise, can also act to partially mitigate conflict [82–86]. In addition, the occurrence of dominance reversal could favour the accumulation of sexually antagonistic alleles on the autosomes, instead of the X chromosome [85]. Although signatures of dominance reversal may be difficult to detect, multiple promising methods have been proposed (reviewed in [86]). Sex-dependent dominance reversal has been identified in a single large-effect locus determining salmon age at maturity [83], in a supergene that mediates trout migration tendency [87] and in a polygenic trait underlying *Drosophila* immunocompetence [88]. Seed beetles also show genome-wide dominance reversals for sexually antagonistic alleles underlying fitness [84]. As noted by Grieshop *et al.* [86], dominance reversals mitigate but do not fully resolve conflict because homozygotes will always exist to some degree in the sex where the genotype is not favoured.

### (vi) The sex chromosomes

Owing to the sexual asymmetry in X (or Z) chromosome inheritance, theory predicts that sex-chromosome linkage can decouple female and male phenotypes and ameliorate sexually antagonistic effects (e.g. [89,90]). Specifically, male-benefit recessive alleles and female-benefit dominant alleles are expected to accumulate on the X (or Z) chromosome [90,91], with dominance effects acting to partially resolve conflict at these antagonistic loci. Complete dimorphism may then be facilitated by sex-specific modifiers that act on these sexually antagonistic loci ([90]; but see [92]).

Precise predictions regarding the accumulation of X-linked sexually antagonistic alleles are complicated by mechanisms of dosage compensation (reviewed in [93]). For example, if dosage compensation involves the inactivation of one X in females, the X-linked male recessive advantage (discussed above) no longer exists [93]. It is also worth noting that the lack of dosage compensation can itself facilitate sexual differentiation: the sexes can differ in gene expression and protein production due to differences in sex-chromosome number, potentially reducing phenotypic conflict [5]. Although there is evidence consistent with the accumulation of sexually antagonistic alleles on the X chromosome (e.g. [94–98]; but see [48]), empirical proxies of sexually antagonistic fitness variation may be biased towards detecting X-linked effects [99]. Furthermore, one unbiased test has revealed no enrichment of sexually antagonistic alleles on the human X chromosome [100], but further studies controlling for X-linked biases in a range of taxa are required.

While the role of X and Z chromosomes in conflict resolution is unclear, W and Y chromosomes provide sex-limited regions of the genome that can accumulate genes underlying sex-specific benefits [89]. Movement of male-benefit alleles to the Y chromosome (or female-benefit alleles to the W in ZW systems) can effectively decouple male and female phenotypes (figure 2e), and Y chromosomes continuously accumulate genes [101–104] with important male-phenotypic effects (e.g. [105–107]). For example, Y-linked additive variation impacts body-size sexual dimorphism by as much as 30% in seed beetles [108]. Studies are also increasingly demonstrating that Y-linked genes have significant epistatic effects (e.g. [109–113]), which could be important for sexual dimorphism. Further work examining the impact of sex determination systems on conflict resolution is required.

### (vii) Sex ratio

In addition to mechanisms that resolve conflict over shared traits, offspring sex-ratio adjustment could mask effects of sexually antagonistic loci ([114]; also see [115]). For example, females might produce excess sons and fewer daughters if they mate

with highly masculinized males, or they may be able to vary the sex ratio based on their own phenotype [116–121]. Male traits targeted by sexual selection, which are usually honest indicators of male quality [120,122–124], provide an obvious means for females to assess whether males carry ‘excess’ male-benefit alleles [125,126]. A single-locus model suggests that sex-ratio adjustment is likely to evolve in response to autosomal antagonistic alleles, but if antagonistic alleles are X-linked, the adjustment should be based on the homogametic genotype [114]. These strategic solutions should be kept in mind when assessing the implications of ongoing conflict over shared traits.

### (viii) Resolution: the overall picture

There is still much progress to be made towards understanding how, and how effectively, the possible mechanisms above uncouple male and female phenotypes to resolve conflict. For example, although links between mechanisms and sexual dimorphism are often clear, impacts on sexual antagonism are rarely quantified. In addition, conflict is often studied in a univariate context, but the maintenance of sexually antagonistic fitness variation likely involves multiple traits linked by networks of interconnected genes, highlighting the need to explore conflict and its resolution in multivariate phenotypic space. It should be noted that pleiotropy and complicated gene networks may hinder conflict resolution in some instances, but they also provide many potential avenues for conflict resolution. Finally, studies across a broad taxonomic range are required, as variation in trait genetic architecture across species may impact the scope for these mechanisms to facilitate the evolution of sexually dimorphic phenotypes.

We also highlight that in the absence of complete conflict resolution, males may often be closer to fitness-maximizing phenotypes than females because of sexual selection. For example, stronger sexual selection on males may mean more masculinized phenotypes contribute disproportionately to the gene pool shared between the sexes [127,128], and there is some evidence that genomes are masculinized in populations where sexual selection is present, compared to those with experimentally enforced monogamy and no sexual selection ([59]; but also see [60,129,130]).

## 3. Empirical concerns and future directions

To fully understand conflict over shared traits, we need greater understanding of three fundamental parameters: sexually antagonistic selection, genetic variance–covariance and intersexual covariance matrices. Without documenting antagonistic selection, it is not possible to know whether conflict over a shared phenotype is present. Without understanding the genetics, we cannot determine the extent to which male and female phenotypes are decoupled or how free they are to respond to sex-specific selection. Unfortunately, it is difficult to comprehensively estimate any of these fundamental parameters with any single approach.

If our goal is to understand how sexual conflict affects natural populations, one could argue that one must go outside to study it; however, studying natural systems is difficult. For example, measures of selection are inevitably incomplete (only partial phenotypes are measured), and estimates often involve the assessment of single or small numbers of selective episodes. Thus, we may not be assessing the true targets of selection and may miss the most significant episodes of selection, including those in which sexual antagonism occurs. Long-term studies can address at least the latter issue, and as one such study shows, long-term relatively weak selection can change direction rapidly and significantly to reverse trait evolution [131]. Nonetheless, studies in natural systems that have focused on antagonistic phenotypes have revealed important aspects of sexual conflict and its resolution (e.g. [83,132]), and others have tracked allele frequency changes to reveal the interplay between selection and sexual conflict [133]. However, estimating  $G$  and  $B$  is difficult, generally requiring large sample sizes and known pedigree structures, and invariably estimates are incomplete. This is also true for field studies.

The power of experimental evolution and artificial selection in understanding conflict over shared traits has been highlighted recently [55,134], and these approaches overcome some of the difficulties mentioned above, namely the complexity of natural environments and the difficulty of multigenerational studies. Examples include sex-limited evolution, where selection on the whole phenome is limited to one sex (e.g. [135–139]), and artificial selection on specific phenotypes in one sex (e.g. [42,140–142]). However, simplistic laboratory environments that make the study of conflict more tractable may inflate sexual conflict and limit responses to it [143–145,145,146]. The very strong selection coefficients in laboratory studies may also bias estimates of genetic architecture underlying sexual conflict. Because of this, experimental evolution may be best suited to studying the rapid resolution of conflict [26,36] or the consequences of its cessation [147]. In addition, laboratory selection studies frequently involve insects and *Wolbachia* infection, which is common in these animals, can inflate estimates of sexual conflict over shared traits [148].

Linking genotype and phenotype is also crucial in understanding sexual conflict over shared traits. Evolve-and-resequence methods (e.g. [149]; discussed in [9,134]) powerfully enable the mapping of phenotypes to genome-wide sequence variation, biological processes or genetic pathways. Experimental manipulation of candidate genes and measurement of sex-specific fitness effects can also be used to verify ongoing antagonism (e.g. [70,150]). Importantly, understanding the biological significance of antagonism requires genetic manipulations in different tissue types and quantification of multivariate phenotypes over various life stages.

There is still much to learn about sexual conflict over shared traits, and critically, there are few documented loci with demonstrated sexually antagonistic effects. Detecting the genomic signatures of these loci is made difficult by the noisy nature of the signals and the complexity of selection acting on any portion of the genome [151]. However, genomic data coupled with phenotypic studies and fitness measures can identify antagonistic loci (e.g. [83,152]) and reveal the longevity of



sexually antagonistic loci. In addition, experimental evolution approaches coupled with more natural levels of environmental heterogeneity [143–145,145,146] will help to disentangle factors influencing conflict. Overall, the wider implications of sexual conflict over shared traits are becoming increasingly clear, from genome architecture (e.g. [11,90,151]) to population viability (e.g. [153–155]) and even human health (e.g. [156–158]). As a result, work clarifying and deepening our understanding of this conflict is likely to deliver benefits on multiple fronts.

**Ethics.** This work did not require ethical approval from a human subject or animal welfare committee.

**Data accessibility.** This article has no additional data.

**Declaration of AI use.** We have not used AI-assisted technologies in creating this article.

**Authors' contributions.** T.M.P.: writing—original draft, writing—review and editing; J.E.M.: conceptualization, writing—original draft, writing—review and editing; S.H.A.: conceptualization, writing—original draft, writing—review and editing; D.J.H.: conceptualization, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

**Conflict of interest declaration.** We declare we have no competing interests.

**Funding.** This work was supported by the National Science Foundation Grant IOS-1655297 and funds from the University of California Santa Cruz (S.H.A.), NSERC and a Canada 150 Research Chair (J.E.M.) and the University of Exeter and the Leverhulme Trust (D.J.H.).

**Acknowledgements.** We thank the Editors for kindly inviting us to contribute to this special issue. We discovered during writing that we actually agree on more about conflict resolution than we thought. We also thank the reviewers and Editor for their helpful comments that enabled us to greatly improve the manuscript and its presentation. We additionally thank the participants of a 2019 workshop at UBC for helping us better understand the issues we discuss here and the Exeter-UBC Partnership Development Fund for supporting that event.

## References

- Parker GA. 1979 Sexual selection and sexual conflict. In *Sexual selection and reproductive competition in insects* (eds MS Blum, NA Blum), pp. 123–166. New York, NY: Academic Press.
- Lande R. 1980 Sexual dimorphism, sexual selection, and adaptation in polygenic characters. *Evolution* **34**, 292–305. (doi:10.1111/j.1558-5646.1980.tb04817.x)
- Rice WR, Holland B. 1997 The enemies within: intergenomic conflict, interlocus contest evolution (ICE), and the intraspecific red queen. *Behav. Ecol. Sociobiol.* **41**, 1–10. (doi:10.1007/s002650050357)
- Arnqvist G, Rowe L. 2005 *Sexual conflict*. Princeton, NJ: Princeton University Press.
- Bonduriansky R, Chenoweth SF. 2009 Intralocus sexual conflict. *Trends Ecol. Evol.* **24**, 280–288. (doi:10.1016/j.tree.2008.12.005)
- van Doorn GS. 2009 Intralocus sexual conflict. *Ann. NY Acad. Sci.* **1168**, 52–71. (doi:10.1111/j.1749-6632.2009.04573.x)
- Pennell TM, Morrow EH. 2013 Two sexes, one genome: the evolutionary dynamics of intralocus sexual conflict. *Ecol. Evol.* **3**, 1819–1834. (doi:10.1002/ece3.540)
- Rice WR, Gavrilets S. 2014 *The genetics and biology of sexual conflict*. New York, NY: Cold Spring Harbor Laboratory Press.
- Schenkel MA, Pen I, Beukeboom LW, Biller JC. 2018 Making sense of intralocus and interlocus sexual conflict. *Ecol. Evol.* **8**, 13035–13050. (doi:10.1002/ece3.4629)
- Chippindale AK, Gibson JR, Rice WR. 2001 Negative genetic correlation for adult fitness between sexes reveals ontogenetic conflict in *Drosophila*. *Proc. Natl Acad. Sci. USA* **98**, 1671–1675. (doi:10.1073/pnas.98.4.1671)
- Price PD, Parkus SM, Wright AE. 2023 Recent progress in understanding the genomic architecture of sexual conflict. *Curr. Opin. Genet. Dev.* **80**, 102047. (doi:10.1016/j.gde.2023.102047)
- Partridge L, Hurst LD. 1998 Sex and conflict. *Science* **281**, 2003–2008. (doi:10.1126/science.281.5385.2003)
- Rowe L, Day T. 2006 Detecting sexual conflict and sexually antagonistic coevolution. *Phil. Trans. R. Soc. B* **361**, 277–285. (doi:10.1098/rstb.2005.1788)
- Lessells CM, Snook RR, Hosken DJ. 2009 The evolutionary origin and maintenance of sperm: selection for a small, motile gamete mating type. In *Sperm biology: an evolutionary perspective* (eds TR Birkhead, DJ Hosken, S Pitnick), pp. 43–67. New York, NY: Academic Press.
- Parker GA, Baker RR, Smith VGF. 1972 The origin and evolution of gamete dimorphism and the male–female phenomenon. *J. Theor. Biol.* **36**, 529–553. (doi:10.1016/0022-5193(72)90007-0)
- Lehtonen J, Kokko H, Parker GA. 2016 What do isogamous organisms teach us about sex and the two sexes *Phil. Trans. R. Soc. B* **371**, 20150532. (doi:10.1098/rstb.2015.0532)
- Parker GA. 1984 Sperm competition and the evolution of animal mating strategies. In *Sperm competition and the evolution of animal mating systems* (ed. RL Smith), pp. 1–60. New York, NY: Academic Press.
- Fisher RA. 1930 *The genetical theory of natural selection*. Oxford, UK: Clarendon Press.
- Andersson MB. 1994 *Sexual selection*. Princeton, NJ: Princeton University Press.
- Hosken DJ, Stockley P, Tregenza T, Wedell N. 2009 Monogamy and the battles of the sexes. *Annu. Rev. Entomol.* **54**, 361–378. (doi:10.1146/annurev.ento.54.110807.090608)
- Fournier D, Estoup A, Orivel J, Foucaud J, Jourdan H, Le Breton J, Keller L. 2005 Clonal reproduction by males and females in the little fire ant. *Nature* **435**, 1230–1234. (doi:10.1038/nature03705)
- Littleford-Colquhoun BL, Clemente C, Thompson G, Cristescu RH, Peterson N, Strickland K, Stuart-Fox D, Frere CH. 2019 How sexual and natural selection shape sexual size dimorphism: evidence from multiple evolutionary scales. *Funct. Ecol.* **33**, 1446–1458. (doi:10.1111/1365-2435.13337)
- Silva NR, Berneck BVM, da Silva HR, Haddad CFB, Zamudio KR, Mott T, Nali RC, Prado CPA. 2020 Egg-laying site, fecundity and degree of sexual size dimorphism in frogs. *Biol. J. Linn. Soc.* **131**, 600–610. (doi:10.1093/biolinnean/blaa126)
- Chelini MC, Brock K, Yeager J, Edwards DL. 2021 Environmental drivers of sexual dimorphism in a lizard with alternative mating strategies. *J. Evol. Biol.* **34**, 1241–1255. (doi:10.1111/jeb.13881)
- Parker GA. 1978 Selection on non-random fusion of gametes during evolution of anisogamy. *J. Theor. Biol.* **73**, 1–28. (doi:10.1016/0022-5193(78)90177-7)
- Stewart AD, Rice WR. 2018 Arrest of sex-specific adaptation during the evolution of sexual dimorphism in *Drosophila*. *Nat. Ecol. Evol.* **2**, 1507–1513. (doi:10.1038/s41559-018-0613-4)
- Corral-Lopez A *et al.* 2024 Functional convergence of genomic and transcriptomic architecture underlying sociability in a live-bearing fish. *Nat. Ecol. Evol.* **8**, 98–110. (doi:10.1038/s41559-023-02249-9)

28. Fairbairn DJ. 2013 *Odd couples: extraordinary differences between the sexes in the animal kingdom*. Princeton, NJ: Princeton University Press.
29. Poissant J, Wilson AJ, Coltman DW. 2010 Sex-specific genetic variance and the evolution of sexual dimorphism: a systematic review of cross-sex genetic correlations. *Evolution* **64**, 97–107. (doi:10.1111/j.1558-5646.2009.00793.x)
30. McGlothlin JW, Cox RM, Brodie ED III. 2019 Sex-specific selection and the evolution of between-sex genetic covariance. *J. Hered.* **110**, 422–432. (doi:10.1093/jhered/esz031)
31. Fairbairn DJ, Roff DA, Wolak ME. 2023 Tests for associations between sexual dimorphism and patterns of quantitative genetic variation in the water strider, *Aquarius remigis*. *Heredity* **131**, 109–118. (doi:10.1038/s41437-023-00626-5)
32. Connallon T, Matthews G. 2019 Cross-sex genetic correlations for fitness and fitness components: connecting theoretical predictions to empirical patterns. *Evol. Lett.* **3**, 254–262. (doi:10.1002/evl3.116)
33. van der Bijl W, Mank JE. 2021 Widespread cryptic variation in genetic architecture between the sexes. *Evol. Lett.* **5**, 359–369. (doi:10.1002/evl3.245)
34. Kaufmann P, Howie JM, Immonen E. 2023 Sexually antagonistic selection maintains genetic variance when sexual dimorphism evolves. *Proc. R. Soc. B* **290**, 20222484. (doi:10.1098/rspb.2022.2484)
35. Khrantsova EA, Davis LK, Stranger BE. 2019 The role of sex in the genomics of human complex traits. *Nat. Rev. Genet.* **20**, 173–190. (doi:10.1038/s41576-018-0083-1)
36. Delph LF, Steven JC, Anderson IA, Herlihy CR, Brodie ED. 2011 Elimination of a genetic correlation between the sexes via artificial correlational selection. *Evolution* **65**, 2872–2880. (doi:10.1111/j.1558-5646.2011.01350.x)
37. Stulp G, Kuijper B, Buunk AP, Pollet TV, Verhulst S. 2012 Intralocus sexual conflict over human height. *Biol. Lett.* **8**, 976–978. (doi:10.1098/rsbl.2012.0590)
38. Sanjak JS, Sidorenko J, Robinson MR, Thornton KR, Visscher PM. 2018 Evidence of directional and stabilizing selection in contemporary humans. *Proc. Natl Acad. Sci. USA* **115**, 151–156. (doi:10.1073/pnas.1707227114)
39. Wyman MJ, Stinchcombe JR, Rowe L. 2013 A multivariate view of the evolution of sexual dimorphism. *J. Evol. Biol.* **26**, 2070–2080. (doi:10.1111/jeb.12188)
40. Arnold SJ, Bürger R, Hohenlohe PA, Ajie BC, Jones AG. 2008 Understanding the evolution and stability of the **G**-matrix. *Evolution* **62**, 2451–2461. (doi:10.1111/j.1558-5646.2008.00472.x)
41. Kirkpatrick M. 2009 Patterns of quantitative genetic variation in multiple dimensions. *Genetica* **136**, 271–284. (doi:10.1007/s10709-008-9302-6)
42. Harano T, Okada K, Nakayama S, Miyatake T, Hosken DJ. 2010 Intralocus sexual conflict unresolved by sex-limited trait expression. *Curr. Biol.* **20**, 2036–2039. (doi:10.1016/j.cub.2010.10.023)
43. Reeve JP, Fairbairn DJ. 2001 Predicting the evolution of sexual size dimorphism. *J. Evol. Biol.* **14**, 244–254. (doi:10.1046/j.1420-9101.2001.00276.x)
44. Milocco L, Salazar-Ciudad I. 2022 Evolution of the **G** matrix under nonlinear genotype-phenotype maps. *Am. Nat.* **199**, 420–435. (doi:10.1086/717814)
45. Lewis Z, Wedell N, Hunt J. 2011 Evidence for strong intralocus sexual conflict in the Indian meal moth, *Plodia interpunctella*. *Evolution* **65**, 2085–2097. (doi:10.1111/j.1558-5646.2011.01267.x)
46. Szepeanacz JL, Houle D. 2019 Cross-sex genetic covariances limit the evolvability of wing-shape within and among species of *Drosophila*. *Evolution* **73**, 1617–1633. (doi:10.1111/evo.13788)
47. Matthews G, Hangartner S, Chapple DG, Connallon T. 2019 Quantifying maladaptation during the evolution of sexual dimorphism. *Proc. R. Soc. B* **286**, 20191372. (doi:10.1098/rspb.2019.1372)
48. Ruzicka F, Hill MS, Pennell TM, Flis I, Ingleby FC, Mott R, Fowler K, Morrow EH, Reuter M. 2019 Genome-wide sexually antagonistic variants reveal long-standing constraints on sexual dimorphism in fruit flies. *PLoS Biol.* **17**, e3000244. (doi:10.1371/journal.pbio.3000244)
49. Cox RM, Costello RA, Camber BE, McGlothlin JW. 2017 Multivariate genetic architecture of the *Anolis* dewlap reveals both shared and sex-specific features of a sexually dimorphic ornament. *J. Evol. Biol.* **30**, 1262–1275. (doi:10.1111/jeb.13080)
50. Iglesias PP, Machado FA, Llanes S, Hasson E, Soto EM. 2023 Opportunities and constraints imposed by the **G** matrix of *Drosophila buzzatii* wings. *Evol. Biol.* **50**, 127–136. (doi:10.1007/s11692-022-09593-x)
51. Wittman TN, Robinson CD, McGlothlin JW, Cox RM. 2021 Hormonal pleiotropy structures genetic covariance. *Evol. Lett.* **5**, 397–407. (doi:10.1002/evl3.240)
52. Gavrilets S. 2000 Rapid evolution of reproductive barriers driven by sexual conflict. *Nature* **403**, 886–889. (doi:10.1038/35002564)
53. Pennell TM, de Haas FJH, Morrow EH, van Doorn GS. 2016 Contrasting effects of intralocus sexual conflict on sexually antagonistic coevolution. *Proc. Natl Acad. Sci. USA* **113**, E978–E986. (doi:10.1073/pnas.1514328113)
54. Kashimada K, Koopman P. 2010 *Sry*: the master switch in mammalian sex determination. *Development* **137**, 3921–3930. (doi:10.1242/dev.048983)
55. Rowe L, Chenoweth SF, Agrawal AF. 2018 The genomics of sexual conflict. *Am. Nat.* **192**, 274–286. (doi:10.1086/698198)
56. Oliva M *et al.* 2020 The impact of sex on gene expression across human tissues. *Science* **369**, eaba3066. (doi:10.1126/science.aba3066)
57. Darolti I, Mank JE. 2023 Sex-biased gene expression at single-cell resolution: cause and consequence of sexual dimorphism. *Evol. Lett.* **7**, 148–156. (doi:10.1093/evlett/qrqad013)
58. Mank JE. 2017 Population genetics of sexual conflict in the genomic era. *Nat. Rev. Genet.* **18**, 721–730. (doi:10.1038/nrg.2017.83)
59. Hollis B, Houle D, Yan Z, Kawecki TJ, Keller L. 2014 Evolution under monogamy feminizes gene expression in *Drosophila melanogaster*. *Nat. Commun.* **5**, 3482. (doi:10.1038/ncomms4482)
60. Veltsos P, Fang Y, Cossins AR, Snook RR, Ritchie MG. 2017 Mating system manipulation and the evolution of sex-biased gene expression in *Drosophila*. *Nat. Commun.* **8**, 2072. (doi:10.1038/s41467-017-02232-6)
61. Wright AE, Fumagalli M, Cooney CR, Bloch NI, Vieira FG, Buechel SD, Kolm N, Mank JE. 2018 Male-biased gene expression resolves sexual conflict through the evolution of sex-specific genetic architecture. *Evol. Lett.* **2**, 52–61. (doi:10.1002/evl3.39)
62. Sayadi A, Martinez Barrio A, Immonen E, Dainat J, Berger D, Tellgren-Roth C, Nystedt B, Arnqvist G. 2019 The genomic footprint of sexual conflict. *Nat. Ecol. Evol.* **3**, 1725–1730. (doi:10.1038/s41559-019-1041-9)
63. Karp NA *et al.* 2017 Prevalence of sexual dimorphism in mammalian phenotypic traits. *Nat. Commun.* **8**, 15475. (doi:10.1038/ncomms15475)
64. Singh P, Ahi EP. 2022 The importance of alternative splicing in adaptive evolution. *Mol. Ecol.* **31**, 1928–1938. (doi:10.1111/mec.16377)
65. Rogers TF, Palmer DH, Wright AE. 2021 Sex-specific selection drives the evolution of alternative splicing in birds. *Mol. Biol. Evol.* **38**, 519–530. (doi:10.1093/molbev/msaa242)
66. Singh A, Agrawal AF. 2023 Two forms of sexual dimorphism in gene expression in *Drosophila melanogaster*: their coincidence and evolutionary genetics. *Mol. Biol. Evol.* **40**, msad091. (doi:10.1093/molbev/msad091)
67. Gallach M, Betrán E. 2011 Intralocus sexual conflict resolved through gene duplication. *Trends Ecol. Evol.* **26**, 222–228. (doi:10.1016/j.tree.2011.02.004)
68. Connallon T, Clark AG. 2011 The resolution of sexual antagonism by gene duplication. *Genetics* **187**, 919–937. (doi:10.1534/genetics.110.123729)
69. Castellanos MDP, Wickramasinghe CD, Betrán E. 2024 The roles of gene duplications in the dynamics of evolutionary conflicts. *Proc. R. Soc. B* **291**, 20240555. (doi:10.1098/rspb.2024.0555)

70. Van Kuren NW, Long M. 2018 Gene duplicates resolving sexual conflict rapidly evolved essential gametogenesis functions. *Nat. Ecol. Evol.* **2**, 705–712. (doi:10.1038/s41559-018-0471-0)
71. Day T, Bonduriansky R. 2004 Intralocus sexual conflict can drive the evolution of genomic imprinting. *Genetics* **167**, 1537–1546. (doi:10.1534/genetics.103.026211)
72. Macias-Velasco JF *et al.* 2022 Parent-of-origin effects propagate through networks to shape metabolic traits. *elife* **11**, e72989. (doi:10.7554/eLife.72989)
73. Wood AJ, Oakey RJ. 2006 Genomic imprinting in mammals: emerging themes and established theories. *PLoS Genet.* **2**, e147. (doi:10.1371/journal.pgen.0020147)
74. Renfree MB, Hore TA, Shaw G, Graves JAM, Pask AJ. 2009 Evolution of genomic imprinting: insights from marsupials and monotremes. *Annu. Rev. Genomics Hum. Genet.* **10**, 241–262. (doi:10.1146/annurev-genom-082908-150026)
75. Matsuura K. 2020 Genomic imprinting and evolution of insect societies. *Pop. Ecol.* **62**, 38–52. (doi:10.1002/1438-390X.12026)
76. Hager R, Cheverud JM, Wolf JB. 2008 Maternal and paternal effects as the cause of parent-of-origin effects that mimic genomic imprinting. *Genetics* **178**, 1755–1762. (doi:10.1534/genetics.107.080697)
77. Gregg C, Zhang J, Butler JE, Haig D, Dulac C. 2010 Sex-specific parent-of-origin allelic expression in the mouse brain. *Science* **329**, 682–685. (doi:10.1126/science.1190831)
78. Faisal M, Kim H, Kim J. 2014 Sexual differences of imprinted genes' expression levels. *Gene* **533**, 434–438. (doi:10.1016/j.gene.2013.10.006)
79. Kincaid-Smith J, Picard MAL, Cosseau C, Boissier J, Severac D, Grunau C, Toulza E. 2018 Parent-of-origin-dependent gene expression in male and female schistosome parasites. *Genome Biol. Evol.* **10**, 840–856. (doi:10.1093/gbe/evy037)
80. Abbott JK, Innocenti P, Chippindale AK, Morrow EH. 2013 Epigenetics and sex-specific fitness: an experimental test using male-limited evolution in *Drosophila melanogaster*. *PLoS One* **8**, e70493. (doi:10.1371/journal.pone.0070493)
81. Tanaka K, Besson V, Rivagorda M, Oury F, Marazzi G, Sassono DA. 2022 Paternally expressed gene 3 (*Pw1/Peg3*) promotes sexual dimorphism in metabolism and behavior. *PLoS Genet.* **18**, e1010003. (doi:10.1371/journal.pgen.1010003)
82. Spencer HG, Priest NK. 2016 The evolution of sex-specific dominance in response to sexually antagonistic selection. *Am. Nat.* **187**, 658–666. (doi:10.1086/685827)
83. Barson NJ *et al.* 2015 Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature* **528**, 405–408. (doi:10.1038/nature16062)
84. Grieshop K, Arnqvist G. 2018 Sex-specific dominance reversal of genetic variation for fitness. *PLoS Biol.* **16**, e2006810. (doi:10.1371/journal.pbio.2006810)
85. Connallon T, Chenoweth SF. 2019 Dominance reversals and the maintenance of genetic variation for fitness. *PLoS Biol.* **17**, e3000118. (doi:10.1371/journal.pbio.3000118)
86. Grieshop K, Ho EK, Kasimatis KR. 2024 Dominance reversals: the resolution of genetic conflict and maintenance of genetic variation. *Proc. R. Soc. B* **291**, 20232816. (doi:10.1098/rspb.2023.2816)
87. Pearse DE *et al.* 2019 Sex-dependent dominance maintains migration supergene in rainbow trout. *Nat. Ecol. Evol.* **3**, 1731–1742. (doi:10.1038/s41559-019-1044-6)
88. Geeta Arun M, Agarwala A, Syed ZA, Kashyap M, Venkatesan S, Chechi TS, Gupta V, Prasad NG. 2021 Experimental evolution reveals sex-specific dominance for surviving bacterial infection in laboratory populations of *Drosophila melanogaster*. *Evol. Lett.* **5**, 657–671. (doi:10.1002/evl3.259)
89. Fisher RA. 1931 The evolution of dominance. *Biol. Rev.* **6**, 345–368. (doi:10.1111/j.1469-185X.1931.tb01030.x)
90. Rice WR. 1984 Sex chromosomes and the evolution of sexual dimorphism. *Evolution* **38**, 735–742. (doi:10.1111/j.1558-5646.1984.tb00346.x)
91. Patten MM. 2019 The X chromosome favours males under sexually antagonistic selection. *Evolution* **73**, 84–91. (doi:10.1111/evo.13646)
92. Connallon T, Clark AG. 2010 Sex linkage, sex-specific selection, and the role of recombination in the evolution of sexually dimorphic gene expression. *Evolution* **64**, 3417–3442. (doi:10.1111/j.1558-5646.2010.01136.x)
93. Fairbairn DJ, Roff DA. 2006 The quantitative genetics of sexual dimorphism: assessing the importance of sex-linkage. *Heredity* **97**, 319–328. (doi:10.1038/sj.hdy.6800895)
94. Gibson JR, Chippindale AK, Rice WR. 2002 The X chromosome is a hot spot for sexually antagonistic fitness variation. *Proc. R. Soc. B* **269**, 499–505. (doi:10.1098/rspb.2001.1863)
95. Pischedda A, Chippindale AK. 2006 Intralocus sexual conflict diminishes the benefits of sexual selection. *PLoS Biol.* **4**, e356. (doi:10.1371/journal.pbio.0040356)
96. Foerster K, Coulson T, Sheldon BC, Pemberton JM, Clutton-Brock TH, Kruuk LEB. 2007 Sexually antagonistic genetic variation for fitness in red deer. *Nature* **447**, 1107–1110. (doi:10.1038/nature05912)
97. Connallon T, Jakubowski E. 2009 Association between sex ratio distortion and sexually antagonistic fitness consequences of female choice. *Evolution* **63**, 2179–2183. (doi:10.1111/j.1558-5646.2009.00692.x)
98. Lucotte EA, Laurent R, Heyer E, Ségurel L, Toupance B. 2016 Detection of allelic frequency differences between the sexes in humans: a signature of sexually antagonistic selection. *Genome Biol. Evol.* **8**, 1489–1500. (doi:10.1093/gbe/evw090)
99. Ruzicka F, Connallon T. 2020 Is the X chromosome a hot spot for sexually antagonistic polymorphisms? Biases in current empirical tests of classical theory. *Proc. R. Soc. B* **287**, 20201869. (doi:10.1098/rspb.2020.1869)
100. Ruzicka F, Connallon T. 2022 An unbiased test reveals no enrichment of sexually antagonistic polymorphisms on the human X chromosome. *Proc. R. Soc. B* **289**, 20212314. (doi:10.1098/rspb.2021.2314)
101. Koerich LB, Wang X, Clark AG, Carvalho AB. 2008 Low conservation of gene content in the *Drosophila* Y chromosome. *Nature* **456**, 949–951. (doi:10.1038/nature07463)
102. Tobler R, Nolte V, Schlötterer C. 2017 High rate of translocation-based gene birth on the *Drosophila* Y chromosome. *Proc. Natl Acad. Sci. USA* **114**, 11721–11726. (doi:10.1073/pnas.1706502114)
103. Cechova M, Vegesna R, Tomaszewicz M, Harris RS, Chen D, Rangavittal S, Medvedev P, Makova KD. 2020 Dynamic evolution of great ape Y chromosomes. *Proc. Natl Acad. Sci. USA* **117**, 26273–26280. (doi:10.1073/pnas.2001749117)
104. Lin Y, Darolti I, Furman BLS, Almeida P, Sandkam BA, Breden F, Wright AE, Mank JE. 2022 Gene duplications to the Y chromosome in *Trinidadian guppies*. *Mol. Ecol.* **31**, 1853–1863. (doi:10.1111/mec.16355)
105. Chippindale AK, Rice WR. 2001 Y chromosome polymorphism is a strong determinant of male fitness in *Drosophila melanogaster*. *Proc. Natl Acad. Sci. USA* **98**, 5677–5682. (doi:10.1073/pnas.101456898)
106. Sandkam BA, Almeida P, Darolti I, Furman BLS, van der Bijl W, Morris J, Bourne GR, Breden F, Mank JE. 2021 Extreme Y chromosome polymorphism corresponds to five male reproductive morphs of a freshwater fish. *Nat. Ecol. Evol.* **5**, 939–948. (doi:10.1038/s41559-021-01452-w)
107. Kaufmann P, Wiberg RAW, Papachristos K, Scofield DG, Tellgren-Roth C, Immonen E. 2023 Y-linked copy number polymorphism of target of rapamycin is associated with sexual size dimorphism in seed beetles. *Mol. Biol. Evol.* (doi:https://doi.org/10.1093/molbev/msad167)
108. Kaufmann P, Wolak ME, Husby A, Immonen E. 2021 Rapid evolution of sexual size dimorphism facilitated by Y-linked genetic variance. *Nat. Ecol. Evol.* **5**, 1394–1402. (doi:10.1038/s41559-021-01530-z)
109. Lemos B, Araripe LO, Hartl DL. 2008 Polymorphic Y chromosomes harbour cryptic variation with manifold functional consequences. *Science* **319**, 91–93. (doi:10.1126/science.1148861)

110. Jiang PP, Hartl DL, Lemos B. 2010 Y not a dead end: epistatic interactions between Y-linked regulatory polymorphisms and genetic background affect global gene expression in *Drosophila melanogaster*. *Genetics* **186**, 109–118. (doi:10.1534/genetics.110.118109)
111. Yee WKW, Rogell B, Lemos B, Dowling DK. 2015 Intergenic interactions between mitochondrial and Y-linked genes shape male mating patterns and fertility in *Drosophila melanogaster*. *Evolution* **69**, 2876–2890. (doi:10.1111/evo.12788)
112. Archer CR, Stephens RM, Sharma MD, Hosken DJ. 2017 The *Drosophila simulans* Y chromosome interacts with the autosomes to influence male fitness. *J. Evol. Biol.* **30**, 1821–1825. (doi:10.1111/jeb.13141)
113. Ågren JA, Munasinghe M, Clark AG. 2020 Mitochondrial-Y chromosome epistasis in *Drosophila melanogaster*. *Proc. R. Soc. B* **287**, 20200469. (doi:10.1098/rspb.2020.0469)
114. Blackburn GS, Albert AYK, Otto SP. 2010 The evolution of sex ratio adjustment in the presence of sexually antagonistic selection. *Am. Nat.* **176**, 264–275. (doi:10.1086/655220)
115. Trivers RL, Willard DE. 1973 Natural selection of parental ability to vary the sex ratio of offspring. *Science* **179**, 90–92. (doi:10.1126/science.179.4068.90)
116. Calsbeek R, Sinervo B. 2004 Within-clutch variation in offspring sex determined by differences in sire body size: cryptic mate choice in the wild. *J. Evol. Biol.* **17**, 464–470. (doi:10.1046/j.1420-9101.2003.00665.x)
117. Calsbeek R, Bonneaud C. 2008 Postcopulatory fertilization bias as a form of cryptic sexual selection. *Evolution* **62**, 1137–1148. (doi:10.1111/j.1558-5646.2008.00356.x)
118. Cox RM, Calsbeek R. 2010 Cryptic sex-ratio bias provides indirect genetic benefits despite sexual conflict. *Science* **328**, 92–94. (doi:10.1126/science.1185550)
119. Roulin A, Altwegg R, Jensen H, Steinsland I, Schaub M. 2010 Sex-dependent selection on an autosomal melanic female ornament promotes the evolution of sex ratio bias. *Ecol. Lett.* **13**, 616–626. (doi:10.1111/j.1461-0248.2010.01459.x)
120. Katsuki M, Harano T, Miyatake T, Okada K, Hosken DJ. 2012 Intralocus sexual conflict and offspring sex ratio. *Ecol. Lett.* **15**, 193–197. (doi:10.1111/j.1461-0248.2011.01725.x)
121. Romano A, Romano M, Caprioli M, Costanzo A, Parolini M, Rubolini D, Saino N. 2015 Sex allocation according to multiple sexually dimorphic traits of both parents in the barn swallow (*Hirundo rustica*). *J. Evol. Biol.* **28**, 1234–1247. (doi:10.1111/jeb.12650)
122. Alatalo RV, Höglund J, Lundberg A. 1988 Patterns of variation in tail ornament size in birds. *Biol. J. Linn. Soc.* **34**, 363–374. (doi:10.1111/j.1095-8312.1988.tb01969.x)
123. Jennions MD, Møller AP, Petrie M. 2001 Sexually selected traits and adult survival: a meta-analysis. *Q. Rev. Biol.* **76**, 3–36. (doi:10.1086/393743)
124. Hernández A, Martínez-Gómez M, Beamonte-Barrientos R, Montoya B. 2021 Colourful traits in female birds relate to individual condition, reproductive performance and mate preferences: a meta-analytic approach. *Biol. Lett.* **17**, 20210283. (doi:10.1098/rsbl.2021.0283)
125. Burley N. 1981 Sex-ratio manipulation and selection for attractiveness. *Science* **211**, 721–722. (doi:10.1126/science.211.4483.721)
126. Burley N. 1986 Sex-ratio manipulation in color-banded populations of zebra finches. *Evolution* **40**, 1191–1206. (doi:10.1111/j.1558-5646.1986.tb05744.x)
127. Shuster SM, Wade MJ. 2003 *Mating systems and strategies*. Princeton, NJ: Princeton University Press.
128. Winkler L, Moiron M, Morrow EH, Janicke T. 2021 Stronger net selection on males across animals. *Elife* **10**, e68316. (doi:10.7554/eLife.68316)
129. Immonen E, Snook RR, Ritchie MG. 2014 Mating system variation drives rapid evolution of the female transcriptome in *Drosophila pseudoobscura*. *Ecol. Evol.* **4**, 2186–2201. (doi:10.1002/ece3.1098)
130. Li Richter XY, Hollis B. 2021 Softness of selection and mating system interact to shape trait evolution under sexual conflict. *Evolution* **75**, 2335–2347. (doi:10.1111/evo.14329)
131. Grant PR, Grant BR. 2002 Unpredictable evolution in a 30-year study of Darwin's finches. *Science* **296**, 707–711. (doi:10.1126/science.1070315)
132. Okhovat M, Berrio A, Wallace G, Ophir AG, Phelps SM. 2015 Sexual fidelity trade-offs promote regulatory variation in the prairie vole brain. *Science* **350**, 1371–1374. (doi:10.1126/science.aac5791)
133. Glaser-Schmitt A, Wittmann MJ, Ramnarine TJS, Parsch J. 2021 Sexual antagonism, temporally fluctuating selection, and variable dominance affect a regulatory polymorphism in *Drosophila melanogaster*. *Mol. Biol. Evol.* **38**, 4891–4907. (doi:10.1093/molbev/msab215)
134. Ruzicka F *et al.* 2020 The search for sexually antagonistic genes: practical insights from studies of local adaptation and statistical genomics. *Evol. Lett.* **4**, 398–415. (doi:10.1002/evl3.192)
135. Rice WR. 1998 Male fitness increases when females are eliminated from gene pool: implications for the Y chromosome. *Proc. Natl Acad. Sci. USA* **95**, 6217–6221. (doi:10.1073/pnas.95.11.6217)
136. Prasad NG, Bedhomme S, Day T, Chippindale AK. 2007 An evolutionary cost of separate genders revealed by male-limited evolution. *Am. Nat.* **169**, 29–37. (doi:10.1086/509941)
137. Bedhomme S, Prasad NG, Jiang PP, Chippindale AK. 2008 Reproductive behaviour evolves rapidly when intralocus sexual conflict is removed. *PLoS One* **3**, e2187. (doi:10.1371/journal.pone.0002187)
138. Morrow EH, Stewart AD, Rice WR. 2008 Assessing the extent of genome-wide intralocus sexual conflict via experimentally enforced gender-limited selection. *J. Evol. Biol.* **21**, 1046–1054. (doi:10.1111/j.1420-9101.2008.01542.x)
139. Nordén AK, Ramm SA, Abbott JK. 2023 Rapid evolution of sex role specialization in a hermaphrodite under sex-limited selection. *Evolution* **77**, 1066–1076. (doi:10.1093/evolut/qpaa025)
140. Berg EC, Maklakov AA. 2012 Sexes suffer from suboptimal lifespan because of genetic conflict in a seed beetle. *Proc. R. Soc. B* **279**, 4296–4302. (doi:10.1098/rspb.2012.1345)
141. Berger D, Berg EC, Widegren W, Arnqvist G, Maklakov AA. 2014 Multivariate intralocus sexual conflict in seed beetles. *Evolution* **68**, 3457–3469. (doi:10.1111/evo.12528)
142. Pick JL, Hutter P, Tschirren B. 2017 Divergent artificial selection for female reproductive investment has a sexually concordant effect on male reproductive success. *Evol. Lett.* **1**, 222–228. (doi:10.1002/evl3.21)
143. Arbuthnott D, Dutton EM, Agrawal AF, Rundle HD. 2014 The ecology of sexual conflict: ecologically dependent parallel evolution of male harm and female resistance in *Drosophila melanogaster*. *Ecol. Lett.* **17**, 221–228. (doi:10.1111/ele.12222)
144. Yun L, Chen PJ, Singh A, Agrawal AF, Rundle HD. 2017 The physical environment mediates male harm and its effect on selection in females. *Proc. R. Soc. B* **284**, 20170424. (doi:10.1098/rspb.2017.0424)
145. García-Roa R, Chirinos V, Carazo P. 2019 The ecology of sexual conflict: temperature variation in the social environment can drastically modulate male harm to females. *Funct. Ecol.* **33**, 681–692. (doi:10.1111/1365-2435.13275)
146. Rodríguez-Exposito E, García-González F. 2021 Metapopulation structure modulates sexual antagonism. *Evol. Lett.* **5**, 344–358. (doi:10.1002/evl3.244)
147. Wyrer CAS, Cator LJ, Hollis B. 2023 Release from sexual selection leads to rapid genome-wide evolution in *Aedes aegypti*. *Curr. Biol.* **33**, 1351–1357. (doi:10.1016/j.cub.2023.02.031)
148. Duffy E, Archer CR, Sharma MD, Prus M, Joag RA, Radwan J, Wedell N, Hosken DJ. 2019 *Wolbachia* infection can bias estimates of intralocus sexual conflict. *Ecol. Evol.* **9**, 328–338. (doi:10.1002/ece3.4744)
149. Dugand RJ, Tomkins JL, Kennington WJ. 2019 Molecular evidence supports a genic capture resolution of the lek paradox. *Nat. Commun.* **10**, 1359. (doi:10.1038/s41467-019-09371-y)
150. Rostant WG, Kay C, Wedell N, Hosken DJ. 2015 Sexual conflict maintains variation at an insecticide resistance locus. *BMC Biol.* **13**, 34. (doi:10.1186/s12915-015-0143-3)
151. Mank JE. 2017 The transcriptional architecture of phenotypic dimorphism. *Nat. Ecol. Evol.* **1**, 6. (doi:10.1038/s41559-016-0006)

152. Rusuwa BB, Chung H, Allen SL, Frentiu FD, Chenoweth SF. 2022 Natural variation at a single gene generates sexual antagonism across fitness components in *Drosophila*. *Curr. Biol.* **32**, 3161–3169. (doi:10.1016/j.cub.2022.05.038)
153. Kokko H, Brooks R. 2003 Sexy to die for? Sexual selection and the risk of extinction. *Ann. Zool. Fenn.* **40**, 207–219.
154. Le Galliard JF, Fitze PS, Ferrière R, Clobert J. 2005 Sex ratio bias, male aggression, and population collapse in lizards. *Proc. Natl Acad. Sci. USA* **102**, 18231–18236. (doi:10.1073/pnas.0505172102)
155. Grieshop K, Berger D, Arnqvist G. 2017 Male-benefit sexually antagonistic genotypes show elevated vulnerability to inbreeding. *BMC Evol. Biol.* **17**, 134. (doi:10.1186/s12862-017-0981-4)
156. Morrow EH. 2015 The evolution of sex differences in disease. *Biol. Sex Differ.* **6**, 1–7. (doi:10.1186/s13293-015-0023-0)
157. Archer CR, Recker M, Duffy E, Hosken DJ. 2018 Intralocus sexual conflict can resolve the male-female health-survival paradox. *Nat. Commun.* **9**, 5048. (doi:10.1038/s41467-018-07541-y)
158. Harper JA, Janicke T, Morrow EH. 2021 Systematic review reveals multiple sexually antagonistic polymorphisms affecting human disease and complex traits. *Evolution* **75**, 3087–3097. (doi:10.1111/evo.14394)