

# UCSF

## UC San Francisco Previously Published Works

### Title

A context-specific circadian clock in adipocyte precursor cells modulates adipogenesis

### Permalink

<https://escholarship.org/uc/item/6nc08355>

### Journal

Adipocyte, 7(4)

### ISSN

2162-3945

### Authors

Jung, Yunshin  
Feldman, Brian J

### Publication Date

2018-10-02

### DOI

10.1080/21623945.2018.1516099

Peer reviewed

COMMENTARY



## A context-specific circadian clock in adipocyte precursor cells modulates adipogenesis

Yunshin Jung<sup>a</sup> and Brian J. Feldman<sup>a,b</sup>

<sup>a</sup>Department of Pediatrics/Endocrinology, Stanford University School of Medicine, Stanford, CA, USA; <sup>b</sup>Program in Regenerative Medicine, Stanford University, Stanford, CA, USA

### ABSTRACT

The circadian clock is an intricate molecular network that paces a variety of physiological process to ~ 24 hour day/night cycles. Whereas the central circadian clock in the brain is primarily entrained by light signals, peripheral circadian clocks, which are in most cells in the body, receive cues not only from the central pacemaker but also endocrine and other systemic and tissue-specific signals. Prior studies have connected peripheral circadian clocks to metabolism, primarily with studies focused on the robust clock in the liver that responds to feeding/fasting cycles. Adipose tissue is also critical for metabolism and adipocytes have circadian clocks. Yet, the role of the circadian clock in adipocytes is poorly understood. Here we describe our studies that revealed components of the circadian clock in primary adipocyte precursor cells (APCs) in mice. We made the surprising discovery of a particularly prominent role for the circadian gene *Period 3* (*Per3*) in the APC clock. Furthermore, we elucidated that *Per3* directly regulates an output pathway of the APC clock to modulate the expression of the Kruppel-like factor 15 (*Klf15*) gene. Finally, we discovered that this clock-Klf15 pathway regulates adipogenesis in APCs. These findings have important implications for our understanding of adipose tissue biology and metabolism and, we speculate, will generate opportunities to develop novel therapeutic strategies based on the context-specific features of the circadian clock in APCs.

### ARTICLE HISTORY

Received 9 July 2018  
Revised 15 August 2018  
Accepted 16 August 2018

### KEYWORDS

Circadian clock; adipocyte precursor cells; *in vivo* adipogenesis; *Per3*; *Klf15*

### Introduction

A circadian clock is an endogenous 24-hr-cycle timekeeping system, controlled by oscillations in molecular components.<sup>1,2</sup> The core circadian clock machinery uses positive feedback components (i.e. *Clock* and *Bmal1*) and negative feedback components (i.e. *Per* genes), to regulate the periodicity of circadian and clock-controlled genes.<sup>1</sup> Although, three *Period* paralogs (*Per 1–3*) are reported in mammals, *Per3* is the least understood and was not regarded as a core circadian clock gene because deleting *Per3* had minimal effect on circadian locomotion in mice.<sup>3,4</sup>

The central circadian clock is located in the suprachiasmatic nuclei of the brain and most cells in the body appear to contain peripheral circadian clocks.<sup>2,5</sup> Interestingly, there are context specific differences in the components used in these peripheral clocks<sup>6</sup> but the implications and mechanisms driving these differences are incompletely understood. Signals from photoreceptors in the retina to the central clock are generated by diurnal light cycles and entrain the central clock.<sup>2,5</sup> The central clock serves as an important pacemaker for peripheral circadian clocks. A number of functions of peripheral tissues are modulated

by central clock pacing including: insulin secretion from the pancreas and sensitivity in muscle.<sup>7,8</sup> However, peripheral clocks are also subject to distinct inputs and contextual regulation that may uncouple oscillations from the central clock.<sup>1,9–12</sup> For example, the timing of food intake can alter peripheral clock oscillations and rhythmicity of bile synthesis in liver.<sup>13–16</sup> These, and a body of other related studies, indicate that peripheral circadian clocks provide a mechanism that links peripheral tissues with systemic metabolism.<sup>17–20</sup>

Another example of a metabolically relevant biological process that is influenced by peripheral circadian clocks is adipogenesis – a differentiation pathway where adipocyte precursor cells develop into mature adipocytes.<sup>21–23</sup> Dysregulated adipogenesis contributes to expansion of adipocyte tissue and thus is often implicated in obesity.<sup>21</sup> Furthermore, disruption of circadian rhythms is a risk-factor for obesity.<sup>1,24</sup> To begin to examine the mechanisms by which the circadian clock is connected to adipogenesis, previous studies employed *in vitro* adipocyte differentiation protocols using mouse embryonic fibroblasts cells (MEFs) and immortalized 3T3-L1 cells, e.g.<sup>25</sup> Recently, we performed studies using primary adipocyte precursor

cells (APCs) and approaches that monitor adipogenesis *in vivo* to establish the presence of peripheral clocks in bona fide APCs and mechanisms by which the circadian clock regulates the process of adipogenesis.<sup>23</sup>

## Results

As defined by a *Scal* (stem cell marker) positive and lineage marker negative population,<sup>26</sup> we isolated primary APCs from subcutaneous adipose depots of circadian clock reporter mice (*mPer2<sup>Luc</sup>*).<sup>27</sup> These studies demonstrated that primary APCs contain a circadian clock.<sup>23</sup> However, to our surprise, our further investigations elucidated that expression of the classic core clock component *Per1* does not significantly oscillate in APCs.<sup>23</sup> Strikingly, we discovered that *Per3*, has the most robust circadian expression profile of the *Per* paralogs in primary APCs.<sup>23</sup> These findings suggested to us that *Per3* might have a particularly prominent role in the context of APC biology. Indeed, deletion of the *Per3* gene altered the circadian cycling of core clock components, supporting our hypothesis that *Per3* activity impacts the clock machinery in APCs.<sup>23</sup>

Intrigued by the above results, we investigated the functional role of *Per3* in APC biology. To probe this question, we compared the level of adipogenesis occurring in *Per3* deficient mice (*Per3<sup>-/-</sup>*) *in vivo* to that of wild type mice using a pulse-chase approach we had previously validated.<sup>23,28,29</sup> Using this protocol, we revealed that *Per3<sup>-/-</sup>* mice have increased levels of adipogenesis *in vivo*.<sup>23</sup> The increased adipogenesis upon *Per3* ablation was also captured in *ex vivo* culture assays by monitoring the levels of spontaneous adipogenesis of isolated primary APCs, indicating that disruption of *Per3* in APCs was sufficient to cell autonomously promote adipogenesis in APCs.<sup>23</sup>


Next, to elucidate the mechanisms by which *Per3* regulates adipogenesis in APCs, we performed unbiased genome-wide expression profiling studies comparing APCs isolated from wild type and *Per3<sup>-/-</sup>* mice. Among the genes with significant expression changes across the genotypes, *Klf15* was previously reported as having a potential role in modulating adipogenesis.<sup>30</sup> Therefore, we decided to examine whether *Klf15* is responsible for connecting circadian clock to *in vivo* adipogenesis. We confirmed that both the messenger and protein expression levels of *Klf15* were significantly increased in APCs from *Per3<sup>-/-</sup>* mice.<sup>23</sup> Conversely, *Per3* overexpression in wild type APCs decreased *Klf15* expression level, suggesting that PER3 inhibits *Klf15* expression.<sup>23</sup> Interestingly, we discovered *Klf15* expression has a circadian oscillation pattern in APCs from wild type mice and,

furthermore, these oscillations were abrogated in the absence of *Per3* gene.<sup>23</sup> Together, these findings indicate that PER3 plays a prominent role in the regulation of *Klf15* expression in APCs. Indeed, our molecular and biochemical studies elucidated that PER3 forms a complex with BMAL1 at an E-box in the *Klf15* gene to regulate expression.<sup>23</sup> Our results suggest that when PER3 levels are low BMAL1 occupancy of this E-box promotes *Klf15* expression and, as PER3 levels increase, PER3 binds to BMAL1 and *Klf15* expression is repressed.<sup>23</sup> This mechanism reveals a direct pathway by which the peripheral circadian clock controls *Klf15* expression, resulting in the circadian oscillations in *Klf15* expression observed in wild type APCs.<sup>23</sup> Finally, we tested if this clock-*Klf15* pathway was relevant for the regulation of adipogenesis. To this end, we found that knocking-down *Klf15* was sufficient to rescue the *Per3<sup>-/-</sup>* adipogenesis phenotype.<sup>23</sup> These results indicate that PER3-KLF15 pathway is critical to regulating adipogenesis.

## Discussion

In conclusion, we discovered that primary APCs contain a peripheral circadian clock. Further, we exposed what appear to be context-specific elements of the APC clock. In particular, the poorly understood *Per3* clock component has a significant role in the APC clock, directly regulating an output pathway that modulates adipogenesis. As the rate of adipogenesis has systemic implications, including for metabolism and body composition, these new insights have broad implications.

In addition, our study further delineates the intriguing relationship between circadian oscillations and metabolic processes. For example, recent studies by others have demonstrated the importance of cycling between feeding and fasting states for maintaining healthy systemic metabolism in humans.<sup>31,32</sup> This is particularly interesting to us because circadian cycles of feeding and fasting provide potent entrainment cues for peripheral clocks. This has most extensively been studied in the liver, where disruption in feeding cycles, as well as direct abrogation of the circadian clock, leads to metabolic derangement.<sup>33-35</sup> To date, the effect of circadian feeding/fasting cycles for the clock in APCs and adipocytes has yet to be extensively explored. We speculate that, like in the liver, disruption of feeding/fasting cycles will have detrimental functional implications in adipose tissue. Indeed, we hypothesize that the rate of adipogenesis might be modulated by altering these circadian cycles by mechanisms that impinge upon the same peripheral clock-*Klf15* pathway exposed in our study (Figure 1).



**Figure 1. Schematic of the central circadian clock and peripheral clocks in the liver and APCs.** The peripheral circadian clock in the liver receives multiple inputs including signals from the central clock and feeding. *Per3* has a context specific significant role in the circadian clock in APCs and regulates an output pathway to *Klf15* that modulates adipogenesis.

However, we also note that, while liver and adipose clocks may share some commonality in receiving entrainment from feeding cycles, there are likely to be important distinctions as well. In particular, our study found that, consistent with a more canonical peripheral clock, hepatocytes maintain robust oscillations in the expression of the core clock component *Per1*, while in APCs, we found that oscillations in *Per1* expression is not significant.<sup>23</sup> This may signal a unique role for *Per3* in APCs and, by extension, adipose tissue biology in general. The logic behind this context difference still needs to be elucidated, but it is consistent with enabling tissue specific input and output responses to systemic changes and signals. If correct, this context-specificity of peripheral clocks would offer opportunities for the development of more targeted tissue-specific therapeutic interventions.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by the National Institutes of Health [R01DK114217]

## References

1. Bass J, Lazar MA. Circadian time signatures of fitness and disease. *Sci*. [Internet]. 2016 Nov 25;354(6315):994. Available from: <http://science.sciencemag.org/content/354/6315/994.abstract>.
2. Reppert SM, Weaver DR. Coordination of circadian timing in mammals. *Nature*. [Internet]. 2002 Aug 29;418:935.
3. Bae K, Jin X, Maywood ES, Hastings MH, Reppert SM, Weaver DR. Differential functions of mPer1, mPer2, and mPer3 in the SCN circadian clock. *Neuron*.

- [Internet]. 2001 May 1;30(2): 525–536. Available from. <http://www.sciencedirect.com/science/article/pii/S0896627301003026>.
4. Shearman LP, Jin X, Lee C, Reppert SM, Weaver DR. Targeted disruption of the mPer3 gene: subtle effects on circadian clock function. *Mol Cell Biol* [Internet]. [2000 Sep 1];20(17):6269–6275. Available from <http://mcb.asm.org/content/20/17/6269.abstract>
5. Schibler U, Gotic I, Saini C, Gos P, Curie T, Emmenegger Y, Sinturel F, Gosselin P, Gerber A, Fleury-Olela F, et al. Clock-talk: interactions between central and peripheral circadian oscillators in mammals. *Cold Spring Harb Symp Quant Biol*. [Internet]. 2015. Jan 1;80: 223–232. Available from. <http://sympo.scripps.edu/content/80/223.abstract>.
6. Mohawk JA, Green CB, Takahashi JS. Central and peripheral circadian clocks in mammals. *Annu Rev Neurosci*. [Internet]. 2012 Jun 20;[cited]. 2018 Jul 4. 35(1): 445–462.
7. Perelis M, Marcheva B, Moynihan Ramsey K, Schipma MJ, Hutchison AL, Taguchi A, Peek CB, Hong H, Huang W, Omura C, et al.  $\beta$  cell enhancers regulate rhythmic transcription of genes controlling insulin secretion. *Sci*. [Internet]. 2015 Nov 6;350(6261): Available from. <http://science.sciencemag.org/content/350/6261/aac4250.abstract>.
8. Hong S, Zhou W, Fang B, Lu W, Loro E, Damle M, Ding G, Jager J, Zhang S, Zhang Y, et al. Dissociation of muscle insulin sensitivity from exercise endurance in mice by HDAC3 depletion. *Nat Med*. [Internet]. 2016 Dec 19;23(223). DOI:10.1038/nm.4245
9. Balsalobre A, Brown SA, Marcacci L, Tronche F, Kellendonk C, Reichardt HM, Schütz G, Schibler U. Resetting of circadian time in peripheral tissues by glucocorticoid signaling. *Sci*. [Internet]. 2000 Sep 29;289(5488): 2344. Available from. <http://science.sciencemag.org/content/289/5488/2344.abstract>.
10. Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev*. [Internet]. 2000 Dec 1;14(23): 2950–2961. Available from. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC317100/>.
11. Vollmers C, Gill S, DiTacchio L, Pulivarthy SR, Le HD, Panda S. Time of feeding and the intrinsic circadian clock drive rhythms in hepatic gene expression. *Proc Natl Acad Sci USA*. [Internet]. 2009 Dec 15;106(50): 21453. Available from. <http://www.pnas.org/content/106/50/21453.abstract>.
12. Mukherji A, Kobiita A, Damara M, Misra N, Meziane H, Champy M-F CP. Shifting eating to the circadian rest phase misaligns the peripheral clocks with the master SCN clock and leads to a metabolic syndrome. *Proc Natl Acad Sci USA*. [Internet]. 2015 Dec 1; 112(48): E6691. Available from. <http://www.pnas.org/content/112/48/E6691.abstract>.
13. Hatori M, Vollmers C, Zarrinpar A, DiTacchio L, Bushong EA, Gill S, Leblanc M, Chaix A, Joens M, Fitzpatrick JA, et al. Feeding without reducing caloric intake prevents metabolic diseases in mice fed a high-fat diet. *Cell Metab*. [Internet]. 2012 Jun 6;15(6): 848–860.

- Available from. <http://www.sciencedirect.com/science/article/pii/S1550413112001891>. .
14. Chaix A, Zarrinpar A, Miu P, Panda S. Time-restricted feeding is a Preventative and therapeutic intervention against diverse nutritional challenges. *Cell Metab.* [Internet]. 2014 Dec 2;20(6): 991–1005. Available from. <http://www.sciencedirect.com/science/article/pii/S1550413114004987>. .
  15. Adamovich Y, Rousoo-Noori L, Zwihaft Z, Neufeld-Cohen A, Golik M, Kraut-Cohen J, Wang M, Han X, Asher G. Circadian clocks and feeding time regulate the oscillations and levels of hepatic triglycerides. [Internet]. *Cell Metab.* cited. 2014 Feb 4;19(2):319–330.
  16. Han S, Zhang R, Jain R, Shi H, Zhang L, Zhou G, Sangwung P, Tugal D, Atkins GB, Prosdocimo DA, et al. Circadian control of bile acid synthesis by a KLF15-Fgf15 axis. *Nat Commun.* [Internet]. 2015 Jun 4;6:7231.
  17. Panda S, Antoch MP, Miller BH, Su AI, Schook AB, Straume M, Schultz PG, Kay SA, Takahashi JS, Hogenesch JB. Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell.* [Internet]. 2002 May 3;109(3): 307–320. Available from. <http://www.sciencedirect.com/science/article/pii/S0092867402007225>. .
  18. Asher G, Schibler U. Crosstalk between components of circadian and metabolic cycles in mammals. *Cell Metab.* [Internet]. 2011 Feb 2;13(2): 125–137. Available from. <http://www.sciencedirect.com/science/article/pii/S1550413111000076>. .
  19. Feng D, Ma L. Clocks, metabolism, and the epigenome. *Mol Cell.* [Internet]. 2012 Jul 27;47(2): 158–167. Available from. <http://www.sciencedirect.com/science/article/pii/S1097276512005515>. .
  20. Bass J, Takahashi JS. Circadian integration of metabolism and energetics. *Sci.* [Internet]. 2010 Dec 3;330(6009): 1349. Available from. <http://science.sciencemag.org/content/330/6009/1349.abstract>. .
  21. Rutkowski JM, Stern JH, Scherer PE. The cell biology of fat expansion. *J Cell Biol.* [Internet]. 2015 Mar 2; 208(5): 501. Available from: <http://jcb.rupress.org/content/208/5/501.abstract>. .
  22. Krueger KC, Costa MJ, Du H, Feldman BJ. Characterization of cre recombinase activity for in vivo targeting of adipocyte precursor cells. *Stem Cell Rep.* [Internet]. 2014 Dec 9;cited. 2018 Jun 1. 3(6): 1147–1158.
  23. Aggarwal A, Costa MJ, Rivero-Gutiérrez B, Ji L, Morgan SL, Feldman BJ. The circadian clock regulates adipogenesis by a Per3 crosstalk pathway to Klf15. *Cell Rep.* [Internet]. 2017 Nov 28;cited. 2018 Jun 2. 21(9): 2367–2375.
  24. Froy O. Metabolism and circadian rhythms—implications for obesity. *Endocr Rev.* [Internet]. 2010 Feb 1; 31(1):1–24.
  25. Grimaldi B, Bellet MM, Katada S, Astarita G, Hirayama J, Amin RH, Granneman JG, Piomelli D, Leff T, Sassone-Corsi P. PER2 controls lipid metabolism by direct regulation of PPAR $\gamma$ . *Cell Metab.* [Internet]. 2010 Nov 3;12(5): 509–520. Available from. <http://www.sciencedirect.com/science/article/pii/S1550413110003578>. .
  26. Rodeheffer MS, Birsoy K, Friedman JM. Identification of white adipocyte progenitor cells in vivo. *Cell.* [Internet]. 2008 Oct 17;135(2): 240–249. Available from. <http://www.sciencedirect.com/science/article/pii/S009286740801194X>.
  27. Yoo S-H, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Slepka SM, Hong H-K, Oh WJ, Yoo OJ, et al. PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci U S A.* [Internet]. 2004 Apr 13;101(15): 5339. Available from. <http://www.pnas.org/content/101/15/5339.abstract>. .
  28. Wong JC, Krueger KC, Costa MJ, Aggarwal A, Du H, McLaughlin TL, Feldman BJ. A glucocorticoid- and diet-responsive pathway toggles adipocyte precursor cell activity in vivo. *Sci Signal.* [Internet]. 2016 Oct 25;9(451): ra103. Available from. <http://stke.sciencemag.org/content/9/451/ra103.abstract>. .
  29. Wong JC, Feldman BJ. Adamts1 responds to systemic cues and gates adipogenesis. *Adipocyte.* [Internet]. 2017 Oct 2;6(4): 293–297.
  30. Mori T, Sakaue H, Iguchi H, Gomi H, Okada Y, Takashima Y, Nakamura K, Nakamura T, Yamauchi T, Kubota N, et al. Role of Krüppel-like factor 15 (KLF15) in transcriptional regulation of adipogenesis. *J Biol Chem.* [Internet]. 2005 Apr 1;280(13): 12867–12875. Available from. <http://www.jbc.org/content/280/13/12867.abstract>. .
  31. Sutton EF, Beyl R, Early KS, Cefalu WT, Ravussin E, Peterson CM. Early time-restricted feeding improves insulin sensitivity, blood pressure, and oxidative stress even without weight loss in men with prediabetes. [Internet]. *Cell Metab.* cited. 2018 Jun 5;27(6):1212–1221.e3.
  32. Longo VD, Panda S. Fasting, circadian rhythms, and time-restricted feeding in healthy lifespan. *Cell Metab.* [Internet]. 2016 Jun 14;23(6): 1048–1059. Available from. <http://www.sciencedirect.com/science/article/pii/S1550413116302509>. .
  33. Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, Laposky A, Losee-Olson S, Easton A, Jensen DR, et al. Obesity and metabolic syndrome in circadian clock mutant mice. *Sci* [Internet]. [2005 May 13];308(5724):1043. Available from <http://science.sciencemag.org/content/308/5724/1043.abstract>
  34. Kohsaka A, Laposky AD, Ramsey KM, Estrada C, Joshu C, Kobayashi Y, Turek FW, Bass J. High-fat diet disrupts behavioral and molecular circadian rhythms in mice. *Cell Metab.* [Internet]. 2007 Nov 7; 6(5): 414–421. Available from. <http://www.sciencedirect.com/science/article/pii/S1550413107002665>. .
  35. Kettner NM, Mayo SA, Hua J, Lee C, Moore DD, Fu L. Circadian dysfunction induces leptin resistance in mice. *Cell Metab.* [Internet]. 2015 Sep 1;22(3): 448–459. Available from. <http://www.sciencedirect.com/science/article/pii/S1550413115002727>.