UC Irvine

UC Irvine Previously Published Works

Title

Communication: Fully coherent quantum state hopping.

Permalink

<https://escholarship.org/uc/item/6nm4t9tj>

Journal

The Journal of chemical physics, 143(14)

ISSN

0021-9606

Author

Martens, Craig C

Publication Date

2015-10-01

DOI

10.1063/1.4933049

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Communication: Fully coherent quantum state hopping

Craig C. Martens^{a)}

University of California, Irvine, California 92697-2025, USA

(Received 25 August 2015; accepted 30 September 2015; published online 9 October 2015)

In this paper, we describe a new and fully coherent stochastic surface hopping method for simulating mixed quantum-classical systems. We illustrate the approach on the simple but unforgiving problem of quantum evolution of a two-state quantum system in the limit of unperturbed pure state dynamics and for dissipative evolution in the presence of both stationary and nonstationary random environments. We formulate our approach in the Liouville representation and describe the density matrix elements by ensembles of trajectories. Population dynamics are represented by stochastic surface hops for trajectories representing diagonal density matrix elements. These are combined with an unconventional coherent stochastic hopping algorithm for trajectories representing off-diagonal quantum coherences. The latter generalizes the binary (0,1) “probability” of a trajectory to be associated with a given state to allow integers that can be negative or greater than unity in magnitude. Unlike existing surface hopping methods, the dynamics of the ensembles are fully entangled, correctly capturing the coherent and nonlocal structure of quantum mechanics. © 2015 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4933049>]

The quantum dynamics of many-body systems is a subject of great current interest. A range of theoretical quantum-classical methods have been proposed to treat such systems. These include trajectory surface hopping,^{1–6} semiclassical initial value representation,^{7–11} and quantum-classical Wigner function-based approaches,^{12–20} to cite just a few.

Probably, the most popular method for combining classical dynamics with quantum transitions is trajectory surface hopping,^{1–6,21} and, in particular, variants of Tully’s fewest switches surface hopping (FSSH).² Together, these methods are well-known for their mistreatment of coherence. The origin of the problem is that each trajectory carries with it its own complete quantum state, which is fully coherent. This is fundamentally incorrect, as quantum effects in a trajectory representation emerge as the inability to separate the unified whole of a quantum state into independent ensemble members. Quantum mechanics *entangles* classical trajectory evolution.^{22,23} From a trajectory ensemble perspective, quantum coherence results from the relationships *between* ensemble members and is not a property of any single trajectory. This perspective has been emphasized in our previous work.^{12–16,22,23} Recently, a great deal of attention has been brought to correct the overcoherence problem while retaining the general surface hopping approach (see, e.g., Ref. 21).

In this paper, we present a fully coherent stochastic quantum state hopping method for simulating quantum dynamics. We consider the simple but unforgiving two-state system evolving coherently in a pure state without external perturbations and then add random noise to the ensemble evolution to demonstrate the method’s ability to accurately capture decoherence effects in both stationary and nonstationary environments.

We formulate our approach in the Liouville representation and describe the quantum evolution by a density matrix.²⁴ The Hamiltonian of the system is given by

$$\hat{H} = \begin{pmatrix} E_1 & V \\ V & E_2 \end{pmatrix}, \quad (1)$$

where the diagonal elements satisfy $E_2 - E_1 = \hbar\omega$. The off-diagonal coupling is taken to be a real constant V . The state of the system is described by the density matrix

$$\hat{\rho}(t) = \begin{pmatrix} \rho_{11}(t) & \rho_{12}(t) \\ \rho_{21}(t) & \rho_{22}(t) \end{pmatrix}, \quad (2)$$

where $\rho_{11}(t)$ and $\rho_{22}(t)$ are the time-dependent populations of states $|1\rangle$ and $|2\rangle$, respectively. The coupling V induces transfer of population between the states. The off-diagonal elements $\rho_{12}(t)$ and $\rho_{21}(t) = \rho_{12}^*(t)$ are the coherences. The density operator $\hat{\rho}(t)$ undergoes pure state evolution under the quantum Liouville equation,

$$i\hbar \frac{d\hat{\rho}(t)}{dt} = [\hat{H}, \hat{\rho}(t)]. \quad (3)$$

This can be broken down into coupled ordinary differential equations for the density matrix elements,

$$\dot{\rho}_{11} = -\frac{2V}{\hbar} \text{Im } \rho_{12}, \quad (4)$$

$$\dot{\rho}_{22} = \frac{2V}{\hbar} \text{Im } \rho_{12}, \quad (5)$$

$$\dot{\rho}_{12} + i\omega\rho_{12} = \frac{iV}{\hbar}(\rho_{11} - \rho_{22}). \quad (6)$$

We represent the time evolution of the density matrix in terms of ensembles of stochastic realizations, which we refer to as *trajectories*. First, consider the evolution of populations, which are represented collectively by an ensemble of N trajectories. Each trajectory carries a discrete random variable $\sigma_j(t)$,

^{a)}Email: cmartens@uci.edu

$j = 1, 2, \dots, N$ which can take on values of either 0 or 1. If $\sigma_j(t) = 1$ at time t , then the j th trajectory occupies state 1, while if $\sigma_j(t) = 0$, the trajectory occupies state 2. The density matrix elements can then be written as sums over the trajectory ensemble,

$$\rho_{11}(t) = \frac{1}{N} \sum_{j=1}^N \sigma_j(t) = \frac{N_1(t)}{N}, \quad (7)$$

$$\rho_{22}(t) = \frac{1}{N} \sum_{j=1}^N (1 - \sigma_j(t)) = \frac{N_2(t)}{N}, \quad (8)$$

where $N_1(t)$ and $N_2(t)$ are the number of trajectories associated with states 1 and 2, respectively, and $N_1(t) + N_2(t) = N$. This conservation of trajectories is honored by the equations of motion, Eqs. (4) and (5), as $\dot{\rho}_{11} + \dot{\rho}_{22} = 0$.

The dynamics of the $\sigma_j(t)$ are represented by a stochastic hopping algorithm. In terms of $N_1(t) = N\rho_{11}(t)$ we have, for a time step Δt ,

$$\Delta N_1(t) = -\frac{2V}{\hbar} \text{Im} \rho_{12}(t) N \Delta t. \quad (9)$$

This gives the number of ensemble members that will join or leave the state 1 portion of the ensemble during the time step Δt . The direction of change depends on the sign of the right side of the equation. In practice, Δt is chosen so that the magnitude of ΔN_1 during the time step is much less than N . The probability that a given member of the ensemble will hop is determined as follows. If the j th trajectory is already a member of the state 1 part of the ensemble with $\sigma_j(t) = 1$ and if $\dot{\rho}_{11}$ is positive, nothing happens to the status of the trajectory. If $\dot{\rho}_{11}$ is negative, the probability that the trajectory will undergo a transition to state 2 is given by

$$P(1 \rightarrow 2) = \frac{\Delta N_1}{N_1} = \frac{2V}{\hbar} \text{Im} \rho_{12} \frac{N}{N_1} \Delta t = \frac{2V}{\hbar} \frac{\text{Im} \rho_{12}}{\rho_{11}} \Delta t. \quad (10)$$

The probability $P(2 \rightarrow 1)$ for hops from state 2 to state 1 is computed in an analogous manner. The numerical implementation is achieved by generating a uniform random number ξ between 0 and 1 for each candidate population ensemble member at each time t and executing the hop if $\xi \leq P$.

The coherence $\rho_{12}(t)$ is also represented by an ensemble of N trajectories. This aspect of the method is novel and differs from previous surface hopping approaches. As we will see, it correctly captures the nature of quantum coherent dynamics.

It is convenient to write the coherence in amplitude-phase form: $\rho_{12}(t) = R(t)e^{i\Phi(t)}$. In terms of the trajectory ensemble, we have

$$R(t)e^{i\Phi(t)} = \frac{1}{N} \sum_{j=1}^N r_j(t)e^{i\phi_j(t)}. \quad (11)$$

This is a complex number that results from summing N complex contributions from each trajectory. The nature of the contributing trajectory amplitudes r_j and phases ϕ_j will be addressed below. We consider the increment of ρ_{12} during a time interval Δt : $\Delta \rho_{12} = \dot{\rho}_{12} \Delta t$. In terms of increments of the amplitudes and phases of the ensemble,

$$\Delta \rho_{12} = (\Delta R + iR\Delta\Phi)e^{i\Phi} = \frac{1}{N} \sum_{j=1}^N (\Delta r_j + ir_j\Delta\phi_j)e^{i\phi_j}. \quad (12)$$

The real and imaginary parts give the relations

$$\Delta R = \frac{1}{N} \sum_{j=1}^N [\Delta r_j \cos(\phi_j - \Phi) - r_j \Delta\phi_j \sin(\phi_j - \Phi)], \quad (13)$$

$$\Delta\Phi = \frac{1}{NR} \sum_{j=1}^N [\Delta r_j \sin(\phi_j - \Phi) + r_j \Delta\phi_j \cos(\phi_j - \Phi)]. \quad (14)$$

From the equations of motion, Eqs. (4)–(6), we have

$$\Delta R = \lambda \sin \Phi \Delta t, \quad (15)$$

$$\Delta\Phi = \left(-\omega + \frac{\lambda}{R} \cos \Phi \right) \Delta t, \quad (16)$$

where $\lambda = \frac{V}{\hbar}(\rho_{11} - \rho_{22})$.

Equating Eqs. (13) to (15) and (14) to (16) gives the conditions for determining the increments Δr_j and $\Delta\phi_j$. This is a highly under-determined set of equations. A great deal of freedom is thus present for creating an algorithm for consistently updating their values. This is reminiscent of the gauge-like freedom of quantum trajectory methods in general that we discussed previously.²⁵ To establish an algorithm, we impose constraints, but note that a different choice of constraints will generate a different dynamics for the trajectory ensemble that nonetheless will be an equivalent representation of the quantum state evolution.

The initial conditions $\sigma_j(0)$ and $(r_j(0), \phi_j(0))$ $j = 1, 2, \dots, N$ are chosen to represent the initial populations $\rho_{11}(0), \rho_{22}(0)$ and coherence $\rho_{12}(0)$, respectively. We restrict the freedom noted above by choosing the initial trajectory phases uniformly between 0 and 2π : $\phi_j(0) = 2\pi j/N$, $j = 1, 2, \dots, N$ and require that the phases evolve, for pure state evolution, as $\Delta\phi_j = -\omega\Delta t$. With that constraint imposed, conditions for Δr_j can be derived by equating Eqs. (13) with (15) and (14) with (16). After some algebra, these can be reduced to

$$\frac{1}{N} \sum_{j=1}^N \Delta r_j \sin \phi_j = \lambda \Delta t, \quad (17)$$

$$\frac{1}{N} \sum_{j=1}^N \Delta r_j \cos \phi_j = 0. \quad (18)$$

A solution for Δr_j is then given by

$$\Delta r_j = \lambda \Delta t \left(\frac{-\langle cs \rangle \cos \phi_j + \langle c^2 \rangle \sin \phi_j}{\langle c^2 \rangle \langle s^2 \rangle - \langle cs \rangle^2} \right), \quad (19)$$

where $\langle c^2 \rangle = \frac{1}{N} \sum_{j=1}^N \cos^2 \phi_j$, $\langle s^2 \rangle = \frac{1}{N} \sum_{j=1}^N \sin^2 \phi_j$, and $\langle cs \rangle = \frac{1}{N} \sum_{j=1}^N \cos \phi_j \sin \phi_j$.

We now implement Eq. (19) with a stochastic algorithm. We interpret Δr_j as a *generalized* probability of change during the time interval Δt . In particular, we generate a random number ξ between 0 and 1 and compare the magnitude of Δr_j with this number. If $|\Delta r_j| \leq \xi$, the trajectory undergoes a coherence transition. This is accomplished by changing the value of r_j by an integer amount. If $\Delta r_j > 0$, then $r_j \rightarrow r_j + 1$. For a negative Δr_j , we let $r_j \rightarrow r_j - 1$. Unlike the population case, the “probability” that a trajectory is in a coherent state is not necessarily a positive number nor is its magnitude bounded by unity. Rather, the coherence state of a given trajectory is an integer: $r_j = \dots$,

FIG. 1. Population of state 1, $\rho_{11}(t)$ (blue dots), and imaginary part of the coherence, $\text{Im } \rho_{12}(t)$ (red dots) compared with exact results (solid lines). (a) $\omega = 0$, $V = 1$. (b) $\omega = 5$, $V = 1$. Ensemble size $N = 10\,000$.

$-3, -2, -1, 0, 1, 2, 3, \dots$. This is reminiscent of the “generalized probability” status of the Wigner function,^{26,27} where its magnitude can be larger than allowed by a classical probability density, while its value can be negative. Restricting the sign or magnitude of r_j in the numerical implementation leads to poor results in practice.

We test the method by treating the coherent evolution of the two-state system with Hamiltonian given by Eq. (1). The dynamics of this system consists of Rabi oscillations induced by the coupling. We consider the initial density matrix with $\rho_{11}(0) = 1$, $\rho_{22}(0) = \rho_{12}(0) = \rho_{21}(0) = 0$. In Fig. 1(a), we show results for the degenerate case $\omega = 0$, $V = 1$, while the dynamics for $\omega = 5$, $V = 1$ case are given in Fig. 1(b). The population $\rho_{11}(t)$ is shown in blue, while the imaginary part of the coherence $\text{Im } \rho_{12}$ is shown in red. These are compared with the exact solution (solid curves). An ensemble of $N = 10\,000$ is used in all cases, which gives effectively quantitative agreement with exact results. (We note that ensembles as small as $N = 100$ give reasonable and stable, if noisy, results.)

In Fig. 2, we show results for the two-state system perturbed by a stationary random environment. This is modeled by adding to ω a fluctuating term $\delta\omega(t)$, characterized by Gaussian white noise with a standard deviation η . This corresponds in the modified Liouville equation governing the quantum system as damping of the coherences, adding a term $-\gamma\rho_{12}(t)$ to the equation for $\rho_{12}(t)$ and a corresponding term for $\rho_{21}(t)$, where $\gamma = \eta^2/2$. In Fig. 2(a), the simulated evolution of $\rho_{11}(t)$ and $\text{Im } \rho_{12}(t)$ is compared with exact results for the case $\omega = 0$, $V = 1$, and $\eta = 0.5$, while Fig. 2(b) gives results for the more strongly damped $\eta = 1.0$ case. Both systems show decaying

coherent oscillations which relax to an incoherent mixture with equal populations in each state.

For both the undamped and damped systems shown above, the coherent surface hopping method gives nearly quantitative agreement with exact results. These are very simple systems that nonetheless provide rigorous tests for our stochastic method for modeling populations and coherence, as coherent quantum effects dominate the dynamics.

We now apply our approach to the nontrivial and largely unexplored problem of quantum dynamics coupled to a *nonequilibrium* environment. We consider a nonstationary stochastic model introduced by us previously to describe the decay of initial coherence coupled to a local nonequilibrium bath, for which analytic results could be obtained.^{28,29}

The model is based on a nonequilibrium local phonon oscillator, which modulates the difference frequency ω by a term $\delta\omega(t) = \delta\omega_o \cos[\Omega(t - t_j) + \theta(t - t_j)]$, where Ω is the local phonon frequency and t_j corresponds to relative time origin for which the oscillator is excited by a quantum transition. The oscillator phase $\theta(t)$ takes on a definite initial value $\theta(0) = \theta_o$, which is related to the mechanical properties of the system and bath and which leads to *classical mechanical* coherence of the ultrafast bath response. As time progresses, the phase $\theta(t)$ undergoes diffusion from the initial value θ_o with diffusion constant D . The interplay of quantum coherence and the classical mechanical coherence of the bath oscillator can influence the quantum dynamics, and the behavior of the system can depend significantly on the control parameter θ_o .^{28,29}

A number of possible realizations of this nonequilibrium model in the context of our stochastic method suggest

FIG. 2. Evolution of $\rho_{11}(t)$ and $\text{Re } \rho_{12}(t)$, as in Fig. 1. (a) $\eta = 0.5$ and (b) $\eta = 1.0$ (right). For both systems, $\omega = 0$, $V = 1$, and $N = 10\,000$.

themselves. In the present paper, our focus is on the quantum simulation method itself, and so, we adopt one simple prescription and leave a detailed analysis of alternatives for future work.

In our approach, each trajectory $(r_j(t), \phi_j(t))$ of the coherence ensemble interacts with its own environmental ensemble member. We include the effect of fluctuations of the instantaneous difference frequency $\delta\omega(t)$ resulting from both stationary Gaussian noise with strength η , as above, and the nonstationary local phonon.

The first hop that creates coherence induces a nonequilibrium bath response and so adds to the fluctuating difference frequency a term $\delta\omega(t) = \delta\omega_o \cos[\Omega(t - t_j) + \theta(t - t_j)]$, where t_j is the time of the transition of r_j . Subsequent transitions of $r_j(t)$ to other nonzero values impulsively add to the response of the evolving local phonon. In general, this will generate a series of trigonometric terms, each with its own relative time zero $t_j^{(k)}$ and stochastic phase history $\theta^{(k)}(t - t_j^{(k)})$. We represent this approximately as a single effective oscillatory term, which simplifies the method while retaining a nonstationary bath with control parameter θ_o .

Immediately after a transition of $r_j(t)$, the term $\delta\omega(t)$ is updated to become $\delta\omega(t) = \delta\omega_o^{(k+1)} \cos[\Omega(t - t_{\text{eff}}^{(k+1)}) + \theta^{(k+1)}(t - t_{\text{eff}}^{(k+1)})]$, where $\theta_o^{(k+1)} = \frac{1}{2}[\theta^{(k)}(t_h^{(k+1)} - t_{\text{eff}}^{(k)}) + \theta_o]$ and $t_{\text{eff}}^{(k+1)} = \frac{1}{2}(t_h^{(k+1)} + t_{\text{eff}}^{(k)})$. The prefactor is updated using $\delta\omega_o^{(k+1)} = \delta\omega_o^{(k)}(1 + \cos(\Omega(t_h^{(k+1)} - t_{\text{eff}}^{(k)})/2))$.

In Fig. 3(a), we show the dynamics of the state 1 population ρ_{11} , while the evolution of the linear entropy $S_L = -\text{Tr}(\rho \ln \rho) = -2(\rho_{11}\rho_{22} - |\rho_{12}|^2)$ is given in Fig. 3(b). We investigate the dependence of the dynamics on the initial phase θ_o for the case $V = 0.5$, $\omega = 1$, $\Omega = 1$, $\delta\omega_o = 0.05$, $D = 0.1$, and $\eta = 0.5$. The trajectory ensemble size is again $N = 10000$. Three initial phase values are shown: $\theta_o = 0$ (blue), $\theta_o = 1.0$ (red), and $\theta_o = \pi$ (green), along with the results for which the oscillator is absent, with only the stationary Gaussian component present (black dashed). The pure dephasing of the coupled system ultimately degrades the Rabi oscillations and leads to equilibration of the populations to $\rho_{11} = 0.5$. The linear entropy S_L measures the purity of the state: a pure state is characterized by $S_L = 0$, while a completely incoherent and equilibrated system decays to $S_L = 0.5$.

A pronounced dependence of the population dynamics on the initial oscillator phase θ_o is revealed in Fig. 3(a). In

particular, the $\theta_o = 0$ case exhibits a significantly slower decay to equilibrium than the purely damped system. The $\theta_o = 1.0$ shows a small positive effect, while $\theta_o = \pi$ leads to an increase in the population transfer at intermediate times, depressing ρ_{11} below the pure dephasing case.

In Fig. 3(b), we show the corresponding time dependence of the linear entropy S_L . The damped system in the absence of the local nonstationary oscillator exhibits a time-dependent growth of S_L from its initial value of $S_L = 0$, indicating a pure state, to the equilibrium value of $S_L = 0.5$. The nonequilibrium system with $\theta_o = 0$ shows a strong suppression of the rate of growth of the linear entropy, especially at intermediate to long times. The effect is less pronounced for $\theta_o = 1.0$. In contrast, the system with $\theta_o = \pi$ exhibits an enhanced rate of increase of S_L .

Persistent nonequilibrium bath coherence is created by and subsequently fully entangled with the quantum evolution. The interplay between quantum system and classical bath has a strong influence on coupled dynamics. As discussed in Refs. 28 and 29, the initial phase θ_o of local environmental oscillator acts as a control parameter for this bath-mediated “coherent control” process. We are unaware of other numerical methods that can simulate fully entangled nonequilibrium system-bath dynamics, which will be explored in more detail in future publications.

In the present method, population and coherence hops are determined stochastically from a global ensemble level representation. The evolution of a given trajectory is “entangled” with that of all the other trajectories. This is consistent with the underlying exact quantum equations of motion, where the *functions* representing the state of the system—the individual density matrix elements—are coupled directly to each other. This entanglement and ensemble level nature of the evolution does not single out a given trajectory’s history as observable. Each member has a definite history, but different stochastic realizations would give different histories without changing anything measurable or observable. Quantum coherence in the present approach is represented at the *ensemble* level through *interrelationship* between contributions of each trajectory that determines the coherence of the state. Energy conservation is also *not* obeyed by individual trajectories, but only holds at the ensemble level. Random perturbations that affect each trajectory in a distinct manner lead to decoherence by eroding systematic relations between different ensemble

FIG. 3. (a) Evolution of $\rho_{11}(t)$ and (b) the linear entropy $S_L(t)$ for the nonequilibrium bath model for two initial bath oscillator phases θ_o , as described in the text.

members. These perturbations, which single out members of the ensemble by the distinctions of each stochastic realization, are essentially acts of measurement, and so modify—and ultimately destroy—the fragile coherences that characterize quantum dynamics. In contrast with the conventional surface hopping methodology, this fundamental characteristic of quantum mechanics is correctly incorporated in the present approach.

We are grateful to Shaul Mukamel, Carter T. Butts, and Rachel Martin for helpful discussions. Stimulating discussions at the Telluride Science Research Center (TSRC) are also gratefully acknowledged. This work is an outgrowth of research supported by the National Science Foundation under Nos. CHE-0614005 and CHE-0802913.

¹J. C. Tully and R. K. Preston, *J. Chem. Phys.* **55**, 562 (1971).

²J. C. Tully, *J. Chem. Phys.* **93**, 1061 (1990).

³J. C. Tully, *J. Chem. Phys.* **137**, 22A301 (2012).

⁴J. E. Subotnik and N. Shenvi, *J. Chem. Phys.* **134**, 024105 (2011).

⁵B. R. Landry and J. E. Subotnik, *J. Chem. Phys.* **135**, 191101 (2011).

⁶T. Nelson, S. Fernandez-Alberti, A. E. Roitberg, and S. Tretiak, *J. Chem. Phys.* **138**, 224111 (2013).

⁷X. Sun and W. H. Miller, *J. Chem. Phys.* **106**, 6346 (1997).

⁸S. Xiong, H. Wang, and W. H. Miller, *J. Chem. Phys.* **109**, 7064 (1998).

⁹X. Sun and W. H. Miller, *J. Chem. Phys.* **110**, 6635 (1999).

¹⁰W. H. Miller, *J. Phys. Chem. A* **105**, 2942 (2001).

¹¹J. M. Moix and E. Pollak, *J. Chem. Phys.* **129**, 064515 (2008).

¹²C. C. Martens and J. Y. Fang, *J. Chem. Phys.* **106**, 4918 (1997).

¹³A. Donoso and C. C. Martens, *J. Phys. Chem. A* **102**, 4291 (1998).

¹⁴A. Donoso and C. C. Martens, *J. Chem. Phys.* **112**, 3980 (2000).

¹⁵A. Donoso, D. Kohen, and C. C. Martens, *J. Chem. Phys.* **112**, 7345 (2000).

¹⁶A. Donoso and C. C. Martens, *Int. J. Quantum Chem.* **87**, 1348 (2002).

¹⁷R. Kapral and G. Ciccotti, *J. Chem. Phys.* **110**, 8919 (1999).

¹⁸R. Kapral, *J. Phys. Chem. A* **105**, 2885 (2001).

¹⁹D. M. Kernan, G. Ciccotti, and R. Kapral, *J. Chem. Phys.* **116**, 2346 (2002).

²⁰G. Hanna and R. Kapral, *Acc. Chem. Res.* **39**, 21 (2006).

²¹N. Shenvi, J. E. Subotnik, and W. Yang, *J. Chem. Phys.* **134**, 144102 (2011).

²²A. Donoso and C. C. Martens, *Phys. Rev. Lett.* **87**, 223202 (2001).

²³A. Donoso, Y. Zheng, and C. C. Martens, *J. Chem. Phys.* **119**, 5010 (2003).

²⁴S. Mukamel, *Principles of Nonlinear Optical Spectroscopy* (Oxford University Press, Oxford, 1995).

²⁵P. Hogan, A. V. Wart, A. Donoso, and C. C. Martens, *Chem. Phys.* **370**, 20 (2010).

²⁶E. P. Wigner, *Phys. Rev.* **40**, 749 (1932).

²⁷M. Hillery, R. F. O'Connell, M. O. Scully, and E. P. Wigner, *Phys. Rep.* **106**, 12 (1984).

²⁸C. C. Martens, *J. Chem. Phys.* **133**, 241101 (2010).

²⁹C. C. Martens, *J. Chem. Phys.* **139**, 024109 (2013).