

UNIVERSITY OF CALIFORNIA

Santa Barbara

A Novel Class of Selective Non-Nucleoside Inhibitors of Human DNA Methyltransferase 3A

A Thesis submitted in partial satisfaction of the
requirements for the degree Master of Science
in Chemistry

by

Sunzeyu Huang

Committee in charge:

Professor Norbert O. Reich, Chair

Professor Trevor W. Hayton

Professor Susannah Scott

Professor Mahdi Abu-Omar

December 2021

The thesis of Sunzeyu Huang is approved.

Trevor W. Hayton

Susannah Scott

Mahdi M. Abu-Omar

Norbert O. Reich, Committee Chair

September 2021

A Novel Class of Selective Non-Nucleoside Inhibitors of Human DNA Methyltransferase 3A

Copyright © 2021

by

Sunzeyu Huang

ACKNOWLEDGEMENTS

First of all, I would like to express my sincere appreciation to my advisor Dr. Norbert Reich for giving me the opportunities to work with him. I still remember the first talk that we had and how his enthusiasm inspired me to join the group. Dr. Reich, to me, you are more than an academic advisor but also a friend, a mentor and a wisdom elder that I admire and respect. To Dr. Greene, Dr. Chen, Dr. Mukherjee, Dr. Sepunaru and Dr. Dahlquist, thank you all for generously sharing your resources and instruments. I would also like to thank my committee members for their support and time.

I would like to have a special thanks to all my group members especially my project partner, Nathaniel Jake. Stillson and Jonathan Sandoval. Thank you, Jake, for being a very helpful, hardworking, smart and creative partner. Thank you, Jonathan, for giving me all the training and help since the first day I joined in the group. Thanks Erin Morgan and Olivia Konttinen for all the great and inspiring discussion.

Furthermore, I give my special thanks to all my friends that I meet in Santa Barbara and old pals in Shenzhen. Thank you Yu-Hsuan, Zhiyuan, Pianpian and Potato for all the company I need whenever I need a break. Thank you, Yihui and Ganghua, for always encourage me whenever I feel frustrated and taking me to gym. Thanks to my roommate Jiaxiang and Wenkai for being the best roommate I can ever think of.

Last but not least, my sincere gratefulness to my parents. Mom and Dad, you guys are my reason that I can start this incredible journey and go on so far. Thank you for always giving me your trust, love and unreasonable support. I share any of my achievement with you. I love you.

ABSTRACT

A Novel Class of Selective Non-Nucleoside Inhibitors of Human DNA Methyltransferase 3A

by

Sunzeyu Huang

Screening of a small chemical library (*Medicines for Malaria Venture* Pathogen Box) identified two structurally related pyrazolone (inhibitor 1) and pyridazine (inhibitor 2) DNMT3A inhibitors with low micromolar inhibition constants. The uncompetitive and mixed type inhibition patterns with DNA and AdoMet suggest these molecules act through an allosteric mechanism, and thus are unlikely to bind to the enzyme's active site. Unlike the clinically used mechanism based DNMT inhibitors such as decitabine or azacitidine that act via the enzyme active site, the inhibitors described here could lead to the development of more selective drugs. Both inhibitors show promising selectivity for DNMT3A in comparison to DNMT1 and bacterial DNA cytosine methyltransferases. With further study, this could form the basis of preferential targeting of *de novo* DNA methylation over maintenance DNA methylation.




TABLE OF CONTENTS

CHAPTER I. A NOVEL CLASS OF SELECTIVE NON-NUCLEOSIDE INHIBITORS OF HUMAN DNA METHYLTRANSFERASE 3A	1
<i>A. INTRODUCTION</i>	1
<i>B. RESULTS AND DISCUSSION</i>	4
<i>C. CONCLUSION</i>	10
<i>D. MATERIAL AND METHODS</i>	10
REFERENCES:.....	14
APPENDIX.....	19

Chapter I. A Novel Class of Selective Non-Nucleoside Inhibitors of Human DNA Methyltransferase 3A

This work represented in this chapter was published on Bioorganic and Medicinal Chemistry Letters. Copyright to Elsevier Ltd.

Citation: Huang S, Stillson NJ, Sandoval JE, Yung C, Reich NO, 'A novel class of selective non-nucleoside inhibitors of human DNA methyltransferase 3A', *Bioorganic & Medicinal Chemistry Letters*, Volume 40, **2021**, 127908

A. Introduction

Epigenetic modifications of proteins and nucleic acids are crucial for normal development.¹⁻³ Human DNA undergoes methylation largely at CpG dinucleotides, and the patterns are developmentally regulated and tissue-specific; these patterns contribute to the epigenetic code, which is essential for viability.³⁻⁶ Aberrant methylation patterns can result in hypermethylation of gene promoters, leading to the silencing of critical tumor suppressor genes, resulting in tumorigenesis.^{5,6} DNA methylation is carried out by a family of enzymes (DNMTs, Figure 1) while demethylation is carried out by the ten-eleven translocation (TET) enzyme family.⁷⁻¹⁰ DNMTs rely on the methyl donor S-adenosyl-l-methionine (AdoMet) and display both random and ordered kinetic mechanisms.¹¹⁻¹⁵ The housekeeping protein DNMT1 primarily acts on hemimethylated DNA, and the two *de novo* methyltransferases DNMT3A and DNMT3B, act predominately on unmethylated DNA.¹⁶⁻¹⁸ The DNMT3s, which also include a catalytically inactive regulatory protein DNMT3L, are mostly expressed during the early development phase

of mammalian germ cells.^{1,4,16,19} DNMT1, meanwhile, is expressed throughout the lifetime of mammalian somatic cells and is localized near replication forks.^{17,20}




Figure 1 – Comparison of the primary structures of human DNMTs. The C-terminal domain contains conserved motifs (I-X) and is active in the absence of the N-terminal domain. The N terminal domain has several conserved segments known to interact with regulatory proteins and histones. The abbreviations used are: DMAPD – DNA methyltransferase associated protein 1 interacting domain, PBD – PCNA binding domain, RFTS – Replication foci targeting domain, BAH – bromo-adjacent homology domain, ADD – ATRX-DNMT3-DNMT3L domain.

All DNMTs share the same domain architecture. The less conserved of their two domains is the N-terminal domain, which contains regulatory segments including the replication fork binding RFD sequence in DNMT1 and the H3 binding ADD sequence in DNMT3s.^{1,4,20,21} The C-terminal, or catalytic domain has the highly conserved methyltransferase motifs (I-X) that are found in both prokaryotic and eukaryotic methyltransferases (see Figure 1).²² These motifs are responsible for cofactor binding and catalysis.^{4,21} DNMT3A forms tetramers with DNA binding occurring along the seam of the dimerization domain (see Figure 2).²¹ Mutations that disrupt the oligomeric state of DNMT3A occur in a number of cancers, and in particular, acute myeloid

leukemia (AML).^{5,23–26} Both catalytically active DNMTs, and in particular, DNMT3A, interact with diverse partners and disruptions to these interactions alter the function of DNMT3A and contribute to tumor-specific changes in methylation patterns.^{5,23,27,28}




Figure 2 – Crystal structure of DNMT3A-3L heterotetrameric complex; the DNA is modeled from the M.HhaI-DNA cocrystal structure (PDB code 6f57).

In the last twenty years, interest in developing drugs that target epigenetic pathways has increased, particularly for histone and DNA modifying enzymes.^{29–31} An obvious feature of these pathways is their inherent reversibility, unlike mutational changes which frequently demand therapeutic strategies leading to cytotoxic interventions. Interestingly, the FDA approved DNMT nucleoside inhibitors, azacytidine and decitabine are highly cytotoxic. These prodrugs are converted to the triphosphates, incorporated into DNA and inhibit DNMTs through the formation of an irreversible suicide complex.^{30,32,33} The dose-limiting toxicity manifested by these drugs has led to the search for non-nucleoside inhibitors; interestingly, many of these act by binding the enzyme active site or act by unknown mechanisms^{29,30}.

B. Results and Discussion

Our interest is to determine if new mechanistic classes of DNMT inhibitors can be identified. The long term goal is to develop therapeutic approaches not hampered by the toxicity and related issues associated with currently used and recently described DNMT inhibitors.^{29,30} For example, there are over 60 known proteins which interact with DNMT3A,³⁴ some of which are implicated in directing DNMT3A to inappropriately methylate and regulate tumor suppressor genes.^{26,27} Moreover, the clinically identified DNMT3A mutations in diverse cancers are known to alter the stability and functional outcomes of the complexes formed between DNMT3A and its partner proteins.^{26,27,35} This network of interactions could be the basis of tumor-specific protein-protein inhibitors (PPIs).^{30,36} Certainly, the recent progress in developing PPIs for diverse therapeutic targets, including leukemia, forms a strong basis for such a strategy.^{37,38} Finally, allosteric enzyme modulators can provide a basis for enhanced selectivity and, potentially, decreased toxicity.³⁹⁻⁴¹

Here we describe our initial compound screening effort, relying on open-source chemical library constructed from the *Medicines for Malaria Venture* (MMV) Pathogen Box. The library consists of 400 drug-like molecules with known activities against targets for neglected tropical diseases. The relative merits of using a library of well-established molecules that show good bio-activity versus other approaches have been well described.⁴² Using 50 compounds of the library, we first determined that a compound concentration of 60 μM resulted in 5% of the molecules showing 90% or more inhibition. We then relied on a modified version of our standard radiochemical assay using tritiated AdoMet,²⁶ which measures DNA methylation (see Methods, Supplementary). The assay uses poly dI-dC which is an excellent DNMT3A substrate, due to the presence of multiple sites for DNMT3A-mediated methylation. The conditions allow for multiple

catalytic turnovers with an excess of DNA.^{26,27} Importantly, many literature reports describing DNMT screens are actually done under conditions which compromise interpretation of any inhibition effects, such as excess enzyme over DNA, or less than a single catalytic cycle.⁴³⁻⁴⁵

The screen of the library generated 12 compounds that showed at least 90% inhibition. The screening assay was repeated on these 12 compounds to verify the inhibitory properties (see Figure 1S), and the results were used to select two structurally similar compounds for further analysis (see compounds 1 and 2, Figure 3). These two compounds were previously identified as potential inhibitors of TbrPDEs, a class of phosphodiesterases found in *T. brucei* – the parasite responsible for trypanosomiasis (*African sleeping sickness*).⁴⁶ Compounds 1 and 2 both show potent and selective inhibition of TbrPDE, good antitrypanosomal effects, and are part of an extensive study of TbrPDE inhibitors involving numerous analogs.⁴⁷

The inhibitory mechanisms of both compounds were examined by varying both substrates. The results were fit to models representing various modes of inhibition (see Methods, Supplemental). The potencies (K_I values) of compounds 1 and 2 against DNMT3A range from 3.7 to 18 μM (AdoMet) and 11 to 41 μM (poly dI-dC), which compare favorably to numerous published efforts.^{29,30} The best fits to the inhibition data for both compounds against poly dI-dC and AdoMet are consistent with mixed type or uncompetitive mechanisms (see Figure 3, Table 1). Importantly, both mechanisms require that compounds 1 and 2 bind allosterically, away from the active site of the enzyme. The mixed type mechanism allows for scenarios in which the inhibitor binds both forms of the enzyme with the pertinent substrate bound, or unbound. In contrast, the uncompetitive mechanisms (Figure 3, Table 1, Compound 2) implicate a mechanism wherein the inhibitor only binds to the form of the enzyme already bound by the

DNA. The mechanisms of other DNMT inhibitors, when reported, often display competitive mechanisms with DNA, AdoMet, or both.^{30,44,48} The simplest interpretation of these mechanisms is that the inhibitor binds the same site as DNA or AdoMet, or, minimally, binds the same form of the enzyme bound by these substrates.⁴⁹

<i>Inhibitor</i>	1		2	
<i>Substrate</i>	AdoMet	poly dI-dC	AdoMet	poly dI-dC
<i>Best fit K_i (μM)</i>	3.70 – 7.06	12.64 – 40.56	9.16-18.85	11.37 – 23.34
<i>Best fit Mechanism</i>	<i>Mixed</i>	<i>Mixed</i>	<i>Uncompetitive</i>	<i>Uncompetitive</i>
<i>Goodness of fit (R^2)</i>	0.957	0.831	0.983	0.891

Table 1 – Values for the various fits of inhibitors with respect to both substrates. Fits were determined using the *Noncompetitive* and *Uncompetitive* nonlinear regression models in *Prism 8.4.3*. The reported bounds define the 95% confidence interval of the K_i value.





Figure 3 – Best fit plots of the inhibition with respect to both substrates, poly dI-dC (A, B) and AdoMet (C, D). Assays were performed with 150 nM DNMT3A with an excess of the substrate being held constant. Radiolabeled ^3H -AdoMet was used to determine product formation. All reactions were assayed for 30 min, then quenched with 0.1% SDS and spotted onto charged nylon membranes for detection. Data was collected with two replicates ($n=2$). Fitting was performed with standard inhibition equations being applied to the whole model. Extracted K_i values are boxed, while corresponding reciprocal plot with

The widespread cellular reliance on AdoMet-dependent methyltransferases suggests that the development of drugs specific for DNA methyltransferases or drugs that distinguish between DNMT1 and DNMT3A will be challenging. This is reflected by the fact that the majority of DNA methyltransferase inhibitors are poorly selective for DNMT3A, likely contributing to the limiting toxicity displayed by these compounds.³⁰ DNMT1 is critical to cell viability and given the prevalence of DNMT1 throughout the lifetime of somatic cells, the selective inhibition of DNMT3A over DNMT1 is important in the development of cancer treatments.⁷ This is especially true of cancers like AML, where prevalence of DNMT3A mutations is particularly high. Further inhibition studies aimed to see if these compounds would affect DNMT1. Additionally, given the implicated allostery, we wanted to investigate if these compounds could inhibit the bacterial cytosine methyltransferase *M. SssI* (see Methods, Supplemental). This protein has a highly conserved active site with respect to DNMT3A but does not share its allosteric structure.²² The results with respect to DNMT3A and DNMT1 found both compounds show some selectivity, with inhibitor 2 being the more selective of the two (see Figure 4). Neither compound shows inhibition of the bacterial DNA cytosine methyltransferase *M.SssI*, even at 60 μ M. Both compounds show little inhibition of DNMT1 at 6 μ M, and compound 2 retains this selectivity even at 60 μ M. It is intriguing that both inhibitors show greater inhibition of the catalytic domain of DNMT3A (residues 628 to 912, see Figure 1) than the full length DNMT3A, suggesting that the large N-terminal segment interferes with the inhibition. The basis of this difference has diverse molecular explanations, which we are actively investigating. Both compound 1 and 2 are still able to modulate the more biologically relevant full-length form of DNMT3A.




Figure 4 – Inhibitors 1 (A) and 2 (B) modulate the activity of human *de novo* DNMT3A_CD (■), DNMT3A_FL (■) and maintenance DNMT1 (■) DNA methyltransferases but not of the bacterial DNA methyltransferase M.SssI (■). To test the specificity of inhibitor 1 and 2 for these distinct types of CpG DNA methyltransferases, assays were performed with 150 nM DNMT3A_CD (■), DNMT3A_FL (■), 133 U/mL DNMT1 (■) and 266 U/mL M.SssI (A and B). Inhibitors 1 (A) and 2 (B) were tested at 6 and 60 μM with poly dI-dC and AdoMet at a fixed concentration of 1.8 ng/μL and 5 μM, respectively. In all reactions (A and B), enzymes and inhibitor 1 or 2 were pre-incubated for 15 minutes at 37 °C in methylation reaction buffer (see methods) prior to initiating the reaction by the addition of DNA. Methylation reactions were assayed for 30 min, quenched with 0.1% SDS and spotted onto charged nylon membranes for detection of radiolabeled ³H-AdoMet DNA. Fold decrease was calculated by dividing the product formed in reactions without compound 1 or 2 by the product formed in reactions with compound 1 or 2. Data reflect the mean ± S.D. of 3 independent experiments; one-way analysis of variance (ANOVA) was used to compare the values

C. Conclusion

In summary, the screening of a small chemical library of known drugs against human DNMT3A identified two non-nucleoside molecules of low micromolar potency. Both molecules inhibit the enzyme by binding outside the active site, and not only selectively inhibit human over bacterial DNMTs, but also shows some promising preferential targeting of *de novo* over maintenance DNA methyltransferases. This highlights the potential use of these molecules for the treatment of malignancies associated with disruptions to DNMT3A activity. The large number of analogs of these two inhibitors which have been described provides a promising basis for further optimization of this new group of DNMT3A inhibitors, with reasonable prospects of showing improved toxicity over known DNA methyltransferase drugs.⁴⁶

D. Material and Methods

Protein expression and purification

The isolated catalytic domain (CD) of DNMT3A (residues 634-912), which was used in all DNMT3A assays, has comparable kinetic parameters (k_{cat} , K_{mDNA} , K_{mAdoMet}) and similar regulatory responses to DNMT3L to the full length protein.^{50,51} The protein was expressed using codon optimized plasmids pET28a-hDNMT3A_CD ($\Delta 1-611$).⁵²

All proteins were expressed in NiCo21 (DE3) competent *E. coli* cells (New England Biolabs). Using LB medium, cell cultures were grown at 37 °C until an $A_{600 \text{ nm}}$ of 0.6 was reached. Following growth, expression was induced at 28 °C using 1 mM isopropyl β -D-thiogalactopyranoside (Gold Biotechnology). This addition marked the beginning of a 6 h induction time that ended with centrifugation and collection of the resulting cell pellet for storage

at -80 °C. Further purification began with cell lysis via sonication in 50 mM HEPES, 500 mM NaCl, 50 mM imidazole, 10% glycerol and 1% phenylmethylsulfonyl fluoride (PMSF) at pH 7.8. The solution was clarified by centrifugation and then loaded into an AKTA start FPLC (GE Healthcare) for purification. This was performed using a nickel-nitrilotriacetic acid column (GE Healthcare), a 50 mM HEPES, 500 mM NaCl, 50 mM imidazole, 10% glycerol and 1% PMSF at pH 7.8 buffer for equilibration and a similar 70 mM imidazole buffer for washing. Elution of column bound protein was triggered with a 500 mM imidazole buffer equivalent. The storage of all collected proteins was done at -80 °C in 50 mM KH₂PO₄/ K₂HPO₄, 20% glycerol buffer at pH 7.8. The activity of obtained DNMT3A was determined by methylation assay (described below).

Methylation Assays

In vitro methylation assays were used to determine the total amount of product (methylated poly dI-dC DNA) generated through the catalytic cycle of active DNMT3A. Reactions with 150 nM DNMT3A tetramer were carried out at 37 °C, pH 7.8 in 50 mM KH₂PO₄/K₂HPO₄, 1 mM DTT, 0.2 mg/mL BSA, 20 mM NaCl and 5 μM AdoMet (composed of 4.5 μM unlabeled and 0.5 μM 3H-methyl labeled cofactor). 20 μL assays were preincubated for 20 min at 37 °C followed by the addition of 1.8 ng/μL poly dI-dC DNA. After the specified assay time, the reactions were quenched with 0.1% SDS and spotted onto Hybond-XL membranes (GE Healthcare). The membranes were washed with both 50 mM KH₂PO₄/ K₂HPO₄ and ethanol. Following a drying period, samples were analyzed using a Beckmann LS6000 liquid scintillation counter.

Library screening

Screening was done in 96 conical well plates (Costar). The average maximum turnover rate from 8 wells with 150 nM DNMT3A WT served as control groups. A master mix contained reaction buffer (50 mM $\text{KH}_2\text{PO}_4/\text{K}_2\text{HPO}_4$, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml of BSA, 20 mM NaCl), and 5 μM AdoMet (from a 50 μM stock composed of 45 μM unlabeled and 5 μM ^3H -methyl labeled at pH 7.8) and 150 nM 3A were mixed and then aliquoted into each well. Chemical library compounds (Pathogen box) were added into each well at 60 μM and incubated at 37 °C for 60 mins after addition of 5 μM poly dI-dC. All wells were quenched with 0.1% SDS (1:1) after 1-hour. Samples (15 μL) were spotted onto Hybond-XL membranes (GE Healthcare), washed, and dried.

Kinetic studies

To determine the dependence of inhibition on the individual cofactors, methylation assays (described above) were performed with inhibitor at three different concentrations (0 μM , $\sim\text{IC}_{50}$, Excess), one cofactor held constant (5 μM AdoMet or 12.0 ng/ μL poly dI-dC), and the other varied. For the variation of AdoMet, concentrations of 10, 4, 1, 0.5, 0.2, 0.16, and 0.08 μM were used, while for the variation of DNA, 0.15, 0.25, 0.3, 0.5, 1, 1.5, +3.0 and 12.0 ng/ μL poly dI-dC DNA were used.

Data Analysis

Using *Prism 8.4.3*, all data was fit both to non-linear models and converted to double-reciprocal format and fitted to linear-models. Using the tabulated Goodness of Fit (R^2) values, the most statistically likely mechanism of inhibition was chosen (Table 1S).

<i>Inhibitor</i>	1		2	
<i>Cofactor</i>	AdoMet	poly dI-dC	AdoMet	poly dI-dC
<i>Competitive Fit</i>	0.8927	0.7690	0.9507	0.5304
<i>Uncompetitive Fit</i>	0.9467	0.8124	0.9833	0.8909
<i>Noncompetitive Fit</i>	0.9571	0.8312	0.9804	0.8572

Table 1S – R^2 values for the non-linear fits of the available inhibition data with respect to three generalized inhibition mechanisms: competitive, uncompetitive and noncompetitive

Specificity of inhibitors to human DNA methyltransferases

To assess the specificity of inhibitor 1 and 2 for DNMT3A, DNA methylation assays (described above) were performed with each inhibitor at two distinct concentrations (6 or 60 μ M) with both poly dI-dC DNA (1.8 ng/ μ L) and AdoMet (5 μ M) held at a constant concentration. The distinct types of DNA methyltransferases and their concentrations were as follows: 150 nM full-length (DNMT3A_FL) or catalytic domain DNMT3A (DNMT3A_CD), 133 U/mL DNMT1 (purchased form NEB) and 266 U/mL M.SssI (purchased form NEB).

REFERENCES:

1. Bergman Y, Cedar H. DNA methylation dynamics in health and disease. *Nat Struct Mol Biol.* 2013;20(3):274-281. doi:10.1038/nsmb.2518
2. Baylin SB, Jones PA. A decade of exploring the cancer epigenome-biological and translational implications. *Nat Rev Cancer.* 2011;11(10):726-734. doi:10.1038/nrc3130
3. Li E, Bestor TH, Jaenisch R. Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell.* 1992. doi:10.1016/0092-8674(92)90611-F
4. Jeltsch A, Jurkowska RZ. Allosteric control of mammalian DNA methyltransferases - A new regulatory paradigm. *Nucleic Acids Res.* 2016;44(18):8556-8575. doi:10.1093/nar/gkw723
5. Wong KK, Lawrie CH, Green TM. Oncogenic Roles and Inhibitors of DNMT1, DNMT3A, and DNMT3B in Acute Myeloid Leukaemia. *Biomark Insights.* 2019;14. doi:10.1177/1177271919846454
6. Vertino PM, Yen RW, Gao J, Baylin SB. De novo methylation of CpG island sequences in human fibroblasts overexpressing DNA (cytosine-5-)-methyltransferase. *Mol Cell Biol.* 1996;16(8):4555-4565. doi:10.1128/mcb.16.8.4555
7. Jeltsch A, Jurkowska RZ. New concepts in DNA methylation. *Trends Biochem Sci.* 2014;39(7):310-318. doi:10.1016/j.tibs.2014.05.002
8. Rasmussen KD, Helin K. Role of TET enzymes in DNA methylation, development, and cancer. *Genes Dev.* 2016;30(7):733-750. doi:10.1101/gad.276568.115
9. Okano M, Bell DW, Haber DA, Li E. DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell.* 1999. doi:10.1016/S0092-8674(00)81656-6
10. Tahiliani M, Koh KP, Shen Y, et al. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science (80-).* 2009. doi:10.1126/science.1170116
11. Gromova ES, Khoroshaev A V. Prokaryotic DNA methyltransferases: The structure and the mechanism of interaction with DNA. *Mol Biol.* 2003;37(2):300-314.
12. Wu JC, Santi D V. Kinetic and catalytic mechanism of HhaI methyltransferase. *J Biol Chem.* 1987;262(10):4778-4786.
13. Bheemanaik S, Reddy YVR, Rao DN. Structure, function and mechanism of exocyclic


- DNA methyltransferases. *Biochem J*. 2006;399(2):177-190. doi:10.1042/BJ20060854
14. Reich NO, Mashhoon N. Kinetic Mechanism of the EcoRI DNA Methyltransferase. *Biochemistry*. 1991;30(11):2933-2939. doi:10.1021/bi00225a029
 15. Yokochi T, Robertson KD. Preferential Methylation of Unmethylated DNA by Mammalian de Novo DNA Methyltransferase Dnmt3a*. doi:10.1074/jbc.M106590200
 16. Lyko F. The DNA methyltransferase family: A versatile toolkit for epigenetic regulation. *Nat Rev Genet*. 2018;19(2):81-92. doi:10.1038/nrg.2017.80
 17. Hermann A, Goyal R, Jeltsch A. DNA Processively with High Preference for Hemimethylated. *J Biol Chem*. 2004.
 18. Pradhan S, Bacolla A, Wells RD, Roberts RJ. Recombinant human DNA (cytosine-5) methyltransferase. I. Expression, purification, and comparison of novo and maintenance methylation. *J Biol Chem*. 1999. doi:10.1074/jbc.274.46.33002
 19. Okano M, Xie S, Li E. Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases [1]. *Nat Genet*. 1998. doi:10.1038/890
 20. Leonhardt H, Page AW, Weier HU, Bestor TH. A targeting sequence directs DNA methyltransferase to sites of DNA replication in mammalian nuclei. *Cell*. 1992. doi:10.1016/0092-8674(92)90561-P
 21. Zhang ZM, Lu R, Wang P, et al. Structural basis for DNMT3A-mediated de novo DNA methylation. *Nature*. 2018;554(7692):387-391. doi:10.1038/nature25477
 22. Lukashevich O V., Cherepanova NA, Jurkovska RZ, Jeltsch A, Gromova ES. Conserved motif VIII of murine DNA methyltransferase Dnmt3a is essential for methylation activity. *BMC Biochem*. 2016. doi:10.1186/s12858-016-0064-y
 23. Brunetti L, Gundry MC, Goodell MA. DNMT3A in Leukemia. *Cold Spring Harb Perspect Med*. 2017;7(2). doi:10.1101/cshperspect.a030320
 24. Khrabrova DA, Loiko AG, Tolkacheva AA, et al. Functional analysis of DNMT3A DNA methyltransferase mutations reported in patients with acute myeloid leukemia. *Biomolecules*. 2020;10(1). doi:10.3390/biom10010008
 25. Holz-Schietinger C, Matje DM, Reich NO. Mutations in DNA methyltransferase (DNMT3A) observed in acute myeloid leukemia patients disrupt processive methylation. *J Biol Chem*. 2012;287(37):30941-30951. doi:10.1074/jbc.M112.366625
 26. Sandoval JE, Huang YH, Muise A, Goodell MA, Reich NO. Mutations in the DNMT3A

- DNA methyltransferase in acute myeloid leukemia patients cause both loss and gain of function and differential regulation by protein partners. *J Biol Chem.* 2019;294(13):4898-4910. doi:10.1074/jbc.RA118.006795
27. Sandoval JE, Reich NO. The R882H substitution in the human de novo DNA methyltransferase DNMT3A disrupts allosteric regulation by the tumor suppressor p53. *J Biol Chem.* 2019;294(48):18207-18219. doi:10.1074/jbc.RA119.010827
 28. Emperle M, Adam S, Kunert S, et al. Mutations of R882 change flanking sequence preferences of the DNA methyltransferase DNMT3A and cellular methylation patterns. *Nucleic Acids Res.* 2019;47(21):11355-11367. doi:10.1093/nar/gkz911
 29. Gros C, Fahy J, Halby L, et al. DNA methylation inhibitors in cancer: Recent and future approaches. *Biochimie.* 2012;94(11):2280-2296. doi:10.1016/j.biochi.2012.07.025
 30. Yu J, Xie T, Wang Z, et al. DNA methyltransferases: emerging targets for the discovery of inhibitors as potent anticancer drugs. *Drug Discov Today.* 2019;24(12):2323-2331. doi:10.1016/j.drudis.2019.08.006
 31. Ganesan A, Arimondo PB, Rots MG, Jeronimo C, Berdasco M. The timeline of epigenetic drug discovery: From reality to dreams. *Clin Epigenetics.* 2019;11(1):1-17. doi:10.1186/s13148-019-0776-0
 32. Santi D V., Norment A, Garrett CE. Covalent bond formation between a DNA-cytosine methyltransferase and DNA containing 5-azacytosine. *Proc Natl Acad Sci U S A.* 1984. doi:10.1073/pnas.81.22.6993
 33. Stresemann C, Lyko F. Modes of action of the DNA methyltransferase inhibitors azacytidine and decitabine. *Int J Cancer.* 2008;123(1):8-13. doi:10.1002/ijc.23607
 34. DNMT3A Result Summary | BioGRID. <https://thebiogrid.org/108125/summary/homo-sapiens/dnmt3a.html>. Accessed August 22, 2020.
 35. Russler-Germain DA, Spencer DH, Young MA, et al. Cancer Cell The R882H DNMT3A Mutation Associated with AML Dominantly Inhibits Wild-Type DNMT3A by Blocking Its Ability to Form Active Tetramers. 2014. doi:10.1016/j.ccr.2014.02.010
 36. Díaz-Eufracio BI, Naveja JJ, Medina-Franco JL. Protein–Protein Interaction Modulators for Epigenetic Therapies. In: *Advances in Protein Chemistry and Structural Biology.* Vol 110. Academic Press Inc.; 2018:65-84. doi:10.1016/bs.apcsb.2017.06.002
 37. Ni D, Lu S, Zhang J. Emerging roles of allosteric modulators in the regulation of protein-

- protein interactions (PPIs): A new paradigm for PPI drug discovery. *Med Res Rev.* 2019;39(6):2314-2342. doi:10.1002/med.21585
38. Cheng SS, Yang GJ, Wang W, Leung CH, Ma DL. The design and development of covalent protein-protein interaction inhibitors for cancer treatment. *J Hematol Oncol.* 2020;13(1):26. doi:10.1186/s13045-020-00850-0
 39. Wenthur CJ, Gentry PR, Mathews TP, Lindsley CW. Drugs for Allosteric Sites on Receptors. *Annu Rev Pharmacol Toxicol.* 2014;54(1):165-184. doi:10.1146/annurev-pharmtox-010611-134525
 40. Dokholyan N V. Controlling Allosteric Networks in Proteins. *Chem Rev.* 2016. doi:10.1021/acs.chemrev.5b00544
 41. Peracchi A, Mozzarelli A. Exploring and exploiting allostery: Models, evolution, and drug targeting. *Biochim Biophys Acta - Proteins Proteomics.* 2011;1814(8):922-933. doi:10.1016/j.bbapap.2010.10.008
 42. Dahlin JL, Walters MA. The essential roles of chemistry in high-throughput screening triage. *Future Med Chem.* 2014;6(11):1265-1290. doi:10.4155/fmc.14.60
 43. Siedlecki P, Boy RG, Musch T, et al. Discovery of two novel, small-molecule inhibitors of DNA methylation. *J Med Chem.* 2006;49(2):678-683. doi:10.1021/jm050844z
 44. Valente S, Liu Y, Schnekenburger M, et al. Selective non-nucleoside inhibitors of human DNA methyltransferases active in cancer including in cancer stem cells. *J Med Chem.* 2014;57(3):701-713. doi:10.1021/jm4012627
 45. Ceccaldi A, Rajavelu A, Ragozin S, et al. Identification of novel inhibitors of dna methylation by screening of a chemical library. *ACS Chem Biol.* 2013;8(3):543-548. doi:10.1021/cb300565z
 46. Blaazer AR, Singh AK, De Heuvel E, et al. Targeting a Subpocket in Trypanosoma brucei Phosphodiesterase B1 (TbrPDEB1) Enables the Structure-Based Discovery of Selective Inhibitors with Trypanocidal Activity. *J Med Chem.* 2018;61(9):3870-3888. doi:10.1021/acs.jmedchem.7b01670
 47. Orrling KM, Jansen C, Vu XL, et al. Catechol pyrazolinones as trypanocidals: Fragment-based design, synthesis, and pharmacological evaluation of nanomolar inhibitors of trypanosomal phosphodiesterase B1. *J Med Chem.* 2012;55(20):8745-8756. doi:10.1021/jm301059b

48. Datta J, Ghoshal K, Denny WA, et al. A new class of quinoline-based DNA hypomethylating agents reactivates tumor suppressor genes by blocking DNA methyltransferase 1 activity and inducing its degradation. *Cancer Res.* 2009;69(10):4277-4285. doi:10.1158/0008-5472.CAN-08-3669
49. Segel IH. *Enzyme Kinetics: Behavior and Analysis of Rapid Equilibrium and Steady-State Enzyme Systems.* Wiley; 1975.
50. Chen T, Ueda Y, Dodge JE, Wang Z, Li E. Establishment and Maintenance of Genomic Methylation Patterns in Mouse Embryonic Stem Cells by Dnmt3a and Dnmt3b. *Mol Cell Biol.* 2003;23(16):5594-5605. doi:10.1128/mcb.23.16.5594-5605.2003
51. Kaneda M, Okano M, Hata K, et al. Essential role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting. *Nature.* 2004;429(6994):900-903. doi:10.1038/nature02633
52. Purdy MM, Holz-Schietinger C, Reich NO. Identification of a second DNA binding site in human DNA methyltransferase 3A by substrate inhibition and domain deletion. *Arch Biochem Biophys.* 2010;498(1):13-22. doi:10.1016/j.abb.2010.03.007

APPENDIX



Secondary Screen Results




Figure 1S – Results from the secondary screen. Compounds that exhibited more than 90% inhibition in the primary screen of the Pathogen Box library were subjected to duplicate (n=2) repetition of the inhibition. Activity assays were performed as described in the Methods. The compounds are designated by their location in the library and reference structures are provided below. Compound 1 is designated AC05 while compound 2 is BC02.