Lawrence Berkeley National Laboratory

Recent Work

Title

Hopf Algebra of Vector Fields on Complex Quantum Groups

Permalink

<https://escholarship.org/uc/item/6pg0g9mz>

Authors

Drabant, B.
Jurco, B.
Schlieker, M.
et al.

Publication Date

1992-05-01

May 1992

MPI-Ph/92-39

LMU-TPW 1992-6

UCB-PTh 92/16

LBL-32354

The Hopf Algebra of Vector Fields on Complex Quantum Groups

*Bernhard Drabant*¹, *Branislav Jurčo*^{2 †}, *Michael Schlieker*³
*Wolfgang Weich*³, *Bruno Zumino*⁴

¹ *Max-Planck-Institut für Physik, Werner-Heisenberg-Institut
Föhringer Ring 6, D-8000 München 40, Germany*

² *Arnold-Sommerfeld-Institut, Technische Universität Clausthal
D-3392 Clausthal-Zellerfeld, Germany*

³ *Sektion Physik der Universität München, Lehrstuhl Professor Wess
Theresienstraße 37, D-8000 München 2, Germany*

⁴ *Department of Physics, University of California
Berkeley, California 94720, USA*

and

*Lawrence Berkeley Laboratory, Theoretical Physics Group
1 Cyclotron Road, Berkeley, California 94720, USA*

Abstract

We derive the equivalence of the complex quantum enveloping algebra and the algebra of complex quantum vector fields for the Lie algebra types A_n , B_n , C_n and D_n by factorizing the vector fields uniquely into a triangular and a unitary part and identifying them with the corresponding elements of the algebra of regular functionals.

[†] *Humboldt-Fellow*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

1. Introduction

In [CDSWZ] we constructed the algebra of vector fields on complex quantum groups [CW, DSWZ, OSWZ, SWZ]. For the special case of A_1 we proved that the algebra of vector fields generates the algebra of regular functionals and coincides in some sense with $U_q sl(2, \mathcal{C})$ thus showing in particular that the algebra of vector fields admits a Hopf-structure. In this paper we want to prove that these facts generalize to A_n, B_n, C_n and D_n .

In the real case it is known that the Hopf algebra of regular functionals is generated by the vector fields which appear in the bicovariant differential calculus on quantum groups [CSWW, Jur, Wor, Zum]. This is proved in [Bur] using the fact that the matrices L^{+i}_j and L^{-i}_j generating the algebra of regular functionals are upper and lower triangular respectively. A similar procedure involving the Gram-Schmidt-orthogonalization scheme, also used for quantum groups in [JS], is applied here to the vector fields of the complex quantum groups [CDSWZ] leading to the result mentioned above that vector fields generate the algebra of regular functionals.

2. The Algebra of Vector Fields

Throughout this paper we are using the notations and conventions of [DSWZ].

Set $(I) := (i, \bar{i})$, $\bar{I} := (\bar{i}, i) = (\bar{i}, i)$, $(i, \bar{i} = 1, \dots, N)$ where $N = n$ for A_{n-1} , $N = 2n+1$ for B_n and $N = 2n$ for C_n, D_n . Define then the $2N \times 2N$ -matrix

$$(T^I_J) := \begin{pmatrix} t & 0 \\ 0 & \hat{t} \end{pmatrix} \quad (2.1)$$

and the $\hat{\mathcal{R}}$ -matrix

$$(\hat{\mathcal{R}}^{IJ}_{KL}) := \begin{pmatrix} \beta \hat{R}_q & 0 & 0 & 0 \\ 0 & 0 & \beta \hat{R}_q & 0 \\ 0 & \beta^{-1} \hat{R}_q^{-1} & 0 & 0 \\ 0 & 0 & 0 & \beta^{-1} \hat{R}_q^{-1} \end{pmatrix} \quad (2.2)$$

with the corresponding $\hat{R}_q$ -matrix [FRT] and with

$$\beta := \begin{cases} q^{-\frac{1}{n}} > 0 & \text{for } A_{n-1} \\ 1 & \text{for } B_n, C_n \text{ and } D_n. \end{cases} \quad (2.3)$$

For convenience we fixed the parameters α_i from [DSWZ] to the special values $\alpha_0 = \alpha_1 = \alpha_2^{-1} = \alpha_3^{-1} = \beta$.

We are considering the following quantum group

$$\mathcal{A} := \mathcal{C} \langle T^I_J \rangle / ((2.5), (2.6), (2.7)). \quad (2.4)$$

where the ideal is generated by

$$I_{ST}^{IJ} := \hat{\mathcal{R}}_q^{IJ}{}_{KL} T^K_S T^L_T - T^I_V T^J_W \hat{\mathcal{R}}_q^{VW}{}_{ST}, \quad (2.5)$$

$$\det(t^i_j) - 1 = \frac{(-1)^{n-1}}{[n]_q!} q^{-\binom{n}{2}} \varepsilon^{k_1 \dots k_n} t^{l_1}_{k_1} \dots t^{l_n}_{k_n} \varepsilon_{l_1 \dots l_n} - 1 \quad \text{for } A_{n-1}, \quad (2.6)$$

$$t^i_s (C^{-1})^{sk} t^l_k C_{lj} - \delta^i_j \mathbf{1}, \quad (C^{-1})^{ik} t^l_k C_{ls} t^s_j - \delta^i_j \mathbf{1} \quad \text{for } B_n, C_n, D_n,$$

$$\det(\hat{t}^{\bar{i}}_{\bar{j}}) - 1 = \frac{(-1)^{n-1}}{[n]_q!} q^{-\binom{n}{2}} \varepsilon^{\bar{k}_1 \dots \bar{k}_n} \hat{t}^{\bar{l}_1}_{\bar{k}_1} \dots \hat{t}^{\bar{l}_n}_{\bar{k}_n} \varepsilon_{\bar{l}_1 \dots \bar{l}_n} - 1 \quad \text{for } A_{n-1}, \quad (2.7)$$

$$\hat{t}^{\bar{i}}_{\bar{s}} (C^{-1})^{\bar{s}\bar{k}} \hat{t}^{\bar{l}}_{\bar{k}} C_{\bar{l}\bar{j}} - \delta^{\bar{i}}_{\bar{j}} \mathbf{1}, \quad (C^{-1})^{\bar{i}\bar{k}} \hat{t}^{\bar{l}}_{\bar{k}} C_{\bar{l}\bar{s}} \hat{t}^{\bar{s}}_{\bar{j}} - \delta^{\bar{i}}_{\bar{j}} \mathbf{1} \quad \text{for } B_n, C_n, D_n$$

with $\varepsilon_{i_1 \dots i_n} = (-1)^{n-1} \varepsilon^{i_1 \dots i_n} = (-q)^{l(\sigma)}$, $l(\sigma)$ is the length (minimal number of transpositions) of the permutation $\sigma = \begin{pmatrix} 1 & \dots & n \\ i_1 & \dots & i_n \end{pmatrix}$, $[n]_q = (q^n - q^{-n})/(q - q^{-1})$, $[n]_q! = [1]_q \dots [n]_q$ [CSWW] and C_{ij} is the usual metric [FRT].

With the involution

$$(t^i_j)^* := \kappa(\hat{t}^{\bar{j}}_{\bar{i}}) \quad (2.8)$$

$\mathcal{A}$ becomes a $*$ -Hopf algebra with comultiplication Φ , counit e and antipode κ [DSWZ].

The dual space $\mathcal{A}^*$ of the Hopf algebra $\mathcal{A}$ is an algebra with the convolution product. One can introduce an antimultiplicative involution $''^*$ on $\mathcal{A}^*$: For $f \in \mathcal{A}^*$ one sets

$$\forall a \in \mathcal{A}: \quad f^*(a) := \overline{f(\kappa^{-1}(a^*))}. \quad (2.9)$$

We define functionals $L^{\pm I}_J \in \mathcal{A}^*$ through their action on the generators of $\mathcal{A}$:

$$\begin{aligned} L^{\pm I}_J(\mathbf{1}) &:= \delta^I_J, \\ L^{\pm I}_J(T^K_L) &:= \hat{\mathcal{R}}_q^{\pm 1 IK}_{LJ} \end{aligned} \quad (2.10)$$

and their comultiplication

$$\forall a, b \in \mathcal{A}: \quad L^{\pm I}_J(ab) = L^{\pm I}_K(a) L^{\pm K}_J(b). \quad (2.11)$$

The algebra $U_{\mathcal{R}}$ of regular functionals on $\mathcal{A}$ is the unital algebra generated by $\{L^{\pm I}_J\}$ [DSWZ]. It is shown in [DSWZ] that $U_{\mathcal{R}}$ is a $*$ -Hopf algebra with comultiplication Δ , counit ϵ and antipode S .

Now we introduce the matrices

$$\begin{aligned} Y &:= L^+ S(L^-) = \begin{pmatrix} y & 0 \\ 0 & \hat{y} \end{pmatrix}, \\ Y^{-1} &:= L^- S(L^+) = \begin{pmatrix} y^{-1} & 0 \\ 0 & \hat{y}^{-1} \end{pmatrix} \end{aligned} \quad (2.12)$$

with the matrix entries Y^I_J and $Y^{-1 I}_J \in U_{\mathcal{R}}$. It follows from the commutation relations of $L^{\pm I}_J$ derived in [DSWZ] that

$$\hat{\mathcal{R}}_q(\mathbf{1} \otimes Y) \hat{\mathcal{R}}_q(\mathbf{1} \otimes Y) = (\mathbf{1} \otimes Y) \hat{\mathcal{R}}_q(\mathbf{1} \otimes Y) \hat{\mathcal{R}}_q. \quad (2.13)$$

The Y^I_J have the comultiplication

$$\Delta(Y^I_J) = \Theta^{+IK}_{LJ} \otimes Y^L_K, \quad (2.14)$$

where

$$\Theta^{+IK}_{LJ} = L^{+I}_L S(L^{-K}_J). \quad (2.15)$$

We set for any matrix M , $M^I_J \in U_{\mathcal{R}}$ the hermitian involution " $\dagger$ " with $M^{\dagger I}_J := (M^J_I)^*$.

Using the involution properties of the $L^{\pm I}_J$ (see (3.13) of [DSWZ]) one obtains

$$Y^{\dagger J}_I = Y^{-1 J}_I. \quad (2.16)$$

For A_{n-1} it is possible to define an invariance relation for antisymmetric n -fold products of the y^i_j and $\hat{y}^{\bar{i}}_{\bar{j}}$ (—from that one gets a modified determinant for the y^i_j and the $\hat{y}^{\bar{i}}_{\bar{j}}$ resp.). By observing that the $t^i_k y^k_j$ behave like the t^i_j under coaction of the group and have the same commutation relations we make the ansatz

$$\varepsilon_{i_1 \dots i_n} t^{i_1}_{k_1} y^{k_1}_{j_1} \dots t^{i_n}_{k_n} y^{k_n}_{j_n} = \varepsilon_{j_1 \dots j_n} \det(ty) \quad (2.17)$$

Formula (3.23) from [CDSWZ] and the relation corresponding to (2.17) for the t^i_j yield the following invariance relation

$$q^{-(n^2-n)} \varepsilon^{i_1 \dots i_n} \cdot e = (y)_1 (\hat{R}_{q2,1}(y)_1 \hat{R}_{q2,1}) \dots \dots (\hat{R}_{qn,n-1} \dots \hat{R}_{q2,1}(y)_1 \hat{R}_{q2,1} \dots \hat{R}_{qn,n-1})^{i_1 \dots i_n}_{j_1 \dots j_n} \varepsilon^{j_n \dots j_1}. \quad (2.18)$$

For the $\hat{y}^{\bar{i}}_{\bar{j}}$ one obtains analogously

$$q^{(n^2-n)} \varepsilon^{\bar{i}_1 \dots \bar{i}_n} \cdot e = (\hat{y})_1 (\hat{R}_q^{-1}{}_{2,1}(\hat{y})_1 \hat{R}_q^{-1}{}_{2,1}) \dots \dots (\hat{R}_q^{-1}{}_{n,n-1} \dots \hat{R}_q^{-1}{}_{2,1}(\hat{y})_1 \hat{R}_q^{-1}{}_{2,1} \dots \hat{R}_q^{-1}{}_{n,n-1})^{\bar{i}_1 \dots \bar{i}_n}_{\bar{j}_1 \dots \bar{j}_n} \varepsilon^{\bar{j}_n \dots \bar{j}_1}. \quad (2.19)$$

The matrices $\hat{R}_q^{\pm 1}{}_{l,l-1}$ are defined as in [FRT]. From (2.18) and (2.19) it is now possible to derive the modified determinant $\text{Det}(y)$ and the $\text{Det}(\hat{y})$ and to express the $(y^{-1})^i_j$ and the $(\hat{y}^{-1})^{\bar{i}}_{\bar{j}}$ in terms of the Y^I_J . If we normalize $\text{Det}(y)$ and $\text{Det}(\hat{y})$ to e we obtain

$$\begin{aligned} \det(ty) &= q^{-(n^2-1)} \det(t) \text{Det}(y), \\ \det(\hat{t}\hat{y}) &= q^{(n^2-1)} \det(\hat{t}) \text{Det}(\hat{y}). \end{aligned} \quad (2.20)$$

For B_n , C_n and D_n we get similar results.

$$\begin{aligned} C^{kt} &= q^{N-1} y^t_a y^s_u \hat{R}_q^{ka}{}_{vs} C^{vu}, \\ C^{\bar{k}\bar{t}} &= q^{-(N-1)} \hat{y}^{\bar{t}}_{\bar{a}} \hat{y}^{\bar{s}}_{\bar{u}} \hat{R}_q^{-1\bar{k}\bar{a}}{}_{\bar{v}\bar{s}} C^{\bar{v}\bar{u}}. \end{aligned} \quad (2.21)$$

From (2.21) one obtains Y^{-1} in terms of Y .

Because of (2.16) the algebra generated by $\{Y^I_J\}$ is closed under the involution " * ".

3. Generalized Schmidt Orthogonalization

With the orthogonalization scheme of Schmidt we decompose now the y -matrix ($\hat{y}$ -matrix) uniquely into the product of an upper (resp. lower) triangular and a unitary matrix ($U^\dagger = U^{-1}$) with entries in the algebra generated by y^a_b and $\hat{y}^{\bar{a}}_{\bar{b}}$. These factors are then identified as L^{+i}_j (resp. $L^{+\bar{i}}_{\bar{j}}$) which is upper (resp. lower) triangular with diagonalizable diagonal elements and $S(L^{-i}_j)$ (resp. $S(L^{-\bar{i}}_{\bar{j}})$) which is unitary.

For simplicity we fix the remaining ambiguity of the parameters α_i (see [DSWZ]) by $\alpha_i > 0$. In all other cases the results can be derived by just the same manner, if we restrict to $\alpha_0 = \alpha_1 = (\alpha_2)^{-1}$.

That L^{+i}_j (resp. $L^{+\bar{i}}_{\bar{j}}$) is upper (respectively lower) triangular has been discussed in [DSWZ]. The diagonal elements L^{+I}_I are algebra homomorphisms acting on $\mathcal{A}$ and

$$L^{+I}_I(T^K_L) = \hat{\mathcal{R}}^{IK}_q{}_{LI} > 0. \quad (3.1)$$

For $S(L^{-i}_j)$ one obtains

$$S(L^{-i}_j)^\dagger = S(L^{-j}_i)^* = (S^{-1}(L^{-j}_i))^* = L^{-\bar{i}}_{\bar{j}} = L^{-i}_j \quad (3.2)$$

using the equations (3.13) and part 3 from *proposition 8* in [DSWZ]. Thus we have proved the unitarity of $S(L^{-i}_j)$. The unitarity of $S(L^{-\bar{i}}_{\bar{j}})$ is proved analogously.

Write the matrix y

$$y = \begin{pmatrix} \vec{y}_1 \\ \vec{y}_2 \\ \vdots \\ \vec{y}_N \end{pmatrix} \quad (3.3)$$

with row vectors y_i .

Define for N -dimensional row vectors $\vec{a}, \vec{b} \in (U_{\mathcal{R}})^N$

$$(\vec{a}, \vec{b}) = \sum_{i=1}^N a^i (b^i)^*. \quad (3.4)$$

The decomposition equation reads

$$\begin{pmatrix} \vec{y}_1 \\ \vec{y}_2 \\ \vdots \\ \vec{y}_N \end{pmatrix} = \begin{pmatrix} T_{11} & \dots & \dots & T_{1N} \\ 0 & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & T_{NN} \end{pmatrix} \begin{pmatrix} \vec{g}_1 \\ \vec{g}_2 \\ \vdots \\ \vec{g}_N \end{pmatrix} \quad (3.5)$$

where the row vectors g_i fulfill an orthonormality condition

$$(g_i, g_j) = \delta_{ij} \quad (3.6)$$

and the diagonal elements T_{ii} being positive and diagonalizable.

This equation leads uniquely to the following recursion relations (Gram-Schmidt)

Starting with $k = N$ going to $k = 1$:

$$T_{kk} \vec{g}_k = \vec{y}_k - \sum_{\nu=k+1}^N (\vec{y}_k, \vec{g}_\nu) \vec{g}_\nu$$

and hence leading to the unique positive root of (3.7)

$$|T_{kk}|^2 = \left| \vec{y}_k - \sum_{\nu=k+1}^N (\vec{y}_k, \vec{g}_\nu) \vec{g}_\nu \right|^2$$

for $i > k$: $T_{ki} = (\vec{y}_k, \vec{g}_i)$.

This proves the uniqueness of the decomposition equation (3.5). One solution to (3.5) is given by

$$\begin{pmatrix} \vec{y}_1 \\ \vec{y}_2 \\ \vdots \\ \vec{y}_N \end{pmatrix} = \begin{pmatrix} T_{11} & \dots & \dots & T_{1N} \\ 0 & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & T_{NN} \end{pmatrix} \begin{pmatrix} \vec{g}_1 \\ \vec{g}_2 \\ \vdots \\ \vec{g}_N \end{pmatrix} = L^+ S(L^-). \quad (3.8)$$

The analogue procedure (where T is lower triangular) leads to the construction of the L^{+i}_j and $S(L^{-i}_j)$ in terms of the y^i_j and $\hat{y}^i_j$.

It is possible to obtain the L^{-I}_J from $S(L^{-K}_L)$ by the following relation

$$L^{-I}_J = \left(S(L^{-J}_I) \right)^* \quad (3.9)$$

because the Y -Algebra is closed under " $*$ ". We have thus shown how the algebra generated by $\{L^{\pm i}_j\}$ is reconstructed from the algebra of vector fields $\{Y^I_J\}$.

It is therefore proved that the algebra of regular functionals generated by $\{L^{\pm I}_J\}$ is equivalent to the algebra of vector fields $\{Y^I_J\}$ for A_n , B_n , C_n and D_n . With eqs. (2.14), (2.15) and the factorization $S(Y^I_J) = S^2(L^{-K}_J)L^{+I}_K$ (and using eq. (3.11) from [DSWZ]) it is obvious that the algebra of vector fields is a $\ast$ -Hopf algebra.

References

- [Bur] N. Burroughs, *Comm. Math. Phys.* **133**, 91 (1987).
- [CDSWZ] C. Chrysomalakos, B. Drabant, M. Schlieker, W. Weich and B. Zumino, preprint UCB 92/03 (1992), to appear in *Comm. Math. Phys.*.
- [CSWW] U. Carow-Watamura, M. Schlieker, S. Watamura and W. Weich, *Comm. Math. Phys.* **142**, 605 (1991).
- [CW] U. Carow-Watamura and S. Watamura, preprint TU-382 (1991).
- [DSWZ] B. Drabant, M. Schlieker, W. Weich and B. Zumino, preprint MPI-PTh/91-75, LMU-TPW 1991-5 (1992), to appear in *Comm. Math. Phys.*.
- [FRT] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, *Algebra and Analysis* **1**, 178 (1987).
- [Jur] B. Jurčo, *Lett. Math. Phys.* **22**, 177 (1991).
- [JS] B. Jurčo and P. Šťovíček, LMU-TPW-1992-3 (1992).
- [OSWZ] O. Ogievetsky, W.B. Schmidke, J. Wess and B. Zumino, MPI-Ph/91-51 (1991).
- [SWZ] W.B. Schmidke, J. Wess and B. Zumino, *Z. Phys. C* **52**, 471 (1991).
- [Wor] S.L. Woronowicz, *Comm. Math. Phys.* **122**, 125 (1989).
- [Zum] B. Zumino, preprint LBL-31432, UCB-PTh-62/91.