UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

UV-excitation from an experimental perspective: frequency resolved.

Permalink

<https://escholarship.org/uc/item/6ps8g3kn>

Journal

Topics in current chemistry, 355

ISSN

0340-1022

Author

de Vries, Mattanjah S

Publication Date

2015

DOI

10.1007/128_2014_560

Peer reviewed

UV-excitation from an experimental perspective I: frequency resolved

Mattanjah S. de Vries

Abstract

Electronic spectroscopy of DNA bases in the gas phase provides detailed information about the electronic excitation, which places the molecule in the Franck-Condon region in the excited state and thus prepares the starting conditions for excited state dynamics. Double resonant or hole burning spectroscopy in the gas phase can provide such information with isomer specificity, probing the starting potential energy landscape as a function of tautomeric form, isomeric structure, or hydrogen bonded or stacked cluster structure. Action spectroscopy, such REMPI, can be affected by excited state lifetimes.

keywords

DNA bases, nucleotides, REMPI, R2PI, hole burning, clusters, conical intersections

2.1. Introduction

Understanding the response of DNA bases to UV radiation is critical for both practical and fundamental reasons.¹⁻⁵ Nucleobase photochemistry following UV absorption constitutes a fundamental step in radiation-induced DNA damage. Short excited state lifetimes are often described as nature's strategy to protect the building blocks of life against UV photodamage. Furthermore, UV photo-selection may have played a key role in prebiotic chemistry on an early earth. This photochemistry schematically involves two steps: absorption of a photon prepares the nucleobase in an excited state, followed by ensuing chemistry. The first step, the excitation, can be probed by various forms of electronic spectroscopy. The second step, the dynamics of the excited state once it is formed, can often be followed in the time domain, which is the topic of the next chapter. This current chapter focuses on the frequency domain, probing primarily the excitation step. However, some time domain information also follows from the frequency domain because both steps are connected by Heisenberg's uncertainty principle: $\Delta E \Delta \tau \geq h/2\pi$ such that a shorter excited state lifetime, τ , corresponds to a larger linewidth, ΔE .

The fate of a molecule upon irradiation depends on the shape of the excited state potential energy landscape⁶. Electronic excitation places the molecule in the Franck-Condon region of the excited state, as the starting point of a number of competing possible processes, such as fluorescence, internal conversion, intersystem crossing or exciplex formation^{7, 8}. The availability of these pathways and their relative rates, or associated excited state lifetimes, determines the outcome of this competition and thus the product of the photochemistry. One molecule can therefore be intrinsically more resistant to damage than another even though they absorb UV light equally. Furthermore, the same molecule can have a different UV response, depending on the molecular environment. Of particular interest in the case of the nucleobases is the rate of internal conversion, compared to other possible processes. Internal conversion returns the molecule to the electronic ground state in a radiationless way, and thus converts electronic energy to heat in the form of ground state vibrational

energy. Heat can subsequently be safely dissipated to the environment. When internal conversion is fast enough to prevent other photochemical reactions from taking place, the molecule will have a very short excited state lifetime, τ^* , and be stable against UV photodamage⁹⁻¹².

Figure 1: Compilation of gas phase excited state lifetimes of selected nucleobases and derivatives.

Rapid internal conversion, when available to a molecule, diffuses electronic energy by converting it to internal energy in the ground state and therefore minimizes access to other photochemical pathways. Generally the biologically most relevant forms of the purines and pyrimidines exhibit the shortest excited state lifetimes, thus selectively minimizing the chances for photochemical damage in the molecular building blocks of life. In stark contrast, many other nucleobase derivatives, even isomers, have orders of magnitude longer lifetimes (See Figure 1). In fact, ultrafast internal conversion — the same property found in many UV sunscreens — is observed for all nucleobases implicated in replication today and, intriguingly, there are no examples of canonical nucleobases that are highly fluorescent¹³. Strengthening the correlation between excited state lifetime and the propensity for photoreaction is the observation that the minor DNA base 5-methylcytosine, with its 10-fold longer lifetime than cytosine, is a hot spot for photodamage¹⁴.

Evolution requires the existence of self-replicating molecules; the selection of the nucleobases as building blocks of those macromolecules would logically have taken place prior to any biological processes. Therefore, if the excited state properties have played a role in a chemical selection of today's nucleobases, they may be a relic of prebiotic chemistry on an early earth.

The intramolecular mechanism governing the ultrafast internal conversion in single purines and pyrimidines is now emerging. The key is the occurrence of conical intersections that connect the excited state potential energy surface, reached by photon absorption, to the ground state energy surface. The dramatic lifetime differences between derivatives of nucleobases appear to be due to variations in the excited state potential surfaces that restrict or slow access to these conical intersections¹⁵.

The excited state potential energy surface, and thus the corresponding electronic absorption and dynamics, is affected by molecular structure and by interactions, such as those with solvent molecules. To disentangle the different effects it is desirable to study isolated molecules, which is possible in the gas phase under collision free conditions. First, this makes it possible to observe intrinsic properties and explore the potential energy

surfaces of the individual molecules. Secondly, by employing double resonant techniques, it is possible to perform isomer specific spectroscopy. Some details of how interactions affect excited states can be studied by cluster spectroscopy in the gas phase. Once intrinsic properties are mapped out, these may be extrapolated to bulk conditions, which are the subject of later chapters.

2.2 Vibronic spectra

The nucleobases are heterocyclic aromatic compounds which absorb to their first excited electronic state in the 31,000-37,000 cm^{-1} range. This range also forms the onset of absorption in the liquid phase. Figure 2 shows resonance enhanced multiphoton ionization (REMPI) spectra of all of the canonical bases and some derivatives, as well as some of their clusters, representing base pair combinations. Asterisks indicate origins of different tautomers for the monomers and different hydrogen bonded structures for the clusters. Table 1 lists the respective $S_1 \leftarrow S_0$ origins.

Levy and coworkers reported the first REMPI spectra for uracil and thymine in 1988 ¹⁶. The spectra were very broad, merely showing an onset of unstructured absorption around 36,000 cm^{-1} for T and somewhat higher energy for U. These results suggested the absence of useful resolved spectroscopy for nucleobases. However that picture changed with the first report of the resolved guanine REMPI spectrum ¹⁷, followed by those of adenine ¹⁸ and cytosine ¹⁹. Observation of these spectra became possible with the employment of laser desorption jet cooling techniques, optimized for the study of these types of molecules ²⁰⁻²³.

The vapor pressure of the nucleobases at room temperature is too low for gas phase spectroscopy and heating to obtain sufficient vapor pressure causes thermal degradation. With some care, adenine can still be sufficiently heated in an oven to form a seeded supersonic beam, but this does not work satisfactorily for the other bases. Pulsed heating at a rate of the order of 1000 degrees in 10 ns allows fragment-free vaporization of all bases and this heating rate can be achieved with pulsed laser desorption. Typical experiments use a Nd:YAG laser at 1064 nm with 10 ns pulses of a fraction of a mJ/cm^2 to desorb from a graphite substrate, which is moved to expose fresh material in subsequent shots ^{21, 24, 25}. To permit spectroscopy, the desorbed molecules are entrained in a pulsed supersonic expansion, cooling them to internal temperatures corresponding to the order of 20 K ²¹. Such temperatures are low enough for the vibronic spectroscopy described in this chapter, but it may be noted that ions in a cold trap can be cooled to lower temperatures by about an order of magnitude and entrainment in helium droplets leads to even lower temperatures ^{26, 27}.

Figure 2: REMPI spectra of the canonical bases and their base pairs. Asterisks indicate origins of different tautomeric or structural isomers.

	Compound	origin cm ⁻¹	remarks	referen ces
nucleobases	guanine	32 864	7H enol <i>syn</i>	28
		33 269	oxo-imino	28
		33 910	oxo-imino	28
		34 755	N9H hydroxyl-amino	28
	adenine	36 105	π - π^*	29, 30
		36 062	n- π^*	29, 30
	cytosine	31 826	keto	31
		~36 000	enol	31
	thymine	36 000	broad onset	16
	uracil	36 700	broad onset	16
nucleosides	guanosine	34 443	enol	32
	2' deoxyguanosine	34 436	enol	32
	3' deoxyguanosine	34 443	enol	32
base-pairs	G-C	33314	C in enol form	33
		~32 800	doubly H-bonded	34
	A-T	35 064		35
	G-G	33 103, 33 282		36
	A-A	35 040		29, 37
	C-C	33 483		31
derivatives	1-methylcytosine	31 908		31
	5-methylcytosine	31 269		31
	5-OH-uracil	34 440		38
	5-NH ₂ -uracil	31 939		38
	9-methylguanine	34 612		36
	2,4-diaminopyrimidine	34 459	diamino	39
	2,6-diaminopurine	34 881 32 215	9H diamino 7H diamino	39
derivative pairs	9-mG-1mC	33 000	broad	34
	C-1mC	33 419		31
	C-5mC	32 500, 32 691, 32 916		31
	5mC-5mC	32 493, 32691, 32 872		31

Table 1: Observed S₁ band origins of nucleobases, nucleosides, base-pairs, and assorted nucleobase derivatives. M (as in 1mC) designates methyl.

Spectroscopic techniques

In molecular beams, direct absorption is generally not measurable because the optical density is too low. Intracavity techniques can sometimes be used, such as cavity ringdown⁴⁰ and microwave spectroscopy⁴¹, but the common approach is a form of action spectroscopy, especially laser induced fluorescence (LIF) and resonant two photon ionization (REMPI). A consequence of this technical limitation is that the excited state becomes part of the detection method and thus the dynamics of the excited state can influence the outcome of the measurement. In particular, both LIF and REMPI are blind for transitions to an excited state with a lifetime significantly shorter than the laser pulse duration. The following sections will discuss some implications of this limitation for understanding excited state behavior.

A major advantage of gas phase spectroscopy is the ability to perform isomer specific measurements by double resonant, or hole-burning techniques. Figure 3 schematically outlines this approach for the case of REMPI. Purple arrows indicate wavelengths that are scanned to obtain a wavelength dependent spectrum. In (a) this produces a vibronic REMPI spectrum. In (b) the blue arrow represents the probe laser, set to a single resonant wavelength in the REMPI spectrum. Its ion signal comes from one specific isomer. The purple arrow represents the burn laser, which precedes the probe laser by typically 100 ns. Whenever the burn laser is resonant with a transition it depletes the ground state and alters the Franck-Condon area, resulting in a dip in the ion signal from the probe laser, but only if both lasers interact with the same isomer population. Therefore a scan with the probe laser produces an ion dip spectrum that amounts to an isomer selected rovibronic spectrum. In (c) the same procedure is followed with an IR laser as probe laser to generate an isomer selected ground state IR spectrum. In Figure 2 asterisks denote origins of different isomers, identified by double resonant spectroscopy.

Figure 3: Schematic diagram of spectroscopic techniques. Purple arrow indicates wavelength that is being scanned to obtain a spectrum. Blue arrows indicate fixed wavelength. Green arrows indicate time delay between two laser pulses. (a) REMPI provides vibronic spectra. (b) UV-UV hole burning separates isomeric contributions to the UV spectrum. (c) IR-UV hole burning provides isomer selected ground state IR spectra.

Electronic spectra essentially contain information about the energy difference between the ground state and the excited state. By reducing the ground state vibrational population to $v=0$ through cooling, vibronic spectral structure can provide additional information on the excited state. IR-UV hole burning on the other hand provides ground state vibrational frequencies which can be used to identify the starting structures of the excitation, especially tautomeric form and cluster structure. Further information can be derived from line contours for identifying excited state symmetries and lifetimes.

Figure 4: Detail of the $34,720\text{ cm}^{-1}$ peak in the 2,4-DAPy REMPI spectrum, compared with simulations of the rotational envelope at three different Lorentzian linewidths ³⁹.

The latter is demonstrated in Figure 4, showing the comparison of a peak in the REMPI spectrum of 2,4-diaminopyrimidine with simulated lineshapes based on different Lorentzian linewidths ³⁹. The upper limit to the linewidth of 0.5 cm^{-1} thus observed in this case corresponds to a lower limit in the excited state lifetime of 10 ps.

Further information about the excited state dynamics may be derived by comparing these electronic spectra, for example by comparing frequency shifts between various isomers and derivatives and by contrasting broad versus sharp spectra. Furthermore, since all of these REMPI spectra were obtained with nanosecond pulses, absences of signal from selected species may point to ultrafast internal conversion.

2.3 Isomers

The origins on the UV spectra, obtained by REMPI, reflect the energies of the S_1 excited state relative to S_0 ground state, which is difficult to obtain computationally with any reasonable accuracy. Figure 2 shows that both the purine and pyrimidine bases absorb in the same frequency range, but significant shifts exist between different related compounds. For example, adenine absorbs 3000 cm^{-1} to the blue of the lowest energy observed tautomer of guanine ^{39, 42}. Adenine, which is 6-aminopurine (6AP), also absorbs about 3500 cm^{-1} to the blue of its isomer 2-aminopurine (2AP). Comparing those two isomers, one notices not only the large difference in origin, but also the fact that the adenine spectrum features only a few lines while the 2-aminopurine spectrum is very extensive and covers a large frequency range ⁴³. These observations are consistent with a model that is

summarized schematically in Figure 5 ⁴². This model assumes excitation to an S_1 state, which interacts with a second excited state through a curve crossing. This picture represents potential curves as a function of a single internuclear coordinate. In the more general case of multidimensional potential surfaces, to be discussed below, the curve crossing will be replaced by a conical intersection. The second excited state may couple with the electronic ground state to quench fluorescence and phosphorescence. In the case of the purines, the S_1 state is likely to be of π - π^* character and the second excited state is likely to be of n - π^* character, to be discussed below. The absorption characteristics of the purines and their excited state lifetimes then depend strongly on two parameters: (i) the relative energies of these two states and (ii) the amount of electronic coupling between them.

Possible arrangements are shown schematically in Figure 5. A curve crossing, as sketched schematically for 6AP, will typically lead to spectra that are diffuse or show a cutoff at absorption energies larger than that of the crossing. If, on the other hand, as sketched for 2AP, the interaction between the two excited states is weak, or if the S_1 state is significantly lower in energy, then we may expect a sharp and extended vibronic spectrum. A lower energy S_1 state is also consistent with the large observed red-shift. Finally, the other extreme is sketched in the inset: If the interaction is strong and the S_1 state is higher in energy, then it is possible to have a barrierless curve crossing, resulting in a completely structureless vibronic spectrum and an extremely short excited state lifetime.

Within this picture the striking photochemical difference of purine and adenine as compared to 2-aminopurines is the result of the relative energies of the π - π^* and the n - π^* states. The fact that the origin for the 2AP spectrum is significantly to the red of the adenine spectrum, implies that the S_1 state is lower in this case. Furthermore, the extent of the spectrum suggest that 2AP does *not* exhibit a curve crossing as sketched in Figure 5 2b, as opposed to adenine. Therefore the gas-phase spectra suggest that the amino substitution in the 2 position of purine lowers the excited state potential of the π - π^* state, while substitution in the 6 position has little effect. This is fully consistent with solution results, as well as with circular dichroism measurements by Holmen et al. which identify a π - π^* state about 4000 cm^{-1} below the n - π^* state for 2AP ⁴⁴.

Figure 5: Left along vertical axis, REMPI spectra of 6-aminopurine (adenine) in red and 2-aminopurine in blue. Right, schematic potential energy diagrams of the excited state as a function one internuclear coordinate, representing a model with a curve crossing between S_1 and another excited state. The inset shows the case without a barrier, which would not produce a discrete spectrum; see text for details.

2.4 Derivatives

When we consider more than one internuclear coordinate we can describe the more general situation as illustrated schematically in Figure 6. The crossings of multidimensional potential surfaces are conical intersections. These features can only occur in regions of the potential energy landscape that represent a deformation of the molecular frame from the ground state equilibrium geometry. The following case study of adenine derivatives serves as an example^{18, 45-57}. For 4-aminopyrimidine (Figure 7 top left), surface hopping calculations identified two conical intersections⁵⁸: deformation at the C2 position (red circle) leads to deactivation of the excited state with a lifetime, τ^* , of 1 ps and deformation at the C5=C6 bond (blue rectangle) leads to deactivation with τ^* of 400 fs. Immobilizing the latter with a 5 membered ring forms adenine with a single conical intersection due to the C2 deformation and τ^* of 1 ps. Figure 7 4 (bottom left) shows the geometry at this conical

intersection as calculated by Marian et al.⁵⁹. Substitution at the C2 position further modifies the excited state potential and eliminates this conical intersection. Consequently, both 2,6-diaminopurine and 2-aminopurine have fluorescent excited states with lifetimes of the order of nanoseconds. In fact, the latter is used as a fluorescent tag in DNA research. In the same way, C5 substituents in pyrimidines alter excited state lifetimes over a range of picoseconds to nanoseconds, by modification of the topography of the potential energy surfaces around C5=C6 torsion and stretching coordinates⁶⁰⁻⁶². Interestingly, the same coordinates are found to play a role in thymine photo-dimerization in DNA⁶³. Other excited states also need to be considered: N9-H motion forms an additional coordinate in adenine that can lead to a conical intersection with a $\pi\sigma^*$ state^{29, 55}. A weakly absorbing $n\pi^*$ state plays a role as well, in addition to the $\pi\pi^*$ state, and can sometimes serve as a “dark state” with an extended lifetime^{30, 64}.

Furthermore, purine, the parent compound of adenine as well as guanine, forms triplet states in high yield⁶⁵. Lifetimes also depend strongly on tautomeric forms, as discussed below in section 2.6^{28, 66-70}. Understanding these mechanisms helps explain how subtle structural differences between substituted nucleobases can produce excited state lifetime differences of orders of magnitude.

A similar strong effect can be observed in uracil (U) derivatives upon substitution in the C5 position³⁸. Two-photon ionization and IR/UV double-resonant spectra of the uracil analogues, 5-OH-U and 5-NH₂-U show that there is only one tautomer present for each with an excited state lifetime of 1.8 ns for 5-OH-Ura and 12.0 ns for 5-NH₂-Ura as determined from pump-probe experiments. These lifetimes are 3 and 4 orders of magnitude longer, respectively, than that for unsubstituted uracil⁷¹. Nachtigallova et al. determined vertical excitation energies, excited state minima, minima on the crossing seam and reaction paths

towards them by means of multi-reference ab initio methods ³⁸. They found sizeable barriers on these paths that provide an explanation for the lifetimes of several nanoseconds observed in the experiment.

Kistler and Matsika reported a similar effect for cytosine derivatives as well ^{61, 72}. For example, 5-methyl-2-pyrimidin-(1*H*)-one (5M2P, with a methyl substituent in the C5 position instead of the amino group in the C4 position) has a fluorescence lifetime that is orders of magnitude longer than the picosecond lifetime of cytosine ^{5, 64, 71, 73}. This excited state lifetime difference correlates directly with a lower S_1 excitation energy for 5M2P, analogous to the energetic considerations for adenine vs. 2-aminopurine.

2.5 Character of the excited state

The character of the first excited states is not always entirely clear but generally the UV spectra contain contributions to two excited states, of $\pi\pi^*$ and $n\pi^*$ character respectively ^{30, 74}. The $\pi\pi^*$ transition carries most oscillator strength and the $n\pi^*$ state can function as a dark state for internal conversion, depending on whether the relative energies lead to the occurrence of conical intersections. In the case of adenine, Lee et al. characterized the excited states based on rotational band contour analyses ³⁰. They showed that the lowest $n\pi^*$ and $\pi\pi^*$ states can be labeled with their excited-state vibronic symmetry, and exhibit a strong $\pi\pi^*$ - $n\pi^*$ vibronic coupling via an out-of-plane vibrational mode. They assigned the band at 36,062 cm⁻¹ as the $n\pi^*$ transition, and the 36,105 cm⁻¹ band as the $\pi\pi^*$ transition by symmetry analysis. The band at 36,248 cm⁻¹ provides evidence of the strong $\pi\pi^*$ - $n\pi^*$ vibronic coupling via an out-of-plane vibrational mode.

Sobolewski and Domcke showed that in the excited state dynamics a $\pi\sigma^*$ can also play a role along a NH nuclear coordinate ⁷⁵. Motion along this coordinate can lead to hydrogen loss and Hunig et al. demonstrated this channel by detecting the dissociating hydrogen using Lyman- α excitation and analyzing the kinetic energy from the Doppler shift ⁵⁵.

2.6 Tautomers

A complication in the study of nucleobase structure and dynamics is the fact that these compounds exhibit multiple tautomeric forms. The electronic spectra, such as those in Figure 2, may contain contributions of all the tautomers present in the beam.

As shown in Figure 2, the $S_1 \leftarrow S_0$ transitions of the two most stable cytosine tautomers, keto and enol, are a remarkable 4000 cm⁻¹ apart, while energetically they differ by only 0.03 eV ^{19, 31}. The third major tautomer, keto-imine, is significantly higher in energy ⁷⁶⁻⁷⁸ and not observed in these spectra. The keto form is the biologically important one, with Watson-Crick base pairing in DNA, and predominant in solution. In matrix isolation Szczesniak et al. observed both keto and enol forms with higher abundances for the latter and small contributions from the imino form ⁷⁹. Brown et al. have obtained rotational

constants for all three tautomeric forms by microwave spectroscopy ⁸⁰. Schiedt et al. also identified the existence of keto and enol tautomers of neutral cytosine in the gas phase by anion spectroscopy ⁸¹.

Kosma et al. reported excited state dynamics by probing with different excitation wavelengths in a range from 260 nm to 290 nm, which allowed them to distinguish between keto only versus mixed populations ⁸². They conclude that the deactivation pathways are quite different for the different tautomers, with three time constants for keto-cytosine and two for the others. Theoretical treatments in the literature had focused on modeling the latter, while the new results point to the importance of tautomeric distinction in these analyses. For the keto form the two faster channels, with femtosecond and picosecond lifetimes are consistent with fast, but not barrierless excited-state relaxation via conical intersections between the $1\pi\pi^*$ and ground state ⁷⁶. The third channel with a lifetime of hundreds of picoseconds at 290 nm is proposed to involve excited-state tautomerization to a low-lying $1n\pi^*$ state of the keto-imino tautomer ⁸³. The authors propose that in the condensed phase this channel would be quenched by rapid back hydrogen-transfer, catalyzed by hydrogen-bound water molecules ^{84, 85}. Consequently this pathway would not alter the inherent photostability.

Figure 8: The major tautomers of cytosine. The IR-UV double resonant experiments observe the keto and enol forms (yellow) but do not distinguish between the two enol tautomers.

In most cases electronic spectra of different tautomers do not exhibit significant shifts and thus vibronic spectra potentially consist of overlapping contributions from a number of tautomers. Such overlapping electronic spectra can be disentangled by IR-UV double

resonant spectroscopy. However, an additional complication with this technique is that it is based on action spectroscopy in which the probe signal is multiphoton ionization. As a result this approach is blind for tautomers for which the excited state life time is significantly shorter than the laser pulse. Such tautomers do not show up in the electronic spectrum to begin with.

Figure 9: Ten of the lowest energy tautomers of guanine. Relative energies in wavenumbers from reference ⁶⁷. Yellow structures are observed in gas phase nanosecond REMPI experiments. Red structures have sub picosecond excited state lifetimes. Structures 1-4 are observed in helium droplets. Keto-N9H is the Watson-Crick structure.

A case in point is the so-called guanine puzzle ^{28, 69}. The REMPI vibronic spectrum consists of the overlapping spectra from four different tautomers, as determined by UV-UV holeburning ^{24, 86, 87}. However the identification of the specific tautomers by IR-UV double resonance spectroscopy proved challenging. Two enol tautomers could easily be assigned based on the strong OH stretch around 3600 cm⁻¹. However the other two species were initially incorrectly assigned as keto tautomers. Figure 9 shows the ten lowest energy tautomers of guanine with relative energies in wavenumbers from reference ⁶⁷. Clearly the two keto tautomers (N7H and N9H) are the lowest energy forms and their IR spectra formed a quite reasonable match with calculated frequencies for these forms, making this a plausible assignment. However, subsequent data from a helium droplet experiment exhibited slightly different IR spectra that formed a better match with the keto forms ²⁷. This led to a re-analysis of the IR-UV data and a reassignment of the two tautomers from the keto to the imino form. This assignment was subsequently confirmed by IR-UV measurements in the imino stretch region below 2000 cm⁻¹ ⁶⁷. Gas phase microwave experiments also found the keto tautomers ⁴¹.

These observations are summarized in Figure 9 in which the red tautomers were observed by the two techniques that involve direct absorption in the ground state (helium droplets and microwave), while the yellow tautomers were observed by the technique that involves action spectroscopy via the electronic excited state (REMPI). This difference suggests that the lowest energy forms are most abundant in the gas phase in all cases, but are not

observed with REMPI because they have short lived excited states. This explanation is confirmed by computations, showing a rapid internal conversion pathway through a conical intersection, which is not accessible for the higher energy enol and imino tautomers⁶⁹.

The tautomeric landscape of adenine is somewhat less varied than in the case of guanine because of the absence of the oxygen. Plützer and Kleinerermanns reported IR-UV double resonance spectroscopy and observed two tautomers⁵². Both tautomers are of the amino form, with the 9H form most abundant and a small presence of the 7H form. This finding is consistent with microwave measurements by Brown et al.⁸⁸. At the conditions of jet cooling the imino form appears to be absent, although in the gas phase at elevated temperature the IR spectra seem to comprise multiple tautomers, including imino.

Microwave measurements of uracil in a heated cell suggest the di-keto form as the most abundant. Brown et al. reported to the first microwave measurements in a seeded molecular beam and also concluded that the di-keto form was predominant^{89,90}. Viant et al. reported to the first rotationally resolved gas phase IR spectra of uracil⁹¹. This work employed a slit nozzle, an IR diode laser, and a multi-pass arrangement to obtain high-resolution IR absorption spectra of the out of phase ν_6 stretching vibration. The rotational analysis unambiguously assigned this species to the diketo tautomer. Brown et al. also observed the diketo form of thymine in a seeded molecular beam, based on hyperfine structure⁹².

2.7. Intermolecular effects – clusters

It is often assumed that the same internal conversion processes govern the subpicosecond de-excitation timescales of nucleobases in solution, albeit modified by solvent interactions. It is reasonable to postulate that the chromophore is affected primarily by the first solvent shell. Therefore many of the details of the dynamics in solution can be elucidated by studying clusters with small numbers of water molecules^{24, 25, 93-98}. It appears that hydrogen bonding can play a role in two ways. First the excited state potential energy surfaces will be modified, thus affecting possible trajectories through conical intersections. Secondly the hydrogen bond itself can provide another coordinate for coupling excited states and the ground state. This effect is not limited to hydrogen bonding with water, but may also play a role in base pairing and in intramolecular hydrogen bonds. This coordinate can be thought of as an analogue of the N-H stretch coordinate correlated with a $\pi\sigma^*$ state, and its associated conical intersections^{75, 99-102}. In the latter case, if the $\pi\sigma^*$ potential surface fails to reconnect with S_0 , the hydrogen can actually dissociate. In the hydrogen bonded case the hydrogen is confined to the system and this coordinate can only mediate a single or double proton transfer.

Figure 10: The lowest energy structures of hydrogen bonded GC clusters with energies in kcal/mol. Yellow structures were observed in REMPI experiments, while red structures were not.

An example is the finding that the specific Watson-Crick (WC) structure in isolated GC nucleobase pairs has a short excited state lifetime that is explained by proton motion in the middle hydrogen bond ^{34, 103-105}. By contrast, in other GC base pair structures this coordinate does not lead to accessible conical intersections ^{34, 106}. Figure 10 shows the eight lowest energy structures of hydrogen bonded GC clusters, with the very lowest energy structure representing the WC structure. The two lowest energy structures, marked in red, were not directly observed in gas phase experiments with nanosecond timescale REMPI detection, while the next two higher energy structures, marked in yellow, were observed and identified by IR-UV hole burning ^{33, 34, 106}.

For AT base pairs the Watson-Crick structure was also not identified, however in that case it is not the lowest energy structure so it is possible that it was insufficiently populated in the molecular beam ³⁵. In REMPI spectra of homo base pair combinations the predicted lowest energy structure was not observed. This absence occurred for GG dimers ¹⁰⁷, CC dimers ³¹, and AA dimers ³⁷. In these homodimers the missing structure is symmetric so it is possible that an excimer state is formed with a considerable splitting causing a shift in the UV band outside of the experimental range. However, it is also possible that the absence of those cluster structures in the REMPI spectra is due to short lived excited states, that are dark in nanosecond timescale REMPI detection. For 9-methyladenine-adenine clusters Plützer et al. identified a stacked structure, which they detected on the mass of protonated adenine. They interpreted their finding as involving hydrogen transfer from the 9H position of the adenine. Schultz and coworkers reported indications of this process in time resolved photoelectron spectroscopy studies of adenine and adenine containing clusters ^{95, 108-110} and in nucleobase pair analogues ^{103, 111, 112}.

Another structure in which hydrogen bonding occurs and possibly affects dynamics is in nucleosides in the *syn* configuration. Nir et al. found by double resonant spectroscopy that isolated guanosine indeed forms an internal hydrogen bond between 5'OH and 3N¹¹³. Nevertheless they found a sharp REMPI spectrum implying a relatively long lived excited state. By contrast, an REMPI spectrum for adenosine has not been observed, suggesting a possibly short excited state lifetime^{114, 115}. It should be noted that for guanosine in the gas phase only the enol tautomer has been observed, so far, which suggests that the keto tautomer is selectively short lived, analogously to the nucleobase. So whether and to what extent the shortened lifetime observed in solution is an intrinsic property of nucleosides remains an open question. Other aspects of the role of the sugar moiety in excited state dynamics may also play a role. It has been proposed that the additional degrees of freedom provide a bath for quenching excited state excitation. Furthermore, in solution the dynamics may be further complicated because the sugar also provides additional hydration sites.

Another intermolecular interaction is base stacking, which provides much of the stability in helices. In the π stacked structural motif, exciplex formation appears to play a major role in the photodynamics^{115, 116}. The overlap of the π clouds depends critically on the relative position of the molecules. For example, Schreier et al. have shown that thymine dimer formation in DNA is determined largely by conformational properties at the instant of excitation¹¹⁷. In solution phase experiments Crespo-Hernández and Kohler and coworkers have shown that base stacking in single- and double-stranded DNA favors non-radiative decay via the formation of charge-transfer or exciplex states¹¹⁸⁻¹²⁰. Such states decay much more slowly than the $\pi\pi^*$ states in single bases (see Figure 11). These considerations are the topic of the next chapter.

Summary

Electronic and vibrational spectroscopy of isolated nucleobases sheds light on excited state dynamics indirectly in a number of ways. UV-VIS excitation spectroscopy probes the potential energy landscape in the Franck-Condon region, which establishes the starting conditions of any ensuing photochemistry. The combination with IR hole burning to obtain vibrational spectroscopy as well, offers the important opportunity to obtain isomer selected data. This approach makes it possible to probe individual tautomers, structural conformers, isomers, and cluster structures. This isomer selectivity as well as the fact that nucleobases can be studied free of interactions provides unique insights in their intrinsic properties, unperturbed by any secondary effects of the biological environment. Therefore such gas phase experiments offer the best data for comparison with computational chemistry at the highest levels. The frequency domain is coupled with the time domain by the inverse relationship between energy and line width. Particularly in the 10-100 ps range excited state lifetimes may be derived from spectra line widths. Shorter lifetimes not only lead to extensive broadening but often render excited states unobservable by action spectroscopy, such as REMPI or LIF, with nanosecond timescale laser pulses. Possibly future experiments in the frequency domain will provide additional insight in excited state dynamics. Examples may include spectroscopy with shorter laser pulses, including a trade-off between time and frequency resolution, spectroscopic characterization of the hot ground state products of internal conversion, and more elaborate studies of clusters with water to detail the role of micro hydration.

Acknowledgement

This material is based upon work supported by the National Science Foundation under CHE-1301305 and by NASA under Grant No. NNX12AG77G.

TOC GRAPHIC:

Following electronic excitation, probed in the frequency domain (blue), internal conversion (red) can compete with fluorescence (green) and other photochemical pathways (black).

References

1. Middleton CT, de La Harpe K, Su C, Law YK, Crespo-Hernandez CE, and Kohler B: *DNA Excited-State Dynamics: From Single Bases to the Double Helix*. Annu. Rev. Phys. Chem., **60**:217-239 (2009).
2. Kleineremanns K, Nachtigallova D, and de Vries MS: *Excited state dynamics of DNA bases*. Int. Rev. Phys. Chem., **32**(2):308-342 (2013).
3. Gonzalez-Ramirez I, Roca-Sanjuan D, Climent T, Serrano-Perez JJ, Merchan M, and Serrano-Andres L: *On the photoproduction of DNA/RNA cyclobutane pyrimidine dimers*. Theor. Chem. Acc., **128**(4-6):705-711 (2011).
4. Serrano-Andres L and Merchan M: *Are the five natural DNA/RNA base monomers a good choice from natural selection? A photochemical perspective*. J Photoch Photobio C, **10**(1):21-32 (2009).
5. Crespo-Hernandez CE, Cohen B, Hare PM, and Kohler B: *Ultrafast excited-state dynamics in nucleic acids*. Chem. Rev., **104**(4):1977-2019 (2004).
6. Plasser F, Barbatti M, Aquino AJA, and Lischka H: *Electronically excited states and photodynamics: a continuing challenge*. Theor. Chem. Acc., **131**(1) (2012).
7. de Vries MS and Hobza P: *Gas-phase spectroscopy of biomolecular building blocks*. Annu. Rev. Phys. Chem., **58**:585-612 (2007).
8. Matsika S: *Radiationless decay of excited states of uracil through conical intersections*. J. Phys. Chem. A, **108**(37):7584-7590 (2004).
9. Otterstedt J-EA: *Photostability and Molecular Structure*. J. Chem. Phys., **58**:5716-5725 (1973).
10. Zgierski MZ, Fujiwara T, and Lim EC: *Conical intersections and ultrafast intramolecular excited-state dynamics in nucleic acid bases and electron donor-acceptor molecules*. Chem. Phys. Lett., **463**(4-6):289-299 (2008).
11. Zgierski MZ, Fujiwara T, Kofron WG, and Lim EC: *Highly effective quenching of the ultrafast radiationless decay of photoexcited pyrimidine bases by covalent modification: photophysics of 5,6-trimethylenecytosine and 5,6-trimethylenauracil*. PCCP, **9**(25):3206-3209 (2007).
12. Zgierski MZ, Patchkovskii S, Fujiwara T, and Lim EC: *On the origin of the ultrafast internal conversion of electronically excited pyrimidine bases*. J. Phys. Chem. A, **109**(42):9384-9387 (2005).
13. Middleton CT, de La Harpe K, Su C, Law YK, Crespo-Hernández CE, and Kohler B: *DNA Excited-State Dynamics: From Single Bases to the Double Helix*. Annu. Rev. Phys. Chem., **60**:217-239 (2009).
14. Tommasi S, Denissenko MF, and Pfeifer GP: *Sunlight induces pyrimidine dimers preferentially at 5-methylcytosine bases*. Cancer Res., **57**(21):4727-4730 (1997).
15. Malone RJ, Miller AM, and Kohler B: *Singlet Excited-state Lifetimes of Cytosine Derivatives Measured by Femtosecond Transient Absorption*. Photochem. Photobiol., **77**(2):158-164 (2003).
16. Brady BB, Peteanu LA, and Levy DH: *The Electronic Spectra of the Pyrimidine Bases Uracil and Thymine in a Supersonic Molecular Beam*. Chem. Phys. Lett., **147**:538-543 (1988).
17. Nir E, Grace L, Brauer B, and de Vries MS: *REMPI spectroscopy of jet-cooled guanine*. J. Am. Chem. Soc., **121**(20):4896-4897 (1999).
18. Plützer C, Nir E, de Vries MS, and Kleineremanns K: *IR-UV double-resonance spectroscopy of the nucleobase adenine*. PCCP, **3**(24):5466-5469 (2001).
19. Nir E, Muller M, Grace LI, and de Vries MS: *REMPI spectroscopy of cytosine*. Chem. Phys. Lett., **355**(1-2):59-64 (2002).
20. Arrowsmith P, de Vries MS, Hunziker HE, and Wendt HR: *Pulsed laser desorption near a jet orifice: concentration profiles of entrained perylene vapor*. Appl. Phys. B, **46**:165-173 (1988).

21. Meijer G, de Vries MS, Hunziker HE, and Wendt HR: *Laser desorption jet-cooling of organic molecules. Cooling characteristics and detection sensitivity*. Appl. Phys. B, **51**:395-403 (1990).
22. Tembreull R and Lubman DM: *Resonant two-photon ionization in biomolecules using laser desorption in supersonic beam-mass spectrometry*. Appl. Spectrosc., **41**:431-436 (1987).
23. Boesl U, Grotemeyer J, Walter K, and Schlag EW: *Resonance ionization and time-of-flight mass spectrometry, high resolution, involatile molecules*. Resonance Ionization Spectroscopy, **369**:223-228 (1986).
24. Piuzzi F, Mons M, Dimicoli I, Tardivel B, and Zhao Q: *Ultraviolet spectroscopy and tautomerism of the DNA base guanine and its hydrate formed in a supersonic jet*. Chem. Phys., **270**(1):205-214 (2001).
25. Saigusa H, Urashima S, and Asami H: *IR-UV Double Resonance Spectroscopy of the Hydrated Clusters of Guanosine and 9-Methylguanine: Evidence for Hydration Structures Involving the Sugar Group*. J. Phys. Chem. A, **113**(15):3455-3462 (2009).
26. Boyarkin OV, Mercier SR, Kamariotis A, and Rizzo TR: *Electronic spectroscopy of cold, protonated tryptophan and tyrosine*. J. Am. Chem. Soc., **128**(9):2816-2817 (2006).
27. Choi MY and Miller RE: *Four tautomers of isolated guanine from infrared laser spectroscopy in helium nanodroplets*. J. Am. Chem. Soc., **128**(22):7320-7328 (2006).
28. Mons M, Piuzzi F, Dimicoli I, Gorb L, and Leszczynski J: *Near-UV resonant two-photon ionization spectroscopy of gas phase guanine: Evidence for the observation of three rare tautomers*. J. Phys. Chem. A, **110**(38):10921-10924 (2006).
29. Kabelac M, Plützer C, Kleineremanns K, and Hobza P: *Isomer selective IR experiments and correlated ab initio quantum chemical calculations support planar H-bonded structure of the 7-methyl adenine - adenine and stacked structure of the 9-methyl adenine - adenine base pairs*. PCCP, **6**(10):2781-2785 (2004).
30. Lee Y, Schmitt M, Kleineremanns K, and Kim B: *Observation of ultraviolet rotational band contours of the DNA base adenine: Determination of the transition moment*. J. Phys. Chem. A, **110**(42):11819-11823 (2006).
31. Nir E, Hünig I, Kleineremanns K, and de Vries MS: *The nucleobase cytosine and the cytosine dimer investigated by double resonance laser spectroscopy and ab initio calculations*. PCCP, **5**(21):4780-4785 (2003).
32. Nir E, Imhof P, Kleineremanns K, and de Vries MS: *REMPI spectroscopy of laser desorbed guanosines*. J. Am. Chem. Soc., **122**(33):8091-8092 (2000).
33. Nir E, Janzen C, Imhof P, Kleineremanns K, and de Vries MS: *Pairing of the nucleobases guanine and cytosine in the gas phase studied by IR-UV double-resonance spectroscopy and ab initio calculations*. PCCP, **4**(5):732-739 (2002).
34. Abo-Riziq A, Grace L, Nir E, Kabelac M, Hobza P, and de Vries MS: *Photochemical selectivity in guanine-cytosine base-pair structures*. Proceedings of the National Academy of Sciences of the United States of America, **102**(1):20-23 (2005).
35. Plützer C, Hünig I, Kleineremanns K, Nir E, and de Vries MS: *Pairing of isolated nucleobases: Double resonance laser spectroscopy of adenine-thymine*. ChemPhysChem, **4**(8):838-842 (2003).
36. Nir E, Plützer C, Kleineremanns K, and de Vries M: *Properties of isolated DNA bases, base pairs and nucleosides examined by laser spectroscopy*. European Physical Journal D, **20**(3):317-329 (2002).
37. Plützer C, Hünig I, and Kleineremanns K: *Pairing of the nucleobase adenine studied by IR-UV double-resonance spectroscopy and ab initio calculations*. PCCP, **5**(6):1158-1163 (2003).
38. Nachtigallova D, Lischka H, Szymczak JJ, Barbatti M, Hobza P, Gengeliczki Z, Pino G, Callahan MP, and de Vries MS: *The effect of C5 substitution on the photochemistry of uracil*. PCCP, **12**(19):4924-4933 (2010).

39. Gengeliczki Z, Callahan MP, Svadlenak N, Pongor CI, Sztaray B, Meerts L, Nachtigallova D, Hobza P, Barbatti M, Lischka H, and de Vries MS: *Effect of substituents on the excited-state dynamics of the modified DNA bases 2,4-diaminopyrimidine and 2,6-diaminopurine*. PCCP, **12**(20):5375-5388 (2010).
40. Chapo CJ, Paul JB, Provencal RA, Roth K, and Saykally RJ: *Is arginine zwitterionic or neutral in the gas phase? Results from IR cavity ringdown spectroscopy*. J. Am. Chem. Soc., **120**(49):12956-12957 (1998).
41. Alonso JL, Pena I, Lopez JC, and Vaquero V: *Rotational Spectral Signatures of Four Tautomers of Guanine*. Angew Chem Int Edit, **48**(33):6141-6143 (2009).
42. Nir E, Kleinermmanns K, Grace L, and de Vries MS: *On the photochemistry of purine nucleobases*. J. Phys. Chem. A, **105**(21):5106-5110 (2001).
43. Seefeld KA, Plützer C, Löwenich D, Häber T, Linder R, Kleinermmanns K, Tatchen J, and Marian CM: *Tautomers and electronic states of jet-cooled 2-aminopurine investigated by double resonance spectroscopy and theory*. PCCP, **7**(16):3021-3026 (2005).
44. Holmen A, Broo A, Albinsson B, and Norden B: *Assignment of electronic transition moment directions of adenine from linear dichroism measurements*. J. Am. Chem. Soc., **119**:12240-12250 (1997).
45. Broo A: *A theoretical investigation of the physical reason for the very different luminescence properties of the two isomers adenine and 2-aminopurine*. J. Phys. Chem. A, **102**:526-531 (1998).
46. Andreasson J, Holmén A, and Albinsson B: *The photophysical properties of the adenine chromophore*. J. Phys. Chem. B, **103**(44):9782-9789 (1999).
47. Kim NJ, Jeong G, Kim YS, Sung J, Kim SK, and Park YD: *Resonant two-photon ionization and laser induced fluorescence spectroscopy of jet-cooled adenine*. J. Chem. Phys., **113**(22):10051-10055 (2000).
48. Mishra SK, Shukla MK, and Mishra PC: *Electronic spectra of adenine and 2-aminopurine: an ab initio study of energy level diagrams of different tautomers in gas phase and aqueous solution*. Spectrochim. Acta, Part A, **56**(7):1355-1384 (2000).
49. Lühns DC, Viallon J, and Fischer I: *Excited state spectroscopy and dynamics of isolated adenine and 9-methyladenine*. Phys. Chem. Chem. Phys., **3**(10):1827-1831 (2001).
50. Kang H, Jung B, and Kim SK: *Mechanism for ultrafast internal conversion of adenine*. J. Chem. Phys., **118**(15):6717-6719 (2003).
51. Sobolewski AL and Domcke W: *On the mechanism of nonradiative decay of DNA bases: ab initio and TDDFT results for the excited states of 9H-adenine*. Eur. Phys. J. D, **20**:369-374 (2002).
52. Plützer C and Kleinermmanns K: *Tautomers and electronic states of jet-cooled adenine investigated by double resonance spectroscopy*. PCCP, **4**(20):4877-4882 (2002).
53. Barbatti M and Lischka H: *Can the Nonadiabatic Photodynamics of Aminopyrimidine Be a Model for the Ultrafast Deactivation of Adenine?* J. Phys. Chem. A, **111**(15):2852-2858 (2007).
54. Marian CM: *A new pathway for the rapid decay of electronically excited adenine*. J. Chem. Phys., **122**:104314 (2005).
55. Hünig I, Plützer C, Seefeld KA, Löwenich D, Nispel M, and Kleinermmanns K: *Photostability of isolated and paired nucleobases: N-H dissociation of adenine and hydrogen transfer in its base pairs examined by laser spectroscopy*. ChemPhysChem, **5**(9):1427-1431 (2004).
56. Crespo-Hernández CE, Cohen B, Hare PM, and Kohler B: *Ultrafast Excited-State Dynamics in Nucleic Acids*. Chem. Rev., **104**:1977-2019 (2004).
57. Zierhut M, Roth W, and Fischer I: *Dynamics of H-atom loss in adenine*. PCCP, **6**(22):5178-5183 (2004).
58. Zechmann G and Barbatti M: *Ab initio study of the photochemistry of aminopyrimidine*. Int. J. Quantum Chem., **108**(7):1266-1276 (2008).

59. Marian CM: *A new pathway for the rapid decay of electronically excited adenine*. J. Chem. Phys., **122**(10):10 (2005).
60. Kistler KA and Matsika S: *Three-state conical intersections in cytosine and pyrimidinone bases*. J. Chem. Phys., **128**(21):- (2008).
61. Kistler KA and Matsika S: *Cytosine in context: A theoretical study of substituent effects on the excitation energies of 2-pyrimidinone derivatives*. J. Phys. Chem. A, **111**(35):8708-8716 (2007).
62. Kistler KA and Matsika S: *Radiationless decay mechanism of cytosine: An ab initio study with comparisons to the fluorescent analogue 5-methyl-2-pyrimidinone*. J. Phys. Chem. A, **111**(14):2650-2661 (2007).
63. Yarasi S, Brost P, and Loppnow GR: *Initial excited-state structural dynamics of thymine are coincident with the expected photochemical dynamics*. J. Phys. Chem. A, **111**(24):5130-5135 (2007).
64. Canuel C, Mons M, Piuze F, Tardivel B, Dimicoli I, and Elhanine M: *Excited states dynamics of DNA and RNA bases: Characterization of a stepwise deactivation pathway in the gas phase*. J. Chem. Phys., **122**(7):074316-074317 (2005).
65. Quinones E and Arce R: *Photochemistry and photophysics of purine free base and 6-methylpurine*. J. Am. Chem. Soc., **111**(21):8218-8223 (1989).
66. Wilson RW and Callis PR: *Fluorescent tautomers and the apparent photophysics of adenine and guanine*. Photochem. Photobiol., **31**:323-327 (1980).
67. Seefeld K, Brause R, Häber T, and Kleinerhanns K: *Imino tautomers of gas-phase guanine from mid-infrared laser spectroscopy*. J. Phys. Chem. A, **111**(28):6217-6221 (2007).
68. Nachtigallova D, Hobza P, and Spirko V: *Assigning the NH stretches of the Guanine tautomers using adiabatic separation: CCSD(T) benchmark calculations*. J. Phys. Chem. A, **112**(9):1854-1856 (2008).
69. Marian CM: *The guanine tautomer puzzle: Quantum chemical investigation of ground and excited states*. J. Phys. Chem. A, **111**(8):1545-1553 (2007).
70. Cerny J, Spirko V, Mons M, Hobza P, and Nachtigallova D: *Theoretical study of the ground and excited states of 7-methyl guanine and 9-methyl guanine: comparison with experiment*. PCCP, **8**(26):3059-3065 (2006).
71. Kang H, Lee KT, Jung B, Ko YJ, and Kim SK: *Intrinsic lifetimes of the excited state of DNA and RNA bases*. J. Am. Chem. Soc., **124**(44):12958-12959 (2002).
72. Kistler KA and Matsika S: *The fluorescence mechanism of 5-methyl-2-pyrimidinone: An ab initio study of a fluorescent pyrimidine analog*. Photochem. Photobiol., **83**(3):611-624 (2007).
73. Ullrich S, Schultz T, Zgierski MZ, and Stolow A: *Electronic relaxation dynamics in DNA and RNA bases studied by time-resolved photoelectron spectroscopy*. PCCP, **6**(10):2796-2801 (2004).
74. Kim NJ, Jeong G, Kim YS, Sung J, Kim SK, and Park YD: *Resonant two-photon ionization and laser induced fluorescence spectroscopy of jet-cooled adenine*. J. Chem. Phys., **113**(22):10051-10055 (2000).
75. Perun S, Sobolewski AL, and Domcke W: *Ab initio studies on the radiationless decay mechanisms of the lowest excited singlet states of 9H-adenine*. J. Am. Chem. Soc., **127**(17):6257-6265 (2005).
76. Tomic K, Tatchen J, and Marian CM: *Quantum chemical investigation of the electronic spectra of the keto, enol, and keto-imine tautomers of cytosine*. J. Phys. Chem. A, **109**(37):8410-8418 (2005).
77. Trygubenko SA, Bogdan TV, Rueda M, Orozco M, Luque FJ, Sponer J, Slavicek P, and Hobza P: *Correlated ab initio study of nucleic acid bases and their tautomers in the gas phase, in a microhydrated environment and in aqueous solution - Part 1. Cytosine*. PCCP, **4**(17):4192-4203 (2002).

78. Kobayashi R: *A CCSD(T) study of the relative stabilities of cytosine tautomers*. J. Phys. Chem. A, **102**(52):10813-10817 (1998).
79. Szczesniak M, Szczepaniak K, Kwiatkowski JS, Kubulat K, and Person WB: *Matrix-isolation infrared studies of nucleic-acid constituents .5. Experimental matrix-isolation and theoretical abinitio scf molecular-orbital studies of the infrared-spectra of cytosine monomers*. J. Am. Chem. Soc., **110**:8319-8330 (1988).
80. Brown RD, Godfrey PD, McNaughton D, and Pierlot A: *Tautomers of Cytosine by Microwave Spectroscopy*. J. Am. Chem. Soc., **111**:2308-2310 (1989).
81. Schiedt J, Weinkauff R, Neumark DM, and Schlag EW: *Anion spectroscopy of uracil, thymine and the amino-oxo and amino-hydroxy tautomers of cytosine and their water clusters*. Chem. Phys., **239**(1-3):511-524 (1998).
82. Kosma K, Schroter C, Samoylova E, Hertel IV, and Schultz T: *Excited-State Dynamics of Cytosine Tautomers*. J. Am. Chem. Soc., **131**(46):16939-16943 (2009).
83. Gonzalez-Vazquez J and Gonzalez L: *A Time-Dependent Picture of the Ultrafast Deactivation of keto-Cytosine Including Three-State Conical Intersections*. ChemPhysChem, **11**(17):3617-3624 (2010).
84. Podolyan Y, Gorb L, and Leszczynski J: *Ab initio study of the prototropic tautomerism of cytosine and guanine and their contribution to spontaneous point mutations*. Int J Mol Sci, **4**(7):410-421 (2003).
85. Gorb L, Podolyan Y, and Leszczynski J: *A theoretical investigation of tautomeric equilibria and proton transfer in isolated and monohydrated cytosine and isocytosine molecules*. Journal of Molecular Structure-Theochem, **487**(1-2):47-55 (1999).
86. Chin W, Mons M, Dimicoli I, Piuze F, Tardivel B, and Elhanine M: *Tautomer contribution's to the near UV spectrum of guanine: towards a refined picture for the spectroscopy of purine molecules*. European Physical Journal D, **20**(3):347-355 (2002).
87. Nir E, Janzen C, Imhof P, Kleinermanns K, and de Vries MS: *Guanine tautomerism revealed by UV-UV and IR-UV hole burning spectroscopy*. J. Chem. Phys., **115**:4604-4611 (2001).
88. Brown RD, Godfrey PD, Mcnaughton D, and Pierlot AP: *A Study of the Major Gas-Phase Tautomer of Adenine by Microwave Spectroscopy*. Chem. Phys. Lett., **156**(1):61-63 (1989).
89. Brown RD, Godfrey PD, Mcnaughton D, and Pierlot AP: *Microwave-Spectrum of Uracil*. J. Am. Chem. Soc., **110**(7):2329-2330 (1988).
90. Vaquero V, Sanz ME, Lopez JC, and Alonso JL: *The structure of uracil: A laser ablation rotational study*. J. Phys. Chem. A, **111**(18):3443-3445 (2007).
91. Viant MR, Fellers RS, McLaughlin RP, and Saykally RJ: *Infrared-Laser Spectroscopy of Uracil in a Pulsed Slit Jet*. J. Chem. Phys., **103**(21):9502-9505 (1995).
92. Brown RD, Godfrey PD, Mcnaughton D, and Pierlot AP: *Microwave-Spectrum of the Major Gas-Phase Tautomer of Thymine*. J Chem Soc Chem Comm (1):37-38 (1989).
93. Chin W, Mons M, Piuze F, Tardivel B, Dimicoli I, Gorb L, and Leszczynski J: *Gas phase rotamers of the nucleobase 9-methylguanine enol and its monohydrate: Optical spectroscopy and quantum mechanical calculations*. J. Phys. Chem. A, **108**(40):8237-8243 (2004).
94. Abo-Riziq A, Crews B, Grace L, and de Vries MS: *Microhydration of guanine base pairs*. J. Am. Chem. Soc., **127**(8):2374-2375 (2005).
95. Smith VR, Samoylova E, Ritze HH, Radloff W, and Schultz T: *Excimer states in microhydrated adenine clusters*. PCCP, **12**(33):9632-9636 (2010).
96. Ritze HH, Lippert H, Samoylova E, Smith VR, Hertel IV, Radloff W, and Schultz T: *Relevance of pi sigma(*) states in the photoinduced processes of adenine, adenine dimer, and adenine-water complexes*. J. Chem. Phys., **122**(22) (2005).

97. Saigusa H, Urashima S, Asami H, and Ohba M: *Microhydration of the Guanine-Guanine and Guanine-Cytosine Base Pairs*. J. Phys. Chem. A, **114**(42):11231-11237 (2010).
98. Saigusa H, Mizuno N, Asami H, Takahashi K, and Tachikawa M: *Ultraviolet Spectroscopy and Theoretical Calculations of Mono- and Dihydrated Clusters of the Guanine Nucleosides: Possibility of Different Hydration Structures for Guanosine and 2'-Deoxyguanosine*. Bull. Chem. Soc. Jpn., **81**(10):1274-1281 (2008).
99. Perun S, Sobolewski AL, and Domcke W: *Photostability of 9H-adenine: mechanisms of the radiationless deactivation of the lowest excited singlet states*. Chem. Phys., **313**(1-3):107-112 (2005).
100. Sobolewski AL and Domcke W: *Photophysics of intramolecularly hydrogen-bonded aromatic systems: ab initio exploration of the excited-state deactivation mechanisms of salicylic acid*. PCCP, **8**(29):3410-3417 (2006).
101. Shemesh D, Hattig C, and Domcke W: *Photophysics of the Trp-Gly dipeptide: Role of electron and proton transfer processes for efficient excited-state deactivation*. Chem. Phys. Lett., **482**(1-3):38-43 (2009).
102. Shemesh D, Sobolewski AL, and Domcke W: *Efficient Excited-State Deactivation of the Gly-Phe-Ala Tripeptide via an Electron-Driven Proton-Transfer Process*. J. Am. Chem. Soc., **131**(4):1374-+ (2009).
103. Schultz T, Samoylova E, Radloff W, Hertel IV, Sobolewski AL, and Domcke W: *Efficient deactivation of a model base pair via excited-state hydrogen transfer*. Science, **306**(5702):1765-1768 (2004).
104. Sobolewski AL and Domcke W: *Ab Initio studies on the photophysics of the guanine-cytosine base pair*. PCCP, **6**:2763-2771 (2004).
105. Perun S, Sobolewski AL, and Domcke W: *Role of Electron-Driven Proton-Transfer Processes in the Excited-State Deactivation of the Adenine-Thymine Base Pair*. J. Phys. Chem. A., **110**:9031 (2006).
106. Nir E, Kleinermaans K, and de Vries MS: *Pairing of isolated nucleic-acid bases in the absence of the DNA backbone*. Nature, **408**(6815):949-951 (2000).
107. Nir E, Janzen C, Imhof P, Kleinermaans K, and de Vries MS: *Pairing of the nucleobase guanine studied by IR-UV double-resonance spectroscopy and ab initio calculations*. PCCP, **4**(5):740-750 (2002).
108. Nolting D, Weinkauff R, Hertel IV, and Schultz T: *Excited-state relaxation of protonated adenine*. ChemPhysChem, **8**(5):751-755 (2007).
109. Samoylova E, Lippert H, Ullrich S, Hertel IV, Radloff W, and Schultz T: *Dynamics of photoinduced processes in adenine and thymine base pairs*. J. Am. Chem. Soc., **127**(6):1782-1786 (2005).
110. Ullrich S, Schultz T, Zgierski MZ, and Stolow A: *Direct observation of electronic relaxation dynamics in adenine via time-resolved photoelectron spectroscopy*. J. Am. Chem. Soc., **126**(8):2262-2263 (2004).
111. Samoylova E, Radloff W, Ritze HH, and Schultz T: *Observation of Proton Transfer in 2-Aminopyridine Dimer by Electron and Mass Spectroscopy*. J. Phys. Chem. A, **113**(29):8195-8201 (2009).
112. Samoylova E, Smith VR, Ritze HH, Radloff W, Kabelac M, and Schultz T: *Ultrafast deactivation processes in aminopyridine clusters: Excitation energy dependence and isotope effects*. J. Am. Chem. Soc., **128**(49):15652-15656 (2006).
113. Nir E, Hünig I, Kleinermaans K, and de Vries MS: *Conformers of guanosines and their vibrations in the electronic ground and excited states, as revealed by double-resonance spectroscopy and ab initio calculations*. ChemPhysChem, **5**(1):131-137 (2004).
114. Nir E and de Vries MS: *Fragmentation of laser-desorbed 9-substituted adenines*. Int. J. Mass spectrom., **219**(1):133-138 (2002).

115. Asami H, Yagi K, Ohba M, Urashima S, and Saigusa H: *Stacked base-pair structures of adenine nucleosides stabilized by the formation of hydrogen-bonding network involving the two sugar groups*. Chem. Phys., **419**:84-89 (2013).
116. Saigusa H and Lim EC: *Excimer formation in van der Waals dimers and clusters of aromatic molecules*. Acc. Chem. Res., **29**(4):171-178 (1996).
117. Schreier WJ, Schrader TE, Koller FO, Gilch P, Crespo-Hernandez CE, Swaminathan VN, Carell T, Zinth W, and Kohler B: *Thymine dimerization in DNA is an ultrafast photoreaction*. Science, **315**(5812):625-629 (2007).
118. Crespo-Hernández CE, Cohen B, and Kohler B: *Base stacking controls excited-state dynamics in A·T DNA*. Nature, **436**:1141-1144 (2005).
119. Takaya T, Su C, de La Harpe K, Crespo-Hernández CE, and Kohler B: *UV excitation of single DNA and RNA strands produces high yields of exciplex states between two stacked bases*. Proc. Natl. Acad. Sci. U.S.A., **105**(July 29, 2008):10285-10290 (2008).
120. Crespo-Hernández CE, de La Harpe K, and Kohler B: *Ground-state recovery following UV excitation is much slower in G•C-DNA duplexes and hairpins than in mononucleotides*. J. Am. Chem. Soc., **130**:10844-10845 (2008).