

eScholarship

Combinatorial Theory

Title

Sets of mutually orthogoal projective and affine planes

Permalink

<https://escholarship.org/uc/item/6q20z7sg>

Journal

Combinatorial Theory, 4(1)

ISSN

2766-1334

Authors

Colbourn, Charles J.

Ingalls, Colin

Jedwab, Jonathan

et al.

Publication Date

2024

DOI

10.5070/C64163845

Supplemental Material

<https://escholarship.org/uc/item/6q20z7sg#supplemental>

Copyright Information

Copyright 2024 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at

<https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

SETS OF MUTUALLY ORTHOGOVAL PROJECTIVE AND AFFINE PLANES

Charles J. Colbourn¹, Colin Ingalls^{*2}, Jonathan Jedwab^{*3}, Mark Saaltink⁴, Ken W. Smith⁵, and
Brett Stevens^{*6}

¹ *School of Computing and Augmented Intelligence, Arizona State University, Tempe, AZ 85287-8809, U.S.A.*
Charles.Colbourn@asu.edu

² *School of Mathematics and Statistics, Carleton University, 1125 Colonel By Drive, Ottawa, ON K1S 5B6, Canada*
coliningalls@cunet.carleton.ca

³ *Department of Mathematics, Simon Fraser University, 8888 University Drive, Burnaby, BC V5A 1S6, Canada*
jed@sfu.ca

⁴ *Independent researcher mark@saaltink.ca*

⁵ *Department of Mathematics and Statistics, Sam Houston State University, Huntsville, TX 77341, U.S.A.*
kenwsmith54@gmail.com

⁶ *School of Mathematics and Statistics, Carleton University, 1125 Colonel By Drive, Ottawa ON K1S 5B6, Canada*
brett@math.carleton.ca

Submitted: Oct 21, 2022; Accepted: Jan 7, 2024; Published: Jun 30, 2024

© The authors. Released under the CC BY license (International 4.0).

Abstract. A pair of planes, both projective or both affine, of the same order and on the same point set are *orthogonal* if each line of one plane intersects each line of the other plane in at most two points. In this paper we prove new constructions for sets of mutually orthogonal planes, both projective and affine, and review known results that are equivalent to sets of more than two mutually orthogonal planes. We also discuss the connection between sets of mutually orthogonal planes and covering arrays.

Keywords. Finite geometry, projective planes, affine planes, covering arrays, orthogonal planes

Mathematics Subject Classifications. 05B25, 05B40, 51E20, 51E21

1. Introduction

In a projective plane of order q an *oval* is a set of $q + 1$ points, no three of which are collinear. A beautiful theorem, independently published multiple times, states that if D is a $(q^2 + q + 1, q + 1, 1)$ -difference set over \mathbb{Z}_{q^2+q+1} , then $-D$ is an oval in the projective plane developed from D [BBEF94, RMS14, Gly78, Bru73]. Since $-D$ is itself also a difference set, it can be developed into a second projective plane on the same ground set, \mathbb{Z}_{q^2+q+1} . Hence we

*supported by NSERC

have constructed the blocks of two projective planes with the property that the lines of one are ovals in the other. We introduce the term *orthogonal* to describe this property.

Definition 1.1. Two planes, both projective or both affine, of the same order and on the same point set are *orthogonal* if a line of one plane intersects any line in the other plane in at most two points, or in other words the blocks of one are ovals in the other. A set of planes is called a *set of mutually orthogonal planes* if the planes are pairwise orthogonal.

We provide two clarifications of the term “oval” in this definition. In the case of a projective plane, we consider an oval as having $q + 1$ points (whereas some authors require $q + 2$ points in even characteristic; others call such a set of $q + 2$ points a hyperoval). In the case of an affine plane, we use “oval” to mean a set of q points no three of which are collinear (although this use is not standard).

The primary goal of this paper is to prove new constructions for sets of mutually orthogonal planes, both projective and affine, and to review known results that are equivalent to sets of more than two mutually orthogonal planes.

A pair of orthogonal projective planes of order q was used by Raaphorst, Moura and Stevens to construct a family of covering arrays of strength 3, $CA(2q^3 - 1; 3, q^2 + q + 1, q)$ (see Definition 4.2) from the circulant matrices of linear feedback shift register sequences over finite fields \mathbb{F}_q . For $q > 3$ these are still the best known strength 3 covering arrays for their number of columns and alphabet sizes. Torres-Jimenez and Izquierdo-Marquez constructed covering arrays $CA(2q^3 - q; 3, q^2 - q + 3, q)$ which are equivalent to deleting a particular set of $2q - 2$ columns of the $CA(2q^3 - 1; 3, q^2 + q + 1, q)$ and then noting that, for every non-zero symbol x , the array that remains has two copies of the all- x row. Deleting one of each of these pairs completes the construction [TJIM18]. Colbourn, Lanus and Sarkar found, via Sherwood covering perfect hash families, a $CA(1016; 3, 64, 8)$ using a conditional expectation search [CLS18]. Investigating its structure they observed that if pairs of orthogonal Desarguesian affine planes could be constructed, then keeping just the columns corresponding to the points of the affine planes from the $CA(2q^3 - 1; 3, q^2 + q + 1, q)$ and deleting one copy of any repeated row, would yield $CA(2q^3 - q; 3, q^2, q)$ and thereby improve Torres-Jimenez and Izquierdo-Marquez’s covering arrays by $q - 3$ columns. A second purpose of this paper is to discuss the connection between sets of mutually orthogonal planes and covering arrays.

A pair of orthogonal affine planes of order 4 was used by Colbourn [Col99] as a key ingredient in disproving a conjecture of Jean Doyen on the automorphisms of Steiner 2-designs.

The union of a set of s mutually orthogonal planes of order q gives a “super-simple” $2-(q^2, q, s)$ design in the affine case, or a super-simple $2-(q^2 + q + 1, q + 1, s)$ design in the projective case. These designs have been studied by Bluskov and Heinrich [BH01] and others. Theorem 1.1 of [BH01] gives the same bound as the Johnson bound appearing in Corollary 1.3 below.

In a set of mutually orthogonal planes, any three points collinear in one are non-collinear in all the others. Therefore, taking the collection of all lines from each plane in a set of s mutually orthogonal planes gives a strength 3 packing design [CD07]. The *packing number* $D(v, k, t)$ of a v -set is the maximum number of k -subsets (blocks) such that every t -subset appears in at most one block. This can be used to calculate an upper bound on s .

Theorem 1.2. *If there exists a set of s mutually orthogonal projective planes of order q then*

$$s(q^2 + q + 1) \leq D(q^2 + q + 1, q + 1, 3).$$

If there exists a set of s mutually orthogonal affine planes of order q then

$$s(q^2 + q) \leq D(q^2, q, 3).$$

The Johnson bound [CD07] on the size of packings is

$$D(v, k, 3) \leq \left\lfloor \frac{v}{k} \left\lfloor \frac{v-1}{k-1} \left\lfloor \frac{v-2}{k-2} \right\rfloor \right\rfloor \right\rfloor.$$

Since the blocks in $AG(2, 2)$ have size two, the set of s copies of $AG(2, 2)$ is trivially mutually orthogonal for every s (similar to the fact that every set of s latin squares of order 1 is mutually orthogonal). Otherwise, the Johnson bound gives the following result.

Corollary 1.3. • *If there exists a set of s mutually orthogonal projective planes of order q , then $s \leq \max\{5, q + 2\}$.*

• *If there exists a set of s mutually orthogonal affine planes of order $q > 2$, then $s \leq \max\{7, q + 2\}$.*

We will either tighten or meet these upper bounds for some small values of q .

In Section 2 we discuss projective planes. There we give a new proof of the existence of a pair of orthogonal projective planes from an algebraic geometric point of view using the Cremona transformation. We also consider if larger sets of such projective planes exist. In Section 3 we turn our attention to affine planes. We prove that for even characteristic there always exists a pair of planes with the desired property. When $\gcd(n, 6) = 1$ we construct a set of three mutually orthogonal affine planes of order 2^n . We then construct sets of size of seven for $q = 4, 8$. In Section 4, we discuss the connections between sets of mutually orthogonal planes and covering arrays. We show that for affine planes we can extend the constructed arrays by two additional columns to construct a $CA(2q^3 - q; 3, q^2 + 2, q)$, and demonstrate the utility of sets of more than two orthogonal planes. In Section 5 we give some final thoughts.

2. Projective planes

2.1. A pair of orthogonal projective planes

We start by giving a new proof of the fact that for every prime power q there exists a pair of Desarguesian orthogonal $PG(2, q)$. This proof uses the Cremona transformation and its algebraic geometric approach adds another interesting point of view of this theorem. For a general introduction to algebraic geometry and its terminologies, see [Har77]. While we believe that the result was broadly speaking known to algebraic geometers we do not know of an explicit articulation or proof of it in the literature.

We express the points of $\text{PG}(2, q)$ with homogeneous coordinates using colons between the coordinates to make them distinct from other numbers that appear in round brackets in this article. The *Cremona transformation*, or *standard quadratic transformation* is the partial rational function $\text{Cr} : \text{PG}(2, q) \dashrightarrow \text{PG}(2, q)$ defined by

$$\text{Cr}((x : y : z)) = (x^{-1} : y^{-1} : z^{-1}).$$

We can extend this partial map and will use the same symbol for this extension,

$$\text{Cr}((x : y : z)) = (xyzx^{-1} : xyzzy^{-1} : xyzzz^{-1}) = (yz : xz : xy).$$

This map is undefined on the intersection points of the pairs of lines yz , xz and xy (variety $V(yz, xz, xy)$), $\{(1 : 0 : 0), (0 : 1 : 0), (0 : 0 : 1)\}$, and is birational since $\text{Cr} \circ \text{Cr} = \text{id}$. We are interested in the image of lines and conics under Cr , which are the varieties of degree one and degree two homogeneous polynomials respectively. For example, if ℓ is a line, we take $\text{Cr}(\ell)$ to mean the set $\{\text{Cr}(p) : p \in \ell\}$. This set might be a single point, a set of collinear points or a set of points lying on a conic. It may not contain all the points of this line or conic because Cr is a partial map. We can “close” the image set with the following procedure. Let $C = V(f(x, y, z))$ in $\text{PG}(2, q)$, not contained in the lines $V(x)$, $V(y)$, or $V(z)$. Let $g(x, y, z) = f(1/x, 1/y, 1/z)x^a y^b z^c$, where a, b and c are minimal such that g is a polynomial. Then the closure of the image of f under the Cremona transformation is $\text{Cr}(C) = V(g(x, y, z))$.

The Cremona transformation has the following properties which are not hard to prove. See [Ful89, Sha13] for examples of proofs of properties of Cr .

Proposition 2.1. • Cr is bijective on $\text{PG}(2, q) - V(xyz)$.

- The lines x , y and z map to points $(1 : 0 : 0)$, $(0 : 1 : 0)$ and $(0 : 0 : 1)$ respectively.
- The line $\beta y + \gamma z$ through $(1 : 0 : 0)$ maps to the line $\gamma y + \beta z$, also through $(1 : 0 : 0)$; and similarly for lines $\alpha x + \gamma z$ and $\alpha x + \beta y$ through $(0 : 1 : 0)$ and $(0 : 0 : 1)$ respectively.
- Each line $ax + by + cz$ that does not contain any of $(1 : 0 : 0)$, $(0 : 1 : 0)$ or $(0 : 0 : 1)$ maps to the irreducible conic $ayz + bxz + cxy$ which contains all three of $(1 : 0 : 0)$, $(0 : 1 : 0)$ and $(0 : 0 : 1)$.

Theorem 2.2 ([BBEF94, RMS14, Gly78, Bru73, She86]). For every prime power q , there exists a pair of Desarguesian orthogoval $\text{PG}(2, q)$.

Proof. Let $\omega \in \mathbb{F}_{q^3}$ so that $\omega \notin \mathbb{F}_q$ and ω is not a root of $x^{3q^2} - 3x^{q^2+q+1} + x^{3q} + x^3$. Since 0 is a root with multiplicity 3 and 1 is also a root, if $q \geq 3$ then such an ω exists by the pigeonhole principle. If $q = 2$ then $\alpha + 1$ satisfies these requirements where α is a root of $x^3 + x + 1$. By these conditions, $(\omega : \omega^q : \omega^{q^2})$, $(\omega^q : \omega^{q^2} : \omega)$ and $(\omega^{q^2} : \omega : \omega^q)$ are not collinear. Let σ be the projective linear transformation of $\text{PG}(2, q^3)$ that maps $(1 : 0 : 0)$, $(0 : 1 : 0)$, $(0 : 0 : 1)$ and $(1 : 1 : 1)$ to $(\omega : \omega^q : \omega^{q^2})$, $(\omega^q : \omega^{q^2} : \omega)$, $(\omega^{q^2} : \omega : \omega^q)$ and $(1 : 1 : 1)$ respectively. We conjugate the Cremona transformation by σ

$$\text{Cr}' = \sigma \circ \text{Cr} \circ \sigma^{-1}$$

and consider its action on $PG(2, q^3)$.

Since Cr maps

$$\alpha(1 : 0 : 0) + \beta(0 : 1 : 0) + \gamma(0 : 0 : 1)$$

to

$$\beta\gamma(1 : 0 : 0) + \alpha\gamma(0 : 1 : 0) + \alpha\beta(0 : 0 : 1),$$

Cr' maps

$$\alpha(\omega : \omega^q : \omega^{q^2}) + \beta(\omega^q : \omega^{q^2} : \omega) + \gamma(\omega^{q^2} : \omega : \omega^q)$$

to

$$\beta\gamma(\omega : \omega^q : \omega^{q^2}) + \alpha\gamma(\omega^q : \omega^{q^2} : \omega) + \alpha\beta(\omega^{q^2} : \omega : \omega^q).$$

Lines not containing any of $(\omega : \omega^q : \omega^{q^2})$, $(\omega^q : \omega^{q^2} : \omega)$, $(\omega^{q^2} : \omega : \omega^q)$ are mapped to irreducible conics that contain all three of these conjugate points.

Suppose that $(x : y : z) \in PG(2, q) \subset PG(2, q^3)$. Express this point as a sum

$$(x : y : z) = \alpha(\omega : \omega^q : \omega^{q^2}) + \beta(\omega^q : \omega^{q^2} : \omega) + \gamma(\omega^{q^2} : \omega : \omega^q).$$

Then the fact that the Frobenius map fixes this point $((x : y : z)^q = (x : y : z))$ implies that

$$\beta = \alpha^q \quad \text{and} \quad \gamma = \alpha^{q^2}. \tag{2.1}$$

Conversely if α, β and γ satisfy Equation (2.1) then

$$\alpha(\omega : \omega^q : \omega^{q^2}) + \beta(\omega^q : \omega^{q^2} : \omega) + \gamma(\omega^{q^2} : \omega : \omega^q)$$

is invariant under the Frobenius map and is therefore in the subplane $PG(2, q)$.

Thus,

$$\begin{aligned} Cr'((x : y : z)) &= \beta\gamma(\omega : \omega^q : \omega^{q^2}) + \alpha\gamma(\omega^q : \omega^{q^2} : \omega) + \alpha\beta(\omega^{q^2} : \omega : \omega^q) \\ &= \alpha^{q+q^2}(\omega : \omega^q : \omega^{q^2}) + \alpha^{1+q^2}(\omega^q : \omega^{q^2} : \omega) + \alpha^{1+q}(\omega^{q^2} : \omega : \omega^q). \end{aligned}$$

The coefficients α^{q+q^2} , α^{1+q^2} and α^{1+q} satisfy the identities in Equation (2.1) and so $Cr'((x : y : z)) \in PG(2, q)$ if and only if $(x : y : z) \in PG(2, q)$.

Thus Cr' is a bijection on $PG(2, q)$ that maps lines to irreducible conics and its image forms a second plane which is orthogoval to the first. \square

2.2. Sets of more than two orthogoval projective planes

A $PG(2, 2)$ is a Steiner triple system (STS) of order 7 and any 3 non-collinear points are an oval. Thus two STS(7) are orthogoval if and only if their blocksets are disjoint. It is known that there does not exist a set of three of STS(7)s with mutually disjoint blocksets [CR99]. When $q = 3$ it is known that the collection of all 715 4-sets of a 13-set can be decomposed into 55 copies of $PG(2, 3)$ [Cho83]. But not all of the lines in these copies will be ovals in the other copies: if a 4-set intersects a line in three points it is not an oval in the plane containing the line. From

Corollary 1.3 we know that if a set of s mutually orthogonal projective planes in $\text{PG}(2, 3)$ exists then

$$13s \leq D(13, 4, 3) = 65,$$

and so $s \leq 5$. Chu and Colbourn have shown that a strength 3 packing of 4-sets on 13 points with a cyclic automorphism group has at most 52 blocks [CC04]. There does exist such a cyclic packing which can be decomposed into four projective planes of order 3. This consists of the standard Desarguesian plane, together with the points of the conics

$$\begin{aligned} z^2 + xy &= 0 \\ z^2 + yz + xy &= 0 \\ z^2 + 2yz + 2x &= 0 \end{aligned}$$

and all the images of these conics in the Singer cycle. In terms of difference sets the blocks of the Desarguesian plane are the development of $D = \{0, 1, 3, 9\}$. The three conics above correspond to difference sets $-D$, $2D$ and $-2D$ respectively. Baker et al. proved that no conics are repeated in this set [BBEF94], and it is easy to check that no triple is repeated in any block. These give the blocks

$$\{0, 1, 3, 9\} + x, \quad \{0, 1, 5, 11\} + x, \quad \{0, 1, 4, 6\} + x, \quad \{0, 1, 8, 10\} + x$$

for $x \in \mathbb{Z}_{13}$.

We now show that there does not exist a set of five orthogonal projective planes of order 3. Suppose otherwise, for a contradiction. Then the set of all their blocks forms a maximum packing with $D(13, 4, 3) = 65$ blocks. Each plane corresponds to a clique of size 13 in the 1-block intersection graph of the packing design. In any such packing with 65 blocks, 26 triples are uncovered, the number of blocks containing a pair of points is 5, and each point is contained in exactly 20 blocks. From this it can be deduced that the 26 missing triples are the blocks of a STS(13). Adjoining one new point to each of these and adding them to the design yields a Steiner quadruple system on 14 points (a SQS(14)). Thus every $v = 13$, $k = 4$, $t = 3$ packing with 65 blocks is obtained from an SQS(14) by the deletion of one point. There are 4 non-isomorphic SQS(14) [MH72]. Using SageMath [The21], the free open-source mathematics software system, for each of these SQS(14) we can delete each point and construct the 1-block intersection graph of the resulting packing. None of these graphs has a clique of size larger than 9 (code available upon request), which gives the required contradiction.

To determine the maximum size of a set of mutually orthogonal projective planes of orders $q = 4$ and $q = 5$ we built the 1-intersection graph of all the ovals in these two projective planes in SageMath. Cliques of size $q^2 + q + 1$ correspond to projective planes with ovals for blocks. A graph is then formed with all these possible cliques as vertices and adjacency determined by a pair being orthogonal. For both $q = 4$ and $q = 5$, no pairs of any of these planes were orthogonal, thus there exists a pair of orthogonal projective planes of orders $q = 4$ and 5 but no set of three. Source code is available upon request.

For $q = 4$ there are 1008 ovals. The 1-intersection graph is regular of degree 475, with 960 cliques of size 21. The 1-intersection graph was computed in 37s; the cliques were found in

another 16s, and checking that no two cliques are orthogoval took 62s for a total of 111 seconds. For $q = 5$ there are 3100 ovals. The 1-intersection graph is regular of degree 1404, and has 12000 cliques of size 31. The 1-intersection graph was computed in 149s; the cliques were found in another 73m, and checking that no two cliques are orthogoval took 2h46m, for a total of just over 4 hours. These timings are for unoptimized code in SageMath 9.4 running on an AMD Ryzen 5600; in particular this code does not use symmetry to reduce the search space.

We summarize the results of this subsection.

Theorem 2.3. *The size of a largest set of mutually orthogoval projective planes in $\text{PG}(2, q)$ is two for $q \in \{2, 4, 5\}$ and is four for $q = 3$.*

3. Affine planes

Because all lines have size two in an affine plane of order 2, a collection of any number of affine planes of order 2 on the same point set is trivially mutually orthogoval. If $q = 3$ then the set of disjoint STS(9) as constructed in [Bay17, Tei73] gives a collection of seven mutually orthogoval affine planes in $\text{AG}(2, 3)$. By Corollary 1.3, this is the most possible. Up to isomorphism there are two distinct sets of disjoint STS(9) [Bay17, BGG03].

Theorem 3.1. *The size of a largest set of mutually orthogoval affine planes in $\text{AG}(2, 3)$ is seven.*

This is the only case of orthogoval affine planes of odd order which we know of, whereas when q is even we can construct pairs or triples of orthogoval affine planes and in some cases larger sets. We identify the points of $\text{AG}(2, q)$ as the subset of points $(x : y : 1)$ from $\text{PG}(2, q)$.

3.1. A pair of orthogoval affine planes

To show that there exists a pair of orthogoval affine planes of order $q = 2^n$, we will need irreducible cubic polynomials in $\mathbb{F}_q[x]$ of the form $x^3 + bx + c$. When n is even, $2^n - 1$ is divisible by 3 and thus there exist elements $b \in \mathbb{F}_{2^n}$ which are not cube roots. For each such b , the polynomial $x^3 - b$ is irreducible. When n is odd, every element is a cube root, $f(x) = x^3$ is a bijection and all polynomials of the form $x^3 + b$ are reducible. However irreducible cubics of the desired form still exist.

Lemma 3.2. *For each n , there exists an irreducible polynomial of the form $x^3 + bx + c$ in $\mathbb{F}_{2^n}[x]$.*

Proof. There exist $q(q - 1)$ monic quadratic polynomials $x^2 + bx + c$ with $c \neq 0$, of which $(q - 1)(q - 2)/2 + (q - 1) = q(q - 1)/2$ are reducible and thus $q(q - 1)/2$ are irreducible. There are a total of $q(q - 1)$ cubic polynomials $x^3 + bx + c$ with $c \neq 0$. If such a cubic factors into linear terms, it must be of the form $(x - r_0)(x - r_1)(x + (r_0 + r_1))$ with $r_0, r_1, r_0 + r_1 \neq 0$. Thus there are $(q - 1)(q - 2)/6$ such reducible cubics. If such a cubic factors into an irreducible quadratic and a linear term, it must be of the form $(x^2 + dx + e)(x + d)$ where $d, e \neq 0$ and the linear term is determined by the quadratic term. Thus the number of these reducible cubics matches the number of irreducible quadratics, $q(q - 1)/2$. Thus there are $(q - 1)(q + 1)/3 > 0$ monic irreducible cubics with no x^2 term and non-zero constant term. \square

For $q = 2^n$, we shall call a conic \mathcal{O} in $\text{PG}(2, q)$ such that $(x_0 + x_1 : y_0 + y_1 : 1) \in \mathcal{O}$ for all $(x_0 : y_0 : 1), (x_1 : y_1 : 1) \in \mathcal{O}$ a *subspace conic*; these (possibly when augmented with their nucleus) are a particular type of *translation oval* [Hir98, GS94]. Specifically, these are translation ovals that are conics and have the line $z = 0$ as their axis. We show that there always exists a pencil of subspace conics in $\text{PG}(2, q)$. For this we need to characterize the subspace conics that may not contain $(1 : 0 : 0), (0 : 1 : 0)$ or $(1 : 1 : 1)$.

Lemma 3.3. *If $q = 2^n$, the conic in $\text{PG}(2, q)$ defined by*

$$ax^2 + by^2 + cz^2 + fyz + gxz + hxy = 0$$

is a subspace conic if and only if $h = c = 0$.

Proof. A subspace conic must contain $(0 : 0 : 1)$ so $c = 0$ is necessary. Since the conic is an oval it contains a pair of points $(x_0 : y_0 : 1)$ and $(x_1 : y_1 : 1)$ such that $x_0y_1 \neq x_1y_0$. Thus $(x_0 + x_1 : y_0 + y_1 : 1)$ is also on the oval and

$$\begin{aligned} ax_0^2 + by_0^2 + fy_0 + gx_0 + hx_0y_0 &= 0, \\ ax_1^2 + by_1^2 + fy_1 + gx_1 + hx_1y_1 &= 0, \\ a(x_0 + x_1)^2 + b(y_0 + y_1)^2 + f(y_0 + y_1) + g(x_0 + x_1) + h(x_0 + x_1)(y_0 + y_1) &= 0. \end{aligned}$$

The sum of these three equations is

$$h(x_0y_1 + x_1y_0) = 0$$

and since $x_0y_1 + x_1y_0 \neq 0$ we must have $h = 0$.

Conversely if $c = h = 0$ and $(x_0 : y_0 : 1)$ and $(x_1 : y_1 : 1)$ are on the conic, then

$$\begin{aligned} ax_0^2 + by_0^2 + fy_0 + gx_0 &= 0, \\ ax_1^2 + by_1^2 + fy_1 + gx_1 &= 0. \end{aligned}$$

The quadratic form evaluated at $(x_0 + x_1 : y_0 + y_1 : 1)$ is

$$\begin{aligned} a(x_0 + x_1)^2 + b(y_0 + y_1)^2 + f(y_0 + y_1) + g(x_0 + x_1) \\ = (ax_0^2 + by_0^2 + fy_0 + gx_0) + (ax_1^2 + by_1^2 + fy_1 + gx_1) = 0 \end{aligned}$$

so the conic is a subspace conic. □

Note that any subspace conic is tangent to the line $z = 0$ and thus the restriction to the affine plane part of $\text{PG}(2, q)$ has q points and so is an oval.

We will call two subspace conics which intersect only in $(0 : 0 : 1)$ *trivially intersecting*.

Corollary 3.4. *For every $q = 2^n$ there exists a trivially intersecting pencil of non-singular subspace conics in $\text{PG}(2, q)$.*

Proof. Let $p = x^3 + bx + c$ be an irreducible cubic equation in $\mathbb{F}_q[x]$ (note $c \neq 0$). We note that $x^3 + b^2x + c^2$ is also irreducible. Define conics

$$\begin{aligned} \phi &: x^2 + yz, \\ \chi &: y^2 + byz + cxz. \end{aligned}$$

Both these satisfy Lemma 3.3 so they are subspace conics. The conics determine the pencil

$$\{\alpha\phi + \beta\chi : (\alpha : \beta) \in \text{PG}(1, q)\}.$$

The partial derivatives of $\alpha\phi + \beta\chi$ are all zero at the point $(\alpha + \beta b : \beta c : 0)$ and this is not on the conic because $x^3 + b^2x + c^2$ is irreducible and thus all conics in the pencil are non-singular.

By the irreducibility of p , the equation $\phi(x, y, z) = \chi(x, y, z) = 0$ in $\text{PG}(2, q)$ holds if and only if $(x : y : z) = (0 : 0 : 1)$. The other three points of intersection are a conjugate triple in $\text{PG}(2, q^3)$. Every conic in the pencil contains these four points, thus the pencil of conics $\{\alpha\phi + \beta\chi : (\alpha : \beta) \in \text{PG}(1, q)\}$ meet inside $\text{PG}(2, q)$ only in $(0 : 0 : 1)$ and are otherwise disjoint. \square

In \mathbb{F}_2^{2n} , a set of n -dimensional spaces which pairwise intersect only at the origin is a *spread*. The set of these spaces and their cosets form an affine plane of order 2^n which is called a *translation plane* because the group of all translations is a transitive automorphism group [Dem97]. The Desarguesian affine plane of order $q = 2^n$ can be constructed this way using the spread of 1-dimensional spaces of \mathbb{F}_q^2 which is isomorphic to \mathbb{F}_2^{2n} as a \mathbb{F}_2 -vector space. We call this the *line spread*. These 1-dimensional spaces correspond to the lines $\alpha x + \beta y$ for $(\alpha : \beta) \in \text{PG}(1, q)$. They alternatively can be constructed from the cyclotomic cosets in $\mathbb{F}_{q^2}^*$. If ω is a primitive element of \mathbb{F}_{q^2} , the set

$$C_0 = \langle \omega^{q+1} \rangle = \{\omega^{j(q+1)} : 0 \leq j < q - 1\}$$

is a multiplicative subgroup of $\mathbb{F}_{q^2}^*$. Its cosets are

$$C_i = \omega^i \langle \omega^{q+1} \rangle \tag{3.1}$$

for $0 \leq i \leq q$. When we adjoin 0 to each coset we get exactly the set of 1-dimensional subspaces of \mathbb{F}_q^2 , a spread. Non-Desarguesian translation planes can also be constructed from spreads [Dem97].

We want to understand when the planes constructed from two spreads are orthogoval. Let $\mathcal{S} = \{S_i\}$ and $\mathcal{T} = \{T_j\}$ be two spreads in \mathbb{F}_2^{2n} . If $x, y, z \in (u + S_i) \cap (v + T_j)$ then $0, y - x, z - x \in S_i \cap T_j$ and conversely. This gives the following result.

Lemma 3.5. *The affine planes constructed from \mathcal{S} and \mathcal{T} are orthogoval if and only if $|S_i \cap T_j| \leq 2$ for all i and j .*

Corollary 3.4 shows that there exists a “spread” of non-singular conics. The translation plane constructed from this spread is Desarguesian.

Definition 3.6. A map f on $\text{AG}(2, q)$ which is an \mathbb{F}_2 -linear bijection and maps lines to ovals is *affine ovalinear*.

Lemma 3.7. *If f is an affine ovalinear map on $\text{AG}(2, 2^n)$ then the affine planes $\text{AG}(2, 2^n)$ and $\text{AG}(2, 2^n)^f$ are orthogoval and both Desarguesian.*

Proof. Because f is a bijection, the two planes $\text{AG}(2, 2^n)$ and $\text{AG}(2, 2^n)^f$ are isomorphic and therefore Desarguesian. Since every line in $\text{AG}(2, 2^n)^f$ is an oval, the two planes are orthogoval. \square

We note that, by applying f^{-1} to both planes, we have that $\text{AG}(2, 2^n)$ and $\text{AG}(2, 2^n)^f$ are orthogoval if and only if $\text{AG}(2, 2^n)$ and $\text{AG}(2, 2^n)^{f^{-1}}$ are orthogoval. We will use Lemma 3.7 to prove the next result and again in Section 3.2.

Theorem 3.8. *The translation plane of order $q = 2^n$ constructed from a trivially intersecting pencil of non-singular conics is Desarguesian and orthogoval to the translation plane constructed from the line spread.*

Proof. Suppose that quadratic forms ϕ and χ generate a trivially intersecting pencil of translation conics. Let ψ be the quadratic form z^2 . Define the map on $\text{PG}(2, q)$, $f((x : y : z)) = (\phi(x, y, z) : \chi(x, y, z) : \psi(x, y, z))$. Clearly f preserves the z -coordinate of all points in $\text{PG}(2, q)$ and is thus \mathbb{F}_2 -linear on $\text{AG}(2, q)$. Suppose that $(x : y : z)$ is on $\alpha\phi + \beta\chi + \gamma\psi$. Then $f((x : y : z))$ is on the line $\alpha x + \beta y + \gamma z$. Since each point can be specified by the lines it is on, f is a bijection which takes ovals to lines. Thus f^{-1} is affine ovalinear and so by Lemma 3.7 $\text{AG}(2, 2^n)$ and $\text{AG}(2, 2^n)^f$ are orthogoval and both Desarguesian. \square

Corollary 3.9. *There exists a pair of Desarguesian orthogoval affine planes of order $q = 2^n$.*

We note that the pair of $\text{PG}(2, q)$ constructed above (one from the lines $\alpha x + \beta y + \gamma z$ and the other from the set of conics $\alpha\phi + \beta\chi + \gamma\psi$) has the following property: any three points that are collinear in one $\text{PG}(2, q)$ are non-collinear in the other, except when the three points are all from the line z . We therefore say that this pair of $\text{PG}(2, q)$ is orthogoval *except* for the line z . We will use this fact when discussing the connections to covering arrays in Section 4.

3.2. Sets of three orthogoval affine planes

For a restricted set of n , we can construct sets of three orthogoval affine planes of order $q = 2^n$. As in the construction of two orthogoval affine planes in Section 3.1, we need to control the roots of specific polynomials.

Lemma 3.10. *If k is relatively prime to n , then for any a and b in \mathbb{F}_{2^n} , the equation $ax^{2^k} + bx = 0$ has at most two roots in \mathbb{F}_{2^n} , unless $a = b = 0$.*

Proof. $x = 0$ is one root; any other must satisfy $ax^{2^k-1} + b = 0$. If $a = 0$ this has no roots unless $b = 0$. For $a \neq 0$, the sole root is the $(2^k - 1)$ th root of $a^{-1}b$. That root exists and is unique since $\text{gcd}(2^n - 1, 2^k - 1) = 2^{\text{gcd}(k, n)} - 1 = 1$. \square

Lemma 3.11. *If $3k$ is relatively prime to n , then the polynomial $x^{2^k+1} + x + 1$ has no roots in \mathbb{F}_{2^n} .*

Proof. Suppose $r \in \mathbb{F}_{2^n}$ is a root of this polynomial and let $m = 2^k$. Then from

$$r^{m+1} = r + 1$$

and taking the m th power (which is a Frobenius automorphism) of both sides we have

$$r^{m^2+m} = r^m + 1$$

so that

$$r^{m^2+m+1} = r^{m+1} + r = 1.$$

Now taking the $(m - 1)$ th power gives $r^{m^3-1} = 1$, so that

$$r \in \mathbb{F}_{m^3} = \mathbb{F}_{2^{3k}}.$$

But as we also have $r \in \mathbb{F}_{2^n}$ we conclude

$$r \in \mathbb{F}_{2^{\gcd(3k,n)}} = \mathbb{F}_2.$$

This is a contradiction, as neither 0 nor 1 is a root. □

A similar argument applies to a second polynomial we will encounter.

Lemma 3.12. *If $3k$ is relatively prime to n , then the polynomial $x^{2^{2k}-1} + x^{2^k-1} + 1$ has no roots in \mathbb{F}_{2^n} .*

Proof. Suppose $r \in \mathbb{F}_{2^n}$ is a root of this polynomial and let $m = 2^k$. Then from $r^{m^2-1} = r^{m-1} + 1$, taking the m th power of both sides we have $r^{m^3-m} = r^{m^2-m} + 1$. Multiplying by r^{m-1} gives $r^{m^3-1} = r^{m^2-1} + r^{m-1} = 1$. So $r \in \mathbb{F}_{m^3}$, but as we also have $r \in \mathbb{F}_{2^n}$ we conclude $r \in \mathbb{F}_{2^{\gcd(3k,n)}} = \mathbb{F}_2$. This is again a contradiction, as neither 0 nor 1 is a root. □

Incidentally, this also shows (because it is \mathbb{F}_2 -linear) that $x^{2^{2k}} + x^{2^k} + x$ is a permutation polynomial for \mathbb{F}_{2^n} when n is relatively prime to $3k$.

Let $\phi_k(x, y) = (x^{2^k} + y, y^{2^k} + x + y)$ be a map defined on $\text{AG}(2, 2^n)$. A simple expansion of terms shows

$$\phi_{2k} = \rho \circ \phi_k \circ \phi_k \circ \rho \tag{3.2}$$

where $\rho(x, y) = (y, x)$ is a linear transformation.

Lemma 3.13. *For all $k \geq 1$, if $3k$ is relatively prime to n , then ϕ_k is a bijection.*

Proof. Suppose $\phi_k(x, y) = \phi_k(s, t)$. Then

$$\begin{aligned} x^{2^k} + y &= s^{2^k} + t \text{ and} \\ y^{2^k} + x + y &= t^{2^k} + s + t, \end{aligned}$$

giving

$$\begin{aligned} (x + s)^{2^k} &= y + t \text{ and} \\ (y + t)^{2^k} &= (x + s) + (y + t). \end{aligned}$$

Substituting $y + t$ from the first equation into the second gives

$$(x + s)^{2^{2k}} = (x + s) + (x + s)^{2^k}.$$

That is, $x + s$ is a root of $z^{2^{2k}} + z^{2^k} + z$. Then either $x = s$, or $x + s$ is a root of $z^{2^{2k}-1} + z^{2^k-1} + 1$, which Lemma 3.12 shows to be impossible. \square

Lemma 3.14. *If $3k$ is relatively prime to n , then ϕ_k is affine ovalinear on \mathbb{F}_{2^n} .*

Proof. It is clear that ϕ_k is \mathbb{F}_2 -linear, and Lemma 3.13 shows it to be a bijection. We first find the images of lines through the origin.

The image of the line $x = 0$ is the set

$$\{(y, y^{2^k} + y) : y \in \mathbb{F}_{2^n}\} \quad (3.3)$$

while the image of the line $y = ux$ for $u \in \mathbb{F}_{2^n}$ is

$$\{(x^{2^k} + ux, u^{2^k}x^{2^k} + (u + 1)x) : x \in \mathbb{F}_{2^n}\}. \quad (3.4)$$

Now we can look at the intersection of these sets with lines through the origin. The line $x = 0$ meets the set (3.3) in the set indexed by $\{y \in \mathbb{F} : y = 0\}$, which has one member. It meets the set (3.4) in the set indexed by

$$\{x \in \mathbb{F} : x^{2^k} + ux = 0\}$$

which Lemma 3.10 shows to have at most 2 members.

For $w \in \mathbb{F}$, the line $y = wx$ meets the set (3.3) in the set indexed by

$$\{y \in \mathbb{F} : y^{2^k} + (w + 1)y = 0\},$$

which Lemma 3.10 shows to have at most 2 members. It meets the set (3.4) in the set indexed by

$$\{x \in \mathbb{F} : (u^{2^k} + w)x^{2^k} + (uw + u + 1)x = 0\}.$$

Lemma 3.10 applies again unless

$$u^{2^k} + w = 0 = uw + u + 1$$

but then $w = u^{2^k}$ and $u^{2^{k+1}} + u + 1 = 0$. So u is a root of $x^{2^{k+1}} + x + 1$, which Lemma 3.11 shows to be impossible. \square

Theorem 3.15. *If $6k$ and n are relatively prime and $q = 2^n$, the standard plane $AG(2, q)$, its image $AG(2, q)^{\phi_k}$, and $AG(2, q)^{\phi_k \circ \phi_k}$ are mutually orthogonal.*

Proof. By Lemmas 3.7 and 3.14, we have that $AG(2, q)$ and $AG(2, q)^{\phi_k}$ are orthogonal. Applying bijection ϕ_k to both of these planes shows that $AG(2, q)^{\phi_k}$ and $AG(2, q)^{\phi_k \circ \phi_k}$ are also orthogonal. It remains to show that the standard plane and $AG(2, q)^{\phi_k \circ \phi_k}$ are orthogonal. Lemmas 3.7 and 3.14 show that $AG(2, q)$ is orthogonal to $AG(2, q)^{\phi_{2k}} = AG(2, q)^{\rho \circ \phi_k \circ \phi_k \circ \rho}$. Since ρ is a bijection and $\rho \circ \rho$ is the identity, we can first apply ρ to those two planes to show that $AG(2, q)^\rho$ is orthogonal to $AG(2, q)^{\rho \circ \phi_k \circ \phi_k}$, and then, as $AG(2, q)^\rho = AG(2, q)$, we have $AG(2, q)$ orthogonal to $AG(2, q)^{\phi_k \circ \phi_k}$. \square

Clearly n must always be relatively prime to 6 for this construction. For $k \in \{2^i 3^j \mid i, j \geq 0\} = \{1, 2, 3, 4, 6, 8, \dots\}$ this is sufficient.

Corollary 3.16. *If n is relatively prime to 6, there exists a set of three mutually orthogonal affine planes of order 2^n .*

If $n > 3$ is prime then every k satisfying $1 \leq k \leq n - 1$ gives a triple of mutually orthogonal affine planes. We do not know whether or not the sets for different k are isomorphic.

3.3. Sets of more than three orthogonal affine planes

We can use the construction of translation affine planes from spreads to construct sets of more than three orthogonal affine planes for $q = 4$ and 8. The line spread \mathcal{C} defined just after Equation (3.1) constructs the standard Desarguesian affine plane on point set $\mathbb{F}_{2^{2n}} \cong \mathbb{F}_2^{2n}$. Other spreads which construct isomorphic translation planes can be generated by applying elements $M \in GL(2n, \mathbb{F}_2)$. By Lemma 3.5, if the intersections of the spaces in the image spread with the spaces from the original line spread all have size no more than two then the pair of affine planes constructed is orthogonal. Furthermore if this is true for M^i for each i satisfying $1 \leq i < s$ then the s spreads $\{M^i(\mathcal{C}) : 0 \leq i < s\}$ generate a set of s mutually orthogonal affine planes.

This gives an efficient method for computationally searching for a set of mutually orthogonal affine planes of $AG(2, 2^n)$. Generate matrices M from $GL(2n, \mathbb{F}_2)$, retaining those for which the plane constructed from $M(\mathcal{C})$ is orthogonal to that built from \mathcal{C} . Given a large set M_1, M_2, \dots, M_r of such matrices, construct a graph with vertex set $M_0 = I_{2n}, M_1, M_2, \dots, M_r$ in which vertices M_i and M_j are joined when $M_i(\mathcal{C})$ is orthogonal to $M_j(\mathcal{C})$. A clique of size c in this graph is then a set of c mutually orthogonal affine planes in $AG(2, 2^n)$. This search yielded a set of seven mutually orthogonal affine planes of order 4 (Example 3.17) and a set of seven mutually orthogonal affine planes of order 8 (Example 3.18).

Example 3.17. Let $q = 4$ and let ω be a root of the primitive polynomial $x^4 + x + 1$. The cyclotomic cosets C_i of \mathbb{F}_{16}^* and the corresponding additive subgroups S_i of $\mathbb{F}_2^4 \cong \mathbb{F}_{16}$ (using the

\mathbb{F}_2 -linear map $(t_0t_1t_2t_3) \rightarrow t_0\omega^3 + t_1\omega^2 + t_2\omega + t_3$) are given by

i	C_i (multiplicative)	S_i (additive)
0	$\{1, \omega^5, \omega^{10}\}$	$\{0000, 0001, 0110, 0111\}$
1	$\{\omega, \omega^6, \omega^{11}\}$	$\{0000, 0010, 1100, 1110\}$
2	$\{\omega^2, \omega^7, \omega^{12}\}$	$\{0000, 0100, 1011, 1111\}$
3	$\{\omega^3, \omega^8, \omega^{13}\}$	$\{0000, 1000, 0101, 1101\}$
4	$\{\omega^4, \omega^9, \omega^{14}\}$	$\{0000, 0011, 1010, 1001\}$.

The 4 lines of the parallel class containing S_0 are the cosets

$$L_{01} = 0000 + S_0 = \{0000, 0001, 0110, 0111\}$$

$$L_{02} = 0010 + S_0 = \{0010, 0011, 0100, 0101\}$$

$$L_{03} = 1000 + S_0 = \{1000, 1001, 1110, 1111\}$$

$$L_{04} = 1010 + S_0 = \{1010, 1011, 1100, 1101\}.$$

The remaining 16 lines of the Desarguesian affine translation plane $\text{AG}(2, 4)$ are formed similarly from S_1, \dots, S_4 .

Let

$$M_4 = \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}.$$

Then $\{M_4^i(\text{AG}(2, 4)) : 0 \leq i < 7\}$ is a set of seven mutually orthogonal affine planes of order 4.

With \mathbb{F}_4 generated by $x^2 + x + 1$ and primitive element $\zeta = \omega^5$, the planes $M_4^i(\text{AG}(2, 4))$ for $1 \leq i < 7$ correspond to the six pencil spreads of conics formed from the pairs

$$\begin{array}{ll} \phi_1 : x^2 + (\zeta + 1)yz + \zeta xz & \chi_1 : y^2 + yz + xz; \\ \phi_2 : x^2 + (\zeta + 1)yz + xz & \chi_2 : y^2 + (\zeta + 1)xz; \\ \phi_3 : x^2 + xz + \zeta xy & \chi_3 : y^2 + \zeta yz + \zeta xz; \\ \phi_4 : x^2 + \zeta yz + \zeta xz & \chi_4 : y^2 + yz + (\zeta + 1)xz; \\ \phi_5 : x^2 + yz + xz & \chi_5 : y^2 + yz + \zeta xz; \\ \phi_6 : x^2 + \zeta yz & \chi_6 : y^2 + xz. \end{array}$$

The union of the lines from all seven of the planes forms a SQS(16). A decomposition of an SQS(16) into affine planes was previously known [Mat97] but this is an interesting method of construction. It is unknown if the 1820 4-subsets of a 16-set can be partitioned into SQS(16)s [EZ20]; this would be a large set of Steiner Quadruple Systems. Because the cyclotomic cosets C_i are all disjoint, the collection of all cyclotomic cosets over these seven spreads forms the blocks of a resolvable Steiner triple system of order 15, a solution to Kirkman's schoolgirl problem of 1850 [Kir50]:

“Fifteen young ladies in a school walk out three abreast for seven days in succession: it is required to arrange them daily, so that no two shall walk twice abreast.”

This design is the derived design of the SQS(16) at the point 0. Because the planes have a transitive automorphism group, deriving at any other point gives an isomorphic resolvable STS(15).

Example 3.18. For $q = 8$, let \mathbb{F}_{64} be generated by a primitive element with minimum polynomial $x^6 + x + 1$. Construct the Desarguesian affine translation plane $AG(2, 8)$. Let

$$M_6 = \begin{bmatrix} 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & 0 \end{bmatrix}.$$

Then $\{M_6^i(AG(2, 8)) : 0 \leq i < 7\}$ is a set of seven mutually orthogonal affine planes of order 8.

Only three of these affine planes, those for $i = 1, 2, 4$, are formed from a pencil spread of conics. With \mathbb{F}_8 generated from $x^3 + x^2 + 1$ and ω as its root, the pencils of conics are formed from

$$\begin{array}{ll} \phi_1 : x^2 + \omega yz + \omega xz & \chi_1 : y^2 + (\omega^2 + \omega + 1)yz + (\omega + 1)xz; \\ \phi_2 : x^2 + (\omega^2 + \omega)yz + xz & \chi_2 : y^2 + yz + (\omega^2 + 1)xz; \\ \phi_4 : x^2 + (\omega^2)yz + (\omega + 1)xz + \omega xy & \chi_4 : y^2 + (\omega^2 + \omega)yz + (\omega^2 + \omega)xz. \end{array}$$

The spreads for $i = 3, 5, 6$ are spreads of conics but are not pencils.

The procedure to find suitable matrices $M \in GL(2n, \mathbb{F}_2)$ was implemented in C, and the clique-finding algorithm was implemented in *Mathematica*. The random search was modified in order to allow the C program to run in reasonable time for $q = 16$. Rather than seeking a suitable matrix M directly, we choose the first $2n - 1$ rows of M randomly and calculate its product with the first $2n - 1$ coordinates of each line S_i . If no four such products from the nonzero points of a line S_i occur in the set of products from a single line MS_i , then we proceed to test all possible final rows for M ; otherwise we discard the partial matrix M .

For $q = 8$, the C program running on a 2013 iMac desktop takes on average less than 0.2 milliseconds to find each suitable matrix M , and a set of 1200 such matrices was sufficient to find the first clique of size 7. We later refined it to the solution given in Example 3.18. No clique of size larger than 7 was found from a set of 50,000 such matrices.

For $q = 16$, the C program takes on average 15 seconds to find each suitable matrix M . A set of 1,000,000 such matrices was collected over a number of months on several machines. This set is too large for the clique-finding algorithm of *Mathematica* to handle, but it is straightforward to search in C for a pair of non-identity matrices M, M' from the set for which $M(AG(2, 16))$ is orthogonal to $M'(AG(2, 16))$; together with the identity matrix I_8 , this would give a set of three mutually orthogonal affine planes of order 16. Unfortunately, no such pair was found. The set of 1,000,000 suitable matrices does not contain an example M for which M^2 is also suitable. It

remains an open question as to whether there exists a set of more than two mutually orthogonal affine planes of orders 2^n with $\gcd(n, 6) > 1$ and $n > 3$.

We summarize the results of this subsection, using Corollary 1.3.

Theorem 3.19. • *The size of a largest set of mutually orthogonal affine planes in $\text{AG}(2, 4)$ is seven.*

• *There exists a set of seven mutually orthogonal affine planes in $\text{AG}(2, 8)$.*

4. Connections to covering arrays

Here we discuss the connection between sets of s mutually orthogonal planes, covering perfect hash families and covering arrays.

Definition 4.1. A *covering perfect hash family* $\text{CPHF}_\lambda(n; t, k, q)$ is an $n \times k$ array of elements from $\mathbb{F}_q^t \setminus \vec{0}$ (equivalently the points of $\text{PG}(t-1, q)$) such that, for each set T of t columns, there exist at least λ rows whose entries in the columns of T are linearly independent. If the vector entries all have non-zero last coordinate (equivalently, they are the points of the affine subspace of $\text{PG}(t-1, q)$) then the CPHF is a *Sherwood covering perfect hash family* $\text{SCPHF}_\lambda(n; t, k, q)$.

Definition 4.2. A *covering array* $\text{CA}_\lambda(N; t, k, v)$ is an $N \times k$ array of elements from a v -set such that, for each set T of t columns and each t -tuple from the v -set, there exist at least λ rows whose entries in the columns of T match the t -tuple.

Given a $\text{CA}_\lambda(N; t, k, v)$, we can add arbitrary rows and drop arbitrary columns to get a $\text{CA}_\lambda(N'; t, k', v)$ when $N \leq N'$ and $k \geq k'$. So we say a $\text{CA}_\lambda(N; t, k, v)$ is better than a $\text{CA}_\lambda(N'; t, k', v)$ if $N \leq N'$ and $k \geq k'$, provided $(N, k) \neq (N', k')$. (In the application of covering arrays to testing we want the smallest number of rows for a given number of columns; the rows are a cost while the columns and strength provide value.) Very few provably optimal covering arrays are known, and we only have the “best currently known” covering array for many parameters t, k , and v .

Covering perfect hash families are used to construct covering arrays. We extend the standard constructions [CLS18, SMC06] by considering the index λ of the constructed covering array.

Theorem 4.3. *Suppose $\lambda' \leq \lambda$. If C is a $\text{CPHF}_\lambda(n; t, k, q)$, there exists a $\text{CA}_{\lambda'}(n(q^t - 1) + \lambda'; t, k, q)$. If C is an $\text{SCPHF}_\lambda(n; t, k, q)$, there exists a $\text{CA}_{\lambda'}(n(q^t - q) + \lambda'q; t, k, q)$*

Proof. See [CLS18, SMC06] for the basic proof. The extension to higher index is straightforward. For a given set T of t columns, any row in which the entries from T are linearly independent contributes one copy of every t -tuple over \mathbb{F}_q to the covering array, so every t -tuple appears at least λ times. Every row of a CPHF contributes an all-zero row to the covering array. Every row of a SCPHF contributes each of the q constant rows to the covering array. In each case all but λ' of them can be deleted. \square

Theorem 4.4. *Suppose that a set of s mutually orthogonal Desarguesian projective (affine) planes exists. Then there exists a $\text{CPHF}_{s-1}(s; 3, q^2 + q + 1, q)$ ($\text{SCPHF}_{s-1}(s; 3, q^2, q)$).*

Proof. Represent the points of $PG(2, q)$ as length 3 vectors in homogeneous coordinates over \mathbb{F}_q and the points of $AG(2, q)$ as length 3 vectors in homogeneous coordinates over \mathbb{F}_q with last entry 1. For each i satisfying $1 \leq i \leq s$, let ϕ_i be an isomorphism of the i th plane in the set onto the first plane.

Let C be the array with s rows and columns indexed by the elements of the common point set of the planes. Define the entries of C by

$$C_{iz} = \phi_i(z).$$

Let T be a set of three columns corresponding to points z_1, z_2 and z_3 . When these three points are non-collinear in the plane i , the corresponding entries in row i are linearly independent. If z_1, z_2 and z_3 appear on a line in one of the planes, then by the definition of orthogoval they must be non-collinear in all other $s - 1$ planes. Thus there are at least $s - 1$ rows with linearly independent entries in columns z_1, z_2 and z_3 . \square

The covering arrays constructed from these Sherwood covering perfect hash families can be extended by an additional two columns.

Theorem 4.5. *Suppose that a set of s mutually orthogoval Desarguesian affine planes exists. Then there exists a $CA_{s-1}(sq^3 - q; 3, q^2 + 2, q)$.*

Proof. The affine plane embeds uniquely into the projective plane so we can extend the set of s mutually orthogoval affine planes to a set of s projective planes which are orthogoval *except* for the line z . From these we construct the $s \times (q^2 + q + 1)$ array C as in the proof of Theorem 4.4 defined by

$$C_{iz} = \phi_i(z).$$

Because the planes have the property that any three points not *all* on the line z which are collinear in one are non-collinear in all the others, if we delete all but two of the columns of C which are indexed by the points on the line z , the resulting array is a $CPHF_{s-1}(s; 3, q^2 + 2, q)$.

When constructing the covering array from this extended SCPHF we observe that the repeated rows arise from the multiplication of r with the entry z in C where the vectors r are the vectors with only the last position non-zero, see [CLS18]. The multiplication of these vectors r with the two points kept from the line z ($(1 : 0 : 0)$ and $(0 : 1 : 0)$ for example) yield all zeros so the rows are still repeated in the covering array extended by two columns and the appropriate number can still be deleted. \square

From our constructions of sets of mutually orthogoval Desarguesian projective planes in Theorems 2.2 and 2.3 we can build the corresponding covering perfect hash families and covering arrays.

Corollary 4.6. • *For all prime powers q there exists a $CA(2q^3 - 1; 3, q^2 + q + 1, q)$.*

• *For $\lambda < 4$ there exists a $CA_\lambda(27\lambda + 26; 3, 13, 3)$.*

From our constructions of sets of mutually orthogoval Desarguesian affine planes summarized in Corollary 3.9 and Theorem 3.19 we can build the corresponding covering perfect hash families and covering arrays.

Corollary 4.7. • *For all prime powers $q = 2^n$ there exists a $CA(2q^3 - q; 3, q^2 + 2, q)$.*

- *For all prime powers $q = 2^n$ with $\gcd(n, 6) = 1$ and $\lambda < 3$, there exists a $CA_\lambda(\lambda q^3 + q^3 - q; 3, q^2 + 2, q)$.*
- *For $\lambda < 7$ there exists a $CA_\lambda(27\lambda + 24; 3, 11, 3)$.*
- *For $\lambda < 7$ there exists a $CA_\lambda(64\lambda + 60; 3, 18, 4)$.*
- *For $\lambda < 7$ there exists a $CA_\lambda(512\lambda + 504; 3, 66, 8)$.*

We believe these are the best known constructions for higher index covering arrays (that is, with $\lambda > 1$) in the instances where the sets of mutually orthogonal planes exist.

5. Final thoughts

In this paper we introduced the concept of orthogonal planes as a generalization of a long-known construction of a pair of projective planes constructed from a difference set and its negative. There are several known constructions for a pair of orthogonal projective planes. We gave a new construction using the Cremona transformation in the proof of Theorem 2.2, illustrating that multiple points of view can lead to a deeper understanding of the richness of this area. A few instances of sets of mutually orthogonal affine planes were previously known; we have derived some specific large sets as well as some infinite classes of pairs or triples. We exhibited a connection between sets of orthogonal affine planes and covering arrays and described the resulting new covering arrays. We believe there is much further potential in taking advantage of the interplay between the various areas of mathematics we have drawn on in this paper.

Many open problems remain. Foremost among them is to determine the largest size of a set of mutually orthogonal planes of order q for all prime powers q . We have some exact values for small q in Theorems 2.3 and 3.19. We have the upper bound, usually $q + 2$, of Theorem 1.2 and the lower bound of 2 from Theorem 2.2 and Corollary 3.9, and for some q we have a lower bound of 3 affine planes from Corollary 3.16. There is a large gap between the lower and upper bounds for all but the smallest q .

A secondary question is how many essentially different maximal sets of mutually orthogonal planes of order q exist. Even in the case of the pairs or triples of mutually orthogonal planes constructed in this paper, we do not know if the constructed sets are non-isomorphic.

It may be possible to use Theorem 2.2 to find sets of more than two mutually orthogonal planes. If we pick two different sets of three conjugate points, we can use each to build other planes on the point set of the standard plane that are each orthogonal to the standard plane. The transformation from the second to third plane is just the composition of the two Cremona transformations that formed each. If this composition is itself a conjugated Cremona then we may have constructed a set of three mutually orthogonal projective planes.

An alternative approach to construct more than two orthogonal projective planes is to use difference sets. Baker et al. [BBEF94] prove that if $D \subset \mathbb{Z}_{n^2+n+1}$ is a difference set, then $-D$, $2D$ and $D/2$ are all difference sets that are conics. These could potentially be used to create a

set of three mutually orthogonal planes of orders other than 3. For example $-D/2 = 2D \neq D$ if and only if -4 is a multiplier of D and 2 is not. In this case, all three of D , $-D$ and $2D$ must be distinct and are each related to each other by multipliers of -1 , 2 or $1/2$ and thus form ovals in the planes generated by the other. Gordon, Mills and Welch showed that the only multipliers of a Desarguesian planar difference set in \mathbb{Z}_{q^2+q+1} for $q = p^e$ are the powers of p in \mathbb{Z}_{q^2+q+1} [GMW62]. Thus if there exists $q = p^e$ such that $p^d \equiv -4 \pmod{q^2 + q + 1}$ and 2 is not generated by p , then there exists a set of three Desarguesian orthogonal projective planes of order q . Unfortunately, 3 is the only prime power up to $10,000,000$ for which this is true.

It is also natural to generalize the definition of orthogonal to other geometric objects. Possibilities that are intriguing include inversive planes and higher dimensional projective and affine spaces.

Acknowledgements

We would like to thank Erin Lanus and Kaushik Sarkar. They found an SCPHF(2; 3, 8, 64) in a conditional expectation search. This turned out to have the structure of a pair of orthogonal affine planes of order 8 and discussion of this structure in several venues was the impetus for the research presented in this article. We thank Stefaan De Winter who pointed out that the set of seven mutually orthogonal affine planes of order 4 constructs a Kirkman triple system of order 15.

References

- [Bay17] S. Bays. Une question de Cayley relative au problème des triades de Steiner. *Enseignement Math.*, 19:57–67, 1917.
- [BBEF94] R. D. Baker, J. M. N. Brown, G. L. Ebert, and J. C. Fisher. Projective bundles. *Bull. Belg. Math. Soc. Simon Stevin*, 1(3):329–336, 1994. doi:10.36045/bbms/1103408578.
- [BGG03] Darryn Bryant, Mike Grannell, and Terry Griggs. Large sets of large sets of Steiner triple systems of order 9. *Util. Math.*, 64:115–118, 2003.
- [BH01] Iliya Bluskov and Katherine Heinrich. Super-simple designs with $v \leq 32$. *Journal of Statistical Planning and Inference*, 95(1):121–131, 2001. doi:10.1016/S0378-3758(00)00282-2.
- [Bru73] R. H. Bruck. Circle geometry in higher dimensions. II. *Geometriae Dedicata*, 2:133–188, 1973. doi:10.1007/BF00147854.
- [CC04] Wensong Chu and Charles J. Colbourn. Optimal $(n, 4, 2)$ -OOC of small orders. *Discrete Math.*, 279(1-3):163–172, 2004. In honour of Zhu Lie. doi:10.1016/S0012-365X(03)00266-8.
- [CD07] Charles J. Colbourn and Jeffrey H. Dinitz, editors. *Handbook of combinatorial designs*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, second edition edition, 2007.

- [Cho83] Leo G. Chouinard, II. Partitions of the 4-subsets of a 13-set into disjoint projective planes. *Discrete Math.*, 45(2-3):297–300, 1983. doi:10.1016/0012-365x(83)90045-6.
- [CLS18] Charles J. Colbourn, Erin Lanus, and Kaushik Sarkar. Asymptotic and constructive methods for covering perfect hash families and covering arrays. *Des. Codes Cryptogr.*, 86(4):907–937, 2018. doi:10.1007/s10623-017-0369-x.
- [Col99] Charles J. Colbourn. A Steiner 2-design with an automorphism fixing exactly $r + 2$ points. *Journal of Combinatorial Designs*, 7(5):375–380, 1999. doi:10.1002/(SICI)1520-6610(1999)7:5<375::AID-JCD6>3.0.CO;2-K.
- [CR99] Charles J. Colbourn and Alexander Rosa. *Triple systems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1999.
- [Dem97] Peter Dembowski. *Finite geometries*. Springer-Verlag, Berlin, 1997.
- [EZ20] Tuvi Etzion and Junling Zhou. An improved recursive construction for disjoint Steiner quadruple systems. *J. Combin. Des.*, 28(8):551–567, 2020. doi:10.1002/jcd.21713.
- [Ful89] William Fulton. *Algebraic curves*. Advanced Book Classics. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989. An introduction to algebraic geometry, Notes written with the collaboration of Richard Weiss, Reprint of 1969 original.
- [Gly78] David G. Glynn. *Finite Projective Planes and Related Combinatorial Systems*. Ph.D., University of Adelaide, Australia, 1978.
- [GMW62] Basil Gordon, W. H. Mills, and L. R. Welch. Some new difference sets. *Canadian J. Math.*, 14:614–625, 1962. doi:10.4153/cjm-1962-052-2.
- [GS94] D.G. Glynn and G.F. Steinke. Pencils of translation ovals in translation planes. *Geom Dedicata*, 51:113–131, 1994. doi:10.1007/BF01265323.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.
- [Hir98] J. W. P. Hirschfeld. *Projective Geometries over Finite Fields*. Oxford Mathematical Monographs. Oxford University Press, Oxford, New York, 1998.
- [Kir50] T. P. Kirkman. Query VI. *The Lady's and Gentleman's Diary*, page 48, 1850.
- [Mat97] Rudolf Mathon. Searching for spreads and packings. In *Geometry, combinatorial designs and related structures (Spetses, 1996)*, volume 245 of *London Math. Soc. Lecture Note Ser.*, pages 161–176. Cambridge Univ. Press, Cambridge, 1997.
- [MH72] N. S. Mendelsohn and Stephen H. Y. Hung. On the Steiner systems $S(3, 4, 14)$ and $S(4, 5, 15)$. *Utilitas Math.*, 1:5–95, 1972.
- [RMS14] Sebastian Raaphorst, Lucia Moura, and Brett Stevens. A construction for strength-3 covering arrays from linear feedback shift register sequences. *Des. Codes Cryptogr.*, 73(3):949–968, 2014. doi:10.1007/s10623-013-9835-2.
- [Sha13] Igor R. Shafarevich. *Basic algebraic geometry. I*. Springer, Heidelberg, third edition, russian edition, 2013. Varieties in projective space.

- [She86] F. A. Sherk. The geometry of $\text{GF}(q^3)$. *Canadian Journal of Mathematics*, 38(3):672–696, 1986. doi:10.4153/CJM-1986-035-2.
- [SMC06] George B. Sherwood, Sosina S. Martirosyan, and Charles J. Colbourn. Covering arrays of higher strength from permutation vectors. *J. Combin. Des.*, 14(3):202–213, 2006. doi:10.1002/jcd.20067.
- [Tei73] Luc Teirlinck. On the maximum number of disjoint Steiner triple systems. *Discret. Math.*, 6:299–300, 1973. doi:10.1016/0012-365x(73)90101-5.
- [The21] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.4)*, 2021. Software and documentation available from <https://www.sagemath.org>.
- [TJIM18] Jose Torres-Jimenez and Idelfonso Izquierdo-Marquez. Covering arrays of strength three from extended permutation vectors. *Des. Codes Cryptogr.*, 86(11):2629–2643, November 2018. doi:10.1007/s10623-018-0465-6.