

Lawrence Berkeley National Laboratory

Nuclear Science

Title

Form Factors and Two-Photon Exchange in High-Energy Elastic Electron-Proton Scattering

Permalink

<https://escholarship.org/uc/item/6qw5c53b>

Authors

Christy, ME

Gautam, T




Ou, L

et al.

Publication Date

2021-03-02

Form factors and two-photon exchange in high-energy elastic electron-proton scattering

M. E. Christy,¹ T. Gautam,¹ L. Ou,² B. Schmookler,² Y. Wang,³ D. Adikaram,⁴ Z. Ahmed,⁵ H. Albatineh,⁶ S. F. Ali,⁷ B. Aljawrneh,^{8,9} K. Allada,² S.L. Allison,¹⁰ S. Alsalmi,¹¹ D. Androic,¹² K. Aniol,¹³ J. Annand,¹⁴ J. Arrington,^{15,16} H. Atac,¹⁷ T. Averett,³ C. Ayerbe Gayoso,³ X. Bai,¹⁸ J. Bane,¹⁹ S. Barcus,³ K. Bartlett,³ V. Bellini,²⁰ R. Beminiwattha,²¹ J. Bericic,⁴ H. Bhatt,²² D. Bhetuwal,²² D. Biswas,¹ E. Brash,²³ D. Bulumulla,¹⁰ C. M. Camacho,²⁴ J. Campbell,²⁵ A. Camsonne,⁴ M. Carmignotto,⁷ J. Castellanos,²⁶ C. Chen,¹ J-P. Chen,⁴ T. Chetry,²⁷ E. Cisbani,²⁸ B. Clary,²⁹ E. Cohen,³⁰ N. Compton,²⁷ J. C. Cornejo,^{3,31} S. Covrig Dusa,⁴ B. Crowe,³² S. Danagoulian,⁸ T. Danley,²⁷ W. Deconinck,³ M. Defurne,³⁰ C. Desnault,²⁴ D. Di,¹⁸ M. Dlamini,²⁷ M. Duer,³⁴ B. Duran,¹⁷ R. Ent,⁴ C. Fanelli,² E. Fuchey,²⁹ C. Gal,¹⁸ D. Gaskell,⁴ F. Georges,³⁵ S. Gilad,² O. Glamazdin,³⁶ K. Gnanvo,¹⁸ A. V. Gramolin,³⁷ V. M. Gray,³ C. Gu,¹⁸ A. Habarakada,¹ T. Hague,¹¹ G. Hamad,²⁷ D. Hamilton,¹⁴ K. Hamilton,¹⁴ O. Hansen,⁴ F. Hauenstein,¹⁰ A. V. Hernandez,⁷ W. Henry,¹⁷ D. W. Higinbotham,⁴ T. Holmstrom,³⁸ T. Horn,⁷ Y. Huang,¹⁸ G.M. Huber ,⁵ C. Hyde,¹⁰ H. Ibrahim,³⁹ N. Israel,²⁷ C-M. Jen,⁴⁰ K. Jin,¹⁸ M. Jones,⁴ A. Kabir,¹¹ B. Karki,²⁷ C. Keppel,⁴ V. Khachatryan,^{41,42} P.M. King,²⁷ S. Li,⁴³ W. Li,⁵ H. Liu,⁴⁴ J. Liu,¹⁸ A. H. Liyanage,¹ D. Mack,⁴ J. Magee,³ S. Malace,⁴ J. Mammei,⁴⁵ P. Markowitz,²⁶ S. Mayilyan,⁴⁶ E. McClellan,⁴ F. Meddi,²⁸ D. Meekins,⁴ K. Mesick,⁴⁷ R. Michaels,⁴ A. Mkrtchyan,⁷ B. Moffit,⁴ R. Montgomery,¹⁴ L.S. Myers,⁴ P. Nadel-Turonski,⁴ S. J. Nazeer,¹ V. Nelyubin,¹⁸ D. Nguyen,¹⁸ N. Nuruzzaman,¹ M. Nycz,¹¹ R.F. Obrecht,²⁹ K. Ohanyan,⁴⁶ C. Palatchi,¹⁸ B. Pandey,¹ K. Park,¹⁰ S. Park,⁴¹ C. Peng,⁴⁸ F. D. Persio,²⁸ R. Pomatsalyuk,³⁶ E. Pooser,⁴ A.J.R. Puckett,²⁹ V. Punjabi,⁴⁹ B. Quinn,³¹ S. Rahman,⁴⁵ M.N.H. Rashad,¹⁰ P.E. Reimer ,¹⁶ S. Riordan,⁴¹ J. Roche,²⁷ I. Sapkota,⁷ A. Sarty,⁵⁰ B. Sawatzky,⁴ N. H. Saylor,⁵¹ M. H. Shabestari,²² A. Shahinyan,⁴⁶ S. Širca,⁵² G.R. Smith,⁴ S. Sooriyaarachchilage,¹ N. Sparveris,¹⁷ R. Spies,⁴⁵ A. Stefanko,³¹ T. Su,¹¹ A. Subedi,²² V. Sulkosky,² A. Sun,³¹ Y. Tan,⁵³ L. Thorne,³¹ N. Ton,¹⁸ F. Tortorici,²⁰ R. Trotta,⁵⁴ R. Uniyal,⁷ G.M. Urciuoli,²⁸ E. Voutier,²⁴ B. Waidyawansa,⁴ B. Wojtsekhowski ,^{4,*} S. Wood,⁴ X. Yan,⁵⁵ L. Ye,²² Z. H. Ye,^{18,56} C. Yero,²⁶ J. Zhang,¹⁸ Y. X. Zhao,⁴¹ and P. Zhu⁵⁷

¹Hampton University, Hampton, Virginia 23669, USA

²Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

³William and Mary, Williamsburg, Virginia 23185, USA

⁴Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

⁵University of Regina, Regina, SK, S4S 0A2 Canada

⁶Texas A & M University, Kingsville, Texas 77843, USA

⁷Catholic University of America, Washington, District of Columbia 20064, USA

⁸North Carolina A&T State University, Greensboro, North Carolina 27411, USA

⁹Al Zaytoonah University of Jordan, Amman 117733, Jordan

¹⁰Old Dominion University, Norfolk, Virginia 23529, USA

¹¹Kent State University, Kent, Ohio 44240, USA

¹²University of Zagreb, Trg Republike Hrvatske 14, 10000, Zagreb, Croatia

¹³California State University, Los Angeles, Los Angeles, California 90032, USA

¹⁴SUPA School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, United Kingdom

¹⁵Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

¹⁶Argonne National Laboratory, Lemont, Illinois 60439, USA

¹⁷Temple University, Philadelphia, Pennsylvania 19122, USA

¹⁸University of Virginia, Charlottesville, Virginia 22904, USA

¹⁹University of Tennessee, Knoxville, Tennessee 37996, USA

²⁰Istituto Nazionale di Fisica Nucleare, Dipt. di Fisica dell Univ. di Catania, I-95123 Catania, Italy

²¹Syracuse University, Syracuse, New York, 13244, USA

²²Mississippi State University, Mississippi State, Mississippi 39762, USA

²³Christopher Newport University, Newport News, Virginia 23606, USA

²⁴Institut de Physique Nucleaire, 15 Rue Georges Clemenceau, 91400 Orsay, France

²⁵Dalhousie University, Nova Scotia, NS B3H 4R2, Canada

²⁶Florida International University, Miami, Florida 33199, USA

²⁷Ohio University, Athens, Ohio 45701, USA

²⁸Istituto Nazionale di Fisica Nucleare - Sezione di Roma, P.le Aldo Moro, 2 - 00185 Roma, Italy

²⁹University of Connecticut, Storrs, Connecticut 06269, USA

³⁰Tel Aviv University, Tel Aviv-Yafo 69978, Israel

³¹Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

³²North Carolina Central University, Durham, North Carolina 27707, USA

³³CEA Saclay, 91191 Gif-sur-Yvette, France

³⁴Tel Aviv University, Tel Aviv-Yafo, Israel

³⁵Ecole Centrale Paris, 3 Rue Joliot Curie, 91190 Gif-sur-Yvette, France

³⁶Kharkov Institute of Physics and Technology, Kharkov 61108, Ukraine

³⁷Boston University, Boston, Massachusetts 02215, USA

³⁸Randolph Macon College, Ashland, Virginia 23005, USA

³⁹Cairo University, Cairo, 12613, Egypt

⁴⁰Virginia Polytechnic Inst. & State Univ., Blacksburg, Virginia 234061, USA

⁴¹Stony Brook, State University of New York, New York 11794, USA

⁴²Cornell University, Ithaca, New York 14853, USA

⁴³University of New Hampshire, Durham, New Hampshire 03824, USA

⁴⁴Columbia University, New York, New York 10027, USA

⁴⁵University of Manitoba, Winnipeg, MB R3T 2N2, Canada

⁴⁶AANL, 2 Alikhanian Brothers Street, 0036, Yerevan, Armenia

⁴⁷Rutgers University, New Brunswick, New Jersey 08854, USA

⁴⁸Duke University, Durham, North Carolina 27708, USA

⁴⁹Norfolk State University, Norfolk, Virginia 23504, USA

⁵⁰Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada

⁵¹Rensselaer Polytechnic Institute, Troy, New York 12180, USA

⁵²Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia

⁵³Shandong University, Shandong, Jinan 250100, China

⁵⁴Duquesne University, Pittsburgh, Pennsylvania 15282, USA

⁵⁵Huangshan University, 44 Daizhen Road, Tunxi District, Huangshan, Anhui Province, China

⁵⁶Tsinghua University, 30 Shuangqing Rd, Haidian District, Beijing 100190, China

⁵⁷University of Science and Technology of China, Hefei, Anhui 230026, China

(Dated: March 23, 2022)

We present new precision measurements of the elastic electron-proton scattering cross section for momentum transfer (Q^2) up to 15.75 (GeV/c)^2 . Combined with existing data, these provide an improved extraction of the proton magnetic form factor at high Q^2 and double the range over which a longitudinal/transverse separation of the cross section can be performed. The difference between our results and polarization data agrees with that observed at lower Q^2 and attributed to hard two-photon exchange (TPE) effects, extending to 8 (GeV/c)^2 the range of Q^2 for which a discrepancy is established at $>95\%$ confidence. We use the discrepancy to quantify the size of TPE contributions needed to explain the cross section at high Q^2 .

PACS numbers: 25.30.Bf, 13.40.Gp, 14.20.Dh

Elastic electron scattering is a key process used in studies of matter across a wide range of energy scales and in many sub-fields of physics. In the one-photon exchange approximation (OPE), first calculated in Ref. [1], the differential electron-nucleon elastic scattering cross section, $d\sigma(\theta_e)/d\Omega_e$, is the product of the cross section for a structureless object and a structure-dependent term that depends on the Sachs magnetic and electric form factors [2], $G_M(Q^2)$ and $G_E(Q^2)$, which encode the spatial distributions of magnetization and charge in the proton:

$$\frac{d\sigma(\theta_e)}{d\Omega_e} = \frac{d\sigma_{Mott}}{d\Omega_e} \cdot \frac{\tau G_M^2(Q^2) + \varepsilon G_E^2(Q^2)}{\varepsilon(1 + \tau)}. \quad (1)$$

In Eq. (1), θ_e is the scattering angle of the electron, $d\sigma_{Mott}/d\Omega_e$ is the cross section for scattering of an electron with incident (scattered) energy E_e (E'_e) from a structureless target, $Q^2 = 4E_e E'_e \sin^2(\theta_e/2)$ is the negative four-momentum transfer squared, $\varepsilon \equiv [1 + 2(1 + \tau) \tan^2(\theta_e/2)]^{-1}$ is the virtual photon polarization parameter, and $\tau \equiv Q^2/4M_p^2$. The structure-dependent term is isolated in the reduced cross section,

$$\begin{aligned} \sigma_R &= \tau G_M^2(Q^2) + \varepsilon G_E^2(Q^2) = \sigma_T + \varepsilon \sigma_L \\ &= G_M^2(Q^2)(\tau + \varepsilon \text{RS}(Q^2)/\mu_p^2), \end{aligned} \quad (2)$$

where σ_L and σ_T are the longitudinal and transverse contributions to the cross section, respectively, $\text{RS} = (\mu_p G_E/G_M)^2$ is the normalized Rosenbluth slope, and μ_p is the proton magnetic moment. The form factors can be extracted using measurements at fixed Q^2 but different values of ε , corresponding to different electron scattering angles. A linear fit to measurements of $\sigma_R(\varepsilon)$ yields an intercept of $\sigma_R(\varepsilon = 0) = \tau G_M^2$, and a slope of $d\sigma_R/d\varepsilon = G_E^2$. This method is commonly known as Rosenbluth or Longitudinal/Transverse (L/T) separation.

Pioneering measurements of elastic electron-proton scattering by R. Hofstadter [3] confirmed the theoretical expectation of linear dependence of σ_R as a function of ε , which supported the use of the OPE approximation. Theoretical studies of the effects beyond OPE were performed soon after that [4, 5]. Early experimental searches using recoil proton polarization [6] and lepton charge asymmetry [7, 8] failed to find significant deviations from the OPE approximation. A number of precision measurements, including one in Ref. [9], extended linearity tests up to $Q^2 = 3 \text{ (GeV/c)}^2$ and demonstrated that G_E and G_M both approximately follow the dipole form $G_D \equiv (1 + Q^2/\Lambda^2)^{-2}$, with $\Lambda^2 = 0.71 \text{ GeV}^2$, yielding form factor scaling: $\mu_p G_E/G_M \approx 1$. At larger Q^2 values, τ enhances the contribution from G_M^2 to the cross section, making it difficult to extract G_E^2 . Analyses at higher Q^2 values [10, 11] extracted G_M under the assumption that $\text{RS} = 1$, and found that $Q^4 G_M(Q^2)$ was Q^2 -independent above 10 (GeV/c)^2 , consistent with pQCD predictions [12].

The reduced sensitivity to G_E^2 at high Q^2 in the Rosen-

* Contact person, bogdanw@jlab.org

bluth method motivated the use of double polarization observables [13], for which the OPE formalism was developed in Refs. [13–16]. Polarization measurements are directly sensitive to the ratio G_E/G_M , but not to the individual form factors. About 20 years ago the first precision measurements of G_E/G_M for Q^2 up to several $(\text{GeV}/c)^2$ were performed using the polarization transfer method [17] and a novel effect was discovered: the form factor ratio (FFR) extracted from polarization data decreased dramatically with Q^2 [17–19].

The decrease of the FFR with increasing Q^2 , an unexpected effect, implied a significant reduction of G_E , with theoretical explanations ranging from the role of quark orbital momentum [20] to the effect of the diquark correlation in the nucleon ground state [21]. In addition, the difference between the FFR extracted from the polarization measurements and from the cross section results was surprising, and requires deeper understanding. This difference is referred to henceforth as the form factor ratio puzzle (FFRP).

A reanalysis of the world data on RS [22], and new measurements of RS values with both scattered electron detection [23] and recoil proton detection [24], confirmed with improved precision the original observation of form factor scaling, enhancing the FFRP. Assuming that there are no unexpected errors with the now extensive body of Rosenbluth and polarization measurements, and that the radiative corrections (RC) applied are complete (except for the excluded hard TPE contributions), the only remaining explanation within the Standard Model is two-photon-exchange (TPE) or higher-order corrections. The hard TPE contributions are defined in this context as the TPE terms omitted in conventional radiative correction procedures which include only the IR-divergent terms, meaning that the definition of hard TPE depends slightly on the RC prescription [25].

An analysis of world data found that non-linearities in the reduced cross section as a function of ε , indicating deviations from the OPE picture, were extremely small [26], although the lack of non-linear contributions does not rule out a change to the slope that could explain the FFRP. At large Q^2 values, where the slope arising from G_E^2 is small, even a tiny change of RS can modify the extraction of G_E/G_M significantly.

TPE cross section contributions have the opposite sign for electron and positron scattering, making a comparison of e^+p and e^-p scattering one of the most direct tests for TPE. A global re-examination of electron/positron scattering comparisons in 2003 showed evidence for TPE [27] at low Q^2 values. After 2010, new experiments were performed to improve the precision and extend the kinematic range of these comparisons [28–30], observing clear hard TPE in the ratio of e^+p and e^-p elastic scattering up to $Q^2 \approx 2 (\text{GeV}/c)^2$. Finally, the contribution of TPE to polarization transfer observables was found to be small [31], as predicted by calculations [32–34]. Given this empirical understanding, the discrepancy (FFRP) is taken in our study as a measure of the TPE impact on the cross section, as in Refs. [22, 35, 36].

While most examinations of the FFRP focus on hard TPE, any ε -dependent correction would contribute to the discrepancy, leading to new examinations of the full radiative correction procedures [25, 37, 38]. The most recent and com-

plete update [38] was applied to SLAC data [9, 39], yielding a reduced discrepancy still providing a clear confirmation of the FFRP for Q^2 from 4-7 $(\text{GeV}/c)^2$. Notwithstanding these experimental and theoretical efforts, a full calculation of the TPE contribution is still not available mostly due to its dependence on the hadron structure of the intermediate states (see reviews [40–42]).

This work provides new experimental data at very large Q^2 , addresses the significance of the FFRP at much higher values of Q^2 than previously investigated, provides reanalyzed RC for six previous experiments, and improves the precision of experimental constraints on TPE effects in elastic $e-p$ scattering. Our new low- ε data, combined with existing high- ε measurements [10, 11, 39, 43], provide new Rosenbluth separations of G_E and G_M above 7 $(\text{GeV}/c)^2$ and significantly improved precision in the extraction of G_M . Our new data also provide an important baseline for high- Q^2 measurements enabled by the 12 GeV upgrade at Jefferson Lab, where precise knowledge of the elastic cross section is needed for experimental normalization and cross checks, and as input to the broader program of high- Q^2 proton and neutron structure measurements.

This experiment, referred to hereafter as GMp12, was performed in Hall A of Jefferson Lab using the basic suite of experimental instrumentation [44]. A 100% duty-factor electron beam with current up to 68 μA and energy from 2.2 to 11 GeV was incident on a 15-cm long liquid hydrogen target. The target operated at a temperature of 19 K, a pressure of 25 psia, and a density of 0.0732 g/cm^3 . The hydrogen target was complemented by a “dummy” target consisting of two aluminum foils, used to measure and subtract events originating from the entrance and exit windows of the hydrogen cell. The target density reduction with increasing beam intensity, due to localized boiling of the cryogen, was found to be 2.7% per 100 μA , [45], with an uncertainty in the variation across the current range of the experiment of 0.35%

The energy of incident electrons was determined using the Hall A ARC energy measurement system, [46], which measures the field integral of the dipoles which bend the beam through 34.257 degrees from the accelerator into Hall A. These results were cross checked with spin precession studies and beam energy measurements in Hall C. The uncertainty in the beam energy was found to be less than 0.1% for all kinematics [47]. The beam current was measured by beam charge monitors (BCMs) [48], which were calibrated against a well-understood Unser monitor [49]. The uncertainty on the beam current and accumulated charge was defined by the accuracy of the BCM calibration. An absolute uncertainty of 0.06 μA stems from the current source utilized to calibrate the Unser monitor. The latter results in an uncertainty of 0.1% at a current of 65 μA , utilized for most of the GMp12 kinematics, and up to a maximum of 0.6% for the lowest current of 10 μA .

The scattered electrons were detected in the left and right Hall A High Resolution Spectrometers (LHRS and RHRS, respectively), with the central momentum of the spectrometers set to detect elastically scattered electrons. The HRSs have a solid angle acceptance of 6.0 msr, momentum acceptance of $\pm 4.5\%$, intrinsic momentum resolution of 2.5×10^{-4} , and an-

gular resolution of 0.6 mrad. The primary trigger was formed as a coincidence of signals in the front and back scintillator planes (separated by two meters) and the gas Cherenkov counter. The trigger efficiency was monitored using a sample of triggers that required only two of these three signals. For this experiment, the tracking system in each HRS was upgraded by adding a three-layer straw tube drift chamber to allow accurate determination of the track reconstruction efficiency [50]. The particle identification detectors included a two-layer shower detector and a gas Cherenkov counter with enhanced light collection efficiency by means of a wavelength shifter [51]. Dead times of the trigger counters, front-end electronics, and DAQ were constantly measured using pulser generated events [45, 52]. The uncorrelated systematic uncertainty of the GMp12 cross section data is 1.2-1.3%, while the overall normalization uncertainty is 1.6% (2.0%) for the LHRS (RHRS) data. A more detailed breakdown and discussion of the main systematic uncertainties is presented in the supplemental material [53].

Full simulations of the incident electron-target interaction and the electron trajectory through the HRS magnets and detectors were performed for each kinematic setting using an updated version of the magnetic optics Monte Carlo code [54] incorporating the HRSs. The event distributions in the detector package were compared with the simulated data and used to fine-tune the model of the HRS optical transport. Radiative processes were implemented using the approach built into the Monte Carlo simulation, described in Ref. [55], based on an updated implementation [9] of the RC formalism of Ref. [56]. The resulting cross section values from GMp12 were then adjusted to account for the difference between the prescription above and the RC calculation of Refs. [25, 37, 38] which has the most accurate evaluation of the internal and external radiation. This is essential for the analysis in the present work, as it resolves some of the discrepancy seen in past comparisons based on older radiative correction procedures.

The kinematics and reduced cross section results from the GMp12 experiment are shown in Table I.

We combine our results with cross sections from several JLab and SLAC experiments [9–11, 23, 39, 43] spanning a Q^2 range of 0.4-31 (GeV/c) 2 in a global fit of the Q^2 and ϵ dependence of the elastic cross section using Eq. 2. These experiments, comprising 121 kinematic points, were chosen because the publications provide sufficient information on their RC procedures and cutoffs to allow us to self-consistently implement the RC modification [38]. The normalizations of the data for the individual experiments were allowed to vary based on their quoted normalization uncertainties, except for the data of Ref. [23], which cover a wide range of Q^2 with the best accuracy. The cross sections were fit in terms of G_M and RS with the following simple parametrization:

$$\begin{aligned} G_M &= \mu_p (1 + a_1 \tau) / (1 + b_1 \tau + b_2 \tau^2 + b_3 \tau^3), \\ RS &= 1 + c_1 \tau + c_2 \tau^2. \end{aligned} \quad (3)$$

The fit gives $\chi^2 = 88.7$ for 107 degrees of freedom; the parameters and uncertainties are given in Tab. II. The cross section database and the full covariance matrix of the fit parameters are given in the supplemental material [53].

TABLE I. Kinematics and reduced cross sections for GMp12, with statistical and point-to-point systematic uncertainties added in quadrature. Points labeled with an asterisk (*) were taken with the RHRS. There is an additional 1.6% (2.0%) normalization uncertainty for the LHRS (RHRS) data.

E_e (GeV)	θ_e (deg)	Q^2 (GeV/c) 2	ϵ	σ_R (Eq. 2)
2.222	42.001	1.577	0.701	$(4.273 \pm 0.040) \times 10^{-2}$
2.222*	48.666	1.858	0.615	$(2.983 \pm 0.057) \times 10^{-2}$
6.427	24.250	4.543	0.826	$(3.813 \pm 0.057) \times 10^{-3}$
6.427	30.909	5.947	0.709	$(1.805 \pm 0.025) \times 10^{-3}$
6.427	37.008	6.993	0.599	$(1.113 \pm 0.016) \times 10^{-3}$
6.427	44.500	7.992	0.478	$(7.289 \pm 0.109) \times 10^{-4}$
8.518	30.909	9.002	0.648	$(5.163 \pm 0.078) \times 10^{-4}$
6.427*	55.900	9.053	0.332	$(4.859 \pm 0.107) \times 10^{-4}$
8.518	34.400	9.807	0.580	$(3.923 \pm 0.059) \times 10^{-4}$
8.518*	42.001	11.19	0.448	$(2.565 \pm 0.041) \times 10^{-4}$
8.518*	48.666	12.07	0.356	$(1.933 \pm 0.043) \times 10^{-4}$
8.518*	53.501	12.57	0.301	$(1.664 \pm 0.053) \times 10^{-4}$
10.587	48.666	15.76	0.309	$(8.405 \pm 0.227) \times 10^{-5}$

TABLE II. Fit parameters and uncertainties (Eq. 3)

a_1	b_1	b_2	b_3	c_1	c_2
0.072(22)	10.73(11)	19.81(17)	4.75(65)	-0.46(12)	0.12(10)

Figure 1 shows the global fit to G_M along with the values extracted from individual cross section measurements using the fit to $RS(Q^2)$ to extrapolate to $\epsilon = 0$. Our new data reduce the high- Q^2 uncertainties on G_M in the global fit by $> 30\%$.

We also performed direct Rosenbluth separations by grouping together points with similar Q^2 values, as indicated by the boxes in the top panel of Fig. 1. The normalization resulting from the global fit was applied to each data set, modifying the cross sections from Table I, and the data in each Q^2 bin were interpolated to a common Q_c^2 value using the global fit [53]. G_E and G_M were then extracted from a linear fit to the ϵ dependence of σ_R for each of the seven Q^2 bins. The results of this extraction are given in Table III. Figure 2 shows \sqrt{RS} (yielding $\mu_p G_E/G_M$ in the OPE) from our global analysis, along with a fit to the polarization data.

While it is conventional to compare measurements by showing $\sqrt{RS} = \mu_p G_E/G_M$, it is more correct to use RS which is the observable most directly extracted from the cross sections. Our quantitative comparisons of the FFRP use RS, as detailed in Ref. [53]. We find that the cross section data, using the best available radiative corrections but excluding hard TPE contributions, show a 2σ discrepancy with the polarization data up to 8 GeV 2 (1σ up to 14 GeV 2).

Accommodating this discrepancy at large Q^2 values requires a TPE contribution that reduces the cross section by $\sim 4\%$ at $\epsilon = 0$, assuming a linear ϵ dependence and a vanishing TPE contribution at $\epsilon = 1$, as detailed in the supplemental material [53], which includes Refs. [57? ? –71]. The cross section has a $\sim 2\%$ variation over the typical ϵ range of the




FIG. 1. (Top) Kinematics of elastic e - p data, Refs. [9–11, 23, 39, 43] and this work, used in the global fit and Rosenbluth separations; boxes (1-7) indicate the groupings of points for the Rosenbluth separations. (Bottom) Effective proton magnetic form factor, normalized by the standard dipole $\mu_p G_D$, obtained from the cross section measurements. The curve shows the result of our global fit, with the gray shaded area indicating the 68% confidence interval.

data. This is qualitatively consistent with some high- Q^2 calculations [33, 41] that predict large deviations from linear ϵ dependence which, however, are most significant below the ϵ range of the current data. Note that without the updated radiative corrections applied in this analysis, the discrepancy would have required TPE with a $\sim 6.5\%$ linear ϵ dependence, consistent with previous estimates [22, 72] based on analyses of data at lower Q^2 values using the older RC procedures.

In summary, the e - p elastic scattering cross section was measured for beam energies in the range of 2.2 - 11 GeV and Q^2 up to 15.75 (GeV/c) 2 . These new, high-precision cross sections provide an important baseline for the future proton and neutron structure investigations in the Jefferson Lab 12 GeV program. Our data were combined with existing cross section measurements [9–11, 39, 43] to perform Rosenbluth separations in a new Q^2 regime. The observed difference between the measured Rosenbluth slope and the OPE expectation, based on G_E/G_M from polarization data, would be resolved with a $\sim 4\%$ contribution to the cross section from hard TPE up to $Q^2 = 8$ GeV 2 , with no indication of significant Q^2 dependence at large Q^2 values.




FIG. 2. Direct Rosenbluth separation results for \sqrt{RS} ($= \mu_p G_E / G_M$ in OPE). The black solid (red dashed) curve shows the results of our fit to the cross section data with (without) the new GMP12 data. The blue dot-dashed curve shows $\mu_p G_E / G_M$ from a fit to the polarization data [53]. The shaded bands show the 68% confidence intervals of the respective fits. We plot $-\sqrt{|RS|}$ for the highest Q^2 point (an open circle), where $RS < 0$.

TABLE III. Rosenbluth separation results for the data groupings shown in the top panel of Fig. 1, after centering to the average Q_c^2 . The quoted values of σ_r and σ_L as defined in Eq. 2, and $G_M / (\mu_p G_D)$ and $\mu_p G_E / G_M$ are obtained assuming validity of the OPE approximation. For the largest Q^2 , where $\sigma_L < 0$, we quote $-\sqrt{|RS|}$.

Q_c^2 (GeV/c) 2	$\sigma_r \times 10^5$	$\sigma_L \times 10^5$	$G_M / (\mu_p G_D)$ (OPE)	$\mu_p G_E / G_M$ (OPE)
5.994	167 ± 4	7.1 ± 4.6	1.000 ± 0.011	0.75 ± 0.25
7.020	104 ± 3	9.3 ± 5.3	0.967 ± 0.015	1.18 ± 0.35
7.943	71.0 ± 2.7	4.1 ± 3.9	0.943 ± 0.018	1.0 ± 0.5
8.994	49.8 ± 1.7	0.7 ± 3.0	0.934 ± 0.016	0.5 ± 1.2
9.840	36.9 ± 2.4	1.9 ± 3.5	0.909 ± 0.029	1.1 ± 1.0
12.249	18.0 ± 0.8	1.2 ± 1.8	0.858 ± 0.019	1.3 ± 1.1
15.721	8.6 ± 0.5	-0.2 ± 1.2	0.840 ± 0.025	(-0.9 ± 2.8)

ACKNOWLEDGMENTS

We thank the Jefferson Lab accelerator and Hall A technical staff for their outstanding support of this experiment, which was part of the first run group to take data at JLab after the accelerator upgrade. We thank N. Kivel for careful reading of the manuscript and his valuable suggestions. Communications with P.N. Kirk, S. Rock, and A. Sill about technical aspects of their experiments are very much appreciated. This work was supported in part by the Science Committee of Republic of Armenia under grant 21AG-1C085, the Natural Sciences and Engineering Research Council of Canada (NSERC), the UK Science and Technology Facilities Council (Grant nos. STFC 57071/1 and STFC 50727/1), the U.S. National Science Foundation grant PHY-1508272, and the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contracts DE-AC02-05CH11231, DE-AC02-06CH11357, and DE-SC0016577, and DOE contract DE-

- [1] M. N. Rosenbluth, *Phys. Rev.* **79**, 615 (1950).
- [2] R. Sachs, *Phys. Rev.* **126**, 2256 (1962).
- [3] R. Hofstadter, *Rev. Mod. Phys.* **28**, 214 (1956).
- [4] S. D. Drell and M. A. Ruderman, *Phys. Rev.* **106**, 561 (1957).
- [5] S. D. Drell and S. Fubini, *Phys. Rev.* **113**, 741 (1959).
- [6] J. C. Bizot, J. M. Buon, J. Lefrançois, J. Perez-y Jorba, and P. Roy, *Phys. Rev.* **140**, B1387 (1965).
- [7] J. Mar, B. C. Barish, J. Pine, D. H. Coward, H. DeStaebler, J. Litt, A. Minten, R. E. Taylor, and M. Breidenbach, *Phys. Rev. Lett.* **21**, 482 (1968).
- [8] B. Bouquet, D. Benaksas, B. Grossetête, B. Jean-Marie, G. Parour, J. Poux, and R. Tchapoutin, *Phys. Lett.* **26**, 178 (1968).
- [9] R. C. Walker *et al.*, *Phys. Rev.* **D 49**, 5671 (1994).
- [10] P. N. Kirk *et al.*, *Phys. Rev.* **D 8**, 63 (1973).
- [11] A. F. Sill *et al.*, *Phys. Rev.* **D 48**, 29 (1993).
- [12] G. P. Lepage and S. J. Brodsky, *Phys. Rev. Lett.* **43**, 545 (1979), [Erratum: *Phys. Rev. Lett.* **43**, 1625 (1979)]; *Phys. Rev.* **D 22**, 2157 (1980).
- [13] A. I. Akhiezer, L. N. Rozentsveig, and I. M. Shumushkevich, *Zh. Eksp. Teor. Fiz.* **33**, 765 (1957), [*Sov. Phys. JETP* **6**, 588 (1958)]; A. I. Akhiezer and M. Rekalov, *Dokl. Akad. Nauk Ser. Fiz.* **180**, 1081 (1968), [*Sov. Phys. Dokl.* **13**, 572 (1968)]; *Fiz. Elem. Chast. Atom. Yadra* **4**, 662 (1973), [*Sov. J. Part. Nucl.* **4**, 277 (1974)].
- [14] G. Frolov, *Sov. Phys. JETP* **7**, 525 (1958).
- [15] J. H. Scofield, *Phys. Rev.* **113**, 1599 (1959); *Phys. Rev.* **141**, 1352 (1966).
- [16] R. G. Arnold, C. E. Carlson, and F. Gross, *Phys. Rev.* **C 23**, 363 (1981).
- [17] M. K. Jones *et al.* (Jefferson Lab Hall A), *Phys. Rev. Lett.* **84**, 1398 (2000), arXiv:nucl-ex/9910005 [nucl-ex]; O. Gayou *et al.* (Jefferson Lab Hall A), *Phys. Rev. Lett.* **88**, 092301 (2002), arXiv:nucl-ex/0111010; V. Punjabi *et al.*, *Phys. Rev.* **C 71**, 055202 (2005), [Erratum: *Phys. Rev.* **C 71**, 069902 (2005)], arXiv:nucl-ex/0501018 [nucl-ex].
- [18] A. J. R. Puckett *et al.*, *Phys. Rev.* **C 85**, 045203 (2012), arXiv:1102.5737 [nucl-ex].
- [19] A. J. R. Puckett *et al.*, *Phys. Rev. Lett.* **104**, 242301 (2010), arXiv:1005.3419 [nucl-ex]; *Phys. Rev.* **C 96**, 055203 (2017), [erratum: *Phys. Rev.* **C 98**, no.1, 019907 (2018)], arXiv:1707.08587 [nucl-ex].
- [20] G. A. Miller and M. R. Frank, *Phys. Rev.* **C 65**, 065205 (2002), arXiv:nucl-th/0201021 [nucl-th]; A. V. Belitsky, X.-d. Ji, and F. Yuan, *Phys. Rev. Lett.* **91**, 092003 (2003), arXiv:hep-ph/0212351 [hep-ph].
- [21] C. D. Roberts, M. S. Bhagwat, A. Holl, and S. V. Wright, *Eur. Phys. J. Special Topics* **140**, 53 (2007), arXiv:0802.0217 [nucl-th]; I. C. Cloet, G. Eichmann, B. El-Bennich, T. Klahn, and C. D. Roberts, *Few Body Syst.* **46**, 1 (2009), arXiv:0812.0416 [nucl-th]; M. Y. Barabanov *et al.*, *Prog. Part. Nucl. Phys.* **116**, 103835 (2021), arXiv:2008.07630 [hep-ph].
- [22] J. Arrington, *Phys. Rev.* **C 68**, 034325 (2003), arXiv:nucl-ex/0305009 [nucl-ex]; *Phys. Rev.* **C 69**, 022201 (2004), arXiv:nucl-ex/0309011.
- [23] M. E. Christy *et al.* (E94110 Collaboration), *Phys. Rev.* **C 70**, 015206 (2004), arXiv:nucl-ex/0401030 [nucl-ex].
- [24] I. A. Qattan *et al.*, *Phys. Rev. Lett.* **94**, 142301 (2005), arXiv:nucl-ex/0410010 [nucl-ex].
- [25] L. C. Maximon and J. A. Tjon, *Phys. Rev.* **C 62**, 054320 (2000), arXiv:nucl-th/0002058 [nucl-th].
- [26] V. Tsvakis, J. Arrington, M. E. Christy, R. Ent, C. E. Keppel, Y. Liang, and G. Vittorini, *Phys. Rev.* **C 73**, 025206 (2006), arXiv:nucl-ex/0511021 [nucl-ex].
- [27] J. Arrington, *Phys. Rev.* **C 69**, 032201(R) (2004), arXiv:nucl-ex/0311019 [nucl-ex].
- [28] I. A. Rachek *et al.*, *Phys. Rev. Lett.* **114**, 062005 (2015), arXiv:1411.7372 [nucl-ex].
- [29] D. Adikaram *et al.* (CLAS Collaboration), *Phys. Rev. Lett.* **114**, 062003 (2015), arXiv:1411.6908 [nucl-ex]; D. Rimal *et al.* (CLAS Collaboration), *Phys. Rev.* **C 95**, 065201 (2017), arXiv:1603.00315 [nucl-ex].
- [30] B. S. Henderson *et al.* (OLYMPUS Collaboration), *Phys. Rev. Lett.* **118**, 092501 (2017), arXiv:1611.04685 [nucl-ex].
- [31] M. Mezziane *et al.*, *Phys. Rev. Lett.* **106**, 132501 (2011).
- [32] D. Borisyuk and A. Kobushkin, *Phys. Rev.* **C 90**, 025209 (2014), arXiv:1405.2467 [hep-ph].
- [33] P. G. Blunden, W. Melnitchouk, and J. A. Tjon, *Phys. Rev.* **C 72**, 034612 (2005), arXiv:nucl-th/0506039 [nucl-th].
- [34] A. V. Afanasev, S. J. Brodsky, C. E. Carlson, Y.-C. Chen, and M. Vanderhaeghen, *Phys. Rev.* **D 72**, 013008 (2005), arXiv:hep-ph/0502013.
- [35] I. A. Qattan, A. Alsaad, and J. Arrington, *Phys. Rev.* **C 84**, 054317 (2011), arXiv:1109.1441 [nucl-ex].
- [36] J. C. Bernauer *et al.* (A1 Collaboration), *Phys. Rev.* **C 90**, 015206 (2014), arXiv:1307.6227 [nucl-ex].
- [37] R. Gerasimov and V. Fadin, *Phys. Atom. Nucl.* **78**, 69 (2015).
- [38] A. V. Gramolin and D. M. Nikolenko, *Phys. Rev.* **C 93**, 055201 (2016), arXiv:1603.06920 [nucl-ex].
- [39] L. Andivahis *et al.*, *Phys. Rev.* **D 50**, 5491 (1994).
- [40] J. Arrington, P. G. Blunden, and W. Melnitchouk, *Prog. Part. Nucl. Phys.* **66**, 782 (2011), arXiv:1105.0951 [nucl-th].
- [41] A. Afanasev, P. Blunden, D. Hasell, and B. Raue, *Prog. Part. Nucl. Phys.* **95**, 245 (2017), arXiv:1703.03874 [nucl-ex].
- [42] Z. Ye, J. Arrington, R. J. Hill, and G. Lee, *Phys. Lett.* **B 777**, 8 (2018), arXiv:1707.09063 [nucl-ex].
- [43] S. Rock *et al.*, *Phys. Rev.* **D 46**, 24 (1992).
- [44] J. Alcorn *et al.*, *Nucl. Instrum. Methods* **A 522**, 294 (2004).
- [45] B. Schmoochler, *Nucleon Structure and Its Modification in Nuclei*, Ph.D. thesis, Massachusetts Institute of Technology (2018).
- [46] J. Berthot and P. Vernin, *Nucl. Phys. News* **9N4**, 12 (1999).
- [47] S. Santiesteban *et al.*, "Precision Beam Energy Determination for the CEBAF 12 GeV Upgrade," arXiv:2110.06281 [physics.acc-ph].
- [48] T. Gautam, *Precision Measurement of the Proton Magnetic Form Factor at high Q^2* , Ph.D. thesis, Hampton University, Hampton, Virginia (2019).
- [49] K. Unser, *Application of accelerators in Research and Industry, Proceedings, 6th conference, Denton, Texas, USA, November 3-5, 1980*, IEEE Trans. Nucl. Sci. **28**, 2344 (1981).
- [50] Y. Wang, *Measurement of the Proton Elastic Cross Section at $Q^2 = 0.66, 1.10, 1.51$ and 1.65 GeV²*, Ph.D. thesis, College of William & Mary (2017).
- [51] K. Allada, C. Hurlbut, L. Ou, B. Schmoochler, A. Shahinyan, and B. Wojtsekhowski, *Nucl. Instrum. Meth.* **A 782**, 87 (2015), arXiv:1502.01772 [physics.ins-det].

- [52] L. Ou, *Precision Measurements of Electron-Proton Elastic Scattering Cross Sections at Large Q^2* , Ph.D. thesis, Massachusetts Institute of Technology (2018).
- [53] “See supplemental material in the source files for the cross-section data base and the global fit of the e-p cross section data, which includes references [57-71].”
- [54] J. Arrington, *Inclusive Electron Scattering From Nuclei at $x > 1$ and High Q^2* , Ph.D. thesis, California Institute of Technology (1998), arXiv:nucl-ex/0608013.
- [55] R. Ent, B. W. Filippone, N. C. R. Makins, R. G. Milner, T. G. O’Neill, and D. A. Wasson, *Phys. Rev. C* **64**, 054610 (2001).
- [56] L. W. Mo and Y.-S. Tsai, *Rev. Mod. Phys.* **41**, 205 (1969).
- [57] Y.-C. Chen, C.-W. Kao, and S.-N. Yang, *Phys. Lett. B* **652**, 269 (2007), arXiv:nucl-th/0703017.
- [58] J. J. Kelly, *Phys. Rev. C* **70**, 068202 (2004).
- [59] K. Makino and M. Berz, *Computational accelerator physics. Proceedings, 8th International Conference, ICAP 2004, St. Petersburg, Russia, June 29-July 2, 2004*, *Nucl. Instrum. Meth. A* **558**, 346 (2006).
- [60] R. C. D. Walker, *A measurement of the proton elastic form factors for $1 \leq Q^2 \leq 3$ (GeV/c)²*, Ph.D. thesis, California Institute of Technology (1989).
- [61] Y.-S. Tsai, *Phys. Rev.* **122**, 1898 (1961).
- [62] Y. S. Tsai, (1971), technical report SLAC-PUB-848.
- [63] G. Ron *et al.* (Jefferson Lab Hall A), *Phys. Rev. C* **84**, 055204 (2011), arXiv:1103.5784 [nucl-ex].
- [64] X. Zhan *et al.*, *Phys. Lett. B* **705**, 59 (2011), arXiv:1102.0318 [nucl-ex].
- [65] M. Paolone *et al.*, *Phys. Rev. Lett.* **105**, 072001 (2010), arXiv:1002.2188 [nucl-ex].
- [66] T. Pospischil *et al.* (A1), *Eur. Phys. J. A* **12**, 125 (2001).
- [67] M. K. Jones *et al.* (Resonance Spin Structure), *Phys. Rev. C* **74**, 035201 (2006), arXiv:nucl-ex/0606015.
- [68] C. B. Crawford *et al.*, *Phys. Rev. Lett.* **98**, 052301 (2007), arXiv:nucl-ex/0609007.
- [69] B. D. Milbrath *et al.* (Bates FPP), *Phys. Rev. Lett.* **80**, 452 (1998), [Erratum: *Phys.Rev.Lett.* 82, 2221 (1999)], arXiv:nucl-ex/9712006.
- [70] O. Gayou *et al.*, *Phys. Rev. C* **64**, 038202 (2001).
- [71] A. V. Gramolin and D. M. Nikolenko, (2016), <https://github.com/gramolin/rosenbluth/>.
- [72] J. Arrington, W. Melnitchouk, and J. Tjon, *Phys. Rev. C* **76**, 035205 (2007), arXiv:0707.1861 [nucl-ex].