UCLA

UCLA Previously Published Works

Title

Is waist-to-height ratio better than body mass index as a predictive indicator of coronary atherosclerosis disease? A cohort study.

Permalink

<https://escholarship.org/uc/item/6rm3f00p>

Journal

Journal of cardiovascular computed tomography, 13(4)

ISSN

1934-5925

Authors

Hou, Zhi-Hui
Lu, Bin
Li, Zhen-Nan
et al.

Publication Date

2019-07-01

DOI

10.1016/j.jcct.2019.06.009

Peer reviewed

Association of Anthropometric Measures and Cardiovascular Biomarkers with Non-Calcified Coronary Artery Plaque Progression in Older Hypogonadal Men Treated with Testosterone: Results from Testosterone Trials

Kashif Shaikh,MD¹, Susan Ellenberg,PhD² Rine Nakanishi,MD,PhD¹, Juhwan Lee, MD,PhD¹, Nanette K. Wenger, MD³, Cora E. Lewis,MD⁴, Ronald S. Swerdloff,MD⁵,Elizabeth Barrett-Connor⁶,MD, Alisa Stephens-Sheilds,PhD², Shalender Bhasin,MD⁶, Suguru Matsumoto,MD¹, Jane A Cauley,DrPH⁷, Jill P Crandall,MD⁸, Glenn R Cunningham,MD⁹, Kristine E Ensrud,MD,MPH^{10,11}, Alvin M Matsumoto,MD¹², Mark E Molich,MD¹³, Venkata m Alla¹⁴, Negin Nezarat,MD¹, Anas Alani,MD¹, Chandana Shekar,MD¹, Anoop Kumar¹, Shone Almeida,MD¹, Salman Sheikh,MD¹, Sajad Hamal,MS¹, Sion K Roy,MD¹, Peter J Snyder,MD¹⁵, Mathew Budoff,MD¹

1Los Angeles Biomedical Research Institute, Division of Cardiology, Harbor-University of California at Los Angeles Medical Center, Torrance, CA, USA

2-Department of Biostatistics and Epidemiology, Perelman School of Medicine at The University of Pennsylvania, Philadelphia, PA, USA

³ Department of Medicine, Division of Cardiology, Emory Heart and Vascular Center Emory University School of Medicine, Atlanta, GA, USA

4-Division of Preventive Medicine, University of Alabama at Birmingham, AL, USA

⁵ Los Angeles Biomedical Research Institute, Division of Endocrinology, Harbor- University of California at Los Angeles Medical Center, Torrance, CA, USA

⁶ Department of Family and Preventive Medicine, Division of Epidemiology, University of California, San Diego School of Medicine, La Jolla, CA, USA

⁷ Department of Epidemiology, University of Pittsburgh, Graduate School of Public Health, Pittsburgh, PA, 1 USA

⁸ Divisions of Endocrinology and Geriatrics, Albert Einstein College of Medicine, Bronx, NY, USA

⁹ Departments of Medicine and Molecular & Cellular Biology, Division of Diabetes, Endocrinology and Metabolism, Baylor College of Medicine and Baylor St. Luke's Medical Center, Houston, TX, USA

¹⁰ Department of Medicine, Division of Epidemiology & Community Health, University of Minnesota, Minneapolis, MN, USA

¹¹ Minneapolis VA Health Care System, Minneapolis, MN, USA

¹²Geriatric Research, Education, and Clinical Center, Department of Veterans Affairs, Puget Sound Health System, and Division of Gerontology and Geriatric Medicine, Department of Internal Medicine, University of Washington School of Medicine, Seattle WA

¹³ Division of Endocrinology, Metabolism and Molecular Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

¹⁴Division of Cardiovascular Diseases, Creighton University School of Medicine, Omaha, Nebraska.

¹⁵ Division of Endocrinology, Diabetes, and Metabolism, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

Correspondence: Matthew J. Budoff, MD FACC
Professor of Medicine
Program Director
Division of Cardiology, Harbor UCLA
Los Angeles Biomedical Research Institute at Harbor-UCLA, Torrance,
California, USA
Tel: +1-310 2224107
Fax: +1-3107829652
Email: mbudoff@labiomed.org

Grants: Dr. Budoff reports grants from NIH, during the conduct of the study; grants from General Electric, outside the submitted work Dr. Swerdloff

reports grants from The Bone Trial of the Testosterone Trial during the conduct of the study; grants and other from Clarus, grants from Lipesene, grants and other from Antares, outside the submitted work; The Testosterone Trials were supported by a grant from the National Institute on Aging, National Institutes of Health (U01 AG030644), supplemented by funds from the National Heart, Lung and Blood Institute, National Institute of Neurological Diseases and Stroke, and National Institute of Child Health and Human Development. AbbVie (formerly Solvay and Abbott Laboratories) generously provided funding, AndroGel, and placebo gel.

Disclosures: Dr. Ellenberg reports grants from National Institutes of Health, grants from AbbVie, Inc, during the conduct of the study; grants from AbbVie, Inc., outside the submitted work; Dr. Lewis reports grants from NIH, grants from AbbVie, during the conduct of the study; Dr. Wenger reports grants from Alnylam Pharmaceuticals, grants and personal fees from Gilead Sciences, grants from NHLBI, grants from Pfizer, grants from Society for Women's Health Research, personal fees from Amgen, personal fees from AstraZeneca, personal fees from Merck, outside the submitted work; Dr. Budoff reports grants from NIH, during the conduct of the study; grants from General Electric, outside the submitted work; Dr. Barrett-Connor has nothing to disclose; Dr. Swerdloff reports grants from The Bone Trial of the Testosterone Trial during the conduct of the study; grants and other from Clarus, grants from Lipesene, grants and other from Antares, outside the submitted work; Dr. Stephens-Shields reports grants from National Institute

on Aging and from AbbVie during the conduct of the study; Dr. Bhasin reports grants from NIA, during the conduct of the study; grants and personal fees from Abbvie, grants and personal fees from Lilly, grants from Transition Therapeutics, grants and personal fees from Regeneron, outside the submitted work; In addition, Dr. Bhasin has a patent Free testosterone calculator pending and has equity interest in FPT, LLC.; Dr. Cauley has nothing to disclose; Dr. Crandall has nothing to disclose; Dr. Cunningham reports personal fees from AbbVie, personal fees from Apricus, personal fees from Besins, personal fees from Clarus Therapeutics, personal fees from Endo Pharma, personal fees from Ferring, personal fees from Lilly, personal fees from Pfizer, personal fees from Repros Therapeutics, outside the submitted work; Dr. Ensrud reports grants from National Institute on Aging, during the conduct of the study; Dr. Gill has nothing to disclose; Dr. Matsumoto reports personal fees from AbbVie, personal fees from Endo, personal fees from Lilly, personal fees from Lipocine, personal fees from Clarus, outside the submitted work; Dr. Molitch reports grants from National Institutes of Health, grants from Abbott Laboratories, during the conduct of the study; personal fees from Abbvie (Abbott Laboratories), personal fees from Eli Lilly & Co., personal fees from Pfizer, outside the submitted work; X. Hou has nothing to disclose; Dr. Snyder reports grants from National Institute on Aging, NIH, grants and non-financial support from AbbVie (formerly Solvay and Abbott Laboratories), during the conduct of the study.

Abstract:

OBJECTIVE/BACKGROUND: Several studies have yielded conflicting results as to whether testosterone treatment increases cardiovascular risk. Recent results from the placebo-controlled Cardiovascular Trial of the Testosterone Trials showed that testosterone treatment of older men with low testosterone was associated with greater progression of non-calcified plaque. We sought to evaluate the effect of anthropometric measures and biomarkers on coronary artery plaque progression in individuals in the Testosterone Trial.

Methods: The Testosterone Trials were a coordinated set of seven double-blinded, placebo-controlled trials. The Cardiovascular Trial included 170 men aged 65 years or older with low testosterone and no contraindication to coronary computed tomographic angiography (CCTA). Participants received testosterone gel dose-adjusted to maintain the testosterone level in the normal range for young men, or placebo gel for 12 months. The primary outcome was change in non-calcified coronary artery plaque volume from baseline to 12 months, as determined by CCTA. We also assayed several markers of cardiovascular risk and analyzed each marker individually in a model that included the binary factors for age and testosterone level, the marker by treatment interaction, and the baseline value of non-calcified plaque as predictive variables and change in non-calcified plaque as the dependent variable.

RESULTS: Of 170 enrollees, 138 (73 testosterone, 65 placebo) completed the study and were available for the primary analysis. Among the 138 men, the mean (SD) age was 71.2 (5.7) years, 81% were white. Of 9 markers evaluated, none showed a significant association with the change in non-calcified plaque volume, but a significant interaction between treatment and waist-hip ratio ($p=0.0014$) indicated that this variable influences the testosterone effect on non-calcified plaque volume. The statistical model indicates that every 0.1 change in the waist-hip ratio increases the

testosterone-induced 12-month change in non-calcified plaque volume by 26.96 mm³ (95% confidence interval 7.72, 46.20).

CONCLUSIONS:

Among older men with low testosterone treated with testosterone for one year, greater waist-hip ratio was associated with greater non-calcified plaque progression, as measured by coronary computed tomographic angiography.

Introduction:

Lower serum testosterone concentration has been associated with adverse cardiovascular disease (CVD) outcomes^{1,2}. There are conflicting reports regarding the effect of testosterone treatment on CVD risk. Some retrospective studies reported more CVD events in men taking testosterone, while others did not³⁻⁷. The Testosterone Trials (TTrials) comprised seven coordinated placebo-controlled clinical trials designed to assess the effects of testosterone treatment in older men who had low testosterone concentrations for no apparent reason other than age⁸. In the Cardiovascular Trial, testosterone treatment for one year compared with placebo was associated with significantly greater progression of coronary artery non-calcified plaque volume measured by serial coronary computed tomography angiography (CCTA)⁹.

Serum markers such as total cholesterol, HDL, LDL and hemoglobin A1C, have been recognized as significant risk factors for developing coronary artery plaque and future CVD events^{10,11}. There are contradictory reports about the association of biomarkers and extent, progression of atherosclerosis and coronary events¹²⁻¹⁴. Inflammatory markers such as CRP have been reported to be associated with plaque progression in some studies^{15,16}, other reports found no association^{17,18}. Anthropometric measures such as Waist-Hip ratio and Waist Circumference are predictors of myocardial infarction risk^{19,20}. Abdominal obesity can lead to increases in insulin and glucose levels and is a central feature of metabolic syndrome.

Several observational studies have shown link of low endogenous sex hormones and metabolic syndrome²¹⁻²³. One large cross-sectional study reported that higher testosterone and sex hormone binding globulin levels in older men were independently associated with reduced risk of metabolic syndrome and higher insulin sensitivity²⁴.

The aim of the current study is to evaluate the impact of baseline anthropometric measures and cardiovascular biomarkers on the progression of coronary artery plaque volume in the 138 men who participated in the Cardiovascular Trial of the TTrial. We also assessed the interaction of anthropometric measures and cardiovascular biomarkers with testosterone treatment and its impact on plaque progression as measured by CCTA to determine whether any of these measures influenced the effect of testosterone on plaque progression.

METHODS

Study Design

The TTrial comprised seven double-blind, placebo-controlled randomized controlled trials. The overall study design of TTrial, as well that of Cardiovascular Trial, has been published^{8,25}. To qualify for the TTrial overall, a participant had to qualify for at least 1 of 3 main trials (Sexual Function

Trial, Physical Function Trial, and Vitality Trial). Qualified men could also participate in any of other trials, if respective criteria were met. The participants were allocated to receive testosterone or placebo gel for 1 year^{8,9}. Institutional review boards of all participating sites approved TTrial and Cardiovascular Trial protocols. All participants provided written consent. Trial conduct and participant safety was supervised by an independent safety and data monitoring board.

Participants

The TTrial included men > 65 years' old who had symptoms and objective evidence of low libido, physical dysfunction and/or low vitality, serum testosterone levels that averaged < 275 ng/dL on 2 morning samples. Men who were at moderate or high risk for prostate cancer, who had had a myocardial infarction within the previous 3 months, or had systolic blood pressure >160 mm Hg or diastolic blood pressure >100 mm Hg, were excluded⁸.

Exclusion criteria specifically for the Cardiovascular Trial included circumstances that either made coronary artery CT angiography (CCTA) technically unfeasible (inability to hold breath for 10 seconds, a prior diagnosis of tachycardia or irregular heart rhythm [e.g., atrial fibrillation], weight >136 kg, or history of coronary artery bypass graft surgery) or

increased risk of performing the CCTA (estimated glomerular filtration rate <60 mL/min/1.73 m² or known allergy to iodinated contrast)^{9,25}.

Testosterone Treatment:

Participants were assigned to receive either testosterone as a 1 % gel in a pump bottle (AndroGel) or placebo gel by a double-blinded method for one year. The initial dose was 5 g/d and was adjusted to maintain the serum concentrations within normal range for young men (280-873 ng/dL) measured at central laboratory (Quest Clinical Trials) at months 1, 2, 3, 6, and 9. Whenever dose adjustments were made in a man receiving testosterone treatment, the dose was changed in a man receiving placebo as well to maintain blinding⁸.

Assessments:

The concentrations of cardiovascular biomarkers were measured on serum samples drawn at baseline and months 3 and 12 and stored at -80 C. These assays were performed at the Laboratory for Clinical Biochemistry Research, University of Vermont and University of Minnesota, as described previously^{7,9}. At months 3, 6, 9, and 12, clinical variables were measured.

Details of coronary artery plaque volume by CCTA assessment have been published²⁵. In brief, coronary artery plaque volume was assessed by CCTA at 9 of the 12 TTrials clinical sites. Pre-contrast scans for evaluation of

coronary artery calcium density and post contrast scans for evaluation of coronary artery plaque volume were performed at baseline and 12 months. Scans were assessed at a central reading center (Harbor-UCLA Medical Center) by readers who were blinded both to treatment group and date of scan. Quantitative plaque assessment was conducted according to a previously defined protocol²⁶ using semi-automated plaque analysis software (QAngioCT Research Edition Version 2.0.5; Medis Medical Imaging Systems). Based on the guidelines of the Society of Cardiovascular Computed Tomography, 17-segment coronary artery model vessels greater than 1.5 mm were evaluated²⁷. The volumes of four types of coronary artery plaque (low attenuation, fibrous-fatty, fibrous, and dense calcified) were calculated by Hounsfield unit threshold. The primary outcome was change in non-calcified plaque volume from baseline to month 12. Non-calcified plaque was defined as the sum of the fibrous, fibrous fatty and low attenuation plaque. secondary outcomes were change in calcified plaque volume, and change in coronary artery score. Details of intra- and inter-observer variability have been published. The intra-class correlations (ICCs) and Coefficient of Variation (CVs) were 0.99 and 7.8 % for intra-observer variability respectively. ICC and CV was 0.95 and 19.9 % for inter-observer variability respectively⁹.

Statistical Analyses

The following markers were available for study: total cholesterol; non-HDL cholesterol; HDL; LDL; triglycerides; HgA1c; glucose, insulin; HOMA; d-dimer; troponin; CRP; IL-6; weight; BMI; waist/hip ratio. We evaluated the intercorrelation of the baseline values of these markers, separately within groups where substantial intercorrelation was expected: lipid markers, metabolic markers, markers of inflammation, and clinical markers. We then excluded from further study the marker showing correlation > 0.5 with the most other markers, and then eliminated any marker with correlation > 0.5 with the selected marker from further consideration. We retained any other markers with correlation < 0.5 with the selected marker. If two markers showed high correlation with the same number of other markers, we selected the one with the lowest correlation with the remaining markers.

We tested each selected marker separately in a regression model, including treatment as a covariate as well as age (over or under 75), baseline testosterone (over or under 200 ng/ml) and an interaction term of the marker with treatment. Any variable showing a significant association with the change in plaque volume after adjusting for multiple comparisons using the Holm procedure²⁸ was to be included in a multivariable model, assessing all potentially predictive variables simultaneously.

Secondary analyses included testing association of the selected markers with change in calcified plaque volume and with coronary artery calcium score, using the same approach as above.

Results

The baseline characteristics of the participants in the Cardiovascular Trial were previously reported (9). At baseline, the mean (SD) age was 71.2 (5.7) years. The majority of participants were white (81%) and had relatively high rates of cardiovascular risk factors, including hypertension, hyperlipidemia, obesity, and diabetes. At baseline the mean BMI was (3.8) in the testosterone group and 30 (3.5) in the placebo group; mean weight was 94 kg and the mean waist-hip ratio was 1.0 in each treatment group. The calculated 10-year risk of cardiovascular events was relatively high as well (a mean risk of 27% [95% CI, 6.4%-47.6%] in the placebo group and 24% [95% CI 2.6%-45.4%] in the testosterone group).

Of the 17 markers initially evaluated, 9 remained for further study after removing those that were highly correlated with other markers, as described above. These 9 markers were HDL cholesterol, non-HDL cholesterol, D-dimer, IL-6, CRP, insulin, HgbA1C, weight and waist-hip ratio (Table-1). Among these 9 measures, only the baseline waist-hip ratio interaction with treatment showed a significant association with the progression of non-calcified plaque volume at 12 months, (Table 2, Figure 1). Because it was the interaction term that met the threshold based on the multiple comparisons adjustment ($p=0.0014$ compared to threshold value from the Holm multiple comparisons procedure of 0.0056), we evaluated waist-hip ratio separately for the two treatment groups. The association was seen only in the testosterone group ($p=0.007$). The model indicates that for every 0.1 change in the waist-hip ratio, the effect of testosterone on the 12-month

change in non-calcified plaque volume would increase by 26.96 mm³ (95% confidence interval 7.72, 46.20). (The baseline values of waist-hip ratio ranged from 0.9 to 1.2).

The only variable to show a significant association with change in calcified plaque volume was D-dimer, and the significant finding was only for the interaction term. Further investigation showed that the interaction effect appeared to be based on one or two extreme values in each treatment group. None of the cardiovascular risk markers were statistically significantly associated with change in CAC score when applying the multiple comparisons correction.

DISCUSSION:

We report that in older hypogonadal men participating in the Cardiovascular Trial of the TTrial there was a significant association between baseline waist-hip ratio and progression of non-calcified coronary artery plaque volume measured by coronary artery CT angiography after one year of testosterone treatment. Larger waist-hip ratios were associated with greater progression of non-calcified plaque in men taking testosterone.

Visceral adipose tissue decreases insulin sensitivity and promotes dyslipidemia and hypertension^{29,30}. Visceral adipose tissue stores can be measured by CT, DXA or MRI but these modalities are too expensive and time consuming for day-to-day use^{31,32}. WHR is closely related to visceral fat

and commonly measured in clinical practice³³. Meta-analyses of 28,114 patients from 15 prospective studies showed that for every 0.01 increase in WHR, there was a 5 % increase in risk of future CVD events ³³. Our data indicate that for every 0.1 increase in waist hip ratio, there was 26 mm³ greater increase in progression of non-calcified plaque volume in patients treated with testosterone replacement therapy.

Non-calcified plaque volumes as assessed by cardiac CCTA has been associated with CVD events. In a large single center trial by Zu et al³⁴, the cumulative probability of 3-year MACE increased across the strata for cardiac CT plaque, with non-calcified plaque having highest probability, 37.7%.

WHR and waist circumference, measures of central obesity or abdominal obesity, have been associated with reduced total testosterone levels^{35,36}. A mechanism that may account for this inverse relationship involves strong association of central obesity and excess secretion of cortisol, potentially mediated through suppression of testosterone production via the hypothalamic-pituitary axis³⁷. Another plausible mechanism of decreased testosterone in obese individuals is increased aromatase activity in visceral adipose tissue, which leads to higher conversion of testosterone to estradiol³⁸. Androgen deprivation therapy, as given to patients with prostate cancer, has shown to significantly increase BMI, total weight, body fat mass and decrease in lean body mass^{39,40}. Hence, several studies have

investigated the hypothesis that testosterone replacement therapy may decrease visceral fat stores and improve the metabolic profile in men. However, there are conflicting reports on effects of testosterone replacement on visceral fat. Some studies reported testosterone replacement therapy decreases visceral fat, while other showed no association^{41,42}. In a study of 261 patients in a prospective longitudinal registry, testosterone replacement was associated with a significant reduction in obesity parameters (e.g. WC, BMI) and cholesterol values over the 5-year study period⁴³. However, randomized controlled clinical trials reported no impact of testosterone replacement on weight, BMI and metabolic syndrome^{41,44}. A previous paper from the TTrialS also did not show any changes in WHR, WC and BMI in men treated with testosterone for 12 months compared to those treated with placebo⁷.

These results warrant further investigation of the interaction of visceral adipose tissue stores and testosterone treatment. To our knowledge, no other studies have examined the interaction of testosterone replacement therapy and central obesity on CVD outcomes. The strengths of our trial included requiring all men to have unequivocally low testosterone at baseline, a placebo-controlled design and blinded central review of baseline and 12 month scans. An important limitation of our study is use of a surrogate marker of heart disease, non-calcified plaque, and not a clinical

outcome. Another limitation is that the results apply only to men >65 with low testosterone⁹.

We conclude that older men with higher WHR receiving testosterone treatment may experience greater increases in noncalcified coronary plaque volume than treated men with lower WHR. Future trials should evaluate the interaction of testosterone treatment and surrogate markers of abdominal obesity and visceral fat stores.

Reference

1. Khaw K-T, Dowsett M, Folkard E, et al. Endogenous testosterone and mortality due to all causes, cardiovascular disease, and cancer in men: European prospective investigation into cancer in Norfolk (EPIC-Norfolk) Prospective Population Study. *Circulation* 2007;116:2694-701.
2. Smith GD, Ben-Shlomo Y, Beswick A, Yarnell J, Lightman S, Elwood P. Cortisol, testosterone, and coronary heart disease: prospective evidence from the Caerphilly study. *Circulation* 2005;112:332-40.
3. Baillargeon J, Urban RJ, Kuo Y-F, et al. Risk of myocardial infarction in older men receiving testosterone therapy. *Annals of Pharmacotherapy* 2014;48:1138-44.
4. Finkle WD, Greenland S, Ridgeway GK, et al. Increased risk of non-fatal myocardial infarction following testosterone therapy prescription in men. *PloS one* 2014;9:e85805.
5. Shores MM, Smith NL, Forsberg CW, Anawalt BD, Matsumoto AM. Testosterone treatment and mortality in men with low testosterone levels. *The Journal of Clinical Endocrinology & Metabolism* 2012;97:2050-8.
6. Vigen R, O'donnell CI, Barón AE, et al. Association of testosterone therapy with mortality, myocardial infarction, and stroke in men with low testosterone levels. *Jama* 2013;310:1829-36.
7. Mohler III ER, Ellenberg SS, Lewis CE, et al. The effect of testosterone on cardiovascular biomarkers in the testosterone trials. *The Journal of Clinical Endocrinology & Metabolism* 2017;103:681-8.
8. Snyder PJ, Ellenberg SS, Cunningham GR, et al. The Testosterone Trials: Seven coordinated trials of testosterone treatment in elderly men. *Clinical trials (London, England)* 2014;11:362-75.
9. Budoff MJ, Ellenberg SS, Lewis CE, et al. Testosterone treatment and coronary artery plaque volume in older men with low testosterone. *Jama* 2017;317:708-16.
10. Cao Q, Yu S, Xiong W, et al. Waist-hip ratio as a predictor of myocardial infarction risk: A systematic review and meta-analysis. *Medicine* 2018;97.
11. Diederichsen SZ, Gronhoj MH, Mickley H, et al. CT-Detected Growth of Coronary Artery Calcification in Asymptomatic Middle-Aged Subjects and Association With 15 Biomarkers. *JACC Cardiovascular imaging* 2017;10:858-66.
12. Anroedh SS, Akkerhuis KM, Oemrawsingh RM, et al. Associations of 26 Circulating Inflammatory and Renal Biomarkers with Near-Infrared Spectroscopy and Long-term Cardiovascular Outcome in Patients Undergoing Coronary Angiography (ATHEROREMO-NIRS Substudy). *Current atherosclerosis reports* 2018;20:52.
13. Battes LC, Cheng JM, Oemrawsingh RM, et al. Circulating cytokines in relation to the extent and composition of coronary atherosclerosis: results from the ATHEROREMO-IVUS study. *Atherosclerosis* 2014;236:18-24.
14. Cheng JM, Oemrawsingh RM, Garcia-Garcia HM, et al. Relation of C-reactive protein to coronary plaque characteristics on grayscale, radiofrequency intravascular ultrasound, and cardiovascular outcome in patients with acute coronary syndrome or stable angina pectoris (from the ATHEROREMO-IVUS study). *The American journal of cardiology* 2014;114:1497-503.
15. Alman AC, Kinney GL, Tracy RP, et al. Prospective association between inflammatory markers and progression of coronary artery calcification in adults with and without type 1 diabetes. *Diabetes care* 2013;36:1967-73.

16. Wadwa RP, Kinney GL, Ogden L, et al. Soluble interleukin-2 receptor as a marker for progression of coronary artery calcification in type 1 diabetes. *The international journal of biochemistry & cell biology* 2006;38:996-1003.
17. Gauss S, Klinghammer L, Steinhoff A, et al. Association of systemic inflammation with epicardial fat and coronary artery calcification. *Inflammation research : official journal of the European Histamine Research Society* [et al] 2015;64:313-9.
18. Kronmal RA, McClelland RL, Detrano R, et al. Risk factors for the progression of coronary artery calcification in asymptomatic subjects: results from the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation* 2007;115:2722-30.
19. Cao Q, Yu S, Xiong W, et al. Waist-hip ratio as a predictor of myocardial infarction risk: A systematic review and meta-analysis. *Medicine* 2018;97:e11639.
20. Tigbe WW, Granat MH, Sattar N, Lean MEJ. Time spent in sedentary posture is associated with waist circumference and cardiovascular risk. *International journal of obesity (2005)* 2017;41:689-96.
21. Barrett-Connor E, Khaw K-T. Endogenous sex hormones and cardiovascular disease in men. A prospective population-based study. *Circulation* 1988;78:539-45.
22. Gyllenberg J, Rasmussen SL, Borch-Johnsen K, Heitmann BL, Skakkeb NE, Juul A. Cardiovascular risk factors in men: the role of gonadal steroids and sex hormone-binding globulin. *Metabolism-Clinical and Experimental* 2001;50:882-8.
23. Simon D, Charles M-A, Nahoul K, et al. Association between plasma total testosterone and cardiovascular risk factors in healthy adult men: The Telecom Study. *The Journal of Clinical Endocrinology & Metabolism* 1997;82:682-5.
24. Muller M, Grobbee DE, Den Tonkelaar I, Lamberts SW, Van Der Schouw YT. Endogenous sex hormones and metabolic syndrome in aging men. *The Journal of Clinical Endocrinology & Metabolism* 2005;90:2618-23.
25. Abd Alamir M, Ellenberg SS, Swerdloff RS, et al. The Cardiovascular Trial of the Testosterone Trials: rationale, design, and baseline data of a clinical trial using computed tomographic imaging to assess the progression of coronary atherosclerosis. *Coron Artery Dis* 2016;27:95-103.
26. Papadopoulou SL, Neefjes LA, Garcia-Garcia HM, et al. Natural history of coronary atherosclerosis by multislice computed tomography. *JACC Cardiovascular imaging* 2012;5:S28-37.
27. Leipsic J, Abbara S, Achenbach S, et al. SCCT guidelines for the interpretation and reporting of coronary CT angiography: a report of the Society of Cardiovascular Computed Tomography Guidelines Committee. *J Cardiovasc Comput Tomogr* 2014;8:342-58.
28. Holm S. A simple sequentially rejective multiple test procedure. *Scandinavian journal of statistics* 1979:65-70.
29. Pouliot M-C, Després J-P, Nadeau A, et al. Visceral obesity in men: associations with glucose tolerance, plasma insulin, and lipoprotein levels. *Diabetes* 1992;41:826-34.
30. Tchernof A, Lamarche B, Prud'homme D, et al. The dense LDL phenotype: association with plasma lipoprotein levels, visceral obesity, and hyperinsulinemia in men. *Diabetes care* 1996;19:629-37.
31. Kamel E, McNeill G, Han T, et al. Measurement of abdominal fat by magnetic resonance imaging, dual-energy X-ray absorptiometry and anthropometry in non-obese men and women. *International journal of obesity* 1999;23:686.

32. Onat A, Avcı GŞ, Barlan M, Uyarel H, Uzunlar B, Sansoy V. Measures of abdominal obesity assessed for visceral adiposity and relation to coronary risk. *International journal of obesity* 2004;28:1018.
33. De Koning L, Merchant AT, Pogue J, Anand SS. Waist circumference and waist-to-hip ratio as predictors of cardiovascular events: meta-regression analysis of prospective studies. *European heart journal* 2007;28:850-6.
34. Hou Z-h, Lu B, Gao Y, et al. Prognostic value of coronary CT angiography and calcium score for major adverse cardiac events in outpatients. *JACC: Cardiovascular Imaging* 2012;5:990-9.
35. Pasquali R, Casimirri F, Cantobelli S, et al. Effect of obesity and body fat distribution on sex hormones and insulin in men. *Metabolism* 1991;40:101-4.
36. Svartberg J. Epidemiology: testosterone and the metabolic syndrome. *International journal of impotence research* 2007;19:124.
37. Rosmond R, Wallerius S, Wanger P, Martin L, Holm G, Björntorp P. A 5-year follow-up study of disease incidence in men with an abnormal hormone pattern. *Journal of internal medicine* 2003;254:386-90.
38. Kalyani RR, Dobs AS. Androgen deficiency, diabetes, and the metabolic syndrome in men. *Current Opinion in Endocrinology, Diabetes and Obesity* 2007;14:226-34.
39. Chen Z, Maricic M, Nguyen P, Ahmann FR, Bruhn R, Dalkin BL. Low bone density and high percentage of body fat among men who were treated with androgen deprivation therapy for prostate carcinoma. *Cancer* 2002;95:2136-44.
40. Smith MR, Finkelstein JS, McGovern FJ, et al. Changes in body composition during androgen deprivation therapy for prostate cancer. *The Journal of Clinical Endocrinology & Metabolism* 2002;87:599-603.
41. Hoyos CM, Yee BJ, Phillips CL, Machan EA, Grunstein RR, Liu PY. Body compositional and cardiometabolic effects of testosterone therapy in obese men with severe obstructive sleep apnoea: a randomised placebo-controlled trial. *European journal of endocrinology* 2012;167:531-41.
42. Lunenfeld B. The relationship between sex hormones and the metabolic syndrome. *Acta Bio Medica Atenei Parmensis* 2010;81:79-84.
43. Yassin DJ, Doros G, Hammerer PG, Yassin AA. Long-term testosterone treatment in elderly men with hypogonadism and erectile dysfunction reduces obesity parameters and improves metabolic syndrome and health-related quality of life. *The journal of sexual medicine* 2014;11:1567-76.
44. Boyanov MA, Boneva Z, Christov VG. Testosterone supplementation in men with type 2 diabetes, visceral obesity and partial androgen deficiency. *The aging male : the official journal of the International Society for the Study of the Aging Male* 2003;6:1-7.