

# UC San Diego

## UC San Diego Previously Published Works

### Title

APOEε4 and slow wave sleep in older adults

### Permalink

<https://escholarship.org/uc/item/6s5387fc>

### Journal

PLOS ONE, 13(1)

### ISSN

1932-6203

### Authors

Tranah, Gregory J  
Yaffe, Kristine  
Nievergelt, Caroline M  
[et al.](#)

### Publication Date

2018

### DOI

10.1371/journal.pone.0191281


Peer reviewed

RESEARCH ARTICLE

# APOE $\epsilon$ 4 and slow wave sleep in older adults

Gregory J. Tranah<sup>1\*</sup>, Kristine Yaffe<sup>2,3,4</sup>, Caroline M. Nievergelt<sup>5</sup>, Neeta Parimi<sup>1</sup>, M. Maria Glymour<sup>4</sup>, Kristine E. Ensrud<sup>6,7,8</sup>, Jane A. Cauley<sup>9</sup>, Sonia Ancoli-Israel<sup>5,10</sup>, Sara Mariani<sup>11</sup>, Susan Redline<sup>12</sup>, Katie L. Stone<sup>1</sup>, the Osteoporotic Fractures in Men Study (MrOS) Research Group<sup>1†</sup>

**1** Research Institute, California Pacific Medical Center, San Francisco, California, United States of America, **2** Departments of Psychiatry and Neurology, University of California San Francisco, San Francisco, California, United States of America, **3** Medical Center, San Francisco VA, San Francisco, California, United States of America, **4** Department of Epidemiology and Biostatistics University of California San Francisco, San Francisco, California, United States of America, **5** Department of Psychiatry, University of California San Diego, La Jolla, California, United States of America, **6** Division of Epidemiology and Community Health, University of Minnesota, Minneapolis, Minnesota, United States of America, **7** Center for Chronic Disease Outcomes Research, Minneapolis VA Medical Center, Minneapolis, Minnesota, United States of America, **8** Department of Medicine, University of Minnesota, Minneapolis, Minnesota, United States of America, **9** Department of Epidemiology, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America, **10** Department of Medicine, University of California San Diego, La Jolla, California, United States of America, **11** Division of Sleep & Circadian Disorders, Brigham & Women's Hospital, Harvard Medical School, Boston, Massachusetts, United States of America, **12** Departments of Medicine, Brigham and Women's Hospital and Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, United States of America



**OPEN ACCESS**

**Citation:** Tranah GJ, Yaffe K, Nievergelt CM, Parimi N, Glymour MM, Ensrud KE, et al. (2018) APOE $\epsilon$ 4 and slow wave sleep in older adults. PLoS ONE 13 (1): e0191281. <https://doi.org/10.1371/journal.pone.0191281>

**Editor:** Stephen D Ginsberg, Nathan S Kline Institute, UNITED STATES

**Received:** August 24, 2017

**Accepted:** January 2, 2018

**Published:** January 25, 2018

**Copyright:** © 2018 Tranah et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data access is legally restricted by the Institutional Review Boards (IRBs) at California Pacific Medical Center and the University of California, San Francisco. Users can access the data through the MrOS Study portal using the following URL: <https://mrosdata.sfcc-cpmc.net/>. Users only have to register for an account and accept the terms of a Data Use Agreement before downloading or exploring clinical data. Additionally, interested researchers may send data access requests to the MrOS Project Director Robin M. Fullman at [rfullman@sfcc-cpmc.net](mailto:rfullman@sfcc-cpmc.net) or to Christina Kouma of

<sup>†</sup> Membership of the (MrOS) Research Group is provided in the Acknowledgments  
\* [gtranah@sfcc-cpmc.net](mailto:gtranah@sfcc-cpmc.net)

## Abstract

Slow wave (or stage N3) sleep has been linked to a variety of cognitive processes. However, the role of stage N3 in the elderly is debated. The link between stage N3 and episodic memory may be weakened or changed in the older adult population, possibly due to several altered mechanisms impacting the cellular structure of the brain. The bases for the age-related dissociation between stage N3 and cognition are not understood. Since APOE $\epsilon$ 4 status is the strongest genetic risk factor for cognitive decline, we assessed whether the  $\epsilon$ 4 allele is associated with stage N3 sleep. Participants were from the population-based Osteoporotic Fractures in Men (MrOS) cohort with polysomnography and APOE $\epsilon$ 4 genotype data (n = 2,302, 100% male, mean age 76.6 years). Sleep stages were objectively measured using overnight in-home polysomnography and central electroencephalogram data were used to score stage N3 sleep. Cognitive function was assessed using the Modified Mini Mental State Exam (3MS). The APOE rs429358 single nucleotide polymorphism, which defines the APOE $\epsilon$ 4 allele, was genotyped using a custom genotyping array. Total time in stage N3 sleep was significantly higher (p<0.0001) among the 40 MrOS participants carrying two copies of the  $\epsilon$ 4 allele (62±5.2 minutes) compared with 43±1.5 minutes for carriers of one  $\epsilon$ 4 allele (n = 515) and 40±0.8 minutes for  $\epsilon$ 4 non-carriers (n = 1747). All results were independent of sleep efficiency, number of sleep cycles, and apnea hypopnea index. These findings support an association between APOE $\epsilon$ 4 genotype and sleep stage N3 in the elderly. Increased total stage N3 duration among  $\epsilon$ 4/ $\epsilon$ 4 carriers does not appear to reflect compensation for prior cognitive decline and may reflect overactive downscaling of

the San Francisco Coordinating Center at [CKouma@sfcc-cpmc.net](mailto:CKouma@sfcc-cpmc.net) (<http://sfcc.ucsf.edu>).

**Funding:** The Osteoporotic Fractures in Men (MrOS) Study is supported by National Institutes of Health (NIH) funding. The following institutes provide support: the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), the National Institute on Aging (NIA), the National Center for Research Resources (NCR), and NIH Roadmap for Medical Research under the following grant numbers: U01 AR45580, U01 AR45614, U01 AR45632, U01 AR45647, U01 AR45654, U01 AR45583, U01 AG18197, U01-AG027810, and UL1 RR024140. The National Heart, Lung, and Blood Institute (NHLBI) provides funding for the MrOS Sleep ancillary study "Outcomes of Sleep Disorders in Older Men" under the following grant numbers: R01 HL071194, R01 HL070848, R01 HL070847, R01 HL070842, R01 HL070841, R01 HL070837, R01 HL070838, and R01 HL070839. NIAMS provides funding for the MrOS ancillary study "Replication of candidate gene associations and bone strength phenotype in MrOS" under the grant number R01-AR051124, which funded the creation and maintenance of MrOS DNA collections. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** This was not an industry supported study. Dr. Ancoli-Israel has been a consultant or on the advisory board of Acadia, Astra Zeneca, Aptalis Pharma, Arena, Merck, NeuroVigil, Inc., Orphagen Pharmaceuticals, and Purdue Pharma LP. This does not alter our adherence to PLOS ONE policies on sharing data and materials. The other authors have indicated no financial conflicts of interest.

synapses during sleep. If confirmed, these results might in part explain the high risk of age-related cognitive decline and AD among APOE  $\epsilon 4/\epsilon 4$  carriers.

## Introduction

Sleep disturbances and specific sleep disorders increase with advancing age [1–6] and elevate the risk for a variety of diseases and conditions [7–22] including cognitive impairment [23–33]. Recent research indicates that sleep may specifically impact the cellular structure of the brain, including the ability of the brain to clear toxic metabolites that accumulate with normal neuronal function [34]. This includes beta-amyloid [34], which accumulates in apolipoprotein E (APOE)  $\epsilon 4$  genotype carriers [35] and contributes to synaptic degeneration and cognitive decline. While the APOE $\epsilon 4$  allele is a well-established risk factor for Alzheimer's disease (AD) and age-related cognitive decline [36–42] as well as a risk factor for sleep apnea [43–51], APOE $\epsilon 4$  has not been assessed in relation to sleep stage distributions. To explore the hypothesis that APOE $\epsilon 4$  influences sleep architecture, we examined the effect of the APOE $\epsilon 4$  allele on sleep stage distribution measured in a large, population-based cohort of elderly participants with a focus on stage N3, which represents the synchronization of cortical EEG activity and has been implicated in memory formation in young populations [52–57]. Given the high risk of cognitive decline and AD among APOE $\epsilon 4$  carriers, identifying a genetic link with stage N3 sleep could advance the development of new approaches for interventions to slow the decline in cognitive function among  $\epsilon 4$  allele carriers and that may also benefit the health and well-being of the elderly.

## Materials and methods

### Participants

Data were collected from participants of the Osteoporotic Fractures in Men (MrOS) Study which included participants who were at least 65 years old and excluded participants who required assistance with ambulation or had undergone bilateral hip replacement. In the MrOS study, 5994 older men were recruited from six geographical areas (Birmingham, AL; the Monongahela Valley near Pittsburgh, PA; Minnaeapolis, MN; Palo Alto, CA; San Diego, CA; and Portland, OR) between 2000 and 2002. The MrOS Sleep Study, an ancillary study of the parent MrOS cohort, was conducted between December 2003 and March 2005 and recruited 3135 MrOS participants (68%) for a comprehensive sleep assessment and assessment of circadian gene polymorphisms. Of the 2,859 men who did not participate in this ancillary study, 344 died before the sleep visit, 36 had already stopped participating in the study, 332 were not invited because recruitment goals had already been met, 150 were not eligible and 1,997 refused. Among the participants of the MrOS Sleep Study, 2480 self-identified white men who had DNA extracted for genetic studies. Details regarding the studies have been published previously [58,59]. All data were collected with written informed consent as approved by the institutional review boards of the clinical sites (University of Alabama at Birmingham; University of Minnesota; University of Pittsburgh; Stanford University; Oregon Health & Science University, and University of California, San Diego) and the coordinating center (University of California, San Francisco). The Outcomes of Sleep Disorders in Older Men Study (MrOS Sleep Study) [60], an ancillary study of the parent MrOS Study, enrolled 3,135 participants from 2003–2005. Of the 3135 enrolled participants, 2911 had usable PSG data and, of these, 2745 (94%) with evaluable sleep staging data. The Modified Mini Mental State Exam [61,62] (3MS),

an expanded 100 point version of the original MMSE, was administered in MrOS at the baseline and sleep exams (~3 years) [63]. The Pittsburgh Sleep Quality Index (PSQI) and Epworth Sleepiness Scale (ESS) questionnaires were administered at the sleep exam.

### Polysomnography

In-home, single night sleep studies using unattended polysomnography (PSG, Safiro, Compumedics, Inc., Melbourne, Australia) were performed as previously described [64,65]. The recording montage consisted of  $C_3/A_2$  and  $C_4/A_1$  electroencephalograms, bilateral electrocuculograms, a bipolar submental electromyogram, thoracic and abdominal respiratory inductance plethysmography, airflow (using nasal-oral thermocouple and nasal pressure cannula), finger pulse oximetry, electrocardiogram, body position (mercury switch sensor), and bilateral leg movements (piezoelectric sensors). Trained certified staff members performed home visits for setup of the sleep study units. After sensors were placed and calibrated, signal quality and impedance were checked, and sensors were repositioned as needed to improve signal quality, replacing electrodes if impedances were elevated, using approaches similar to those in the Sleep Health Heart Study [66]. After studies were downloaded, they were transferred to the Case Western Reserve University Reading Center (Cleveland, OH) for centralized scoring of sleep stages and sleep disorders by a trained technician using standard criteria [67,68]. Sleep was staged in 30-sec scoring epochs and electroencephalogram data were used to score stage N3 using the criteria of Rechtschaffen and Kales [67]. Total time (minutes) and percent time [ $100 \times (\text{Number of minutes scored as stage 3-4}) / (\text{total sleep time})$ ] in stage N3 were calculated for these analyses. The apnea hypopnea index (AHI) was calculated as the total number of apneas and hypopneas per hour of sleep (4% desaturation) [69]. Sleep cycles were defined using a modified version of the rules detailed by Feinberg and Floyd [70]. A cycle was defined as a period of at least 15 minutes of NREM sleep starting with stage N2 or N3 and terminated by the end of a period of REM sleep lasting at least 5 minutes for all cycles except cycle 1, or a period of wake or stage N1 lasting 15 minutes or longer. Two epochs of REM sleep separated by no more than 15 minutes of NREM sleep or wake were merged into a single REM period. Average absolute spectral power density in  $\log_{10}(\mu V^2/\text{Hz})$  was calculated for each sleep cycle for the following bands: Slow oscillation: 0.25–1 Hz, Delta: 1.25–4 Hz, Theta: 4.25–8 Hz, Alpha: 8.25–12 Hz, Sigma: 12.25–15 Hz, and Beta: 15.25–20 Hz. There was excellent inter-scoring reliability for the key sleep parameters, including stage N3 (ICCs >0.92). Several additional sleep parameters were scored, including: wake after sleep onset time and sleep efficiency.

### APOE genotyping

The *APOE* rs429358 SNP, which defines the *APOE*ε4 allele, was genotyped using a custom Illumina Golden Gate assay (Illumina, San Diego, CA, USA) that included 529 SNPs as previously described [71]. All individuals in this study are of European ancestry and are unrelated. Of 2,745 MrOS participants with PSG, 2,302 were genotyped for the rs429358 C/T alleles, which define *APOE*ε4 status by the presence of the C allele. All participants in this analysis are of European ancestry.

### Statistical analyses

Associations between *APOE*ε4 status and sleep staging were based on 2,302 MrOS participants with both PSG and rs429358 genotype data. All PSG and cognition measures were treated as continuous variables in analyses examining both additive and dominant genetic models to examine *APOE*ε4 associations for total time and percent time spent in stage N3. All regression

models were adjusted for age and clinic site with additional models including measures of sleep duration (TST), quality (WASO time and sleep efficiency) and AHI. Cognitive function change (~3 years) from the visit prior to sleep assessment was calculated and used to stratify stage N3-cognition analyses into high and low decline groups. All analyses were analyzed using SAS version 9.4 (SAS Institute Inc, Cary, NC).

## Results

Differences in age, cognition, sleep performance and stage N3 across three *APOEε4* categories are shown in Table 1. Total stage N3 time was significantly higher among 40 MrOS participants carrying two copies of the ε4 allele (62±5.2 minutes) compared with 43±1.5 minutes for carriers of one ε4 allele (n = 515) and 40±0.8 minutes for ε4 non-carriers (n = 1747). The *APOEε4* association with stage N3 time (p = 0.0002) was not altered by adjustment for 3MS (p = 0.0001). We examined the *APOEε4*-stage N3 association among MrOS participants who had experienced either ≥2 point decline or <2 point decline (based on median change) from the previous 3MS assessment to the 3MS measured at the sleep visit. Total stage N3 time was significantly higher among participants carrying two copies of the ε4 allele when compared to the other genotype categories in both the group with ≥2 point 3MS decline (p<0.0001) and the group with <2 point decline (p = 0.0002) and the test for interaction between *APOE* allele and 3MS decline was not significant (p = 0.21). In addition, the *APOEε4* association with 3MS

**Table 1. Characteristics of MrOS participants with PSG sleep staging and *APOE* genotyping.**

	<i>APOEε4</i> alleles					
	0	1	2	p-value <sup>a</sup>	p-value <sup>b</sup>	
n (%)	1747 (75.8)	515 (22.5)	40 (1.7)			
	mean (SE)	mean (SE)	mean (SE)			
Age, y	76.6 (0.1)	76.5 (0.2)	75.6 (0.9)	0.48	0.13	
3MS (0 to 100)*	93.4 (0.1)	92.4 (0.2)	90.2 (0.8)	< .0001	0.003	
Change 3MS <sup>c</sup>	-1.01 (0.1)	-1.47 (0.2)	-3.34 (0.7)	0.0025	0.0026	
Time in stage N3 (min)*	40 (0.8)	43 (1.5)	62 (5.2)	< .0001	< .0001	
	Change 3MS ≥ -2	41 (1.0)	42 (1.8)	61 (7.4)	0.03	0.008
	Change 3MS < -2	38 (1.4)	45 (2.5)	64 (7.5)	0.0005	0.002
Time in REM (min)*	70 (0.7)	68 (1.2)	65 (4.5)	0.21	0.13	
Time in stage N1 (min)*	24 (0.3)	24 (0.6)	22 (2.1)	0.68	0.48	
Sleep efficiency (%)*	76 (0.3)	76 (0.5)	81 (1.9)	0.05	0.01	
Wake after sleep onset (min)	115 (1.6)	116 (2.8)	86 (10)	0.02	0.005	
Apnea hypopnea index (4% desat)*	13 (0.3)	12 (0.6)	14 (2.1)	0.57	0.56	
Number of sleep cycles*	4.7 (0.03)	4.6 (0.1)	4.6 (0.2)	0.50	0.97	
Spectral power density- SOsc (log <sub>10</sub> (uV <sup>2</sup> /Hz))* <sup>d</sup>	1.9 (0.01)	1.9 (0.02)	2.0 (0.6)	0.15	0.06	
Arousal index	24 (0.3)	23 (0.5)	24 (1.9)	0.10	0.93	
PSQI*	5.6 (0.1)	5.5 (0.1)	5.3 (0.5)	0.69	0.55	
ESS*	6.1 (0.1)	6.1 (0.2)	5.7 (0.6)	0.76	0.47	
Hypnotics, n	38	10	0			

\* Adjusted for age

<sup>a</sup> Linear p-value across three genotypes

<sup>b</sup> P-value comparing carriers of two ε4 alleles to all others

<sup>c</sup> Baseline to sleep visit ~ 3 years.

<sup>d</sup> Spectral power density of the C3-A2 EEG in the slow oscillation (0.25–1 Hz) band for the final sleep cycle.

PSQI = Pittsburgh Sleep Quality Index; ESS = Epworth Sleepiness Scale

<https://doi.org/10.1371/journal.pone.0191281.t001>

( $p < 0.0001$ ) was not altered by adjustment for stage N3 time ( $p < 0.0001$ ) and the test for interaction between APOE $\epsilon 4$  and stage N3 time in the association with 3MS was statistically significant ( $p = 0.006$ ). The APOE  $\epsilon 4/\epsilon 4$  genotype was associated with higher sleep efficiency and lower WASO but not the PSQI or ESS assessments. Spectral power density of the C3-A2 EEG in the slow oscillation (0.25–1 Hz) band was calculated for the final sleep cycle and did not differ among APOE alleles (Table 1). The total number of sleep cycles did not differ among APOE alleles (Table 1). The APOE associations with total stage N3 time did not change after additional adjustment for sleep efficiency ( $\epsilon 4/\epsilon 4 = 59 \pm 5.2$  minutes,  $p = 0.0003$ ), wake after sleep onset ( $\epsilon 4/\epsilon 4 = 60 \pm 5.2$  minutes,  $p = 0.0002$ ), number of sleep cycles ( $\epsilon 4/\epsilon 4 = 64 \pm 5.6$  minutes,  $p = 0.0001$ ), or AHI ( $\epsilon 4/\epsilon 4 = 62 \pm 5.1$  minutes,  $p < 0.0001$ ).

## Discussion

In the present study of >2,300 older men, we confirmed a significant association between APOE $\epsilon 4$  status and lower cognitive function scores and unexpectedly found that APOE $\epsilon 4$  was associated with increased stage N3 duration. Average total stage N3 time was >50% longer among elderly APOE  $\epsilon 4/\epsilon 4$  carriers (those who at the highest risk of developing AD and age-related cognitive decline [36–42]) when compared to those with one or no copies of the  $\epsilon 4$  allele. While those with the APOE  $\epsilon 4/\epsilon 4$  genotype had more stage N3 time, stage N3 was not associated with cognitive decline in MrOS [72], even after adjustment for APOE $\epsilon 4$  status. Consistent with increased stage N3 time among  $\epsilon 4/\epsilon 4$  genotype carriers, the APOE  $\epsilon 4/\epsilon 4$  genotype was also associated with higher sleep efficiency and decreased WASO, indicating overall less arousability. This finding was unexpected but could reflect a compensatory mechanism wherein the consolidation process is upregulated in reaction to declining cognition. In order to test this possibility we examined the APOE $\epsilon 4$ -stage N3 association among APOE  $\epsilon 4/\epsilon 4$  participants who had experienced either high or low ~3-year decline from the previous 3MS assessment. Total stage N3 time did not differ between participants who had experienced either high or low declines suggesting that the longer stage N3 time among  $\epsilon 4/\epsilon 4$  carriers is not compensating for prior cognitive decline. It has been suggested that the negative correlations previously observed between stage N3 and subsequent learning in older adults is unlikely due to memory consolidation [73], and might be explained by another function of stage N3 such as synaptic downscaling or pruning [74]. During waking hours, encoding leads to increased synaptic weights which eventually results in the saturation of synaptic networks. The downscaling of certain synapses which takes place during stage N3 likely serves to ensure the maintenance of balanced synaptic input. The negative correlations observed between stage N3 and memory in older adults, however, may reflect an altered balance between daytime encoding and nighttime downscaling wherein older adults engage in less daytime encoding than younger adults [75] but over-downscale if they are still gaining relatively high amounts of stage N3. Thus, an age-related proportional increase in downscaling would presumably become detrimental to memory functioning if synapses that could otherwise help encode new memories are pruned. Such sleep-dependent synaptic downscaling has been observed in *Drosophila* [76,77].

Synaptic plasticity and scaling are crucial mechanisms in memory flexibility and AD onset [73,78,79] with experimental downscaling of AMPA receptors contributing directly to AD pathology [78]. In light of the genetic association data presented herein, it is possible that APOE  $\epsilon 4/\epsilon 4$  carriers are experiencing “overactive downscaling” [73]. APOE is produced by the glia in the central nervous system [80], while APOE receptors are expressed on the neuronal cell surface where they mediate lipoprotein uptake and signal transduction [81,82]. It has been proposed that APOE isoforms may differentially accelerate synapse dysfunction and AD



dementia by interfering with APOE receptor-dependent neuromodulation [81]. Of the three APOE isoforms ( $\epsilon 2$ ,  $\epsilon 3$ , and  $\epsilon 4$ ), APOE $\epsilon 4$  selectively impairs synaptic plasticity and NMDA receptor phosphorylation by Reelin, a modulator of synaptic strength [79]. This selective impairment of APOE receptor signaling by the APOE $\epsilon 4$  isoform leads to a decrease in synaptic responsiveness [79]. The APOE $\epsilon 4$  isoform also reduces glutamate receptor function and synaptic plasticity by selectively impairing APOE receptor recycling in primary neurons [79].

Sleep also impacts the cellular structure of the brain, including the ability of the brain to clear beta-amyloid, contributing to synaptic degeneration and cognitive decline [34]. For example, chronic sleep restriction increases amyloid-B plaque formation in the amyloid precursor protein transgenic mouse [34]. Sleep also impacts the ability of the brain to remove potentially toxic biomolecules that accumulate with normal neuronal function, including beta-amyloid [34], which accumulates in APOE $\epsilon 4$  genotype carriers [35]. This "glymphatic system" is strongly stimulated by sleep and is associated with an increase in interstitial volume, possibly by shrinkage of astroglial cells, leading to clearance activity. Clearance during sleep is as much as two-fold faster than during waking hours [34]. Thus, glymphatic dysfunction may play a role in the pathogenesis of neurodegeneration as well as maintenance of cognition. However, it is also possible that accumulation of beta-amyloid and other toxic metabolites may influence synaptic connectivity, and thus influence neurophysiological mechanisms of sleep, altering sleep state distributions [83].

Few studies have examined the role of APOE genotype in both sleep and cognitive performance [84,85]. In a study of 126 cognitively normal adults (30–70 years), chronic daytime somnolence was associated with a distinctive decline in verbal memory among APOE  $\epsilon 4/\epsilon 4$  carriers as compared to APOE $\epsilon 4$  heterozygotes and noncarriers [84]. A second study examined how sleep parameters in AD patients change over time as a function of APOE $\epsilon 4$  status [85]. Among forty-four community-dwelling adults with a diagnosis of probable AD there was greater deterioration on sleep parameters (increased WASO time and decreased TST and sleep efficiency) in non-APOE $\epsilon 4$  carriers when compared with positive APOE $\epsilon 4$  carriers [85]. Associations between sleep performance and cognition also differed between APOE $\epsilon 4$  carriers and non-carriers [85]. The APOE $\epsilon 4$  allele has also been identified in studies as a risk factor for sleep apnea, although the associations have not been consistent [43–51]. Despite these previous APOE $\epsilon 4$ -sleep associations, the APOE $\epsilon 4$ -stage N3 results reported herein were independent of objective measures of sleep continuity as well as apnea hypopnea index.

In conclusion, these findings support a role for APOE $\epsilon 4$  genotype in the regulation of stage N3 in the elderly. Despite the established role of stage N3 in executive function [52,53] and memory consolidation [54–57] in younger and middle-aged adults, the link between stage N3 and cognitive function may be weakened or changed in older adult population [73,86–89] (including the MrOS study [72]). The negative correlations between stage N3 and subsequent learning in older adults could involve several altered mechanisms [74,90] (including memory consolidation, synaptic downscaling, or other macromolecular changes during stage N3 [83]). As demonstrated in previous studies, increased stage N3 might predict worse episodic memory consolidation in the elderly [88] and may be detrimental to morning recall [73,91–94]. Similar changes to the sleep–memory link have also been reported for old rodents [95,96] showing that normal hippocampal reactivation is not preserved in older rats and that deep sleep was associated with worse memory performance as measured by performance on the Morris Water Maze task. Further research is needed to confirm the APOE $\epsilon 4$ -stage N3 associations and to identify the mechanism(s) by which APOE $\epsilon 4$  alterations impact sleep architecture. Quantitative EEG analysis will be useful also to elucidate whether APOE influences homeostatic drive and sleep dynamics, and whether such changes associate with cognition. The important strengths of this study are the large number of participants with overnight PSG,

longitudinal cognitive function measures, and *APOE*ε4 genotyping. An additional study strength is the use of a large cohort of older adults who were not selected on the basis of sleep or cognition characteristics. This study, however, was limited to white men and may not be generalizable to other populations or women. If it is confirmed that increased total stage N3 time among *APOE* ε4/ε4 carriers reflects an overactive downscaling of synapses during sleep, these results might in part explain the high risk of AD and age-related cognitive decline among *APOE* ε4/ε4 carriers.

## Acknowledgments

The Osteoporotic Fractures in Men (MrOS) Study is supported by National Institutes of Health (NIH) funding. The following institutes provide support: the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), the National Institute on Aging (NIA), the National Center for Research Resources (NCRR), and NIH Roadmap for Medical Research under the following grant numbers: U01 AR45580, U01 AR45614, U01 AR45632, U01 AR45647, U01 AR45654, U01 AR45583, U01 AG18197, U01-AG027810, and UL1 RR024140. The National Heart, Lung, and Blood Institute (NHLBI) provides funding for the MrOS Sleep ancillary study “Outcomes of Sleep Disorders in Older Men” under the following grant numbers: R01 HL071194, R01 HL070848, R01 HL070847, R01 HL070842, R01 HL070841, R01 HL070837, R01 HL070838, and R01 HL070839. NIAMS provides funding for the MrOS ancillary study “Replication of candidate gene associations and bone strength phenotype in MrOS” under the grant number R01-AR051124, which funded the creation and maintenance of MrOS DNA collections.

The Osteoporotic Fractures in Men (MrOS) Study Research Group includes: SR Cummings as lead author (California Pacific Medical Center Research Institute, San Francisco, CA, USA, [scummings@sfcc-cpmc.net](mailto:scummings@sfcc-cpmc.net)); E Barrett-Connor University of California, San Diego, San Diego, CA, USA; JA Cauley University of Pittsburgh, Pittsburgh, PA, USA; PM Cawthon, California Pacific Medical Center Research Institute, San Francisco, CA, USA; KE Ensrud Department of Medicine, University of Minnesota, Minneapolis, MN, USA; DC Bauer University of California, San Francisco, San Francisco, CA, USA; JM Shikany University of Alabama at Birmingham, Birmingham, AL, USA; AR Hoffman Stanford University, Stanford, CA, USA; DM Kado University of California, San Diego, San Diego, CA, USA; ML Stefanick Stanford University, Stanford, CA, USA; ES Orwoll Oregon Health & Science University, Portland, OR, USA; JM Zmuda University of Pittsburgh, Pittsburgh, PA, USA; CE Lewis University of Alabama at Birmingham, Birmingham, AL, USA.

## Author Contributions

**Conceptualization:** Gregory J. Tranah, Kristine Yaffe, Caroline M. Nievergelt, M. Maria Glymour, Kristine E. Ensrud, Jane A. Cauley, Sonia Ancoli-Israel, Susan Redline, Katie L. Stone.

**Data curation:** Gregory J. Tranah, Caroline M. Nievergelt, Neeta Parimi, Sara Mariani, Katie L. Stone.

**Formal analysis:** Gregory J. Tranah, Neeta Parimi.

**Funding acquisition:** Gregory J. Tranah, Kristine E. Ensrud, Jane A. Cauley, Sonia Ancoli-Israel, Katie L. Stone.

**Investigation:** Gregory J. Tranah, Caroline M. Nievergelt, M. Maria Glymour, Kristine E. Ensrud, Jane A. Cauley, Sonia Ancoli-Israel, Susan Redline, Katie L. Stone.



**Methodology:** Gregory J. Tranah, Caroline M. Nievergelt, M. Maria Glymour, Jane A. Cauley, Sonia Ancoli-Israel, Sara Mariani, Susan Redline, Katie L. Stone.

**Project administration:** Katie L. Stone.

**Resources:** Kristine E. Ensrud, Jane A. Cauley, Sara Mariani, Katie L. Stone.

**Supervision:** Gregory J. Tranah, Kristine E. Ensrud, Jane A. Cauley, Katie L. Stone.

**Writing – review & editing:** Katie L. Stone.

**Writing – review & editing:** Kristine Yaffe.

## References

1. Foley DJ, Monjan AA, Brown SL, Simonsick EM, Wallace RB, et al. (1995) Sleep complaints among elderly persons: an epidemiologic study of three communities. *Sleep* 18: 425–432. PMID: [7481413](#)
2. Ancoli-Israel S, Klauber MR, Stepnowsky C, Estline E, Chinn A, et al. (1995) Sleep-disordered breathing in African-American elderly. *Am J Respir Crit Care Med* 152: 1946–1949. <https://doi.org/10.1164/ajrccm.152.6.8520760> PMID: [8520760](#)
3. Ancoli-Israel S, Kripke DF, Klauber MR, Mason WJ, Fell R, et al. (1991) Sleep-disordered breathing in community-dwelling elderly. *Sleep* 14: 486–495. PMID: [1798880](#)
4. Ancoli-Israel S, Kripke DF, Klauber MR, Parker L, Stepnowsky C, et al. (1993) Natural history of sleep disordered breathing in community dwelling elderly. *Sleep* 16: S25–29. PMID: [8178017](#)
5. Ancoli-Israel S, Alessi C (2005) Sleep and aging. *Am J Geriatr Psychiatry* 13: 341–343. <https://doi.org/10.1176/appi.ajgp.13.5.341> PMID: [15879581](#)
6. Ancoli-Israel S, Kripke DF (1991) Prevalent sleep problems in the aged. *Biofeedback Self Regul* 16: 349–359. PMID: [1760457](#)
7. Al-Delaimy WK, Manson JE, Willett WC, Stampfer MJ, Hu FB (2002) Snoring as a risk factor for type II diabetes mellitus: a prospective study. *Am J Epidemiol* 155: 387–393. PMID: [11867347](#)
8. Ayas NT, White DP, Al-Delaimy WK, Manson JE, Stampfer MJ, et al. (2003) A prospective study of self-reported sleep duration and incident diabetes in women. *Diabetes Care* 26: 380–384. PMID: [12547866](#)
9. Elmasry A, Janson C, Lindberg E, Gislason T, Tageldin MA, et al. (2000) The role of habitual snoring and obesity in the development of diabetes: a 10-year follow-up study in a male population. *J Intern Med* 248: 13–20. PMID: [10947876](#)
10. Ficker JH, Dertinger SH, Siegfried W, Konig HJ, Pentz M, et al. (1998) Obstructive sleep apnoea and diabetes mellitus: the role of cardiovascular autonomic neuropathy. *Eur Respir J* 11: 14–19. PMID: [9543264](#)
11. Grunstein RR, Stenlof K, Hedner J, Sjoström L (1995) Impact of obstructive sleep apnea and sleepiness on metabolic and cardiovascular risk factors in the Swedish Obese Subjects (SOS) Study. *Int J Obes Relat Metab Disord* 19: 410–418. PMID: [7550526](#)
12. Jennum P, Sjol A (1993) Snoring, sleep apnoea and cardiovascular risk factors: the MONICA II Study. *Int J Epidemiol* 22: 439–444. PMID: [8359959](#)
13. Newman AB, Enright PL, Manolio TA, Haponik EF, Wahl PW (1997) Sleep disturbance, psychosocial correlates, and cardiovascular disease in 5201 older adults: the Cardiovascular Health Study. *J Am Geriatr Soc* 45: 1–7. PMID: [8994480](#)
14. Newman AB, Nieto FJ, Guidry U, Lind BK, Redline S, et al. (2001) Relation of sleep-disordered breathing to cardiovascular disease risk factors: the Sleep Heart Health Study. *Am J Epidemiol* 154: 50–59. PMID: [11434366](#)
15. Newman AB, Spiekerman CF, Enright P, Lefkowitz D, Manolio T, et al. (2000) Daytime sleepiness predicts mortality and cardiovascular disease in older adults. The Cardiovascular Health Study Research Group. *J Am Geriatr Soc* 48: 115–123. PMID: [10682939](#)
16. Shahar E, Whitney CW, Redline S, Lee ET, Newman AB, et al. (2001) Sleep-disordered breathing and cardiovascular disease: cross-sectional results of the Sleep Heart Health Study. *Am J Respir Crit Care Med* 163: 19–25. <https://doi.org/10.1164/ajrccm.163.1.2001008> PMID: [11208620](#)
17. Ancoli-Israel S, Kripke DF, Klauber MR, Fell R, Stepnowsky C, et al. (1996) Morbidity, mortality and sleep-disordered breathing in community dwelling elderly. *Sleep* 19: 277–282. PMID: [8776783](#)
18. Kripke DF, Garfinkel L, Wingard DL, Klauber MR, Marler MR (2002) Mortality associated with sleep duration and insomnia. *Arch Gen Psychiatry* 59: 131–136. PMID: [11825133](#)

19. Manabe K, Matsui T, Yamaya M, Sato-Nakagawa T, Okamura N, et al. (2000) Sleep patterns and mortality among elderly patients in a geriatric hospital. *Gerontology* 46: 318–322. <https://doi.org/10.1159/000022184> PMID: 11044786
20. Mant A, King M, Saunders NA, Pond CD, Goode E, et al. (1995) Four-year follow-up of mortality and sleep-related respiratory disturbance in non-demented seniors. *Sleep* 18: 433–438. PMID: 7481414
21. Rockwood K, Davis HS, Merry HR, MacKnight C, McDowell I (2001) Sleep disturbances and mortality: results from the Canadian Study of Health and Aging. *J Am Geriatr Soc* 49: 639–641. PMID: 11380758
22. Stone KL, Ewing SK, Ancoli-Israel S, Ensrud KE, Redline S, et al. (2009) Self-reported sleep and nap habits and risk of mortality in a large cohort of older women. *J Am Geriatr Soc* 57: 604–611. <https://doi.org/10.1111/j.1532-5415.2008.02171.x> PMID: 19220560
23. Blackwell T, Yaffe K, Ancoli-Israel S, Schneider JL, Cauley JA, et al. (2006) Poor sleep is associated with impaired cognitive function in older women: the study of osteoporotic fractures. *J Gerontol A Biol Sci Med Sci* 61: 405–410. PMID: 16611709
24. Cricco M, Simonsick EM, Foley DJ (2001) The impact of insomnia on cognitive functioning in older adults. *J Am Geriatr Soc* 49: 1185–1189. PMID: 11559377
25. Meguro K, Ueda M, Kobayashi I, Yamaguchi S, Yamazaki H, et al. (1995) Sleep disturbance in elderly patients with cognitive impairment, decreased daily activity and periventricular white matter lesions. *Sleep* 18: 109–114. PMID: 7792490
26. Ohayon MM, Vecchierini MF (2002) Daytime sleepiness and cognitive impairment in the elderly population. *Arch Intern Med* 162: 201–208. PMID: 11802754
27. Bonnet MH (1993) Cognitive effects of sleep and sleep fragmentation. *Sleep* 16: S65–67. PMID: 8178030
28. Crenshaw MC, Edinger JD (1999) Slow-wave sleep and waking cognitive performance among older adults with and without insomnia complaints. *Physiol Behav* 66: 485–492. PMID: 10357438
29. Dealberto MJ, Pajot N, Courbon D, Alperovitch A (1996) Breathing disorders during sleep and cognitive performance in an older community sample: the EVA Study. *J Am Geriatr Soc* 44: 1287–1294. PMID: 8909342
30. Edinger JD, Glenn DM, Bastian LA, Marsh GR (2000) Slow-wave sleep and waking cognitive performance II: Findings among middle-aged adults with and without insomnia complaints. *Physiol Behav* 70: 127–134. PMID: 10978487
31. Engleman HM, Kingshott RN, Martin SE, Douglas NJ (2000) Cognitive function in the sleep apnea/hypopnea syndrome (SAHS). *Sleep* 23 Suppl 4: S102–108.
32. Granat P (1990) Sleep loss and cognitive performance. *J Fam Pract* 30: 632.
33. McCann UD, Penetar DM, Shaham Y, Thorne DR, Gillin JC, et al. (1992) Sleep deprivation and impaired cognition. Possible role of brain catecholamines. *Biol Psychiatry* 31: 1082–1097. PMID: 1356024
34. Xie L, Kang H, Xu Q, Chen MJ, Liao Y, et al. (2013) Sleep drives metabolite clearance from the adult brain. *Science* 342: 373–377. <https://doi.org/10.1126/science.1241224> PMID: 24136970
35. Lim YY, Mormino EC, Alzheimer's Disease Neuroimaging I (2017) APOE genotype and early beta-amyloid accumulation in older adults without dementia. *Neurology*.
36. Berlau DJ, Corrada MM, Head E, Kawas CH (2009) APOE epsilon2 is associated with intact cognition but increased Alzheimer pathology in the oldest old. *Neurology* 72: 829–834. <https://doi.org/10.1212/01.wnl.0000343853.00346.a4> PMID: 19255410
37. Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, et al. (1993) Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 261: 921–923. PMID: 8346443
38. Burt TD, Agan BK, Marconi VC, He W, Kulkarni H, et al. (2008) Apolipoprotein (apo) E4 enhances HIV-1 cell entry in vitro, and the APOE epsilon4/epsilon4 genotype accelerates HIV disease progression. *Proc Natl Acad Sci U S A* 105: 8718–8723. <https://doi.org/10.1073/pnas.0803526105> PMID: 18562290
39. Mahley RW, Rall SC Jr. (2000) Apolipoprotein E: far more than a lipid transport protein. *Annu Rev Genomics Hum Genet* 1: 507–537. <https://doi.org/10.1146/annurev.genom.1.1.507> PMID: 11701639
40. Weisgraber KH, Innerarity TL, Newhouse YM, Young SG, Arnold KS, et al. (1988) Familial defective apolipoprotein B-100: enhanced binding of monoclonal antibody MB47 to abnormal low density lipoproteins. *Proc Natl Acad Sci U S A* 85: 9758–9762. PMID: 3200853
41. Cosentino S, Scarmeas N, Helzner E, Glymour MM, Brandt J, et al. (2008) APOE epsilon 4 allele predicts faster cognitive decline in mild Alzheimer disease. *Neurology* 70: 1842–1849. <https://doi.org/10.1212/01.wnl.0000304038.37421.cc> PMID: 18401023

42. Mahley RW, Huang Y (2006) Apolipoprotein (apo) E4 and Alzheimer's disease: unique conformational and biophysical properties of apoE4 can modulate neuropathology. *Acta Neurol Scand Suppl* 185: 8–14. <https://doi.org/10.1111/j.1600-0404.2006.00679.x> PMID: 16866905
43. Cosentino FI, Bosco P, Drago V, Prestianni G, Lanuzza B, et al. (2008) The APOE epsilon4 allele increases the risk of impaired spatial working memory in obstructive sleep apnea. *Sleep Med* 9: 831–839. <https://doi.org/10.1016/j.sleep.2007.10.015> PMID: 18083630
44. Craig D, Hart DJ, Passmore AP (2006) Genetically increased risk of sleep disruption in Alzheimer's disease. *Sleep* 29: 1003–1007. PMID: 16944667
45. Foley DJ, Masaki K, White L, Redline S (2001) Relationship between apolipoprotein E epsilon4 and sleep-disordered breathing at different ages. *Jama* 286: 1447–1448. PMID: 11572727
46. Gozal D, Capdevila OS, Kheirandish-Gozal L, Crabtree VM (2007) APOE epsilon 4 allele, cognitive dysfunction, and obstructive sleep apnea in children. *Neurology* 69: 243–249. <https://doi.org/10.1212/01.wnl.0000265818.88703.83> PMID: 17636061
47. Kadotani H, Kadotani T, Young T, Peppard PE, Finn L, et al. (2001) Association between apolipoprotein E epsilon4 and sleep-disordered breathing in adults. *Jama* 285: 2888–2890. PMID: 11401610
48. Larkin EK, Patel SR, Redline S, Mignot E, Elston RC, et al. (2006) Apolipoprotein E and obstructive sleep apnea: evaluating whether a candidate gene explains a linkage peak. *Genet Epidemiol* 30: 101–110. <https://doi.org/10.1002/gepi.20127> PMID: 16374834
49. Saarelainen S, Lehtimaki T, Kallonen E, Laasonen K, Poussa T, et al. (1998) No relation between apolipoprotein E alleles and obstructive sleep apnea. *Clin Genet* 53: 147–148. PMID: 9611077
50. Gottlieb DJ, DeStefano AL, Foley DJ, Mignot E, Redline S, et al. (2004) APOE epsilon4 is associated with obstructive sleep apnea/hypopnea: the Sleep Heart Health Study. *Neurology* 63: 664–668. PMID: 15326239
51. Kalra M, Pal P, Kaushal R, Amin RS, Dolan LM, et al. (2008) Association of ApoE genetic variants with obstructive sleep apnea in children. *Sleep Med* 9: 260–265. <https://doi.org/10.1016/j.sleep.2007.05.001> PMID: 17658295
52. Aeschbach D, Cutler AJ, Ronda JM (2008) A role for non-rapid-eye-movement sleep homeostasis in perceptual learning. *J Neurosci* 28: 2766–2772. <https://doi.org/10.1523/JNEUROSCI.5548-07.2008> PMID: 18337406
53. Landsness EC, Crupi D, Hulse BK, Peterson MJ, Huber R, et al. (2009) Sleep-dependent improvement in visuomotor learning: a causal role for slow waves. *Sleep* 32: 1273–1284. PMID: 19848357
54. Wilson MA, McNaughton BL (1994) Reactivation of hippocampal ensemble memories during sleep. *Science* 265: 676–679. PMID: 8036517
55. Yaroush R, Sullivan MJ, Ekstrand BR (1971) Effect of sleep on memory. II. Differential effect of the first and second half of the night. *J Exp Psychol* 88: 361–366. PMID: 4326302
56. Diekelmann S, Wilhelm I, Born J (2009) The whats and whens of sleep-dependent memory consolidation. *Sleep Med Rev* 13: 309–321. <https://doi.org/10.1016/j.smrv.2008.08.002> PMID: 19251443
57. Walker MP (2009) The role of sleep in cognition and emotion. *Ann N Y Acad Sci* 1156: 168–197. <https://doi.org/10.1111/j.1749-6632.2009.04416.x> PMID: 19338508
58. Cummings SR, Black DM, Nevitt MC, Browner W, Cauley J, et al. (1993) Bone density at various sites for prediction of hip fractures. The Study of Osteoporotic Fractures Research Group. *Lancet* 341: 72–75. PMID: 8093403
59. Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12: 189–198. PMID: 1202204
60. Rao MN, Blackwell T, Redline S, Stefanick ML, Ancoli-Israel S, et al. (2009) Association between sleep architecture and measures of body composition. *Sleep* 32: 483–490. PMID: 19413142
61. Teng EL, Chui HC, Schneider LS, Metzger LE (1987) Alzheimer's dementia: performance on the Mini-Mental State Examination. *J Consult Clin Psychol* 55: 96–100. PMID: 3571664
62. Teng EL, Chui HC (1987) The Modified Mini-Mental State (3MS) examination. *J Clin Psychiatry* 48: 314–318. PMID: 3611032
63. Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12: 189–198. PMID: 1202204
64. Blackwell T, Redline S, Ancoli-Israel S, Schneider JL, Surovec S, et al. (2008) Comparison of sleep parameters from actigraphy and polysomnography in older women: the SOF study. *Sleep* 31: 283–291. PMID: 18274276
65. Mehra R, Stone KL, Blackwell T, Ancoli Israel S, Dam TT, et al. (2007) Prevalence and correlates of sleep-disordered breathing in older men: osteoporotic fractures in men sleep study. *J Am Geriatr Soc* 55: 1356–1364. <https://doi.org/10.1111/j.1532-5415.2007.01290.x> PMID: 17767677

66. Redline S, Sanders MH, Lind BK, Quan SF, Iber C, et al. (1998) Methods for obtaining and analyzing unattended polysomnography data for a multicenter study. Sleep Heart Health Research Group. *Sleep* 21: 759–767. PMID: [11300121](https://pubmed.ncbi.nlm.nih.gov/11300121/)
67. Hori T, Sugita Y, Koga E, Shirakawa S, Inoue K, et al. (2001) Proposed supplements and amendments to 'A Manual of Standardized Terminology, Techniques and Scoring System for Sleep Stages of Human Subjects', the Rechtschaffen & Kales (1968) standard. *Psychiatry Clin Neurosci* 55: 305–310. <https://doi.org/10.1046/j.1440-1819.2001.00810.x> PMID: [11422885](https://pubmed.ncbi.nlm.nih.gov/11422885/)
68. (1992) EEG arousals: scoring rules and examples: a preliminary report from the Sleep Disorders Atlas Task Force of the American Sleep Disorders Association. *Sleep* 15: 173–184. PMID: [11032543](https://pubmed.ncbi.nlm.nih.gov/11032543/)
69. Khan A, Harrison SL, Kezirian EJ, Ancoli-Israel S, O'Hearn D, et al. (2013) Obstructive sleep apnea during rapid eye movement sleep, daytime sleepiness, and quality of life in older men in Osteoporotic Fractures in Men (MrOS) Sleep Study. *J Clin Sleep Med* 9: 191–198. <https://doi.org/10.5664/jcsm.2474> PMID: [23493839](https://pubmed.ncbi.nlm.nih.gov/23493839/)
70. Feinberg I, Floyd TC (1979) Systematic trends across the night in human sleep cycles. *Psychophysiology* 16: 283–291. PMID: [220659](https://pubmed.ncbi.nlm.nih.gov/220659/)
71. Evans DS, Parimi N, Nievergelt CM, Blackwell T, Redline S, et al. (2013) Common Genetic Variants in ARNTL and NPAS2 and at Chromosome 12p13 are Associated with Objectively Measured Sleep Traits in the Elderly. *Sleep* 36: 431–446. <https://doi.org/10.5665/sleep.2466> PMID: [23449886](https://pubmed.ncbi.nlm.nih.gov/23449886/)
72. Blackwell T, Yaffe K, Ancoli-Israel S, Redline S, Ensrud KE, et al. (2011) Associations between sleep architecture and sleep-disordered breathing and cognition in older community-dwelling men: the Osteoporotic Fractures in Men Sleep Study. *J Am Geriatr Soc* 59: 2217–2225. <https://doi.org/10.1111/j.1532-5415.2011.03731.x> PMID: [22188071](https://pubmed.ncbi.nlm.nih.gov/22188071/)
73. Scullin MK (2013) Sleep, memory, and aging: the link between slow-wave sleep and episodic memory changes from younger to older adults. *Psychol Aging* 28: 105–114. <https://doi.org/10.1037/a0028830> PMID: [22708533](https://pubmed.ncbi.nlm.nih.gov/22708533/)
74. Tononi G, Cirelli C (2003) Sleep and synaptic homeostasis: a hypothesis. *Brain Res Bull* 62: 143–150. PMID: [14638388](https://pubmed.ncbi.nlm.nih.gov/14638388/)
75. Cirelli C (2012) Brain plasticity, sleep and aging. *Gerontology* 58: 441–445. <https://doi.org/10.1159/000336149> PMID: [22398514](https://pubmed.ncbi.nlm.nih.gov/22398514/)
76. Donlea JM, Ramanan N, Shaw PJ (2009) Use-dependent plasticity in clock neurons regulates sleep need in *Drosophila*. *Science* 324: 105–108. <https://doi.org/10.1126/science.1166657> PMID: [19342592](https://pubmed.ncbi.nlm.nih.gov/19342592/)
77. Gilestro GF, Tononi G, Cirelli C (2009) Widespread changes in synaptic markers as a function of sleep and wakefulness in *Drosophila*. *Science* 324: 109–112. <https://doi.org/10.1126/science.1166673> PMID: [19342593](https://pubmed.ncbi.nlm.nih.gov/19342593/)
78. Chang EH, Savage MJ, Flood DG, Thomas JM, Levy RB, et al. (2006) AMPA receptor downscaling at the onset of Alzheimer's disease pathology in double knockin mice. *Proc Natl Acad Sci U S A* 103: 3410–3415. <https://doi.org/10.1073/pnas.0507313103> PMID: [16492745](https://pubmed.ncbi.nlm.nih.gov/16492745/)
79. Chen Y, Durakoglugil MS, Xian X, Herz J (2010) ApoE4 reduces glutamate receptor function and synaptic plasticity by selectively impairing ApoE receptor recycling. *Proc Natl Acad Sci U S A* 107: 12011–12016. <https://doi.org/10.1073/pnas.0914984107> PMID: [20547867](https://pubmed.ncbi.nlm.nih.gov/20547867/)
80. Mahley RW, Weisgraber KH, Huang Y (2006) Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc Natl Acad Sci U S A* 103: 5644–5651. <https://doi.org/10.1073/pnas.0600549103> PMID: [16567625](https://pubmed.ncbi.nlm.nih.gov/16567625/)
81. Herz J, Chen Y (2006) Reelin, lipoprotein receptors and synaptic plasticity. *Nat Rev Neurosci* 7: 850–859. <https://doi.org/10.1038/nrn2009> PMID: [17053810](https://pubmed.ncbi.nlm.nih.gov/17053810/)
82. Chen Y, Beffert U, Ertunc M, Tang TS, Kavalali ET, et al. (2005) Reelin modulates NMDA receptor activity in cortical neurons. *J Neurosci* 25: 8209–8216. <https://doi.org/10.1523/JNEUROSCI.1951-05.2005> PMID: [16148228](https://pubmed.ncbi.nlm.nih.gov/16148228/)
83. Mackiewicz M, Shockley KR, Romer MA, Galante RJ, Zimmerman JE, et al. (2007) Macromolecule biosynthesis: a key function of sleep. *Physiol Genomics* 31: 441–457. <https://doi.org/10.1152/physiolgenomics.00275.2006> PMID: [17698924](https://pubmed.ncbi.nlm.nih.gov/17698924/)
84. Caselli RJ, Reiman EM, Hentz JG, Osborne D, Alexander GE, et al. (2002) A distinctive interaction between memory and chronic daytime somnolence in asymptomatic APOE e4 homozygotes. *Sleep* 25: 447–453. PMID: [12071547](https://pubmed.ncbi.nlm.nih.gov/12071547/)
85. Yesavage JA, Friedman L, Kraemer H, Tinklenberg JR, Salehi A, et al. (2004) Sleep/wake disruption in Alzheimer's disease: APOE status and longitudinal course. *J Geriatr Psychiatry Neurol* 17: 20–24. <https://doi.org/10.1177/0891988703261994> PMID: [15018693](https://pubmed.ncbi.nlm.nih.gov/15018693/)
86. Pace-Schott EF, Spencer RM (2011) Age-related changes in the cognitive function of sleep. *Prog Brain Res* 191: 75–89. <https://doi.org/10.1016/B978-0-444-53752-2.00012-6> PMID: [21741545](https://pubmed.ncbi.nlm.nih.gov/21741545/)

87. Spiegel R, Koberle S, Allen SR (1986) Significance of slow wave sleep: considerations from a clinical viewpoint. *Sleep* 9: 66–79. PMID: [3961369](#)
88. Rauchs G, Schabus M, Parapatics S, Bertran F, Clochon P, et al. (2008) Is there a link between sleep changes and memory in Alzheimer's disease? *Neuroreport* 19: 1159–1162. <https://doi.org/10.1097/WNR.0b013e32830867c4> PMID: [18596620](#)
89. Peters KR, Ray L, Smith V, Smith C (2008) Changes in the density of stage 2 sleep spindles following motor learning in young and older adults. *J Sleep Res* 17: 23–33. <https://doi.org/10.1111/j.1365-2869.2008.00634.x> PMID: [18275552](#)
90. Soudmand R, Ward LC, Swift TR (1982) Effect of height on nerve conduction velocity. *Neurology* 32: 407–410. PMID: [7199664](#)
91. Cai ZJ (1995) An integrative analysis to sleep functions. *Behav Brain Res* 69: 187–194. PMID: [7546310](#)
92. Mazzone G, Gori S, Formicola G, Gneri C, Massetani R, et al. (1999) Word recall correlates with sleep cycles in elderly subjects. *J Sleep Res* 8: 185–188. PMID: [10476004](#)
93. Feinberg I (1974) Changes in sleep cycle patterns with age. *J Psychiatr Res* 10: 283–306. PMID: [4376564](#)
94. Foulkes D (1993) Symposium: Normal and abnormal REM sleep regulation: Dreaming and REM sleep. *J Sleep Res* 2: 199–202. PMID: [10607095](#)
95. Gerrard JL, Burke SN, McNaughton BL, Barnes CA (2008) Sequence reactivation in the hippocampus is impaired in aged rats. *J Neurosci* 28: 7883–7890. <https://doi.org/10.1523/JNEUROSCI.1265-08.2008> PMID: [18667620](#)
96. Buechel HM, Popovic J, Searcy JL, Porter NM, Thibault O, et al. (2011) Deep sleep and parietal cortex gene expression changes are related to cognitive deficits with age. *PLoS One* 6: e18387. <https://doi.org/10.1371/journal.pone.0018387> PMID: [21483696](#)