

UCLA

UCLA Previously Published Works

Title

Complete genomes of two Variovorax endophytes isolated from surface-sterilized alfalfa nodules.

Permalink

<https://escholarship.org/uc/item/6sb8f6pm>

Authors

Hirsch, Ann M

Humm, Ethan

Rubbi, Mila

et al.

Publication Date

2024-07-05

DOI

10.1128/mra.00336-24

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Complete genomes of two *Variovorax* endophytes isolated from surface-sterilized alfalfa nodules

Ann M. Hirsch,¹ Ethan Humm,² Mila Rubbi,¹ Giorgia del Vecchio,¹ Sung Min Ha,³ Matteo Pellegrini,^{1,4} Robert P. Gunsalus^{2,4}

AUTHOR AFFILIATIONS See affiliation list on p. 2.

ABSTRACT *Variovorax* species catabolize a wide range of natural and industrial products and have been shown to be integral rhizosphere inhabitants. Here, we report the complete genomes of *V. paradoxus* 2u118 and *V. sp.* SPNA7, which were isolated from alfalfa root nodules and possess plant growth-promoting properties.

KEYWORDS endophytes, plant growth-promotion, alfalfa nodule, bioremediation, *Variovorax*

The genus *Variovorax* is known to metabolize a wide range of substrates, including pesticides (1, 2), acyl homoserine lactones (3), and acrylamide (4). *Variovorax* species have the potential as plant growth-promoting bacteria via several strategies including lowering plant ethylene levels (5) and remediating metal-contaminated soil (6), and have been identified as keystone species for maintaining root growth in *Arabidopsis* (7). In a study of alfalfa nodule-associated bacteria, two *Variovorax* isolates were collected from an alfalfa field at CalPoly Pomona (34.045075, -117.812530). Surface-sterilized nodules were crushed with a mortar and pestle, serial dilutions were plated on LB agar, and single colonies were picked after incubation at 30°C for 1 week and streaked to obtain pure cultures. DNA was extracted using a Quick-DNA HMW Magbead Kit (Zymo Research) per the manufacturer's instructions, and fragmented using Covaris gTubes following instructions from the manufacturer (4 passes at 7,000 rpm through the gTube orifice). The average size of the sheared gDNA was checked at the TapeStation 4200 (Agilent). Multiplexed microbial libraries were prepared using the PacBio SMRTbell prep kit 3.0 together with the SMRTbell barcoded adapters 3.0 according to the PacBio protocol. Final whole genome libraries were not size-selected but simply purified via a standard procedure using 1× SMRTbell clean-up beads. DNA sequencing was performed using the PacBio Sequel IIe platform. Demultiplexing and adapter trimming were done using Lima v2.9.0 (<https://github.com/pacificbiosciences/barcoding>). All reads were then targeted for genome assembly by Canu v2.2 (8) and the assembled genomes were further refined by Circlator v1.5.5 (9) to identify circular contigs, remove redundant non-circular contigs, and rotate circular contigs to start with *dnaA*. This resulted in circular genomes and plasmids (Table 1). A completeness check was performed by CheckM v1.0.18 (10) and the N50 value was determined by Assembly stats ver1.01 (<https://github.com/sanger-pathogens/assembly-stats>). High-quality reads, completeness, and N50 quality values for each strain were as follows: *V. sp.* SPNA7: 38,196, 100%, and 5,887,536 bp; *V. paradoxus* 2u118: 23,568, 100%, and 5,622,806 bp. Genome ORF calling and annotation were performed by NCBI's PGAP v6.6 (11) and the IMG Annotation Pipeline v.5.1.17 (12). All software tools used default parameters that were stated in each tool's manual.

Properties of the finished genomes of each *Variovorax* strain are summarized in Table 1. All 16S sequences had greater than 99.45% similarity to the published 16S sequences of *V. paradoxus* NBRC 15149^T (13, 14). ANI values against *V. paradoxus* NBRC 15149^T were 93.94% and 95.27% for *V. sp.* SPNA7 and *V. paradoxus* 2u118, suggesting that SPNA7

Editor Irene L. G. Newton, Indiana University, Bloomington, Indiana, USA

Address correspondence to Ann M. Hirsch, ahirsch@ucla.edu.

The authors declare no conflict of interest.

See the funding table on p. 2.

Received 1 April 2024

Accepted 13 June 2024

Published 5 July 2024

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

TABLE 1 *V. sp.* SPNA7 and *V. paradoxus* 2u118 genome information

Strain	Contig	Topology	Size (bp)	GC%	Coverage	Protein coding	# 16S	# tRNA
SPNA7	#1	Circular	5,887,536	67.5	39.0×	5,459	2	46
	#2	Circular	1,192,185	67.5	39.0×	1,112	0	10
	Total	n/a ^a	7,079,721	67.5	39.0×	6,571	2	56
2u118	#1	Circular	5,622,806	67.5	37.0×	5,238	2	46
	#2	Circular	1,298,350	67	37.0×	1,241	0	0
	Total	n/a	6,921,156	67.5	37.0×	6,479	2	46

^an/a: not available.

could potentially represent a new species. ANI was calculated using contigs and the Ezbiocloud ANI Calculator (15).

Both genomes are enriched in genes related to heavy metal resistance (MerR family copper efflux transcriptional regulators, copper-responsive two-component system CusR/CusS, arsenate reductase, etc.) and xenobiotics degradation (including genes related to degradation of chloroalkanes, dioxins, styrene, toluene, xylene, etc.). Xenobiotics degradation/metabolism accounts for 3.73%–3.79% of genes assigned to KEGG categories. Additionally, both genomes encode the enzyme ACC deaminase (*acdS*), genes for indole-3-acetic acid (IAA) biosynthesis, and acetoin and trehalose biosynthesis, all of which may contribute to plant growth promotion.

ACKNOWLEDGMENTS

We thank the UCLA Institute for Quantitative and Computational Biosciences (QCB) for their support and Prof. David Still of the California Polytechnic Institute, Pomona campus for access to sample collection sites in an alfalfa field.

This work was supported by the National Science Foundation Grant No. NSF 1911781 and the Department of Energy BER Award DE-FC02-02ER63421 to the UCLA DOE Institute.

AUTHOR AFFILIATIONS

¹Department of Molecular, Cell, and Developmental Biology, University of California, Los Angeles, California, USA

²Department of Microbiology, Immunology, and Molecular Genetics, University of California, Los Angeles, California, USA

³Department of Integrative Biology and Physiology, University of California, Los Angeles, California, USA

⁴UCLA DOE Institute, University of California, Los Angeles, California, USA

AUTHOR ORCIDs

Ann M. Hirsch  <http://orcid.org/0000-0002-9633-1538>

Ethan Humm  <http://orcid.org/0000-0002-9727-6809>

FUNDING

Funder	Grant(s)	Author(s)
National Science Foundation (NSF)	1911781	Robert P. Gunsalus
National Science Foundation (NSF)	DE-FC02-02ER63421	Robert P. Gunsalus

DATA AVAILABILITY STATEMENT

The complete genome sequences of *V. paradoxus* 2u118 and *V. sp.* SPNA7 have been deposited in IMG/M under the taxon IDs [8045543215](https://img.jgi.doe.gov/cgi-bin/seqs.cgi?taxon=8045543215) and [8045536498](https://img.jgi.doe.gov/cgi-bin/seqs.cgi?taxon=8045536498), respectively. The assembled genomes are listed under the GenBank accession numbers [CP138515](https://www.ncbi.nlm.nih.gov/nuccore/CP138515) and

CP138516 for 2u118, and CP138513 and CP138514 for SPNA7. The raw sequencing reads have been deposited under the NCBI BioProject numbers PRJNA1026576 for 2u118 and PRJNA1026575 for SPNA7.

REFERENCES

- Dejonghe W, Berteloot E, Goris J, Boon N, Crul K, Maertens S, Höfte M, De Vos P, Verstraete W, Top EM. 2003. Synergistic degradation of linuron by a bacterial consortium and isolation of a single linuron-degrading *Variovorax* strain. *Appl Environ Microbiol* 69:1532–1541. <https://doi.org/10.1128/AEM.69.3.1532-1541.2003>
- Zhang HJ, Zhou QW, Zhou GC, Cao YM, Dai YJ, Ji WW, Shang GD, Yuan S. 2012. Biotransformation of the neonicotinoid insecticide thiacloprid by the bacterium *Variovorax boronicumulans* strain J1 and mediation of the major metabolic pathway by nitrile hydratase. *J Agric Food Chem* 60:153–159. <https://doi.org/10.1021/jf203232u>
- Leadbetter JR, Greenberg EP. 2000. Metabolism of acyl-homoserine lactone quorum-sensing signals by *Variovorax paradoxus*. *J Bacteriol* 182:6921–6926. <https://doi.org/10.1128/JB.182.24.6921-6926.2000>
- Liu ZH, Cao YM, Zhou QW, Guo K, Ge F, Hou JY, Hu SY, Yuan S, Dai YJ. 2013. Acrylamide biodegradation ability and plant growth-promoting properties of *Variovorax boronicumulans* CGMCC 4969. *Biodegradation* 24:855–864. <https://doi.org/10.1007/s10532-013-9633-6>
- Belimov AA, Safronova VI, Sergeyeva TA, Egorova TN, Matveyeva VA, Tsyganov VE, Borisov AY, Tikhonovich IA, Kluge C, Preisfeld A, Dietz KJ, Stepanok VV. 2001. Characterization of plant growth promoting rhizobacteria isolated from polluted soils and containing 1-aminocyclopropane-1-carboxylate deaminase. *Can J Microbiol* 47:642–652. <https://doi.org/10.1139/w01-062>
- Abdelkrim S, Jebara SH, Saadani O, Chiboub M, Abid G, Mannai K, Jebara M. 2019. Heavy metal accumulation in *Lathyrus sativus* growing in contaminated soils and identification of symbiotic resistant bacteria. *Arch Microbiol* 201:107–121. <https://doi.org/10.1007/s00203-018-1581-4>
- Finkel OM, Salas-González I, Castrillo G, Conway JM, Law TF, Teixeira PJPL, Wilson ED, Fitzpatrick CR, Jones CD, Dangl JL. 2020. A single bacterial genus maintains root growth in a complex microbiome. *Nature* 587:103–108. <https://doi.org/10.1038/s41586-020-2778-7>
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>
- Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>
- Chen IA, Chu K, Palaniappan K, Ratner A, Huang J, Huntemann M, Hajek P, Ritter SJ, Webb C, Wu D, Varghese NJ, Reddy TBK, Mukherjee S, Ovchinnikova G, Nolan M, Seshadri R, Roux S, Visel A, Woyke T, Eloe-Fadrosh EA, Kyrpides NC, Ivanova NN. 2023. The IMG/M data management and analysis system v.7: content updates and new features. *Nucleic Acids Res* 51:D723–D732. <https://doi.org/10.1093/nar/gkac976>
- Lee I, Chalita M, Ha SM, Na SI, Yoon SH, Chun J. 2017. ContEst16S: an algorithm that identifies contaminated prokaryotic genomes using 16S RNA gene sequences. *Int J Syst Evol Microbiol* 67:2053–2057. <https://doi.org/10.1099/ijsem.0.001872>
- Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67:1613–1617. <https://doi.org/10.1099/ijsem.0.001755>
- Yoon SH, Ha SM, Lim J, Kwon S, Chun J. 2017. A large-scale evaluation of algorithms to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* 110:1281–1286. <https://doi.org/10.1007/s10482-017-0844-4>