UC Irvine

UC Irvine Previously Published Works

Title

On the variability of the exponent in the power law depth dependence of POC flux estimated from sediment traps

Permalink

<https://escholarship.org/uc/item/6sj4874q>

Journal

Deep-Sea Research Part I: Oceanographic Research Papers, 53(8)

ISSN

0967-0637

Author

Primeau, F

Publication Date

2006-08-01

DOI

10.1016/j.dsr.2006.06.003

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

On the variability of the exponent in the power law depth dependence of POC flux estimated from sediment traps

François Primeau*

Department of Earth System Science, University of California at Irvine, Croul Hall, Irvine, CA 92697, USA

Received 30 August 2005; received in revised form 6 June 2006; accepted 9 June 2006

Abstract

Using the power law relationship $J(z) = J_0(z/z_0)^{-b}$ relating the particulate organic carbon (POC) flux in the water column at depth z to the export flux J_0 at $z_0 = 100$ m, I reanalyzed the sediment-trap data set compiled by Berelson [2001. The flux of particulate organic carbon into the ocean interior: a comparison of four U.S. JGOFS regional studies. *Oceanography* 14, 59–67]. The goal is to better understand the variability of the exponent b . I show that the usual approach of estimating parameters for each station separately and then pooling the estimates from all stations confounds within-station estimation errors with true regional variability and so tends to inflate the apparent variance of b . Furthermore, I show that the correlation between b and J_0 observed by Berelson [2001. The flux of particulate organic carbon into the ocean interior: a comparison of four U.S. JGOFS regional studies. *Oceanography* 14, 59–67] is spurious and attributable to the fact that the estimation errors for b and J_0 are correlated. A two level mixed-effects regression model is introduced to properly take into account the contribution to the variability of the POC fluxes of within- and between-station effects. The analysis shows that the variability of b across stations is only a small contributor to the POC flux variance in the sediment trap data set. In fact the additional POC flux variance captured by a model with regionally varying b in comparison to a simple model with a fixed b is not statistically significant. The best estimate for the fixed b applicable to all stations is 0.70 ± 0.08 (95% C.I. 90 d.f.).

© 2006 Elsevier Ltd. All rights reserved.

1. Introduction

The depth dependence of the downward flux of particulate organic carbon (POC) in the ocean, $J(z)$, is often parameterized by a power law

$$J(z) = J_0(z/z_0)^{-b}, \quad (1)$$

where z_0 is some reference depth, typically near 100 m, at which the export flux $J_0 \equiv J(z_0)$ can be explicitly modeled or measured. The functional

form in Eq. (1) was initially proposed by Martin et al. (1987) as a way of summarizing sediment-trap data, but has since been adopted as a useful parameterization of carbon export by sinking particles in many global biogeochemistry and ecosystem models (e.g. Sarmiento et al., 1993; Yamanaka and Tajika, 1996; Najjar and Orr, 1998). The parameter b controls the depth of particle remineralization. For large values of b there is a rapid decrease of the particle flux with depth, and most of the remineralization happens in the top of the water column just below the euphotic zone. For smaller values of b the particle flux decreases

*Tel.: +1 949 824 9435; fax: +1 949 824 3874.

E-mail address: fprimeau@uci.edu.

more slowly, and a larger fraction of the flux reaches the sea floor. In models, the efficiency of the biological pump and the resulting partition of carbon between the atmosphere and ocean is sensitive to b (Kwon and Primeau, 2006). Constraining b from data is therefore important.

In a summary of JGOFS sediment-trap data using Eq. (1), Berelson (2001) found variability in the estimates of b from different stations and suggested that this variability might point to regional differences in the biogeochemical controls of the attenuation of the POC fluxes with depth. Furthermore, he noted that a significant fraction of the variability of $\hat{b}$ (hats are used to indicate estimates of the “true” value) can be explained by variations in $\hat{J}_0$. Taken at face value, this result implies that regions of higher export have a higher fraction of the flux remineralizing in the upper water column. A similar observation was made by Lutz et al. (2002), who suggested that the flux of POC to depth was more efficient during periods of low primary production after noting a correlation between the export flux J_0 and the ratio, $J(z)/J_0$ (what they call the s_{ratio}). Berelson suggests several possible mechanisms that might explain the correlation: (1) varying rate constants for POC degradation associated with differences in molecular compositions; (2) varying kinetics due to different oxygen, temperature and mineral distributions in the water column; (3) differences in microbial and heterotrophic communities; and (4) variations in settling velocities, all of which could plausibly correlate with the spatially varying biological production in the top 100 m.

Here I show that estimating pairs of b and J_0 values separately for each station and then pooling the resulting estimate pairs as was done by Berelson (2001) and previously by Martin et al. (1987) tends to inflate the apparent station-to-station variability of b because it confounds errors in the estimates of b with true regional differences. I show also that such an analysis produces estimates of b and J_0 with correlated errors, making it difficult to interpret the apparent covariance between J_0 and b noted by Berelson (2001). One of the goals of this article is therefore to introduce an improved estimation method that can properly distinguish within-station errors from across-station variability in b and J_0 . I introduce a two-level mixed-effects regression model (Raudenbush and Bryk, 2002) to properly

take into account the combined contribution of within- and between-station effects to the variability of the POC flux measurements. Within-station effects include measurement errors and other effects that contribute to deviations from the power-law flux profile. Between-station effects include differences in J_0 and b from station to station. A statistical analysis based on the new model shows that with some caveats concerning the North Atlantic Bloom Experiment (NABE) station, the JGOFS sediment-trap data set compiled by Berelson (2001) is consistent with the hypothesis that there is no detectable covariance between b and J_0 . Moreover, the analysis shows that a model in which b varies from station to station does not account for a significantly larger fraction of the POC flux variance than a simpler model with a fixed b .

2. Expected variance and covariance between the estimates of the power-law exponent and POC export flux

Although Berelson (2001) fitted the power law directly to the flux measurements, it is useful first to transform the measured fluxes to a logarithmic scale so that the resulting model becomes linear. Following Martin et al. (1987) and Armstrong (2002), we take the logarithm of (1) to obtain

$$\log J(z) = \log J_0 - b \log z/z_0. \quad (2)$$

Applying Eq. (2) to the discrete set of sediment-trap data from one particular site yields a linear estimation problem of the form

$$X\beta = y + e, \quad (3)$$

where

$$X = \begin{bmatrix} 1 & -\log z_1/z_0 \\ 1 & -\log z_2/z_0 \\ \vdots & \vdots \\ 1 & -\log z_n/z_0 \end{bmatrix}, \quad \beta = \begin{bmatrix} \log J_0 \\ b \end{bmatrix},$$

$$y = \begin{bmatrix} \log J(z_1) \\ \log J(z_2) \\ \vdots \\ \log J(z_n) \end{bmatrix} \quad \text{and} \quad e = \begin{bmatrix} e_1 \\ e_2 \\ \vdots \\ e_n \end{bmatrix}, \quad (4)$$

where z_i are the sediment-trap depths, and e_i are the errors assumed to be independent and identically

distributed with variance σ^2 . The least squares solution of Eq. (3) is

$$\hat{\beta} = (X^T X)^{-1} X^T (y + e), \quad (5)$$

and the expected variance–covariance matrix of the parameter estimates is $\sigma^2 (X^T X)^{-1}$ (e.g. Faraway, 2005). Explicitly, the variance–covariance matrix for the parameter estimates is

$$\begin{bmatrix} \text{Var}[\log \hat{J}_0] & \text{Cov}[\hat{b} \log \hat{J}_0] \\ \text{Cov}[\hat{b} \log \hat{J}_0] & \text{Var}[\hat{b}] \end{bmatrix} = \frac{\sigma^2}{\Delta} \begin{bmatrix} \sum_{i=1}^n (\log z_i / z_0)^2 & \sum_{i=1}^n \log z_i / z_0 \\ \sum_{i=1}^n \log z_i / z_0 & n \end{bmatrix}, \quad (6)$$

where

$$\Delta = n \sum_{i=1}^n (\log z_i / z_0)^2 - \left(\sum_{i=1}^n \log z_i / z_0 \right)^2. \quad (7)$$

When parameter estimates obtained from different locations are combined, (e.g. Berelson, 2001; Martin et al., 1987) it is important to take into account the within-station sampling variability due to random errors in the sediment-trap flux measurements before attributing the variability of $\hat{b}$ or the covariance of $\hat{b}$ with $\hat{J}_0$ to any systematic differences in the processes happening in the water column at

different geographical locations. The variance–covariance one computes by combining estimates $\hat{b}_j$ and $\hat{J}_{0j}$ obtained from different station is the sum of the across-station variance–covariance plus the within-station variance–covariance of the estimates. Only the across-station variance–covariance represents true geographical differences in b and J_0 . The within-station variance–covariance, given by the average across all stations of the matrix expression in (6), represents a bias due to within-station estimation errors.

From the off-diagonal terms in (6) we see that the sign of the bias in the covariance depends only on the depths of sediment traps in relation to the reference depth z_0 . Because the fluxes are referenced to the export flux at 100 m near the top of the water column, the depths z_i are generally much greater than z_0 , and as a result, the expected covariance is positive for all sites. Importantly, the bias does not decrease as the number of stations in the ensemble increases. It decreases only as the number of flux measurements within each station increases. Because the number of flux measurements at any given site in the Berelson (2001) data set is small (n ranges from 3 to 12), the bias is potentially significant.

In Fig. 1, I show plots of the parameter estimates for the data set compiled by Berelson (2001) separated by region, but estimated separately for

Fig. 1. Plot of the parameter estimates $\hat{b}$ and $\hat{J}_0$ together with their 95% confidence intervals separated by region. The units of $\hat{J}_0$ are in $\text{mmol C m}^{-2} \text{ day}^{-1}$. Each point was estimated separately for each station by the ordinary least squares formula. The POC flux data set compiled by Berelson (2001) was used for the estimation. The data set includes sediment-trap fluxes from the Southern Ocean (SO), Equatorial Pacific (EqPac), North Atlantic Bloom Experiment (NABE), and Arabian Sea (AS).

each station by the least squares formula. The data set comprises sediment-trap and thorium-based POC flux estimates from the North Atlantic (NABE: 1 station), Equatorial Pacific (EqPac: 7 stations), Arabian Sea (AS: 5 stations) and Southern Ocean (SO, 4 stations). The data set contains stations with POC flux measurements at shallow, intermediate and deep levels and is uniquely well suited for estimating the exponent b . For references to the original publication of the data, I refer the reader to Berelson (2001). Fig. 1 also shows the 95% confidence ellipses. The tilting of the confidence ellipses relative to the parameter axes in Fig. 1 indicates that the uncertainties of $\hat{b}$ are correlated with those of $\log \hat{J}_0$. From Eq. (6), the slope of the tilt is given by

$$\frac{\text{Cov}[\hat{b} \log \hat{J}_0]}{\text{Var}[\log \hat{J}_0]} = \frac{\sum_{i=1}^n \log(z_i/z_0)}{\sum_{i=1}^n (\log z_i/z_0)^2}, \quad (8)$$

which is fairly uniform for all stations even though the trap deployment depths vary somewhat between stations. Because the uncertainties in the parameter estimates are positively correlated at each site, we expect the combined parameter estimates from all four regions (17 stations) to be positively correlated even if the true parameters are independent. Berelson (2001) computed a regression line for the scatter plot of his estimates of b against J_0 and found what appeared to be a statistically significant positive slope. However, Eq. (6) shows that the errors in the estimates are not independent and the usual significance tests are not applicable. The slope of a regression line through a scatter plot is given by (e.g. Janke and Tinsley, 2005)

$$m = \frac{\widehat{\text{Cov}}[\hat{b}_j \log \hat{J}_{0j}]}{\widehat{\text{Var}}[\log \hat{J}_{0j}]}, \quad (9)$$

where $\widehat{\text{Cov}}$ and $\widehat{\text{Var}}$ are used to denote the sample covariance and variance of the 17 pairs of parameter estimates. If the true b and J_0 are independent, the numerator in (9) will be given by the average over the 17 stations of the off-diagonal term in (6). As already noted, this term is positive for all stations, so the average will also be positive. That a positive slope in the scatter plot of b vs. J_0 is clearly visible in Berelson's (2001) analysis is expected from the correlation of the errors. This feature of the covariance of b and $\log J_0$ demonstrates the need for a more careful analysis to ascertain whether b is spatially varying and whether it co-varies with J_0 .

It is easy to understand the origin of the biased covariance between $\hat{b}$ and $\hat{J}_0$. A random error that leads to an over prediction in the export flux at the surface leads to a positive error in the b -estimate because the POC flux profile has to decrease more rapidly in order to be close to the deeper flux measurements. Similarly, a random error that leads to an under-prediction in export production at the surface leads to a negative error in the b -estimate because in this case the flux profile has to decrease more slowly if it is to be close to the deep flux measurements. Mathematically this effect is captured by the non-orthogonality of the columns in a regression design matrix X in Eq. (4). By choosing a deeper reference depth, ($z_0 \approx 670$ m) it is possible to make the columns of X nearly orthogonal, but the model is then conceptually less useful because it relates the sediment flux profile to a depth that is far from where the particles originate. More fundamentally, the approach of first estimating parameters separately for each station and then combining the resulting estimates to study their variability confounds within-station errors with the "true" regional variability of the parameters. Unless the within-station error variance is much smaller than the sought after variance-covariance of the "true" parameters, the usual estimation approach is inadequate for studying the spatial variability of POC flux profiles.

In the next section I introduce a two-level mixed-effects model (Raudenbush and Bryk, 2002) to properly take into account the contribution of within-station and between-station effects to the variability of the POC flux measurements. Using the two-level model, I reanalyze the sediment-trap data set used by Berelson (2001) to: (i) test if there is a statistically significant correlation between b and J_0 , and (ii) test how much of the often noted station-to-station variability in the estimates of b is due to true geographical differences in b .

3. Two-level mixed effects model

To properly treat the uncertainty associated with modeling POC fluxes measured from sediment traps using Eq. (1), we introduce a two-level mixed-effects regression model. A comprehensive introduction to this type of model can be found in Raudenbush and Bryk (2002). The two-level mixed-effects model I use takes into account uncertainty in the predicted fluxes due to the across-station variability of b and J_0 in addition to uncertainty due to the usual

within-station effects such as measurement errors. In the usual parameter estimation problem based on ordinary least squares (OLS) the coefficients b and J_0 are assumed fixed, whereas here we treat them as random variables. Mixed-effects models have become well established in several research fields and routines for estimating parameters are standard in many statistical software packages. Mixed-effects models are also known as random-effects models in the biometric literature, as random-coefficients regression models in the econometrics literature and as covariance components models in the statistical literature (Raudenbush and Bryk, 2002).

To keep the notation manageable I denote the i th POC flux measurement at the j th station by J_{ij} and define

$$\begin{aligned} Y_{ij} &\equiv \log J_{ij}, \\ Z_{ij} &\equiv \log z_{ij}/z_0. \end{aligned} \quad (10)$$

The first level of the model (level-1) describes the within-station flux variability

$$Y_{ij} = \beta_{0j} + \beta_{1j}Z_{ij} + r_{ij}, \quad (11)$$

where r_{ij} is the within-station error term and where β_{0j} denotes the logarithm of the 100 m reference POC flux for the j th station, and where β_{1j} denotes the b exponent for the j th station.

At the second level (level-2), the model describes the across-station variability of the β coefficients (i.e. the station-to-station variability of $\log J_0$ and b):

$$\begin{aligned} \beta_{0j} &= \gamma_{00} + u_{0j}, \\ \beta_{1j} &= \gamma_{10} + u_{1j}, \end{aligned} \quad (12)$$

where γ_{00} and γ_{10} are, respectively, the across-station mean $\log J_0$ and b values and where u_{0j} and u_{1j} are random effects that vary from station to

station with variance–covariance matrix

$$\text{Var} \begin{bmatrix} u_{0j} \\ u_{1j} \end{bmatrix} = \begin{bmatrix} \tau_{00} & \tau_{01} \\ \tau_{10} & \tau_{11} \end{bmatrix}. \quad (13)$$

τ_{00} and τ_{11} represent, respectively, the variances of $\log J_0$ and b across stations, i.e. the “true” station-to-station variance of the parameters as opposed to the variance of the *estimates*. Similarly $\tau_{01} = \tau_{10}$ represents the “true” covariance between $\log J_0$ and b .

Inserting Eqs. (12) for the level-2 effects into the level-1 model equation (11) gives

$$Y_{ij} = \gamma_{00} + \gamma_{10}Z_{ij} + u_{0j} + u_{1j}Z_{ij} + r_{ij}. \quad (14)$$

Note that estimating the parameters by ordinary least squares is not appropriate, because the error terms, $u_{0j} + u_{1j}Z_{ij} + r_{ij}$, do not have equal variance either within or across stations and moreover are not independent within stations. A proper estimation of γ_{10} (i.e. the mean b value) in Eq. (14) takes into account the different precisions with which b is determined at each station. It thus produces a better estimate than simply averaging the b values obtained from different stations (e.g. Berelson, 2001) or as in the case of Martin et al. (1987) by pooling all the measurements and estimating only a fixed J_0 and b .

I estimated the parameters in (14) using a maximum likelihood approach as implemented in the subroutine lme (Bates and Pinheiro, 1998) in the statistical software package R. The parameter estimates are listed in the row labeled *full* in Table 1. The numbers in parenthesis are the 95% confidence intervals for the estimated parameters. The estimate for the fixed b value is $\hat{\gamma}_{10} = 0.72 \pm 0.1$ (95% C.I.), and the estimated standard deviation of b across stations is $\text{sd}[u_{1j}] = 0.14$, with lower and upper 95% confidence limits of 0.06 and 0.31. In comparison,

Table 1
Parameters estimated from the Berelson (2001) POC flux data set for the two-level mixed-effects regression model

Model	$\hat{\gamma}_{00}$ fixed $\log J_0$	$\hat{\gamma}_{10}$ fixed $\hat{b}$	$[\hat{\tau}_{00}]^{1/2}$ $\text{sd}[\log J_0]$	$[\hat{\tau}_{11}]^{1/2}$ $\text{sd}[b]$	$\hat{\tau}_{10}/[\hat{\tau}_{00}\hat{\tau}_{11}]^{1/2}$ $\text{Corr}[b \log J_0]$	$\text{sd}[r_{ij}]$ $\text{sd}[\text{residuals}]$	Likelihood ratio test
<i>full</i>	1.4 (1.1,1.8)	0.72 (0.62,0.82)	0.67 (0.44,1.0)	0.14 (0.06,0.31)	0.67 (0.03,0.92)	0.53 (0.46,0.63)	<i>full</i> vs. <i>null</i>
<i>null</i>	1.4 (1.1,1.7)	0.70(0.62,0.78)	0.53 (0.36,0.78)	n.a.	n.a.	0.57 (0.50,0.66)	$p = 0.1547$
<i>full*</i>	1.3 (1.0,1.6)	0.68 (0.60,0.76)	0.51 (0.30,0.86)	0.07 (0.01,0.45)	0.13 (−0.87,0.92)	0.50 (0.43,0.59)	<i>full*</i> vs. <i>null*</i>
<i>null*</i>	1.3 (1.0,1.6)	0.68 (0.61,0.75)	0.51 (0.34,0.76)	n.a.	n.a.	0.51 (0.44,0.60)	$p = 0.7807$

The *full* and *null* model parameters were estimated from the full data set. The *full** and *null** model parameters were estimated from the data set in which the two shallow POC fluxes at the NABE station (Buesseler et al., 1992) were excluded. In the *null* and *null** models, b is fixed and is not allowed to vary across stations. J_0 is expressed in $\text{mmol C m}^{-2} \text{day}^{-1}$. The quantities in the parentheses are the lower and upper limits of a 95% confidence interval. The full data set contains 17 stations and a total of 108 flux measurements.

the standard deviation of the b values estimated in Section 2 by ordinary least squares is 0.22, approximately 60% higher. This is consistent with the fact that the variance of the estimates obtained by ordinary least squares is equal to the sum of the estimation error variance plus the across-station variance of the true b . The variance of the OLS estimates of b is 0.047 and the across-station average of the estimation error variance for b from the OLS formula is 0.031. If we try to correct the OLS variance by subtracting the average of the estimated within-station error variance we obtain an across-station standard deviation of $\text{sd}[b] = \sqrt{0.047 - 0.031} = 0.13$ which is closer to our estimate based on the mixed-effects model, but appears to be biased low. The plot of $\hat{b}_j \equiv \hat{\gamma}_{10} + \hat{u}_{1j}$ vs. $\hat{J}_{0j} \equiv \exp\{\hat{\gamma}_{00} + \hat{u}_{0j}\}$ estimated from the two-level mixed-effects model is shown in Fig. 2.

Since the standard deviation of b must be positive, the confidence interval for the across-station standard deviation of b excludes zero. To test the statistical significance of including the across-station deviations in b , we cannot simply check to see if its confidence interval excludes zero. Instead we construct a *null* model in which across-station deviations of b are set identically to zero (i.e. $u_{1j} \equiv 0$) and compare it to a *full* model by a likelihood ratio test (e.g. Faraway, 2005). This test yielded a p -value of 0.15, indicating that there is a 0.15 probability that the additional POC-flux

variance captured by the *full* model is due to chance alone. Furthermore, a closer look at the data shows that the already weak evidence for a variable b is strongly influenced by a single station. The point for the North Atlantic Bloom Experiment labeled NABE in Fig. 2 stands out as a clear outlier. At NABE, two shallow flux estimates, at 150 and 300 m, from Buesseler et al. (1992) based on the thorium method are a factor of 3.5 and 6.5 greater than those obtained by Martin et al. (1993) from floating traps at the same depths. To test the influence of these two flux measurements I repeated the estimation after excluding them. Without the two shallow thorium based fluxes at the NABE station I estimated ($\hat{b} = 0.68$, $\text{sd}[b] = 0.07$) and ($\hat{b} = 0.68$) for the full and null models (denoted *full** and *null** in Table 1). A likelihood ratio test, comparing the *null** and *full** models yielded a p -value of 0.7807. Without the Buesseler et al. (1992) shallow NABE fluxes based on the thorium method there is no evidence for rejecting the *null* model with a fixed b .

A plot of the b and J_0 values estimated from the data set excluding the two shallow thorium-based fluxes at NABE is shown in Fig. 3. Without the high shallow fluxes, the NABE station no longer stands out as an outlier. I hypothesize that including the Buesseler et al. (1992) thorium-derived flux data at NABE without having thorium-based estimates of the flux deeper in the water column biases the exponent towards a higher value because the power

Fig. 2. Plot of the exponent $\hat{b}$ and export flux $\hat{J}_0$ estimated from the Berelson (2001) data set and the two-level mixed-effects model. The units of $\hat{J}_0$ are $\text{mmol C m}^{-2} \text{ day}^{-1}$. The outlier labeled NABE corresponds to the North Atlantic Bloom Experiment. Note that $\hat{J}_0$ is plotted on a logarithmic scale.

Fig. 3. Same as Fig. 2 except that the two shallow NABE fluxes from Buesseler et al. (1992) measured by the thorium method were excluded from Berelson's (2001) data set for the estimation. Note that the axis scales are different than Fig. 2.

law fit has to decrease more rapidly to approach the lower floating trap numbers (assuming that thorium-based fluxes below 300 m would be higher than the floating trap based fluxes in accord with the higher thorium-based fluxes measured at 150 and 300 m). This hypothesis is consistent with the findings of Yu et al. (2001) and Scholten et al. (2001), who estimated the trapping efficiency of sediment traps using ^{230}Th calibration; both conclude that traps near 1000 m at the NABE site considerably underestimate the sediment flux. For example, Yu et al. (2001) report a trapping efficiency in the range of 38–43% for a trap at 1110 m and 70–88% for a trap at 3734 m.

On the grounds of simplicity, the best model appears to be the *null* model in which b is fixed. The best estimate for the fixed b is 0.70 ± 0.08 (95% C.I. 90 d.f.). The value is robust to the inclusion or exclusion of the level-2 random effects for b or to the exclusion of either the shallow NABE fluxes based on the thorium method or the NABE station data altogether; all cases produced fixed b estimates well within the quoted uncertainty (Table 1).

From the *full* model I also estimate $\tau_{01} \equiv \tau_{10} = 0.67$ with a 95% confidence interval ($0.03 < \tau_{01} < 0.92$). Recall that τ_{01} is the true across station covariance between $\log J_0$ and b . Since the confidence interval is positively bounded, there appears to be some evidence that stations with high 100-m export fluxes tend to have higher b exponents as observed by Berelson (2001). However, the correlation is entirely due to the influence of the two shallow fluxes based on the thorium method at the NABE station. This is evident from Figs. 2 and 3, where the apparent correlation disappears when the shallow thorium-based fluxes are excluded from the NABE station. Without the two shallow fluxes based on the thorium method at the NABE site (*null** in Table 1), the estimate for the covariance is $\tau_{01} = 0.13$, with a $(-0.86, 0.92)$ 95% confidence interval, confirming that the correlation is not robust. The tendency for faster remineralization with depth in regions of high export flux is not systematic across the entire data set and is sensitive to the possible inconsistency of using different flux measurement methods at different depths.

I have highlighted the possible inconsistency of mixing POC fluxes based on sediment traps and the thorium method at the NABE site. One should not infer from this that the NABE site was exceptional in Berelson's (2001) data set for having thorium-based shallow fluxes; in fact all stations in the data

set included thorium-based shallow POC fluxes. I have no evidence for discounting the thorium-based shallow fluxes at the NABE site in favor of the sediment-trap measurements. The point I wish to make is that the apparent correlation between $\log J_0$ and $\hat{b}$ is sensitive to inconsistencies in the shallow vs. deep flux measurements, and that the correlation noted by Berelson (2001) is not a systematic feature of the parameters once the errors in these parameters are properly represented.

4. Discussion

I have shown that the usual approach of fitting power law exponents to the vertical profile of POC flux measured from sediment traps overestimates the variability of the exponent because it confounds estimation errors with regional differences in the true b . I introduced a two-level mixed-effects regression model to properly model the uncertainty associated with predicting POC fluxes with a power-law profile. The mixed-effects model takes into account uncertainty associated with the across-station variability of b and J_0 as well as the usual within-station uncertainty associated with measurement errors. Contrary to previous studies I found that a model with a fixed exponent b applicable at all sites adequately fits the data. The additional variance in the POC flux captured by a model in which b varies across stations is not statistically significant. Our best estimate for the exponent is $b = 0.70 \pm 0.08$ (95% C.I. 90 d.f.).

There is of course little reason to expect that a universal b value would be applicable at all stations. In fact the existence of a universal b would be surprising given the complex transformations that affect particles as they rain down from the surface. Our result should not be interpreted as meaning that there is a universal b value. What it tells us is that the variability in the POC flux data set collected from sediment traps is dominated by effects other than the variability of b across stations. This is an important point to bear in mind if, as suggested by Berelson (2001), differences in estimated b values are to be used as guidance for seeking regional differences in the processes that might systematically affect the attenuation of POC fluxes with depth. To detect significant differences in the power law exponent a larger number of measurements per station is needed than was typical in the Berelson (2001) data set. Another possibility for decreasing the exponent's estimation variance would be to pool

stations into subgroups and test for statistically significant differences between the groups. Finally, working directly with the time-series data as opposed to the annually averaged fluxes might reveal differences in b as a function of time. For the purpose of the analysis I presented here, any such temporal variability was lumped into the within-station measurement error.

Our estimated b -value is lower than the average value ($b = 0.82 \pm 0.16$ 1 s.d.) quoted in Berelson (2001). The reason for this is that our estimation procedure based on the two-level mixed-effects model explicitly takes into account the differences in precision with which b is determined at each station. Bishop (1989) reviews several parameterizations of POC flux, and our estimate of b falls in the range of those that are cast in terms of the power law profile ($\bar{b} = 0.628$ – 1.0). Without proper uncertainty estimates for the b values estimated in previous studies it is difficult to judge if the differences in b values are significant. For example, our estimate is lower than the Martin et al. (1987) open-ocean composite (OOC), $b = 0.858$, but without a reliable uncertainty estimate for the OOC it is difficult to assess the significance of the difference. Based on the published results of Martin et al. (1987), I estimated the uncertainty of b based on two methods: one from the quoted r^2 value for the OOC and one from the standard deviation of the b values fitted for each station separately. Appendix A show how I backed out a formal standard error for the residual of the OOC using the quoted r^2 value in Martin et al. (1987). The result is $\sigma_{\text{OOC}} = 0.059$. Which yields a 95% confidence interval of ($0.7395 < b < 0.9765$). This overlaps with our 95% confidence interval ($0.62 < b < 0.78$). However, the formal standard error based on the OLS formula is likely to be an underestimate because it is based on the *incorrect* assumption that the OOC residuals are independent. As our two-level mixed-effects model makes explicit, the residuals for the OOC fit depend on a variable J_0 value that varies across stations but not within stations. Martin et al. (1987) analysis for the OOC assumes a fixed J_0 value for the pooled data. This suggests that the difference is probably not significant. At the other extreme, if I use the standard deviation of the b values for the individual stations that went into the OOC, I obtain a standard error of 0.12, which would suggest that the difference between the OOC and our estimate is insignificant. However, this uncertainty is an overestimate since it does not properly take into account

the larger number of degrees of freedom in the pooled data set. While inconclusive the above considerations make it clear that proper uncertainty estimates are needed in order to detect real regional differences in the power-law exponent. Multi-level mixed effects regression models are ideally suited for this task.

The 95% confidence interval for our estimate, ($0.62 < b < 0.78$), is relatively large suggesting that carbon cycle models have some latitude in tuning the b value in order to improve the model performance. Global models might be able to provide additional constraints on the power law exponent by optimizing the exponent so as to best reproduce global dissolved inorganic carbon, phosphate or alkalinity fields (e.g. Usbeck et al., 2003). Ultimately a better understanding of the processes that affect POC fluxes in the water column should produce a mechanistically based model to replace the empirical power law parameterization (e.g. the mineral-ballast model of Armstrong et al., 2002). I believe that mixed-effects regression models can also be usefully applied to test the statistical significance of parameters in the mineral-ballast model since it can easily be extended to take into account within- and across-station differences in mineral fluxes.

Acknowledgments

I thank Will Berelson for kindly providing the combined POC sediment trap data set. Comments from three anonymous reviewers and the editors of DSR Part I are also acknowledged. I also would like to acknowledge useful discussion of this work with Keith Moore. Support from the National Science Foundation Grant OCE 02215156 is also acknowledged.

Appendix A

To compare the statistical significance of the difference of the fixed b value that I estimated and Martin et al. (1987) OOC we need the standard error of both estimates. Unfortunately Martin et al. (1987) did not quote a standard error for their OOC. For the ordinary least squares approximation, the standard error for b is given by

$$\hat{\sigma}_b = \hat{\sigma} \sqrt{\frac{n}{A}}, \quad (15)$$

where $\hat{\sigma}$ is the standard deviation of the residuals and where A is defined in Eq. (7). The standard

deviation of the residual can be backed out from the quoted r^2 value of 0.81, and the approximate depths, z_i of the 48 sediment traps used in the OOC, from the following relationships

$$b = \frac{\widehat{\text{Var}}[\log J(z)]}{\widehat{\text{Var}}[\log(z/z_0)]} \quad (16)$$

and

$$r^2 \equiv 1 - \frac{\hat{\sigma}^2}{\widehat{\text{Var}}[\log J(z)]}, \quad (17)$$

from which we find

$$\hat{\sigma}^2 = (1 - r^2)b\widehat{\text{Var}}[\log(z/z_0)]. \quad (18)$$

I obtained approximate trap depths within ± 25 m from Fig. 5 in Martin et al. (1987). The resulting standard error for the OOC b value is $\hat{\sigma}_b \approx 0.059$. I tested that the estimate is not very sensitive to errors in our estimation of the OOC trap depths by adding noise with variance of 635 m^2 to the z_i values and repeating the calculation.

References

- Armstrong, R.A., Lee, C., Hedges, J.I., Honjo, S., Wakeham, S.G., 2002. A new, mechanistic model for organic carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals. *Deep-Sea Research II* 49, 219–236.
- Bates, D.M., Pinheiro, J.C., 1998. Computational methods for multilevel models. Available in PostScript or PDF formats at (<http://franz.stat.wisc.edu/pub/NLME/>).
- Berelson, W.M., 2001. The flux of particulate organic carbon into the ocean interior: a comparison of four U.S. JGOFS regional studies. *Oceanography* 14, 59–67.
- Bishop, J.K.B., 1989. Regional extremes in particulate matter composition and flux: effects on the chemistry of the ocean interior. In: Berger, W.H., Smetacek, V.S., Wefer, G. (Eds.), *Productivity of the Ocean: Present and Past*. Wiley, New York, pp. 117–137.
- Buesseler, K.O., Bacon, M.P., Cochran, J.K., Livingston, H.D., 1992. Carbon and nitrogen export during the JGOFS North Atlantic bloom experiment estimated from ^{234}Th : ^{238}U disequilibria. *Deep Sea Research* 39 (7/8), 1115–1137.
- Faraway, J.J., 2005. *Linear Models with R*. Chapman & Hall/CRC Texts in Statistical Science Series, vol. 63, 229pp.
- Janke, S.J., Tinsley, F.C., 2005. *Introduction to Linear Models and Statistical Inference*. Wiley, New York, 584pp.
- Kwon, E.Y., Primeau, F., 2006. Sensitivity and optimization study using an implicit biogeochemistry Ocean model and in situ phosphate, alkalinity, and DIC data. 2006 Ocean Sciences Meeting Conference Proceedings.
- Lutz, M., Dunbar, R., Caldeira, K., 2002. Regional variability in the vertical flux of particulate organic carbon in the ocean interior. *Global Biogeochemical Cycles* 16, doi: 10.1029/2000GB001383.
- Martin, J.W., Knauer, G.A., Karl, D.M., Broenkow, W.W., 1987. VERTEX: carbon cycling in the NE Pacific. *Deep-Sea Research* 34, 267–285.
- Najjar, R., Orr, J., 1998. Design of OCMIP-2 simulations of chlorofluorocarbons, the solubility pump and common biogeochemistry.
- Raudenbush, S.W., Bryk, A.S., 2002. *Hierarchical Linear Models Applications and Data Analysis Methods*, second ed. Sage Publications, London, 483pp.
- Sarmiento, J.L., Slater, R.D., Fasham, M.J.R., Ducklow, H.W., Toggweiler, J.R., 1993. A seasonal three-dimensional ecosystem model of nitrogen cycling in the North Atlantic photic zone. *Global Biogeochemical Cycles* 8, 417–450.
- Usbeck, R., Schlitzer, R., Fischer, G., Wefer, G., 2003. Particle fluxes in the ocean: comparison of sediment trap data with results from inverse modeling. *Journal of Marine Systems* 39, 167–183.
- Yamanaka, Y., Tajika, E., 1996. The role of the vertical fluxes of particulate organic matter and calcite in the ocean carbon cycle: studies using an ocean biogeochemical general circulation model. *Global Biogeochemical Cycles* 10, 361–382.