



# Microbes trading electricity in consortia of environmental and biotechnological significance

Amelia-Elena Rotaru<sup>1</sup>, Mon Oo Yee<sup>2</sup> and Florin Musat<sup>3</sup>

Favorable interspecies associations prevail in natural microbial assemblages. Some of these favorable associations are co-metabolic dependent partnerships in which extracellular electrons are exchanged between species. For such electron exchange to occur, the cells must exhibit electroactive interfaces and get involved in direct cell-to-cell contact (**D**irect **I**nterspecies **E**lectron **T**ransfer/**DIET**) or use available conductive mineral grains from their environment (**C**onductive-particle-mediated **I**nterspecies **E**lectron **T**ransfer/**CIET**). This review will highlight recent discoveries and knowledge gaps regarding DIET and CIET interspecies associations in artificial co-cultures and consortia from natural and man-made environments and emphasize approaches to validate DIET and CIET. Additionally, we acknowledge the initiation of a movement towards applying electric syntrophies in biotechnology, bioremediation and geoengineering for natural attenuation of toxic compounds. Next, we have highlighted the urgent research needs that must be met to develop such technologies.

## Addresses

<sup>1</sup> University of Southern Denmark, Campusvej 55, Odense, Denmark

<sup>2</sup> Lawrence Berkley National Laboratories, Cyclotron Rd. 1, Berkeley, CA, United States

<sup>3</sup> Helmholtz Center for Environmental Research, Permoserstraße 15, Leipzig, Germany

Corresponding author: Rotaru, Amelia-Elena ([arotaru@biology.sdu.dk](mailto:arotaru@biology.sdu.dk))

**Current Opinion in Biotechnology** 2021, **67**:119–129

This review comes from a themed issue on **Environmental biotechnology**

Edited by **Robbert Kleerebezem** and **Diana Z Sousa**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 2nd February 2021

<https://doi.org/10.1016/j.copbio.2021.01.014>

0958-1669/© 2021 Elsevier Ltd. All rights reserved.

## Introduction

We live on a microbial planet — a planet where microbes control the distribution of nearly all life's essential elements. Recent estimates place prokaryotes as the second most abundant organisms of all Earth's biomass (bacteria  $\approx 12.7\%$  and archaea  $\approx 1.3\%$  of  $\approx 550$  Gt bound-C). Although largely surpassed by plants in terrestrial

environments, prokaryotes dominate the subterrestrial (90%) and oceanic realms (70%) [1].

Prokaryotes do not live isolated and typically establish associations between species or with eukaryotes in the environment. Interactions between microbial species could be favorable, like mutualism, or unfavorable, like competition. Favorable interspecies associations based on cross-feeding prevail in natural microbial assemblages, as shown by a thorough survey of 800 microbial communities [2]. During favorable interspecies associations, prokaryotes synchronize their activity and growth via an array of information exchange strategies like quorum sensing, membrane vesicles, intercellular junctions, or intercellular membrane nanotubes [3]. Remarkably, the exchange of cellular material between species can even implicate the entire cytoplasm leading to hybrid cells that can reproduce – a possible unexplored driver of evolutionary diversification. The latter has been recently investigated in a *Clostridium ljungdahlii* and *Clostridium acetobutylicum* co-culture, which exchanged RNA and proteins [4].


Favorable interactions between prokaryotes involve the exchange of cell material (**Figure 1**), including:

- 1 exchange of small metabolites (e.g. H<sub>2</sub>, formate)
- 2 exchange of electrons (e.g. via shuttles, conductive materials, or native redox-active cell-surface molecules) and
- 3 exchange of other small molecules and cytoplasmic material (e.g., iron, vitamins, amino acids, antibiotic resistance proteins).

The first two are also known as syntrophy or metabolic cross-feeding - a cooperative interaction in which two species, (A) a syntroph/electron-donating species and (B) an electron-accepting partner, survive environmental conditions that would benefit neither species alone. In environments without soluble electron acceptors, syntrophs carry out energetically unfavorable reactions, like organic matter oxidation, by releasing reducing equivalents outside of the cell as electrons or small metabolites. These are scavenged by the accepting partner and used as electron donors for their metabolism (**Figure 1**). Without their partner, syntrophs experience catabolite repression. Without the syntroph, the electron-accepting partner experiences famine. Thus, only together, they prevail.

Syntrophic interactions via diffusible chemicals (H<sub>2</sub> or formate) or mediated by electron shuttle molecules

Figure 1



Examples of favorable interactions between prokaryotic species based on intercellular material exchange: (a) via diffusible molecules (e.g. H<sub>2</sub> and formate see Ref. [9]); (b) via an electron shuttle (e.g. via flavins see Ref. [73]); (c) by direct cell-to-cell contacts (e.g. pili [14\*] and outer membrane c-type cytochromes [15]); (d) via conductive particles (e.g. magnetite [28]). The first four (a–d) are typical interactions based on extracellular electron transfer. However, cells can also transfer larger cellular material by (e) membrane fusion (e.g. between two species of *Clostridium* [4]); (f) vesicles (e.g. interspecies iron delivery [74]) or nanotubes (e.g. interspecies amino acid transfer to compensate for amino acid auxotrophies [75]). ED – electron donor; EA – electron acceptor; ES – electron shuttle; ox – oxidized; red – reduced.

(e.g. cysteine, flavin, quinones) have been described elsewhere [5–7].

In this review, we will focus on ‘electric’ syntrophy, established either by relying on direct cell-to-cell electrical contacts (DIET – Direct Interspecies Electron Transfer) or mediated by electrically conductive materials (CIET – Conductive-particle-mediated Interspecies Electron Transfer) in artificial co-cultures and consortia from natural and man-made environments. A timeline of the discoveries in this research field is highlighted in Figure 2. We will highlight methods to validate DIET


and CIET in such environments. Finally, we will provide a list of open questions regarding the ecology of electric syntrophies and their role in future technology applications.

### Direct interspecies electron transfer in artificial co-cultures

#### DIET between *Geobacter metallireducens* and *Geobacter sulfurreducens*

DIET was first demonstrated in an artificial *Geobacter* co-culture provided with ethanol as the electron donor and fumarate as an electron acceptor [8]. Neither

Figure 2



Timeline of discoveries regarding direct interspecies electron transfer (DIET – above the arrow) and conductive mineral mediated interspecies electron transfer (CIET – below the arrow). GM; *Geobacter metallireducens*, GS; *Geobacter sulfurreducens*, GH; *Geobacter hydrogenophilus*, MB; *Methanosarcina barkeri*, MH; *Methanosaeta harundinacea*, PA; *Prosthecochloris aestuarii*, PA\*; *Pseudomonas aeruginosa*, PC; *Pelobacter carbinolicus*, RF; *Rhodoferrax ferrireducens*, SA; *Syntrophus aciditrophicus*, AOM; anaerobic oxidation of methane, OMC; outer membrane cytochrome, HIET; interspecies hydrogen transfer, NP; nanoparticles, IET; interspecies electron transfer, AQDS; anthraquinone-2,6-disulfonate, EET; extracellular electron transfer, ANME; anaerobic methanotrophic archaea. Additional references that have not been discussed in the manuscript text [76–82].

partner could use the ethanol-fumarate energy sources alone. When the ethanol-oxidizing *G. metallireducens* was placed together with the fumarate-reducing *G. sulfurreducens*, they formed a metabolically co-dependent consortium [8].

The partnership did not require enzymes for the metabolism of formate or H<sub>2</sub> [9]. Instead, it required a distinct apparatus for extracellular electron release in the donor strain (*G. metallireducens*) or extracellular electron uptake in the electron-accepting partner (*G. sulfurreducens*) (Figure 1). *G. metallireducens* required electrically conductive pili (*e*-pili) [10,11] and outer-membrane multiheme *c*-type cytochromes [10] to release electrons extracellularly. Conversely, to accept extracellular electrons, *G. sulfurreducens* did not require *e*-pili [11]. Still, it required an outer-membrane multiheme cytochrome (OMC) — OmcS [8] — an OMC, which could self-assemble into electrically conductive cytochrome-chains [12,13].

### DIET between *Geobacter metallireducens* and *Methanosarcinales*

In methanogenic environments, syntrophy is the key process in organic matter decomposition [6]. Thus, methanogens were expected to play the role of electron-accepting partners for syntrophs like *Geobacter metallireducens*. Indeed, DIET was possible between the alcohol-utilizing *G. metallireducens* and *Methanosarcinales* [14\*,15,16\*,17\*\*,18], including strict non-H<sub>2</sub> consumers [14\*,16\*,17\*\*,18]. Instead, *G. metallireducens* could not interact syntrophically with strict H<sub>2</sub> or formate-consuming methanogens [14\*,17\*\*]. It was conceivable that strict acetoclastic methanogens like *Methanosaeta harundinacea*, were only transferring acetate. However, expression analyses coupled with stoichiometry and <sup>14</sup>CO<sub>2</sub>-radiolabeling incubations showed that CH<sub>4</sub> was generated from CO<sub>2</sub> and not acetate alone. Additionally, incubations of *G. metallireducens* – *M. harundinacea* co-cultures with long-chain alcohols (e.g. butanol) that

cannot split into acetate led to DIET-based co-cultures, independent of acetate-transfer [18]. In these co-cultures, long-chain alcohols were oxidized to their respective long-chain fatty acids (e.g. butyrate). All these results confirmed that *Methanosaeta* was exchanging electrons directly with *G. metallireducens*. Nevertheless, the electron uptake mechanisms in *Methanosaeta* remains enigmatic.

During DIET with *Methanosarcinales*, *G. metallireducens* required conductive pili [14\*,15] and outer membrane multiheme *c*-type cytochromes (OMCs) [15]. The process of electron uptake usually involves OMCs in many autotrophs that accept extracellular electrons [19]. Therefore, *Methanosarcinales* were expected to retrieve electrons similarly as the only methanogens with *c*-type cytochromes [20]. However, not all *Methanosarcinales* capable of DIET flaunted multiheme *c*-type cytochromes (MHC) in their genomes [17\*\*]. Besides, one *Methanosarcina* (*M. mazei*), which contains a MHC (Mma\_0663), did not require it for growth with extracellular electrons from DIET partners or electrodes [17\*\*]. Therefore, it appears that *Methanosarcinales* may use unprecedented electron uptake mechanisms, which are profoundly unexplored.

#### Other DIET co-cultures

The diversity of DIET syntrophic interactions in co-cultures is expanding (Figure 2), beyond typical electroactive species. Typically, effective electrogens [22,23] play the role of the electron-donating strains to DIET-accepting partners but not HIET-partners (H<sub>2</sub>-based interspecies electron transfer). These electrogens belong to the genera *Geobacter* and *Rhodospirillum rubrum* [17\*\*,21\*,22], namely, *G. metallireducens*, *G. sulfurreducens*, *G. hydrophilus* and *R. ferrireducens*. Contrariwise, non-electrogenic (*G. bemidjensis*) or poor electrogenic *Geobacter* (*G. bremensis*, *G. uraniiireducens*, *G. humireducens*, *G. chapelei*) could not interact syntrophically with DIET-partners [22].

However, recent studies appear to challenge the hypothesis that effective electrogens are better DIET-ers [24\*,25], indicating that DIET relationships may occur between unexpected partners and under unusual conditions. For example, one interaction occurred only under light conditions, between the acetate-oxidizing *G. sulfurreducens* and the CO<sub>2</sub>-reducing phototrophic partner *Prosthecochloris aestuarii* [21\*].

Semenec *et al.* paired a formate-oxidizing *Pseudomonas aeruginosa* with the fumarate-reducing *G. sulfurreducens* as the electron-accepting partner. They showed that the interaction was dependent on multiheme cytochromes [25]. Although *P. aeruginosa* is capable of extracellular electron transfer (EET), it does so with the aid of self-secreted phenazine shuttles retained in a network of extracellular DNA [26]. Yet, *Pseudomonas*' phenazines were not required for its interaction with *G. sulfurreducens*.

Walker *et al.* showed that the typical H<sub>2</sub>-producing syntroph (*Syntrophus aciditrophicus*) — never characterized as an electrogen — harbored *e*-pili and switched to DIET when a DIET option was available [24\*]. This DIET interaction was demonstrated by placing *Syntrophus* with a partner incapable of H<sub>2</sub> and formate uptake — a *G. sulfurreducens*, which lacked a subunit for formate dehydrogenase and one for hydrogenase [24\*]. Besides, *Syntrophus* is not the only syntrophic bacterium encoding *e*-pili in its genome, hinting at a potential option for other syntrophs to do DIET [24\*].

Moreover, DIET does not always correlate with electroactivity in methanogens.

For example, a *Methanosarcina horonobensis* could not use a cathode as the electron donor but could form DIET consortia with *G. metallireducens* [16\*].

DIET was also indicated as a mode of interaction for a new *Methanobacterium* isolate (and strict formate utilizer) co-cultured with *G. metallireducens* [27]. Astoundingly, the *Methanobacterium-Geobacter* co-culture was independent of *e*-pili. It remains to be determined whether and how this *Methanobacterium* receives extracellular electrons.


Mostly, effective electroactive microorganisms interact by DIET. So, what makes 'non-electroactive species' capable of DIET? Furthermore, how come that archetypal syntrophs interact by DIET-syntrophy in the absence of a possibility for H<sub>2</sub>/formate-transfer? What ecological advantages might they have when switching from H<sub>2</sub>/formate-transfer to DIET and vice-versa? These questions remain open to future investigations.

#### Conductive materials accelerate DIET co-culture metabolism

The metabolism of DIET co-cultures is accelerated by electrically conductive particles (iron-oxide minerals [28] and carbon-based materials [15,16\*,17\*\*,29–31]). Such cell-particle-cell interactions are not strictly speaking DIET. When minerals mediate the interaction between species (CIET), cells are not in direct contact and genes typically involved in DIET are significantly downregulated. For example, a *Geobacter* co-culture amended with a semi-conductive iron-oxide (magnetite) downregulated the expression of OmcS, which was not required for the mineral-mediated interaction, but it required *e*-pili [28].

Moreover, conductive materials are sometimes essential for the syntrophy between partners otherwise incapable of DIET, as was the case for an acetate-nitrate fed *Geobacter sulfurreducens* – *Thiobacillus denitrificans* co-culture, which can only grow together in the presence of iron-oxide minerals [32] or redox-active humic substances [33].

Figure 3



Methods to validate DIET in a co-culture. **(a)** Establishing cross-feeding interactions to ensure substrate selectivity of each member, **(b)** Ensuring the incapability of alternative interspecies electron transfer (e.g. via hydrogen or formate) **(c)** Monitoring expression profiles of electron transfer conduit proteins, **(d)** Validating the absence of possible exogenous and endogenous electron shuttles in the culture media and **(e)** Deletion studies targeting genes involved in extracellular electron transfer (e.g. pili, outer membrane cytochromes).

### Approaches to validate DIET in co-cultures

Precise validation of DIET in artificial co-cultures requires a polyphasic approach (Figure 3). This approach includes: i) determining the potential to form a cross-feeding interaction (with DIET and non-DIET partners) along with ii) the syntrophic consortia's physiology, iii) genomics to document the potential absence of alternative electron transfer strategies, iv) gene expression and v) targeted gene-deletion studies. For example, the incapacity to exchange electrons via H<sub>2</sub>/formate was tested with the help of a donor strain incapable of H<sub>2</sub>/formate transfer [14<sup>\*</sup>,17<sup>\*\*</sup>]. In instances where the donor strain can oxidize their substrate to H<sub>2</sub>/formate [17<sup>\*\*</sup>,22,24<sup>\*</sup>], researchers tested first if the donor strain was unsuccessful at establishing co-dependent interactions with H<sub>2</sub>/formate-utilizing partners [14<sup>\*</sup>,17<sup>\*\*</sup>] and second if it was successful with acceptor strains unable of H<sub>2</sub>/formate uptake (naturally or artificially by gene deletion) [9,16<sup>\*</sup>,24<sup>\*</sup>]. Additional tests are needed to exclude other electron transfer possibilities between species via self-generated shuttles or other redox-active compounds (e.g. flavins or cysteine, respectively). For example, cysteine could be transiently excluded from the media [15], or co-cultures could be spiked with spent cell filtrate, which may stimulate metabolism if rich in shuttles [34<sup>\*\*</sup>].

### Evidence for DIET-syntrophy in environmental dual-species consortia

Recent investigations indicate that DIET is a relevant electron transfer process in microbial consortia catalyzing the anaerobic oxidation of methane (AOM) and higher gaseous alkanes, both coupled with sulfate-reduction. Sulfate-dependent AOM is a process with broad climate impact, controlling methane emissions to the atmosphere. AOM-mediating consortia are abundant in various methane-rich habitats [35], while archaea oxidizing higher alkanes appear widespread in hydrocarbon-impacted sediments [36,37].

### DIET in anaerobic methane-oxidizing consortia

Sulfate-dependent AOM consortia consist of anaerobic methanotrophic (ANME) Archaea tightly packed with partner sulfate-reducing bacteria (SRB). Reducing equivalents from methane oxidation are transferred from ANME-Archaea to the partner SRB, which reduces sulfate to sulfide [35]. Two studies indicated that ANME-SRB interaction is based on DIET [38<sup>\*</sup>,39<sup>\*</sup>]. As determined by stable isotope assimilation [38<sup>\*</sup>] and confirmed by modeling [40], the distribution of metabolically active cells within natural ANME-2-SRB aggregates from cold seeps could only be explained by an interspecies



association dependent on electrically conductive conduits between cells, similar to DIET.

Additionally, ANME-2 genomes contain large multiheme cytochromes (MHC) similar to those in electrogens like *Geobacter* [38<sup>\*</sup>]. Researchers identified probable electroactive interfaces in cellular membranes and the interstitial space between cells via heme staining [38<sup>\*</sup>]. Besides, they identified MHC genes in the genomes of both partners of thermophilic AOM consortia, ANME-1 and HotSeep-1 SRB, enriched from hot seeps [39<sup>\*</sup>]. Moreover, HotSeep-1 encoded type IV pili proteins. MHC and pili genes were specifically overexpressed under methane-oxidizing conditions, and nanowire-like structures were observed in consortia's intercellular space, indicating DIET coupling [39<sup>\*</sup>].

#### DIET in anaerobic butane-oxidizing and ethane-oxidizing consortia

Recently, DIET interactions have been proposed for thermophilic archaea candidate lineages oxidizing butane (*Ca. Syntrophoarchaeum*) or ethane (*Ca. Ethanoperedens*) in consortia with SRB of the HotSeep-1 clade [36,41<sup>\*\*</sup>]. The SRB partners of both *Ca. Syntrophoarchaeum* [41<sup>\*\*</sup>], and *Ca. Ethanoperedens* [36] encode and express MHC and type IV pili, and nanowire-like structures have been observed connecting cells within consortia [41<sup>\*\*</sup>]. A representative of the HotSeep-1 clade (*Ca. Desulfosarcina auxilii*) was enriched without its archaeal partner and shown to be a chemolithoautotrophic H<sub>2</sub>-oxidizer [42]. Together with the detection of H<sub>2</sub> in thermophilic AOM cultures [39<sup>\*</sup>], this raised the prospect of a hydrogen-based coupling of alkane oxidation to sulfate reduction. However, after specific inhibition of the SRB partner, the H<sub>2</sub> concentrations in AOM and butane-oxidizing consortia were far too low to explain the measured sulfate reduction rates, leaving DIET as the only reasonable electron transfer mechanism [39<sup>\*</sup>,41<sup>\*\*</sup>]. Nevertheless, direct proofs for pili and MHCs being undoubtedly linked to DIET in such consortia is yet to be determined.

#### Proposed alternative mechanism via zero-valent sulfur (S<sup>0</sup>)

Chemical imaging of AOM consortia of ANME-2 and *Desulfosarcina*-SRB showed a high abundance of S<sup>0</sup> in the archaeal cells [43]. S<sup>0</sup>-abundance was corroborated with physiology experiments and immunolabelling of canonical enzymes and interpreted as interspecies electron transfer mediated by S<sup>0</sup>-based (polysulfides) compounds. This model had one major drawback, the reliance on a hypothetical, cryptic sulfate-reduction pathway producing S<sup>0</sup>/polysulfides in archaea whose enzymes were never identified. Recently, archaea with high S<sup>0</sup> content have been identified in an ethane-oxidizing culture [44]. Like ANME archaea, the ethane-oxidizing archaea (*Ca. Argoarchaeum ethanivorans*) also depend on partner SRB, but they do not form aggregates and do not exhibit nanowire-like structures.

These recent findings revived the idea that in some consortia, alkane-oxidation may be coupled to sulfate reduction via S<sup>0</sup>-mediated IET and not DIET [44].

#### Evidence for CIET-syntrophy in environmental communities

In the environment, syntrophic partners may interact via conductive minerals [45]. Conductive minerals are often present in natural environments (e.g. in coastal sediments, rice paddies, hydrothermal vents) [46–48], and their absence during laboratory incubations severely impacts species distribution and survival [34<sup>\*\*</sup>,49,50].

Interactions dependent on conductive particles (CIET) are more straightforward to investigate than DIET. This is because we can use conductive minerals to specifically enrich CIET-partners from environmental communities where partners may rely on conductive minerals to interact with each other. Under such enrichment conditions, non-syntrophic species fade out. For example, a *Geobacter* - *Methanosarcina* consortium from Baltic Sea sediments required the presence of conductive materials (iron-oxides or activated carbon) to carry out syntrophic acetate oxidation [34<sup>\*\*</sup>]. Without conductive minerals, syntrophic acetate oxidation ceased, and both groups went extinct. Without conductive minerals, a less abundant and metabolically ineffective species took over acetate turnover via acetoclastic methanogenesis [34<sup>\*\*</sup>]. Stable isotope analyses of fresh sediments showed that acetate was processed via syntrophic acetate oxidation coupled with CO<sub>2</sub> reductive methanogenesis [34<sup>\*\*</sup>,51], likely relying on the conductive iron-minerals abundant in marine sediments [34<sup>\*\*</sup>,49,50].

#### Approaches to validate DIET and CIET in natural guilds

The application of the polyphasic approach mentioned above (Figure 3) to confirm DIET and CIET in environmental communities is not always possible because some syntrophic partners cannot be separated or genetically manipulated, especially those in obligate syntrophic interactions like the ANME-SRB consortia.

Secondly, significant concerns have been raised at describing 'electric'-syntrophy based on metabolism stimulation by conductive materials [52<sup>\*\*</sup>], because conductive materials enhance the metabolism of some methanogens (e.g. carbon nanotubes [53]) independent of being coupled with an 'electric'-syntroph.

Besides, investigations of environmental DIET/CIET associations based on the mere presence of the DNA/RNA of 'electric'-syntrophs (see references in [17<sup>\*\*</sup>]) are not ideal since species abundance (e.g. *Geobacter*) or expression of a certain protein does not necessarily mean they perform DIET/CIET in the environment.

With many *Geobacter*-species incapable of establishing syntrophic associations [22] and many uncharacterized ‘electric’-syntrophs out there [49,50], novel investigation strategies are needed. A combination of tools must be employed after case-by-case adjustment to the process and the environment considered. For example, to demonstrate CIET-dependent interactions, we must verify the strict dependence on conductive minerals over non-conductive materials [34<sup>••</sup>,49]. For both DIET and CIET, expression studies can inform on MHC and pili content [54]. However, these alone cannot tell whether two species are coupled metabolically. For this we need to monitor metabolites to inform on consortia’s stoichiometry, and determine species co-occurrence. Plus, specific inhibition of the donating and accepting partners could tell whether the oxidation and reduction processes are co-dependent [34<sup>••</sup>,49,50]. The role of H<sub>2</sub>/formate or shuttles/enzymes as interspecies intermediates can be excluded by following their impact on the consortia’s metabolism [55]. For example, H<sub>2</sub>-additions to a consortium relying on H<sub>2</sub>-transfer would block the syntroph’s metabolism by feedback inhibition [56]. Or suppose the interaction depends on shuttles/enzymes generated by the consortia. In that case, the shuttles/enzymes in the spent media would facilitate the extracellular electron exchange. Such experiments testing the spent media, are typical when investigating electron uptake during Fe<sup>0</sup>-biocorrosion [57,58]. Additionally, we can apply electrochemical methods like cyclic voltammetry to determine the presence of active redox molecules or enzymes in the live/heat-killed spent culture media [59].

DIET and CIET interactions may have specific isotopic signatures, specific microscopic distribution patterns, molecular and elemental signatures (e.g. high metal-content on cell surfaces) compared to H<sub>2</sub>/formate IET. None are understood or explored sufficiently. Therefore, innovative methods to simplify the verification of these processes in the environment require immediate attention.

### Ecological and biotech ramifications of electric syntrophies

It is apparent that cooperative metabolic dependencies greatly influence environmental chemistry and, consequently, impact our health, climate, and industries (Figure 4). Because we lack tools to study DIET and CIET in the environment, our understanding of how interspecies interactions impact environmental processes is in its infancy. Nevertheless, we mentioned studies that showed how both methane production and methane consumption in marine environments appear to be controlled by DIET/CIET interactions, possibly influencing the release of this greenhouse gas in the atmosphere. Therefore, it is imperative to understand better the triggers and controls for these processes of climate relevance.

Biotechnologies dependent on DIET and CIET are budding, with DIET-syntrophs and conductive materials often applied to stimulate industrial processes like anaerobic digestion. Several recent reviews summarized the implications of electric syntrophies in anaerobic digestion and demanded the development of suitable detection methods (extensively reviewed in Refs. [52<sup>••</sup>,60]).

Another role for DIET and CIET is the bioremediation of toxic compounds from industry off streams, or already released in the environment. Recent studies investigated the possibility to apply CIET to improve the degradation of toxic compounds from the effluents of various industrial processes like: nitrobenzene – found in herbicides, insecticides and pharmaceuticals [61], azo dyes from the textile industry [62], solvents from the printing industry [63], chlorinated compounds (e.g. Refs. [64,65]) generally used as precursors for PVC-production, and petroleum hydrocarbons [66–69].


Geoengineering approaches using CIET to stimulate the attenuation and degradation of contaminants and decontaminate sediments are now under consideration [70]. Two recent studies showed that the addition of activated carbon stimulated polycyclic aromatic hydrocarbon degradation under anaerobic conditions when CIET was possible, but not under aerobic conditions [71,72]. However, the actual implications of DIET and CIET in environmental decontamination remains to be verified. It is advisable to proceed stepwise because adding conductive minerals to contaminated soils could significantly enhance CIET and methane production, possibly enhancing methane emissions to the atmosphere. Thus, it is paramount that primary tests are carried out to investigate the effect of such materials on communities through the sediment depth and verify the effect on microorganisms along the entire spectrum of electron acceptors.

### Conclusion

DIET and CIET have been intensely studied in laboratory co-cultures, natural dual-species consortia and enriched environmental consortia. However, methods to easily fingerprint DIET/CIET associations in the environment are lacking. Here we indicate a polyphasic approach to study such associations in environmental samples and call for additional tools to be developed.

The significance of electric associations along other types of interspecies associations in natural processes is ambiguous. Thus, it is imperative to understand the role of ‘electric’ syntrophies in global element cycles, especially in the interplay between the iron and methane cycles. Climate change has led to increased erosion and input of rock and mineral particles in our oceans, possibly enhancing CIET interactions and the release

Figure 4



Ecological relevance and potential applications of DIET and CIET interactions.

and perhaps consumption of the potent greenhouse gas -methane. Overall, this significantly influences our present-day climate models since we do not comprehend potential novel methane sources and sinks in natural environments. The wastewater and anaerobic digestion industries are now investing resources to determine DIET and CIET implications in speeding up organic matter decomposition. Additionally, geoen-gineering approaches are being sought considering conductive mineral particle additions to contaminated environments to induce bio-attenuation of pollutants. It

is time we, as a scientific community, come together to cover these knowledge gaps.

**Conflict of interest statement**

Nothing declared.

**CRedit authorship contribution statement**

**Amelia-Elena Rotaru:** Conceptualization, Writing - original draft, Writing - review & editing. **Mon Oo Yee:**



Visualization, Writing - review & editing. **Florin Musat:**  
Writing - original draft, Writing - review & editing.

## Acknowledgement

This is a contribution to a Sapere Aude Danish Research Council grant to AER with grant number 4181-00203.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Bar-On YM, Phillips R, Milo R: **The biomass distribution on Earth.** *Proc Natl Acad Sci U S A* 2018, **115**:6506-6511.
2. Zelezniak A, Andrejev S, Ponomarova O, Mende DR, Bork P, Patil KR: **Metabolic dependencies drive species co-occurrence in diverse microbial communities.** *Proc Natl Acad Sci U S A* 2015, **112**:6449-6454.
3. Sime-Ngando T, Bertrand J-C, Bogusz D, Brugère J-F, Franche C, Fardeau M-L, Froussart E, Geiger A, Goñi-Urriza MS, Ollivier B: **The evolution of living beings started with prokaryotes and in interaction with prokaryotes.** *Prokaryotes and Evolution.* Springer; 2018:241-338.
4. Charubin K, Modla S, Caplan JL, Papoutsakis ET: **Interspecies microbial fusion and large-scale exchange of cytoplasmic proteins and RNA in a syntrophic *Clostridium* coculture.** *mBio* 2020, **11**:1-20.
5. Morris BEL, Henneberger R, Huber H, Moissl-Eichinger C: **Microbial syntrophy: interaction for the common good.** *FEMS Microbiol Rev* 2013, **37**:384-406.
6. Shrestha PM, Rotaru A-E: **Plugging in or going wireless: strategies for interspecies electron transfer.** *Front Microbiol* 2014, **5**:237.
7. Schink B, Montag D, Keller A, Müller N: **Hydrogen or formate - alternative key players in methanogenic degradation.** *Environ Microbiol Rep* 2017, **00**.
8. Summers ZM, Fogarty HE, Leang C, Franks AE, Malvankar NS, Lovley DR: **Direct exchange of electrons within aggregates of an evolved syntrophic coculture of anaerobic bacteria.** *Science (80-)* 2010, **330**:1413-1415.
9. Rotaru A-E, Shrestha PM, Liu F, Ueki T, Nevin K, Summers ZM, Lovley DR: **Interspecies electron transfer via hydrogen and formate rather than direct electrical connections in cocultures of *Pelobacter carbinolicus* and *Geobacter sulfurreducens*.** *Appl Environ Microbiol* 2012, **78**:7645-7651.
10. Shrestha PM, Rotaru A-E, Summers ZM, Shrestha M, Liu F, Lovley DR: **Transcriptomic and genetic analysis of direct interspecies electron transfer.** *Appl Environ Microbiol* 2013, **79**:2397-2404.
11. Ueki T, Nevin KP, Rotaru A-E, Wang LY, Ward JE, Woodard TL, Lovley DR: **Geobacter strains expressing poorly conductive pili reveal constraints on direct interspecies electron transfer mechanisms.** *mBio* 2018, **9**:1-10.
12. Filman DJ, Marino SF, Ward JE, Yang L, Mester Z, Bullitt E, Lovley DR, Strauss M: **Cryo-EM reveals the structural basis of long-range electron transport in a cytochrome-based bacterial nanowire.** *Commun Biol* 2019, **2**:219.
13. Wang F, Gu Y, O'Brien JP, Yi SM, Yalcin SE, Srikanth V, Shen C, Vu D, Ing NL, Hochbaum AI *et al.*: **Structure of microbial nanowires reveals stacked hemes that transport electrons over micrometers.** *Cell* 2019, **177**:361-369.e10.
14. Rotaru A-E, Shrestha PM, Liu F, Shrestha M, Shrestha D, Embree M, Zengler K, Wardman C, Nevin KP, Lovley DR: **A new model for electron flow during anaerobic digestion: direct interspecies electron transfer to *Methanosaeta* for the reduction of carbon dioxide to methane.** *Energy Environ Sci* 2014, **7**:408-415.
15. Rotaru A-E, Shrestha PM, Liu F, Markovaite B, Chen S, Nevin KP, Lovley DR: **Direct interspecies electron transfer between *Geobacter metallireducens* and *Methanosaeta barkeri*.** *Appl Environ Microbiol* 2014, **80**:4599-4605.
16. Yee MO, Snoeyenbos-West OL, Thamdrup B, Ottosen LDM, Rotaru A-E: **Extracellular electron uptake by two *Methanosarcina* species.** *Front Energy Res* 2019, **7**:1-10.
17. Yee MO, Rotaru A-E: **Extracellular electron uptake in *Methanosarcinales* is independent of multiheme c-type cytochromes.** *Sci Rep* 2020, **10**:372.
18. Wang L, Nevin KP, Woodard TL, Mu B, Lovley DR: **Expanding the diet for DIET: electron donors supporting Direct Interspecies Electron Transfer (DIET) in defined co-cultures.** *Front Microbiol* 2016, **7**:1-7.
19. Gupta D, Guzman MS, Bose A: **Extracellular electron uptake by autotrophic microbes: physiological, ecological, and evolutionary implications.** *J Ind Microbiol Biotechnol* 2020, **47**:863-876.
20. Thauer RK, Kaster A-K, Seedorf H, Buckel W, Hedderich R: **Thaerogenic archaea: ecologically relevant differences in energy conservation.** *Nat Rev Microbiol* 2008, **6**:579-591.
21. Ha PT, Lindemann SR, Shi L, Dohnalkova AC, Fredrickson JK, Madigan MT, Beyenal H: **Syntrophic anaerobic photosynthesis via direct interspecies electron transfer.** *Nat Commun* 2017, **8**:1-7.
22. Rotaru A-E, Woodard TL, Nevin KP, Lovley DR: **Link between capacity for current production and syntrophic growth in *Geobacter* species.** *Front Microbiol* 2015, **6**:744.
23. Chaudhuri SK, Lovley DR: **Electricity generation by direct oxidation of glucose in mediatorless microbial fuel cells.** *Nat Biotechnol* 2003, **21**:1229-1232.
24. Walker DJF, Nevin KP, Holmes DE, Rotaru A-E, Ward JE, Woodard TL, Zhu J, Ueki T, Nonnenmann SS, McInerney MJ *et al.*: **Syntrophic conductive pili demonstrate that common hydrogen-donating syntrophs can have a direct electron transfer option.** *ISME J* 2020, **14**:837-846.
25. Semenech L, Laloo AE, Schulz BL, Vergara IA, Bond PL, Franks AE: **Deciphering the electric code of *Geobacter sulfurreducens* in cocultures with *Pseudomonas aeruginosa* via SWATH-MS proteomics.** *Bioelectrochemistry* 2018, **119**:150-160.
26. Saunders SH, Edmund CM, Yates MD, Otero FJ, Trammell SA, Stemp EDA, Barton JK, Tender LM, Newman DK: **Extracellular DNA promotes efficient extracellular electron transfer by pyocyanin in *Pseudomonas aeruginosa* biofilms.** *Cell* 2020, **182**:919-932.
27. Zheng S, Liu F, Wang B, Zhang Y, Lovley DR: **Methanobacterium capable of direct interspecies electron transfer.** *Environ Sci Technol* 2020, **54**:15347-15354.
28. Liu F, Rotaru A-E, Shrestha PM, Malvankar NS, Nevin KP, Lovley DR: **Magnetite compensates for the lack of a pilin-associated c-type cytochrome in extracellular electron exchange.** *Environ Microbiol* 2015, **17**:648-655.

29. Liu F, Rotaru A-E, Shrestha PM, Malvankar NS, Nevin KP, Lovley DR: **Promoting direct interspecies electron transfer with activated carbon.** *Energy Environ Sci* 2012, **5**:8982.
30. Chen S, Rotaru A-E, Liu F, Phillips J, Woodard TL, Nevin KP, Lovley DR: **Carbon cloth stimulates direct interspecies electron transfer in syntrophic co-cultures.** *Bioresour Technol* 2014, **173**:82-86.
31. Chen S, Rotaru A-E, Shrestha PM, Malvankar NS, Liu F, Fan W, Nevin KP, Lovley DR: **Promoting interspecies electron transfer with biochar.** *Sci Rep* 2014, **4** 5019.
32. Kato S, Hashimoto K, Watanabe K: **Microbial interspecies electron transfer via electric currents through conductive minerals.** *Proc Natl Acad Sci U S A* 2012, **109**:10042-10046.
33. Zheng Y, Kappler A, Xiao Y, Yang F, Mahadeva GD, Zhao F: **Redox-active humics support interspecies syntrophy and shift microbial community.** *Sci China Technol Sci* 2019, **62**:1695-1702.
34. Rotaru A-E, Calabrese F, Stryhanyuk H, Musat F, Shrestha PM, Weber HS, Snoeyenbos-West OLO, Hall POJ, Richnow HH, Musat N *et al.*: **Conductive particles enable syntrophic acetate oxidation between *Geobacter* and *Methanosarcina* from coastal sediments.** *mBio* 2018, **9**:1-14
- This study is the first polyphasic substantiation of CIET interactions in environmental consortia, where the consortium's metabolism was strictly dependent on mineral additions. A coherent approach on how to identify CIET partners and processes in environmental samples has also been highlighted.
35. Knittel K, Boetius A: **Anaerobic oxidation of methane: progress with an unknown process.** *Annu Rev Microbiol* 2009, **63**:311-334.
36. Hahn CJ, Laso-Pérez R, Vulcano F, Vaziourakis KM, Stokke R, Steen IH, Teske A, Boetius A, Liebeke M, Amann R *et al.*: **"*Candidatus Ethanoperedens*," a thermophilic genus of archaea mediating the anaerobic oxidation of ethane.** *mBio* 2020, **11**.
37. Wang Y, Feng X, Natarajan VP, Xiao X, Wang F: **Diverse anaerobic methane-and multi-carbon alkane-metabolizing archaea coexist and show activity in Guaymas Basin hydrothermal sediment.** *Environ Microbiol* 2019, **21**:1344-1355.
38. McGlynn SE, Chadwick GL, Kempes CP, Orphan VJ: **Single cell activity reveals direct electron transfer in methanotrophic consortia.** *Nature* 2015, **526**:531-535
- By stable isotope probing coupled with single-cell chemical imaging by nanoSIMS, the authors explained the distribution of biosynthetic activity of cells in AOM-dual-species consortia to be most likely dependant on a direct exchange of electrons. DIET involvement was supported by detecting multi-haem cytochromes in the genomes of ANME Archaea and by redox-dependent staining of the aggregate's intercellular space.
39. Wegener G, Krukenberg V, Riedel D, Tegetmeyer HE, Boetius A: **Intercellular wiring enables electron transfer between methanotrophic archaea and bacteria.** *Nature* 2015, **526**:587-590
- This study showed that AOM-consortia express multiheme cytochromes and pili (which were visualized in the interstitial space of the consortia) only under anaerobic methane oxidizing conditions. Additional physiology experiments excluded the role of H<sub>2</sub> as a mediator of electron transfer. This work consolidated the current view that AOM is governed by DIET.
40. He X, Chadwick G, Kempes C, Shi Y, McGlynn S, Orphan V, Meile C: **Microbial interactions in the anaerobic oxidation of methane: model simulations constrained by process rates and activity patterns.** *Environ Microbiol* 2019, **21**:631-647.
41. Laso-Pérez R, Wegener G, Knittel K, Widdel F, Harding KJ, Krukenberg V, Meier DV, Richter M, Tegetmeyer HE, Riedel D *et al.*: **Thermophilic archaea activate butane via alkyl-coenzyme M formation.** *Nature* 2016, **539**:396-401
- This study reports the discovery of Archaeal-Bacterial consortia oxidizing butane using biochemical principles known hitherto only from AOM. The Archaea oxidizing butane, *Candidatus Syntrophoarchaeum*, transferred reducing equivalents to partner SRB. These organisms express multiheme cytochromes and pili during the interaction, and nanowire-like structures have been observed in the intercellular space of aggregates.
- Based on these observations and additional physiology experiments, a DIET-based interaction was proposed.
42. Krukenberg V, Harding K, Richter M, Glöckner FO, Gruber-Vodicka HR, Adam B, Berg JS, Knittel K, Tegetmeyer HE, Boetius A: ***Candidatus Desulfofervidus auxilii*, a hydrogenotrophic sulfate-reducing bacterium involved in the thermophilic anaerobic oxidation of methane.** *Environ Microbiol* 2016, **18**:3073-3091.
43. Milucka J, Ferdelman TG, Polerecky L, Franzke D, Wegener G, Schmid M, Lieberwirth I, Wagner M, Widdel F, Kuypers MMM: **Zero-valent sulphur is a key intermediate in marine methane oxidation.** *Nature* 2012, **491**:541-546.
44. Chen S-C, Musat N, Lechtenfeld OJ, Paschke H, Schmidt M, Said N, Popp D, Calabrese F, Stryhanyuk H, Jaekel U *et al.*: **Anaerobic oxidation of ethane by archaea from a marine hydrocarbon seep.** *Nature* 2019, **568**:108-111.
45. Nielsen LP, Risgaard-Petersen N, Fossing H, Christensen PB, Sayama M: **Electric currents couple spatially separated biogeochemical processes in marine sediment.** *Nature* 2010, **463**:1071-1074.
46. Canfield DE: **Reactive iron in marine sediments.** *Geochim Cosmochim Acta* 1989, **53**:619-632.
47. Morse JW, Cornwell JC: **Analysis and distribution of iron sulfide minerals in recent anoxic marine sediments.** *Mar Chem* 1987, **22**:55-69.
48. Kappler A, Straub KL: **Geomicrobiological cycling of iron.** *Rev Miner Geochem* 2005, **59**:85-108.
49. Rotaru A-E, Posth NR, Loscher CR, Miracle MR, Vicente E, Cox RP, Thompson J, Poulton SW, Thamdrup B, Löscher C *et al.*: **Interspecies interactions mediated by conductive minerals in the sediments of the iron rich meromictic lake La Cruz, Spain.** *Limnetica* 2019, **38**:21-40.
50. Kato S, Igarashi K: **Enhancement of methanogenesis by electric syntrophy with biogenic iron-sulfide minerals.** *Microbiology open* 2019, **8**:e00647.
51. Beulig F, Roy H, McGlynn SE, Jørgensen BB: **Cryptic CH<sub>4</sub> cycling in the sulfate-methane transition of marine sediments apparently mediated by ANME-1 archaea.** *ISME J* 2019, **13**:250-262.
52. Martins G, Salvador AF, Pereira L, Alves MM: **Methane production and conductive materials: a critical review.** *Environ Sci Technol* 2018, **52**:10241-10253
- This review article is a comprehensive report of iron- and carbon-based additions to stimulate cMIET in engineered systems. The authors also discuss the controls used by previous researchers when investigating the mechanism of electron transfer.
53. Salvador AF, Martins G, Melle-franco M, Serpa R, Stams AJM, Cavaleiro AJ, Pereira MA, Alves MM: **Carbon nanotubes accelerate methane production in pure cultures of methanogens and in a syntrophic coculture.** *Environ Microbiol* 2017, **19**:2727-2739.
54. Holmes DE, Shrestha PM, Walker DJF, Dang Y, Nevin KP, Woodard TL, Lovley DR: **Metatranscriptomic evidence for direct interspecies electron transfer between *Geobacter* and *Methanotherix* species in methanogenic rice paddy soils.** *Appl Environ Microbiol* 2017, **83** AEM.00223-17.
55. Morita M, Malvankar NS, Franks AE, Summers ZM, Giloteaux L, Rotaru A-E, Rotaru C, Lovley DR: **Potential for direct interspecies electron transfer in methanogenic wastewater digester aggregates.** *mBio* 2011, **2**:e00159-11.
56. Wu WM, Jain MK, Hickey RF, Zeikus JG: **Perturbation of syntrophic isobutyrate and butyrate degradation with formate and hydrogen.** *Biotechnol Bioeng* 1996, **52**:404-411.
57. Deutzmann JS, Sahin M, Spormann AM: **Extracellular enzymes facilitate electron uptake in biocorrosion and bioelectrosynthesis.** *mBio* 2015, **6**:1-8.
58. Palacios PA, Snoeyenbos-West O, Löscher CR, Thamdrup B, Rotaru A-E: **Baltic Sea methanogens compete with acetogens for electrons from metallic iron.** *ISME J* 2019, **13**:3011-3023.

59. Yee MO, Deutzmann J, Spormann A, Rotaru A-E: **Cultivating electroactive microbes-from field to bench.** *Nanotechnology* 2020, **31**.
60. Van Steendam C, Smets I, Skerlos S, Raskin L: **Improving anaerobic digestion via direct interspecies electron transfer requires development of suitable characterization methods.** *Curr Opin Biotechnol* 2019, **57**:183-190.
61. Zhang D, Shen J, Shi H, Su G, Jiang X, Li J, Liu X, Mu Y, Wang L: **Substantially enhanced anaerobic reduction of nitrobenzene by biochar stabilized sulfide-modified nanoscale zero-valent iron: process and mechanisms.** *Environ Int* 2019, **131** 105020.
62. Wang Z, Yin Q, Gu M, He K, Wu G: **Enhanced azo dye Reactive Red 2 degradation in anaerobic reactors by dosing conductive material of ferrous oxide.** *J Hazard Mater* 2018, **357**:226-234.
63. Ferrero P, Izquierdo M, Álvarez-Hornos FJ, Peña-Roja JM, Martínez-Soria V: **Influence of activated carbon on performance and microbial communities in the treatment of solvent pollutant mixtures in a continuous stirred tank reactor.** *Environ Sci Water Res Technol* 2020, **6**:1445-1455.
64. Leitão P, Aulenta F, Rossetti S, Nouws HPA, Danko AS: **Impact of magnetite nanoparticles on the syntrophic dechlorination of 1, 2-dichloroethane.** *Sci Total Environ* 2018, **624**:17-23.
65. Aulenta F, Fazi S, Majone M, Rossetti S: **Electrically conductive magnetite particles enhance the kinetics and steer the composition of anaerobic TCE-dechlorinating cultures.** *Process Biochem* 2014, **49**:2235-2240.
66. Zhuang L, Tang Z, Ma J, Yu Z, Wang Y, Tang J: **Enhanced anaerobic biodegradation of benzoate under sulfate-reducing conditions with conductive iron-oxides in sediment of Pearl river estuary.** *Front Microbiol* 2019, **10**:374.
67. Kato S, Wada K, Kitagawa W, Mayumi D, Ikarashi M, Sone T, Asano K, Kamagata Y: **Conductive iron oxides promote methanogenic acetate degradation by microbial communities in a high-temperature petroleum reservoir.** *Microbes Environ* 2019, **34**:95-98.
68. Ye Q, Zhang Z, Huang Y, Fang T, Cui Q, He C, Wang H: **Enhancing electron transfer by magnetite during phenanthrene anaerobic methanogenic degradation.** *Int Biodeterior Biodegradation* 2018, **129**:109-116.
69. Aulenta F, Tucci M, Cruz Viggli C, Dolfing J, Head IM, Rotaru A-E: **An underappreciated DIET for anaerobic petroleum hydrocarbon-degrading microbial communities.** *Microb Biotechnol* 2020 <http://dx.doi.org/10.1111/1751-7915.13654>. Epub ahead of print.
70. Pagnozzi G, Carroll S, Reible DD, Millerick K: **Biological natural attenuation and contaminant oxidation in sediment caps: recent advances and future opportunities.** *Curr Pollut Rep* 2020.
71. Bonaglia S, Broman E, Brindefalk B, Hedlund E, Hjorth T, Rolff C, Nascimento FJA, Udekwi K, Gunnarsson JS: **Activated carbon stimulates microbial diversity and PAH biodegradation under anaerobic conditions in oil-polluted sediments.** *Chemosphere* 2020, **248** 126023.
72. Pagnozzi G, Carroll S, Reible DD, Millerick K: **Powdered Activated Carbon (PAC) amendment enhances naphthalene biodegradation under strictly sulfate-reducing conditions.** *Environ Pollut* 2020.
73. Light SH, Su L, Rivera-Lugo R, Cornejo JA, Louie A, Iavarone AT, Ajo-Franklin CM, Portnoy DA: **A flavin-based extracellular electron transfer mechanism in diverse Gram-positive bacteria.** *Nature* 2018, **562**:140-144.
74. Wang M, Nie Y, Wu XL: **Membrane vesicles from a *Dietzia* bacterium containing multiple cargoes and their roles in iron delivery.** *Environ Microbiol* 2020, **00**:1-11.
75. Pande S, Shitit S, Freund L, Westermann M, Bertels F, Colesie C, Bischofs IB, Kost C: **Metabolic cross-feeding via intercellular nanotubes among bacteria.** *Nat Commun* 2015, **6**.
76. Shrestha PM, Rotaru A-E, Aklujkar M, Liu F, Shrestha M, Summers ZM, Malvankar N, Flores DC, Lovley DR: **Syntrophic growth with direct interspecies electron transfer as the primary mechanism for energy exchange.** *Environ Microbiol Rep* 2013, **5**:904-910.
77. Rowe AR, Xu S, Gardel E, Bose A, Girguis P, Amend JP, El-Naggar MY: **Methane-linked mechanisms of electron uptake from cathodes by *Methanosarcina barkeri*.** *mBio* 2019, **10**:1-12.
78. Holmes DE, Rotaru A-E, Ueki T, Shrestha PM, Ferry JG, Lovley DR: **Electron and proton flux for carbon dioxide reduction in *Methanosarcina barkeri* during direct interspecies electron transfer.** *Front Microbiol* 2018, **9**:3109.
79. Kato S, Hashimoto K, Watanabe K: **Methanogenesis facilitated by electric syntrophy via (semi)conductive iron-oxide minerals.** *Environ Microbiol* 2012, **14**:1646-1654.
80. Smith JA, Nevin KP, Lovley DR: **Syntrophic growth via quinone-mediated interspecies electron transfer.** *Front Microbiol* 2015, **6**:1-8.
81. Wang O, Zheng S, Wang B, Wang W: **Necessity of electrically conductive pili for methanogenesis with magnetite stimulation.** *PeerJ* 2018, **2**:1-14.
82. Zhang X, Xia J, Pu J, Cai C, Tyson GW, Yuan Z, Hu S: **Biochar-mediated anaerobic oxidation of methane.** *Environ Sci Technol* 2019, **53**:6660-6668.