UC Irvine

UC Irvine Previously Published Works

Title

Short Enantioselective Total Syntheses of Cheloviolenes A and B and Dendrillolide C via Convergent Fragment Coupling Using a Tertiary Carbon Radical.

Permalink

<https://escholarship.org/uc/item/6vh8n3kv>

Journal

The Journal of organic chemistry, 83(13)

ISSN

0022-3263

Authors

Garnsey, Michelle R
Slutskyy, Yuriy
Jamison, Christopher R
et al.

Publication Date

2018-07-01

DOI

10.1021/acs.joc.7b02458

Peer reviewed

Short Enantioselective Total Syntheses of Cheloviolenes A and B and Dendrillolide C via Convergent Fragment Coupling Using a Tertiary Carbon Radical

Michelle R. Garnsey,^{†,§} Yuriy Slutskyy,^{†,§} Christopher R. Jamison,^{†,§} Peng Zhao,[†] Juyeol Lee,[†] Young Ho Rhee,[‡] and Larry E. Overman^{*,†,§}

[†]Department of Chemistry, University of California, Irvine, California 92697-2025, United States

[‡]Department of Chemistry, Pohang University of Science and Technology, Hyoja-dong San 31, Pohang, Kyungbook 790-784, Republic of Korea

S Supporting Information

ABSTRACT: The development of a convergent fragment coupling strategy for the enantioselective total syntheses of a group of rearranged spongian diterpenoids that harbor the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one unit is described. The key bond disconnection relies on a late-stage fragment coupling between a tertiary carbon radical and an electron-deficient alkene to unite two ring systems and form two new stereocenters, one of which is quaternary, in a stereoselective and efficient manner. This strategy is applied toward scalable 14–15 step syntheses of three rearranged spongian diterpenoids, cheloviolenes A and B, and dendrillolide C.

INTRODUCTION

Synthetic strategies that rely on independent syntheses of fragments, followed by their late-stage union are key to efficient preparation of complex molecules. The major drawback of this convergent approach is the small pool of privileged reactions that are reliable, chemoselective, and high-yielding. Some of the most widely used methods having these properties are Diels–Alder reactions, transition metal-catalyzed cross couplings, Nozaki–Hiayama–Kishi couplings, and olefin metathesis.¹ Bimolecular late-stage fragment couplings that form two new stereocenters are especially challenging, particularly in cases when one of the newly formed centers is quaternary.² Formation of such sterically congested sp^3 – sp^3 σ bonds by intermolecular addition of nucleophilic tertiary radicals to electron deficient olefins is attractive for several reasons: the early transition state with a long forming bond (~ 2.5 Å) reduces the enthalpic penalty of bringing two bulky fragments together;³ the high rates of addition⁴ and stereoselection^{4a,5} realized in additions of tertiary radicals to Michael acceptors often leads to efficient fragment-couplings with reliable stereochemical outcomes. Despite the aforementioned appealing features of the Giese reaction of tertiary carbon radicals, which were known since the 1980s,^{4a,6} bimolecular radical reactions have not been used to unite complex fragments because of the large excesses (commonly 3–10-fold) of one of

the coupling components often required. The recent advent of visible-light photoredox catalysis to generate carbon radicals under mild conditions,⁷ a method that is compatible with most polar functional groups, led to a resurgence of radical-based approaches for convergent construction of complex molecules.⁸ An overarching goal of our research has been to capitalize on the appealing features of the Giese reaction of tertiary carbon radicals to perform a late-stage union of two ring systems in 1:1 stoichiometry. One example of this approach leading to short enantioselective total syntheses of a family of rearranged spongian diterpenoids is the subject of this report.

BACKGROUND

More than 100 natural products harbor a *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one (1) fragment (Figure 1).⁹ The marine diterpenoids dendrillol 1 (2),¹⁰ gracilin C (3),¹¹ and darwinolide (4)¹² exemplify members of this group in which the dioxabicyclooctanone unit is fused to a larger polycyclic ring system. In others, such as the fungal sesquiterpenoid 5,¹³ the dioxabicyclic unit is isolated and joined by a single bond to a second cyclic or polycyclic fragment. Among these is a subset of

Special Issue: Synthesis of Antibiotics and Related Molecules

Received: September 27, 2017

Published: November 13, 2017

A. The *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one ring system and representative structurally diverse natural products harboring this structural fragment.

B. Rearranged spongian diterpenoids having a *cis*-dioxabicyclo[3.3.0]octan-3-one fragment attached to a quaternary carbon of a hydrocarbon fragment.

C. Proposed biogenesis of rearranged spongian diterpenoids such as **6–11** from a precursor having the spongian diterpenoid skeleton.

Figure 1.

rearranged spongian diterpenoids of marine origin, illustrated by **6–11**, having the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one fragment joined at C-6 to a 14-carbon bicyclic hydrocarbon fragment.¹⁴ The first member of this diterpenoid group to be reported was norrisolide (**6**), whose structure and relative configuration were established by single-crystal X-ray diffraction analysis.¹⁵ In the more common members of this group depicted in Figure 1B, the dioxabicyclo[3.3.0]octan-3-one fragment is joined by a single bond to a quaternary carbon of the hydrocarbon unit. The hydrocarbon fragment of these diterpenoids can reside on either the convex or concave face of the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one fragment, as exemplified respectively by cheloviolene A (**7**)^{16,17} and macfarlandin C (**9**).¹⁸ A freely rotating single bond having three staggered conformers of roughly similar energies characterizes this group of rearranged spongian diterpenoids. As a result of this structural feature, relating the configurations of the two chiral bicyclic fragments is notably challenging in the absence of X-ray structures. Only the relative configurations of macfarlandin C (**9**) and cheloviolene A (**7**) are known with certainty by virtue of single-crystal X-ray analyses.^{16,18} The relative configurations of the other diterpenoids in this family are proposed to be as depicted in Figure 1B as a result of their presumed biosynthesis from precursors having a spongian skeleton (see Figure 1C).^{11,19} Absolute configurations for these natural products have not been established experimentally and are also proposed on the basis of this biosynthetic hypothesis.²⁰

The biological activity of rearranged spongian diterpenoids of marine origin has been little explored.^{17,18,21} Our recent interest in the group of natural products exemplified in Figure 1B was stimulated by Sütterlin's observations from a screen of small molecule marine natural products that macfarlandin E (**13**,

Figure 2), a rearranged spongian diterpenoid harboring a 2,7-dioxabicyclo[3.2.1]octan-3-one fragment, exhibits unique

Figure 2.

Golgi-altering activity.²² Macfarlandin E (**13**) induces irreversible fragmentation of the Golgi apparatus with the fragments remaining in the pericentriolar region of the cell. This phenotype contrasts with the effects of other natural products such as brefeldin A,²³ ilimaquinone,²⁴ and norrisolide (**6**),²⁵ which cause Golgi fragmentation with the resulting fragments

being delocalized throughout the cytosol. In our initial studies, we prepared racemic *tert*-butyl analogues **15**–**18** of the rearranged spongian diterpenoids macfarlandin **E** (**13**), aplyviolene (**14**), and dendrillolide **A** (**10**) and explored their chemical reactivity and effects on the Golgi apparatus (Figure 2).^{22,26} Both the bridged and fused dioxabicyclooctanone ring systems were found to react with primary amines to form pyrrole products (e.g., **20**), presumably via the intermediacy of 1,4-dialdehyde **19** generated upon cleavage of the anomeric acetoxy group. Under physiologically relevant conditions, *tert*-butyl analogues **15**–**18** reacted with lysine chains of hen egg white lysozyme (HEWL) to form pyrrole conjugates **21**, a reaction that potentially could be the origin of the effects of these agents on the Golgi. The presence of an acetoxy substituent adjacent to the lactone carbonyl group in analogues **16** and **18** increased the extent of the lysine to pyrrole conversion and was important for induction of the macfarlandin **E** Golgi phenotype.²⁶ To further elucidate the mechanism by which dioxabicyclooctanones **15**–**18** trigger Golgi fragmentation, a more efficient preparation of these molecules was necessary as our initial route required many steps.²⁶

As analogues having the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one ring system were expected to be easier to access by chemical synthesis than their 2,7-dioxabicyclo[3.2.1]octan-3-one counterparts, our recent studies in this area focused on developing a short and versatile synthesis of 6-substituted *cis*-2,8-dioxabicyclo[3.3.0]octan-3-ones. Our plan was to employ fragment coupling reactions of tertiary carbon radicals generated from alcohol or carboxylic acid precursors by visible-light photoredox catalysis²⁷ to unite tertiary carbon fragments with an unsubstituted 5-alkoxybutenolide **22a** or a butenolide, such as **22b**, containing the additional two carbons of the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one product **24** (Scheme 1). In the former instance, alkylation of the coupled

Scheme 1

product **23a** with a haloacetate electrophile would be employed to append the additional two carbons. In either case, coupling of a tertiary radical with butenolides **22a** and **22b** was expected to proceed with high stereoselectivity from the face opposite to the 5-alkoxy substituent.²⁸ The task of relating the configurations of the hydrocarbon and dioxabicyclooctanone fragments in total syntheses of the rearranged spongian diterpenoids depicted in Figure 1B would be addressed by uniting enantiopure butenolide and tertiary hydrocarbon fragments to form the demanding bond joining the two attached ring systems.^{29,30} The sequence delineated in Scheme 1 could potentially access a wide variety of dioxabicyclooctanones **24** in 3–4 steps, representing a significant improvement on our earlier synthesis of **17** which required 14 steps.²⁶ The successful development of the synthetic strategy outlined in Scheme 1 and its use to complete short enantioselective total

syntheses of cheloviolenes **A** (**7**) and **B** (**8**), and dendrillolide **C** (**11**) are the subject of this report.^{31,32}

RESULTS AND DISCUSSION

Initial Exploratory Investigations. We examined the proposed sequence initially with 5-methoxybutenolide **28** containing all the carbon atoms of the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one ring system (Scheme 2). A ring closing

Scheme 2

metathesis route was developed for preparing racemic **28**.³³ The sequence starts with itaconic acid (**25**), which after converting selectively to the monomethyl ester by a known procedure,³⁴ was allylated to yield diester **26**. After examining several metathesis catalysts, the Stewart–Grubbs catalyst was found to be uniquely proficient at promoting the desired transformation to form butenolide **27**.³⁵ Optimization studies showed that reaction concentrations as high as 0.03 M could be employed before side products resulting from bimolecular metathesis were observed. The 5-methoxy substituent was introduced by initial bromination of **27** with NBS, followed by methanolysis³⁶ to deliver butenolide **28** in five steps from commercially available itaconic acid (**25**).

The fragment coupling was explored initially with the *N*-hydroxyphthalimide (NHP) ester of pivalic acid **29a** as the precursor of *tert*-butyl radical (Scheme 3). Using the modification of Okada's conditions³⁷ developed during our first studies in this area,³⁰ the Ru(bpy)₃-catalyzed reaction of equimolar amounts of activated ester **29a** and butenolide **28** in the presence of low-intensity blue LEDs, 1.5 equiv of a Hantzsch ester (diethyl 1,4-dihydro-2,6-dimethylpyridine-3,5-

Scheme 3

^aDetermined by ¹H NMR integration of the crude reaction mixture.

^bIsolated yield.

dicarboxylate), and 1 equiv of *i*-Pr₂NEt proceeded to give the expected stereoisomeric products **30a** and **31** in a 10:1 ratio favoring the formation of the desired all-*trans* adduct **30a** (isolated in 72% yield).^{4a,38} The relative configuration of these stereoisomers was assigned initially on the basis of vicinal coupling constants and ¹H NOE data.^{39,40} In addition to products **30a** and **31**, lactone **32**, which would arise from base-promoted isomerization of the double bond prior to radical coupling, was formed in ~5% yield. As would be expected, increasing the amount of *i*-Pr₂NEt enhanced the formation of byproduct **32**, whereas replacing *i*-Pr₂NEt with *i*-Pr₂NEt·HBF₄ led to **32** being produced in trace amounts only. To our initial surprise, stereoselection was reduced significantly (to 2:1) in coupling reactions conducted in the presence of *i*-Pr₂NEt·HBF₄. Although not understood at the time these experiments were carried out, our recent investigations suggest that in the presence of the basic amine the α -acyloxy radical produced upon conjugate addition is terminated by single-electron transfer (SET) followed by protonation of the lactone enolate, whereas hydrogen-atom transfer (HAT) predominates in the presence of *i*-Pr₂NEt·HBF₄.⁴¹ It would not be surprising that the former termination process is more stereoselective.⁴² Control experiments established that product ratios did not change with time and that adducts **30a** and **31** did not equilibrate at room temperature in the presence of excess *i*-Pr₂NEt.⁴³ Many reaction variables are reported to affect stereoselection in HAT.⁴⁴ We examined a number of these in the hope of enhancing the formation of the minor adduct **31**; however, no conditions were identified that led to this product in useful yield.

The salient results of our efforts to explore the scope of the fragment coupling step and to elaborate the conjugate-addition products to *cis*-2,8-dioxabicyclo[3.3.0]octan-3-ones are summarized in Scheme 4. The NHP esters **29b** and **29c**, the latter

Scheme 4

harboring a side chain chosen for use in future studies to pursue Golgi molecular targets, underwent stereoselective (*dr* = 5–10:1) reductive photoredox coupling with butenolide **28** in the presence of 1 equiv of *i*-Pr₂NEt to yield the all-*trans* trisubstituted products **30b** and **30c** in respectively 53% and 52% yield after purification on silica gel. In initial scouting studies, butyrolactone **30a** was allowed to react with 1 equiv of (*i*-Bu)₂AlH (DIBALH) at –78 °C in the hope that the lactone

carbonyl could be selectively reduced to give, after intramolecular lactonization, *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one **34a**. However, this reaction did not afford **34a**, but gave rise to a 1:1 mixture of the starting lactone ester **30a** and dioxabicyclic lactol **33a**, suggesting that intramolecular lactonization of the diisobutylaluminum lactol intermediate was faster than reduction of the starting butyrolactone. After screening several reductants and reaction conditions with only modest success, we turned to accomplishing the desired conversion in two steps. A quick survey showed that several oxidants, including Br₂, PCC, acetone (via Oppenauer oxidation), and Ag₂CO₃ supported on Celite could transform **33a** to dioxabicyclooctanone **34a**, the latter leading to the highest yield of the desired product. The final sequence involved reduction of the coupled products **30** with 2.1 equiv of (*i*-Bu)₂AlH at –78 °C and isolation of the crude mixture of lactol epimers **33a–c** after workup with an aqueous solution of Rochelle's salt. This crude mixture of lactol intermediates was then directly oxidized with Ag₂CO₃/Celite in refluxing toluene to give *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one **34a–c** in 40–64% yield.

Although the convergent construction of *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one was appealing, the yield of the all-*trans* product from the addition of tertiary radicals to butenolide **28** was compromised by the formation of the two additional isomeric products **31** and **32** depicted in Scheme 3. As a result, we examined the alternate sequence in which an unsubstituted 5-alkoxybutenolide would be the radical acceptor, an approach that could be advantageous because of the commercial availability of several butenolides of this type in high enantiomeric purity.⁴⁵ As summarized in Scheme 5, reductive

Scheme 5

photoredox-catalyzed coupling of the *tert*-NHP ester **29a** with racemic 5-methoxybutenolide (**35**) gave exclusively the *trans* product **36** in 73% yield. The reaction proceeded in identical yield, also with complete stereoselectivity, when *i*-Pr₂NEt·HBF₄ was substituted for *i*-Pr₂NEt. Alkylation of the lithium enolate of **36** with methyl bromoacetate occurred stereoselectively to give a single product **30a** in 56% yield. Although the transformations summarized in Schemes 3–5 likely could be further optimized, we chose to move forward to explore the application of this chemistry for the synthesis of rearranged-spongian diterpenoids that harbor a *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one fragment.

First-Generation Total Synthesis of (+)-Cheloviolene

A. The plan for synthesis of (+)-cheloviolene **A** (**7**) using the approach developed in our exploratory studies is outlined in retrosynthetic format in Scheme 6. The decisive step would be the Giese coupling of *cis*-perhydroazulene radical **38** with butenolide (*S*)-**35** to form tricyclic adduct **37**. The configuration of the C-8 and C-14 stereocenters of the attached chiral bicyclic rings of (+)-cheloviolene **A** would derive from the

Scheme 6

expected high preference for this union to take place from the convex face of the *cis*-perhydroazulene radical³⁰ and the face opposite the methoxy substituent of butenolide (S)-35 to form coupled product 37.

The pivotal radical coupling was examined initially using the enantiopure *cis*-perhydroazulene NHP ester 39, which we had prepared earlier in 14 steps from (+)-fenchone.³⁰ Exposure of the activated ester 39 and butenolide 35 to Ni-catalyzed Giese reaction conditions did not lead to the formation of the desired product 37.^{46,47} In contrast, when equimolar amounts of coupling partners 39 and 35 were subjected to photoredox reaction conditions the desired lactone 37 was obtained as a single stereoisomer (eq 1). However, the efficiency of the

reaction was poor, resulting in a modest 30% yield of coupled product 37. The major byproducts were derived from premature reduction of the tertiary radical generated upon reductive cleavage of *N*-acyloxyphthalimide 39. We hypothesized that the presence of Hantzsch ester, a known hydrogen atom donor,⁴⁸ was largely responsible for the undesired reduction of the intermediate tertiary radical. To circumvent premature reduction of the tertiary radical, we examined the coupling of *cis*-perhydroazulene carboxylic acid 40 with 1 equiv of (S)-35 using the Ir(III)-catalyzed photoredox conditions developed by MacMillan⁴⁹ in which no external reductants or hydrogen atom sources are required. In this case, the desired product 37 was isolated in 44% yield. Once again premature reduction of the tertiary radical was observed,⁵⁰ suggesting that a more activated acceptor would be required to efficiently trap sterically encumbered *cis*-perhydroazulene radical 38.

Despite modest yields of the key coupling step, we turned our attention to elaboration of the lactone fragment of addition product 37 to a *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one to conclude a first-generation total synthesis of (+)-chelviolene A (7) (Scheme 7). Deprotonation of lactone 37 with LDA at $-78\ ^\circ C$, followed by addition of methyl iodoacetate furnished

Scheme 7

methyl ester 41, as a single stereoisomer, in 45% yield. Next, reduction of 41 with excess DIBALH at $-78\ ^\circ C$, followed by direct oxidation of the resulting epimeric mixture of dioxabicyclic lactols under Fétizon oxidation conditions provided tetracyclic intermediate 42 in 80% yield. Finally, hydrolysis of 42 with dilute aqueous HCl at $40\ ^\circ C$ furnished (+)-chelviolene A (7), mp: $157-158\ ^\circ C$, in 70% yield. Synthetic 7 exhibits a higher optical rotation from the one reported in literature, synthetic: $[\alpha]_D^{22} +49$ ($c\ 0.11, CHCl_3$), literature: $[\alpha]_D +4.5$ ($c\ 0.11, CHCl_3$).¹⁶ However, the spectroscopic data for synthetic 7 compared well with those reported for the natural product isolated from the New Zealand sponge *Chelonaphysilla violacea*, leaving little doubt as to their identity.¹⁷ In addition, the structure of synthetic (+)-chelviolene A was confirmed by single-crystal X-ray analysis.⁴⁰

Development of a More Concise Second-Generation Total Synthesis Strategy. Completion of the synthesis of (+)-chelviolene A (7) validated our strategy and identified several problems that needed to be addressed to secure a concise approach to the family of spongian diterpenoids containing a *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one fragment. First, the efficiency of the radical fragment-coupling step had to be improved, which would likely necessitate use of a more activated butenolide radical acceptor. Second, a shorter synthesis of a precursor of the *cis*-perhydroazulene tertiary radical 38 would be needed, as our earlier preparation of carboxylic acid 40 required 13 steps.³⁰ The recent development of convenient methods to generate tertiary radicals from tertiary alcohols^{51,52} directed our attention to the use of a tertiary *cis*-perhydroazulene alcohol as the radical precursor.

Efficient Synthesis of *cis*-Perhydroazulenol 54. A *cis*-perhydroazulene tertiary alcohol, which at the time was assigned structure A (Scheme 8), was an intermediate in our earlier synthesis of *cis*-perhydroazulene carboxylic acid 40.³⁰ As this alcohol was accessed on gram-scale in only eight steps from (+)-fenchone, we initially examined whether it, or potentially an exomethylene analogue, could be utilized in the crucial radical coupling event. To test the feasibility of this approach, alcohol 43 was condensed with methyl chlorooxacetate and the resulting diester selectively saponified with 0.95 equiv of LiOH to provide *cis*-perhydroazulene lithium oxalate salt 44 in 85% yield. Upon attempted coupling of oxalate salt 44 with (R)-butenolide (R)-35 using the Ir(III)-catalyzed photoredox conditions developed earlier,^{51,52} no coupling product was observed, rather a tricyclic lactone 45, whose structure was unambiguously established via single-crystal X-ray analysis of the 4-nitrobenzoate derivative, was isolated in 71% yield.⁴⁰ To our surprise, the perhydroazulene fragment of 45 is *trans*-fused and the configuration at C-1 is opposite to that expected from a

Scheme 8

precursor of structure A. Single-crystal X-ray analysis of the phenylhydrazone derivative of 43 confirmed that tertiary alcohol 43 has the stereostructure depicted in Scheme 8, with the C-1 and C-8a stereocenters opposite to those found in the originally misassigned structure A.⁴⁰ To the best of our knowledge, the addition of an alkoxyacyl radical to a ketone carbonyl group is without precedent.

Brief comment on our original assignment of structure A to the perhydroazulenol intermediate used to prepare *cis*-perhydroazulene carboxylic acid 40 is warranted. First, this structural misassignment was not detected in our earlier studies because the next step in the synthesis of 40, dehydration of tertiary alcohol 43 to form the corresponding conjugated enone, removed the C-1 and C-8a stereocenters of 43.³⁰ In this earlier synthesis, the perhydroazulene ring was constructed by intramolecular nitrile oxide cycloaddition of nitro precursor 46 to yield an isoxazoline intermediate, whose structure was assigned as 47 on the basis of excellent precedent.⁵³ Extensive 2D NMR experiments confirm the relative configuration of cycloadduct 47. Thus, epimerization of the ring fusion and alcohol stereocenter in route to 43 must occur during reduction of isoxazoline 47 under acidic conditions.⁵⁴ This epimerization, which undoubtedly arises by a retro aldol/aldol sequence, has precedent in the unmasking of related tricyclic isoxazolines in the presence of boric acid.^{55–57}

It was clear at this point that a new route to a *cis*-perhydroazulenol intermediate would be required. We conjectured in addition that the tertiary alcohol should reside on the convex face of the *cis*-perhydroazulene ring to avoid the possibility of unwanted cyclization of a transiently generated alkoxyacyl radical onto a proximal carbonyl or exomethylene functional group. Our development of a short enantioselective route to access such an intermediate, *cis*-perhydroazulenol 54, is summarized in Scheme 9. The synthesis began with inexpensive (+)-fenchone, whose oxime derivative underwent a known Beckmann fragmentation when heated at reflux in aqueous sulfuric acid to give cyclopentene nitrile 49 and its $\Delta^{1,5}$ isomer in near-equal amounts.⁵⁸ Although these double-bond regioisomers can be separated by careful chromatography on silver nitrate-impregnated silica gel,³⁰ we found it more

Scheme 9

convenient on scale to selectively epoxidize the less-hindered $\Delta^{1,5}$ isomer of this mixture thereby allowing pure 49 to be obtained reliably on 10 g scales by simple flash chromatography. Conventional reduction of nitrile 49 and Wittig olefination of the aldehyde product gave dienyl nitrile 50 in 87% yield over two steps. As the prelude to forming the seven-membered ring, the trisubstituted double bond of 50 was selectively epoxidized by reaction with 1 equiv of *m*-CPBA at –10 °C in CH₂Cl₂ to give an 8:1 mixture of stereoisomers from which the major isomer 51 was isolated after chromatographic purification in 81% yield. Deprotonation of 51 with 1 equiv of lithium 2,2,6,6-tetramethylpiperide (LiTMP) induced stereospecific cyclization to form *cis*-perhydroazulenol 52 in 85% yield.^{59,60} To our knowledge, this is the first example of forming a seven-membered ring by Stork epoxy-nitrile cyclization,⁶¹ an outcome undoubtedly assisted by the *cis*-double bond in the tether and the presence of *gem*-dimethyl substitution. After examining several nonconventional methods for transforming the nitrile substituent to an exomethylene group in one step, this transformation was ultimately realized in high yield by a two-step sequence. First, reaction of 52 with Raney-Ni and hydrogen (50 atm) in the presence of paraformaldehyde delivered amino alcohol 53 in 91% yield. Formation of the corresponding *N*-oxide and heating this crude intermediate to 120 °C in DMF occasioned clean Cope elimination to provide alcohol 54 in 77% yield.⁶²

Optimization in a Model Series of the Fragment Coupling Step and Elaboration of the Coupled Product to a 6-Substituted *cis*-2,8-Dioxabicyclo[3.3.0]octan-3-one. In addition to developing an expedited route to a precursor of *cis*-perhydroazulene radical 38, both the conjugate addition step and the elaboration of the butenolide fragment of the coupled product to a *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one would need to be optimized significantly in order to define an efficient route to (+)-cheloviolene A (7) and congeners. Initially we wished to evaluate whether incorporation of a radical-stabilizing group at the α -position of a 5-alkoxybutenolide would increase the efficiency of the fragment-coupling step. An obvious choice would be a chlorine substituent.^{30,63} Undoubtedly reflecting the sensitivity of the allylic acetal functionality, our attempts to directly introduce chlorine at C-3 of a 5-alkoxybutenolide led to either decomposition or, when attempted with enantiopure 5-alkoxybutenolides, partial racemization.⁶⁴ As a result, our efforts shifted to the development of

a new route to 3-chloro-5-alkoxybutenolides that would exploit Rhee's recent method for enantioselective synthesis of acetals (Scheme 10).⁶⁵ The sequence began with palladium-catalyzed

Scheme 10

enantioselective alkoxylation of (*D*)-menthol-derived allene **56** with allylic alcohol **55** to deliver mixed acetal **57** in 99% yield as a single detectable diastereomer by ^1H NMR analysis.⁶⁶ Subjection of diene **57** to the Hoveyda-Grubbs second-generation catalyst in toluene at 60°C gave rise to dihydrofuran **58** in excellent yield on gram-scale. Incorporation of the phenyl group in the alkenyl chloride fragment was crucial to the success of this ring-closing metathesis as reported by Dorta.^{67,68} After much experimentation, we found that the demanding allylic oxidation of **58** could be accomplished with CrO_3 and *tert*-butylhydroperoxide⁶⁹ to reliably give butenolide **59** in 32% yield.^{70–72} The opposite enantiomer of butenolide **59** was readily accessible by the same sequence, starting from (*L*)-menthol and employing the opposite enantiomer of the Trost ligand.

Our exploratory studies, summarized in Scheme 11, aimed at developing a one-pot sequence for directly transforming a

Scheme 11

tertiary alcohol to a coupled product and optimizing the coupling with a 5-alkoxybutenolide. We eventually found that reaction of 1-methylcyclohexanol (**60**) with 1 equiv of oxalyl chloride at room temperature in DME, followed by addition of water and 3 equiv of K_2HPO_4 cleanly generated the potassium hemioxalate intermediate **61** *in situ*. Direct addition of 1 equiv of 5-methoxybutenolide (**35**), followed by photoredox-catalyzed coupling as described previously⁵¹ provided *trans* adduct **63** as a single stereoisomer in 58% yield.⁷³ Using menthyloxybutenolide **62**,⁷⁴ coupled product **64** was generated, again as a single stereoisomer, in this case in 60% yield. The analogous coupling with butenolide **59** harboring a 3-chloro

substituent was more efficient delivering a 3:1 mixture of **65** and dechlorinated analogue **64** in 78% combined yield. Addition of 10 equiv of tri-*n*-butylamine to the reaction mixture following the initial fragment coupling with butenolide **59** and allowing the subsequent photocatalytic dechlorination⁷⁵ to proceed for 4 h gave conjugate addition product **64** in 80% yield in one-step from alcohol **60**.

In our first-generation synthesis of (+)-cheloviolene A, a four-step sequence proceeding in 25% yield was used to fashion the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one moiety from the butyrolactone fragment of the coupled product (Scheme 7). We anticipated that this elaboration could be shortened by one step by introducing the acetic ester side chain as a *tert*-butyl ester. This sequence in the 1-methylcyclohexyl model series is summarized in Scheme 12. Enolization of coupled product **64**

Scheme 12

with LiHMDS, followed by trapping with *tert*-butyl bromoacetate took place stereoselectively to give *tert*-butyl ester **66** in high yield. As expected, the steric bulk of the *tert*-butyl ester simplified chemoselective reduction of the lactone carbonyl group of **66** such that reaction with 2.4 equiv of (*i*-Bu)₂AlH in toluene at -78°C delivered lactol **67** in 72% yield. Finally, exposure of this intermediate to 2 M HCl formed the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one moiety and cleaved the menthyl acetal to provide tricyclic product **68** in 61% yield. This expedited sequence furnished **68** in 38% yield over three steps from coupled product **64**.

Total Syntheses of (+)-Cheloviolene B, (+)-Dendrillo-lide C and Second-Generation Total Synthesis of (+)-Cheloviolene A. Our studies began by optimizing the efficiency of the fragment-coupling reaction between butenolide **ent-59** and *cis*-perhydroazulene oxalate salts **69**. In this study, the oxalate salt intermediates were generated by selective hydrolysis of the mixed oxalate diester formed from *cis*-perhydroazulenol **54** and methyl chlorooxalate (Table 1).⁵¹ An initial solvent screen revealed DME and THF to be superior to other solvents, with the Ir(III)-catalyzed visible-light photocatalytic reaction yielding mixtures of coupled product **70** and its dechlorinated analogue **71** in ~40% combined yield (entries 1–4).⁷⁶ We utilized THF in our further optimization experiments because of the overall cleaner reaction profile and lower amounts of dechlorinated product **71** in this solvent. Increasing the amount of water in the reaction was deleterious to reaction efficiency (entry 5). The yield of the transformation could be increased somewhat by using 1.5 equiv of the butenolide radical acceptor (entry 6). However, since our objective was to optimize the coupling step using equimolar amounts of the addends, 1 equiv of the butenolide was used in subsequent experiments.

The yield of the conjugate addition was dramatically enhanced when the concentration of the reaction was increased

Table 1. Optimization of the Fragment Coupling Between **69** and *ent*-**59**

Reaction scheme showing the coupling of **69** (1 equiv) and *ent*-**59** (1 equiv) under conditions: 2 mol % Ir[dF(CF₃)ppy]₂(dtbbpy)PF₆, 2 X 34 W blue LEDs. The products are **70** (R = Cl) and **71** (R = H).

entry	M	conditions	70 , yield (%) ^a	71 , yield (%) ^a
1	Li	MeCN (0.05 M), H ₂ O (10 equiv)	0	0
2	Li	DMF (0.05 M), H ₂ O (10 equiv)	24	0
3	Li	DME (0.05 M), H ₂ O (10 equiv)	29	13
4	Li	THF (0.05 M), H ₂ O (10 equiv)	34	5
5	Li	THF (0.05 M), H ₂ O (100 equiv)	17	12
6 ^b	Li	THF (0.05 M), H ₂ O (10 equiv)	48	0
7	Li	THF (0.6 M), H ₂ O (5 equiv)	73 ^c	<5%
8 ^d	Li	THF (0.6 M), H ₂ O (5 equiv)	0	75 ^c
9	K	THF (0.6 M), H ₂ O (5 equiv)	72 ^c	<5%

^aDetermined by ¹H NMR integration relative to an internal standard (1,2-dibromo-4,5-methylenedioxybenzene). ^b1.5 equiv of butenolide *ent*-**59** was utilized. ^cIsolated yield. ^d*n*-Bu₃N (10 equiv) was added after coupling.

from 0.05 to 0.6 M, providing coupled product **70** in 73% isolated yield (entry 7).⁷⁷ Notably, only minor amounts of the dechlorinated product **71** were formed under these reaction conditions. Resubjection of **70** to the reaction conditions in the presence of *n*-Bu₃N led to a quantitative conversion of **70** to **71**.⁷⁵ Finally, we were able to perform the desired radical fragment coupling and dechlorination in one step to deliver **71** in 75% isolated yield by adding *n*-Bu₃N after 18 h and allowing the subsequent dechlorination to proceed for 4 h (entry 8). As expected from our earlier studies,⁵¹ the yield of the Ir(III) photoredox-mediated fragment coupling under optimized conditions was essentially identical when the oxalate counterion was switched from Li to K. (entry 9).

With reaction conditions for the pivotal fragment coupling step optimized, we turned to investigate accomplishing this union in one step from *cis*-perhydroazulenol **54** and butenolide *ent*-**59** (Scheme 13). To this end, a THF solution of tertiary alcohol **54** was allowed to react with 1 equiv of oxalyl chloride at room temperature for 6 h and then water and 3 equiv of K₂HPO₄ were added to generate potassium oxalate intermediate **72**. Addition of butenolide *ent*-**59** (1 equiv), the photocatalyst and irradiation with high-intensity blue LEDs for 18 h at 60 °C, followed by addition of excess *n*-Bu₃N and irradiation for an additional 4 h gave the desired product **71** in 68% yield after purification. This radical fragment coupling is noteworthy for several reasons: (1) it is the first example of a one-step coupling of an alcohol-derived tertiary radical with a Michael acceptor; (2) equimolar amounts of the two coupling partners are utilized; (3) the desired product **71** is obtained as a single diastereomer at both newly formed stereocenters; and (4) the reaction conditions allowed for selective isolation of either the direct coupling product, or its dehalogenated congener **71**.⁴⁰

Elaboration of coupled product **71** to cheloviolene B (**8**) began with alkylation of the lactone fragment of **71** with *tert*-butyl bromoacetate to deliver ester **73** in 91% yield. In contrast to our earlier results in the model series, reduction of **73** with 2.2 equiv of DIBALH at –78 °C in toluene did not lead exclusively to the desired tricyclic lactol product **75**, but rather to a 2:1 mixture of **75** and **76**, both as mixtures of lactol epimers. Attempts to modify the reduction conditions to furnish **75** selectively were met with no success. To converge these products, **76** was oxidized with PCC to give tetracyclic product **77** harboring the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one moiety. At this point, **75** and **77** were combined and exposed to 2 M HCl in THF/H₂O at room temperature to give what turned out to be (+)-cheloviolene B (**8**), mp: 189–190 °C, [α]_D²¹ +26.6, in 53% overall yield from intermediate **73**.

As the oxidation step in route to (+)-cheloviolene B (**8**) could not be avoided by introducing the two-carbon side chain as a *tert*-butyl ester, and processing both intermediates **75** and **76** was cumbersome, a more efficient route was developed to access (+)-cheloviolene B. In this sequence, the fragment-coupling product **71** was alkylated in high yield with methyl

Scheme 13

bromoacetate to give lactone ester **74**. In this methyl ester series, lactonization of the initially formed tricyclic lactol alkoxide intermediate generated upon exposure of **74** to excess DIBALH was sufficiently rapid at -78°C that tetracyclic lactol product **76** was formed in high yield. Without purification, this mixture of lactol epimers was directly oxidized with PCC to give **77** in 79% yield over the two steps. Exposure of **77** to 2 M HCl in THF/H₂O then afforded (+)-cheloviolene B. Using this four-step sequence, fragment-coupling product **71** was transformed to (+)-cheloviolene B (**8**) in 62% overall yield.

The relative configuration of synthetic (+)-cheloviolene B (**8**) was confirmed by single-crystal X-ray analysis,⁴⁰ whereas its absolute configuration follows rigorously from the absolute configuration of precursors **54** and *ent*-**59**. Structure **8** was originally proposed by Bobzin and Faulkner for a diterpenoid isolated from the marine sponge *Chelonaplysilla* sp. collected in Pohnpei, Federated States of Micronesia, and called chelonaplysins B.¹⁷ Two years later, Taylor and co-workers reported that the ¹H NMR spectra of so-called chelonaplysins B was identical to that of (+)-cheloviolene A (**7**), one of a series of diterpenoids isolated from the sponge *Chelonaplysilla violacea* collected from coastal waters of New Zealand.¹⁶ In addition to (+)-cheloviolene A (**7**), whose structure was confirmed by X-ray analysis, these workers isolated a related diterpenoid, (+)-cheloviolene B, which they assigned as the lactol epimer of cheloviolene A **78** (Figure 3).^{16b} It is this sponge isolate whose reported ¹H and ¹³C NMR spectra are indistinguishable from synthetic **8**. The structure of (+)-cheloviolene B must therefore be revised to be **8**.

Figure 3.

As first pointed out by Faulkner, a diterpenoid of structure **8** would be an outlier in the group of rearranged spongian diterpenoids exemplified in Figure 1B, because the relative configuration of its attached carbons, C-8 and C-14, differs at C-14 from that expected from a precursor having the spongian skeleton **12** (see Figure 1).¹⁷ They suggested that the unexpected *S* configuration at C-14 of a diterpenoid of structure **8** might arise by hydration of the corresponding enol ether double bond of another spongian diterpenoid, dendrillolide C, which this research group had isolated earlier from the sponge *Dendrilla* sp. obtained from a marine lake of Palau and assigned structure **11**.⁷⁸ To confirm the structure of dendrillolide C, establish its absolute configuration, and pursue Faulkner's suggestion for the origin of the unexpected stereostructure of cheviolene B (**8**), we carried out the experiments summarized in Scheme 14. Reaction of **8** with 2.5 equiv of MsCl and excess Et₃N in toluene at 90°C provided (+)-dendrillolide C (**11**), $[\alpha]_{\text{D}}^{21} +133$, in 77% yield. ¹H NMR and optical rotation data of synthetic dendrillolide C (**11**) were indistinguishable from those reported for the diterpenoid isolated from the sponge *Dendrilla* sp.⁷⁷ Exposing synthetic dendrillolide C (**11**) to 2 M HCl in THF/H₂O at 40°C led to the formation of a 1.2:1 mixture of cheloviolene B (**8**) and

Scheme 14

tricyclic furan **79**. Careful analysis of the ¹H NMR spectra of this crude product mixture showed that stereoisomers of **8** were not present in significant amounts.⁴⁰ Other acidic conditions (both Lewis and protic) that we investigated resulted in exclusive formation of furan **79** or intractable mixtures of products. That dendrillolide C (**11**) undergoes protonation at C14 preferentially from the *Si* face is consistent with the Faulkner initial proposal and with torsional effects dictating the stereochemical outcome of the hydration reaction *in vitro*.⁷⁹

Besides efficiency, the ability to access analogues by varying the structure of late-stage fragments is a distinct advantage of convergent synthesis strategies. The second-generation total synthesis of (+)-cheloviolene A (**7**) illustrates this point (Scheme 15). In this case, the fragment coupling step employed butenolide **59**, which trapped the tertiary radical generated from *cis*-pehydroazulenol **54** to give tricyclic lactone **80** as the only detectable stereoisomer in 76% yield. The 10% higher yield of this step than the analogous one employing *ent*-**59** (Scheme 13) reflects a match of the chirality of the enantiopure fragments combining to yield **80**. Processing of this product by the identical four-step sequence used to prepare (+)-cheloviolene B (**8**) gave (+)-cheloviolene A (**7**) in 70% overall yield from intermediate **80**.

CONCLUSION

Enantioselective total syntheses of rearranged spongian diterpenoids **7**, **8**, **11** exemplify advantages of convergent synthesis strategies based upon late-stage fragment coupling between a tertiary carbon radical and an electron-deficient alkene to unite two ring systems and form two new stereocenters, one of which is quaternary, in a stereoselective and efficient manner. (+)-Cheloviolene A (**7**) and (+)-cheloviolene B (**8**) were prepared in 11 steps from the known cyclopentene nitrile **49**⁵⁸ in respectively 22% and 18% overall yield, and in 14 steps and 5–7% overall yield from (+)-fenchone. These short synthetic sequences are made possible in part by the one-step generation of tertiary radical **38** from tertiary alcohol **54** and its *in situ* trapping with chlorobutenolides **59** and *ent*-**59**. Of critical importance, these fragment unions were accomplished using equimolar amounts of the two coupling partners. It should be noted that while this strategy has allowed us to access diterpenoids such as (+)-cheloviolenes A (**7**) and B (**8**) that bear the 14-carbon hydrophobic fragment on the convex face of the *cis*-2,8-dioxabicyclo[3.3.0]octan-3-one unit, we have so far been unsuccessful in developing a complementary stereoselective approach to diterpenoids such as dendrillolide A (**11**) in which the *cis*-perhydroazulene resides on the concave face.⁴⁰ We

Scheme 15

anticipate that the convergent strategy described in this article, namely late-stage union of a structurally complex tertiary carbon radical with an acceptor, will find applications in future syntheses of a variety of stereochemically elaborate natural products.

EXPERIMENTAL SECTION

Unless stated otherwise, reactions were conducted in oven-dried glassware under an atmosphere of nitrogen or argon. Tetrahydrofuran (THF), 1,2-dimethoxyethane (DME), dimethylformamide (DMF), toluene, dichloromethane, methanol (MeOH), *N,N*-diisopropylethylamine (*i*-Pr₂NEt), and triethylamine (Et₃N) were dried by passage through activated alumina. Methyl bromoacetate and *tert*-butyl bromoacetate were distilled under reduced pressure and stored in a Schlenk flask. All other commercial reagents were used as received unless otherwise noted. Reaction temperatures were controlled using a temperature modulator, and unless stated otherwise, reactions were performed at room temperature (rt, approximately 23 °C). Thin-layer chromatography (TLC) was conducted with silica gel 60 F254 precoated plates, (0.25 mm) and visualized by exposure to UV light (254 nm) or by *p*-anisaldehyde, ceric ammonium molybdate, and potassium permanganate staining (KMnO₄). Silica gel 60 (particle size 0.040–0.063 mm) was used for flash column chromatography. ¹H NMR spectra were recorded at 500 or 600 MHz and are reported relative to deuterated solvent signals. Data for ¹H NMR spectra are reported as follows: chemical shift (δ ppm), multiplicity, coupling constant (Hz), and integration. ¹³C NMR spectra were recorded at 126 or 151 MHz. Data for ¹³C NMR spectra are reported in terms of chemical shift. IR spectra were recorded on a FT-IR spectrometer and are reported in terms of frequency of absorption (cm^{−1}). High-resolution mass spectra were obtained with an LCT spectrometer. Optical rotation readings were obtained using JASCO P-1010 polarimeter. Kessil KSH150B LED Grow Light 150, Blue (34 W blue LED lamps) was purchased from <http://www.amazon.com>. Low-intensity blue LEDs (30 cm, 1 W) were purchased from <http://www.creativelightings.com> (product code CL-FRS5050–12WP-12 V) and were powered by 8 AA batteries. See JOC Standard Abbreviations and Acronyms for abbreviations (available at <http://pubs.acs.org/paragonplus/submission/joc/abbreviations.pdf>).

Preparation of Ester 26. A round-bottom flask was charged with 4-methyl 2-methylenesuccinate (**S1**) (3.0 g, 21 mmol, 1.0 equiv),³⁴ DMF (15 mL, 2.0 M), and a magnetic stir bar under an atmosphere of argon. Next, sodium bicarbonate (3.53 g, 42.0 mmol, 2.0 equiv) was added portionwise to the solution and the resulting heterogeneous mixture was stirred at rt for 1 h. After 1 h, allyl bromide (2.7 mL, 31 mmol, 1.5 equiv) was added dropwise to the mixture that was then stirred vigorously at rt for 12 h. After 12 h, H₂O (50 mL) was added to the reaction mixture and the resulting solution was transferred to a separatory funnel and extracted with CH₂Cl₂ (2 × 50 mL). The combined organic layers were washed with H₂O (3 × 50 mL), dried over MgSO₄ and concentrated by use of a rotary evaporator. The residue was purified by flash chromatography on silica gel using 10:90 ethyl acetate:hexanes to yield ester **26** as a clear oil (3.0 g, 16 mmol,

78% yield): R_f 0.35 (10:90 ethyl acetate:hexanes); ¹H NMR (500 MHz, CDCl₃) δ 6.38 (s, 1H), 5.94 (dddd, *J* = 22.4, 16.2, 10.7, 5.4 Hz, 1H), 5.75 (s, 1H), 5.34 (d, *J* = 17.2 Hz, 1H), 5.26 (d, *J* = 10.5 Hz, 1H), 4.68 (d, *J* = 5.5 Hz, 2H), 3.71 (s, 3H), 3.37 (s, 2H); ¹³C NMR (126 MHz, CDCl₃) δ 171.3, 165.9, 133.8, 132.0, 128.9, 118.3, 65.8, 52.2, 37.7; IR (thin film) 1744, 1642 cm^{−1}; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₉H₁₂O₄Na 207.0633; Found 207.0637.

Preparation of Butenolide 27. A round-bottom flask was charged with diene **26** (800 mg, 4.3 mmol, 1.0 equiv), toluene (440 mL, 0.01 M), and a magnetic stir bar under an atmosphere of argon. Next, Stewart-Grubbs catalyst (50 mg, 0.09 mmol, 0.02 equiv)³⁵ was added to the reaction mixture and the resulting solution was heated to 110 °C for 18 h. After 18 h, the solution was allowed to cool to rt and concentrated by use of a rotary evaporator to yield a brown oil. The residue was purified by flash column chromatography on silica gel using 20:80 ethyl acetate:hexanes → 40:60 ethyl acetate:hexanes as eluent to yield butenolide **27** as a brown oil (500 mg, 3.2 mmol, 74% yield): R_f 0.29 (50:50 ethyl acetate:hexanes, stained with KMnO₄); ¹H NMR (600 MHz, CDCl₃) δ 7.52 (s, 1H), 4.87 (s, 2H), 3.75 (s, 3H), 3.38 (s, 2H); ¹³C NMR (151 MHz, CDCl₃) δ 173.7, 170.1, 148.1, 127.1, 70.8, 52.5, 30.5; IR (thin film) 2955, 2871, 1737, 1657, 1438, 1349, 1223, 1078, 1049 cm^{−1}; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₇H₈O₄Na 179.0320; Found 179.0312.

Preparation of Butenolide 28. A round-bottom flask was charged with butenolide **27** (490 mg, 3.1 mmol, 1.0 equiv), CCl₄ (21 mL, 0.15 M), *N*-bromosuccinimide (1.7 g, 9.4 mmol, 3.0 equiv), AIBN (52 mg, 0.31 mmol, 0.1 equiv), and a magnetic stir bar under an atmosphere of argon. The mixture was irradiated with a 100 W compact fluorescent light bulb and heated to 75 °C for 3 h. After 3 h, additional *N*-bromosuccinimide (560 mg, 3.1 mmol, 1.0 equiv) and AIBN (52 mg, 0.31 mmol, 0.1 mmol) were added. The mixture was stirred at 75 °C for 1 h, at which point TLC analysis indicated complete consumption of starting material **27**. The reaction was then cooled to 0 °C, vacuum filtered and concentrated by use of a rotary evaporator to yield the crude 5-bromo analogue as a yellow oil. Diagnostic ¹H NMR shifts (600 MHz, CDCl₃) δ 7.59 (d, *J* = 1.2 Hz, 1H), 6.90 (d, *J* = 1.2 Hz, 1H), 3.76 (s, 3H), 3.44 (br s, 2H). A round-bottom flask was charged with the crude intermediate, MeOH (7 mL, 0.45 M), and a magnetic stir bar under an atmosphere of argon. The solution was heated to 70 °C for 12 h. After 12 h the solution was allowed to cool to rt and concentrated by use of a rotary evaporator. The residue was purified by flash chromatography on silica gel using 10:90 ethyl acetate:hexanes → 25:75 ethyl acetate:hexanes as eluent to yield butenolide **28** as a yellow oil (210 mg, 1.1 mmol, 36% yield over 2 steps): R_f 0.28 (30:70 ethyl acetate:hexanes); ¹H NMR (600 MHz, CDCl₃) δ 7.16 (d, *J* = 1.2 Hz, 1H), 5.83 (d, *J* = 1.2 Hz, 1H), 3.74 (s, 3H), 3.57 (s, 3H), 3.38 (d, *J* = 1.2 Hz, 2H); ¹³C NMR (151 MHz, CDCl₃) δ 170.7, 169.5, 145.6, 131.2, 103.0, 57.1, 52.6, 30.4; IR (thin film) 1780, 1750 cm^{−1}; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₈H₁₀O₅Na 209.0426; Found 209.0418.

Preparation of Carboxylic Acid 52. A round-bottom flask was charged with *i*-Pr₂NH (9.0 mL, 68 mmol, 2.6 equiv), THF (200 mL, 0.13 M), and a magnetic stir bar under an atmosphere of argon. After cooling the solution to −78 °C, 2.4 M *n*-BuLi in hexanes (27 mL, 65

mmol, 2.5 equiv) was added dropwise. The resulting solution was then warmed to 0 °C stirred for 30 min. Next, isobutyric acid (2.4 mL, 26 mmol, 1.0 equiv) was added dropwise at 0 °C. The reaction was maintained at 0 °C for 20 min, followed by a dropwise addition of a solution of 1-iodo-3,6,9,12-tetraoxapentadec-14-yne (10.2 g, 28.6 mmol, 1.1 equiv)⁸⁰ in THF (60 mL, 0.43 M). The resulting heterogeneous mixture was allowed to warm to rt and stirred vigorously for 12 h. After 12 h, the reaction was quenched via addition of H₂O (50 mL). The resulting biphasic mixture was transferred to a separatory funnel and extracted with CH₂Cl₂ (2 × 200 mL). The aqueous layer was acidified with sat. NH₄Cl (aq) and extracted with CH₂Cl₂ (2 × 300 mL). The combined organic layers, resulting from extracting the acidified aqueous layer, were dried over MgSO₄ and concentrated by use of a rotary evaporator to yield **S2** as a clear oil (3.8 g, 13 mmol, 48% yield); ¹H NMR (500 MHz, CDCl₃) δ 4.21 (d, *J* = 2.3 Hz, 2H), 3.73–3.56 (m, 14H), 2.43 (br s, 1H), 1.87 (t, *J* = 5.9 Hz, 2H), 1.23 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ 181.5, 79.7, 74.7, 70.9, 70.6, 70.4, 70.3, 70.2, 69.2, 67.8, 58.5, 40.4, 40.0, 25.5; IR (thin film) 3251, 1728, 1700, 1103 cm⁻¹; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₅H₂₆O₆Na 325.1627; Found 325.1636.

Preparation of NHP Ester 29c. A round-bottom flask was charged with acid **S2** (1.9 g, 7.4 mmol, 1.0 equiv), CH₂Cl₂ (50 mL, 0.2 M), DCC (2.0 g, 9.6 mmol, 1.3 equiv), DMAP (180 mg, 1.5 mmol, 0.2 equiv), and a magnetic stir bar under an atmosphere of argon. The resulting heterogeneous mixture was stirred at rt for 15 min. Next, *N*-hydroxyphthalimide (1.4 g, 8.8 mmol, 1.2 equiv) was added in one portion and the mixture was stirred at rt for 16 h. After 16 h, the reaction was quenched via addition of sat. NH₄Cl (aq). The resulting biphasic mixture was transferred to a separatory funnel and extracted with CH₂Cl₂ (3 × 50 mL). The combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The residue was purified by flash chromatography on silica gel using 50:50 ethyl acetate:hexanes as eluent to yield *N*-acyloxyphthalimide **29c** as a clear oil (1.6 g, 4.0 mmol, 54% yield): *R*_f = 0.35 (50:50 ethyl acetate:hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.88 (dd, *J* = 5.6, 4.7 Hz, 2H), 7.79 (dd, *J* = 5.4, 3.2 Hz, 2H), 4.20 (d, *J* = 2.4 Hz, 2H), 3.71–3.61 (m, 14H), 2.43–2.42 (m, 1H), 2.06 (t, *J* = 7.0 Hz, 2H), 1.44 (s, 6H); ¹³C NMR (126 MHz, CDCl₃) δ 173.8, 162.2, 134.8, 129.2, 124.0, 79.8, 74.6, 70.8, 70.7, 70.5, 70.4, 69.2, 67.7, 58.5, 40.9, 39.7, 25.6; IR (thin film) 3273, 1782, 1743 cm⁻¹; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₂₃H₂₉NO₈Na 470.1791; Found 470.1777.

General Procedure for the Photoredox-mediated Couplings of 29a–c with 28. A 1-dram scintillation vial was charged with *N*-acyloxyphthalimide **29a–c** (1.0 equiv), butenolide **28** (1.0 equiv), Hantzsch ester (1.5 equiv),⁸¹ [Ru(bpy)₃](PF₆)₂ (0.01 equiv), followed by either *i*-Pr₂NEt (1.0 equiv) or *i*-Pr₂NEt·HBF₄ (2.2 equiv),⁸² and a magnetic stir bar. The vial was sealed with a screw cap bearing a Teflon septum and CH₂Cl₂ (0.15 M, sparged with argon for 10 min) was added. Next, the vial was placed in the center of a 30 cm-loop of low-intensity blue LEDs. The heterogeneous reaction mixture was irradiated by low intensity blue LEDs and stirred vigorously at rt for 18 h. The mixture was filtered over silica gel using Et₂O as eluent and concentrated by use of a rotary evaporator to yield a yellow residue that was further purified by flash chromatography on silica gel.

Preparation of Lactone 30a. Following the general procedure, *N*-acyloxyphthalimide **29a** (27 mg, 0.11 mmol, 1.0 equiv) and butenolide **28** (20 mg, 0.11 mmol, 1.0 equiv) were coupled in the presence of *i*-Pr₂NEt (19 μL, 0.11 mmol, 1.0 equiv), [Ru(bpy)₃](PF₆)₂ (1 mg, 0.001 mmol, 0.01 equiv), and Hantzsch ester (41 mg, 0.16 mmol, 1.5 equiv) in CH₂Cl₂ (0.75 mL). Purification of the crude residue by flash chromatography on silica gel using 10:90 ethyl acetate:hexanes as eluent yielded **30a** as a clear oil (19 mg, 0.076 mmol, 72% yield): *R*_f = 0.45 (10:90 ethyl acetate:hexanes, stained with KMnO₄); ¹H NMR (500 MHz, CDCl₃) δ 5.17 (d, *J* = 2.3 Hz, 1H), 3.71 (s, 3H), 3.51 (s, 3H), 2.85–2.73 (m, 3H), 1.95 (app s, *J* = 1.9 Hz, 1H), 0.94 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 177.8, 171.3, 106.5, 57.2, 56.6, 52.1, 38.7, 36.9, 31.8, 27.1; IR (thin film) 1644, 1633 cm⁻¹; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₂H₂₀O₅Na 267.1208; Found 267.1209.

The reaction performed in the presence of *i*-Pr₂NEt·HBF₄ (53 mg, 0.24 mmol, 2.2 equiv) in place of *i*-Pr₂NEt led to product **30a** (17 mg, 0.69 mmol, 66% yield) that was isolated after purification by flash chromatography on silica gel. The product ratios shown in Scheme 3 arise from ¹H and ¹³C NMR analysis of crude reaction mixtures.

Lactone **31** was isolated in 22% yield from the above reaction when *i*-Pr₂NEt·HBF₄ was used: *R*_f = 0.47 (10:90 ethyl acetate:hexanes, stained with KMnO₄); ¹H NMR (500 MHz, CDCl₃) δ 5.27 (s, 1H), 3.74 (s, 3H), 3.63–3.59 (m, 1H), 3.48 (s, 3H), 2.97 (dd, *J* = 17.5, 6.1 Hz, 1H), 2.74 (dd, *J* = 17.5, 9.3 Hz, 1H), 2.36 (d, *J* = 8.2 Hz, 1H), 0.97 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 177.8, 172.0, 105.9, 56.6, 54.2, 52.3, 38.3, 32.3, 32.0, 28.3; IR (thin film) 2956, 2922, 1783, 1742 cm⁻¹; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₂H₂₀O₅Na 267.1208; Found 267.1201.

Lactone **32** was isolated in 1% yield as a 1.5:1 mixture of diastereomers from the above reaction when *i*-Pr₂NEt was used: *R*_f = 0.42 (10:90 ethyl acetate:hexanes, stained with KMnO₄); ¹H NMR (600 MHz, CDCl₃) δ 5.40 (app t, *J* = 5.8 Hz, 1H), 5.36 (dd, *J* = 6.9, 5.2 Hz, 1H), 3.73 (s, 3H), 3.66 (s, 3H), 3.57 (s, 3H), 3.54 (s, 3H), 3.11 (ddd, *J* = 11.8, 10.5, 9.3 Hz, 1H), 3.01 (d, *J* = 3.1 Hz, 1H), 2.88 (dd, *J* = 9.5, 3.0 Hz, 1H), 2.69 (ddd, *J* = 13.9, 9.0, 5.2 Hz, 1H), 2.57 (ddd, *J* = 13.7, 10.4, 6.2 Hz, 1H), 2.45 (d, *J* = 10.3 Hz, 1H), 2.38 (ddd, *J* = 14.6, 9.3, 5.7 Hz, 1H), 2.00 (app dt, *J* = 12.0, 6.9 Hz, 1H), 1.06 (s, 9H), 1.03 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 177.0, 175.2, 173.8, 172.6, 104.5, 103.9, 58.1, 57.7, 55.8, 54.1, 51.6, 51.5, 41.3, 39.8, 35.9, 33.1, 32.3, 28.6, 28.5, 15.4. IR (thin film) 1770, 1644 cm⁻¹; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₂H₂₀O₅Na 267.1208; Found 267.1199.

Preparation of Lactone 30b. Synthesized according to the general procedure from the *N*-acyloxyphthalimide **29b** (77 mg, 0.27 mmol, 1.0 equiv),⁴¹ butenolide **28** (50 mg, 0.27 mmol, 1.0 equiv), *i*-Pr₂NEt (47 μL, 0.27 mmol, 1.0 equiv), [Ru(bpy)₃](PF₆)₂ (2 mg, 0.003 mmol, 0.01 equiv) and Hantzsch ester (100 mg, 0.4 mmol, 1.5 equiv) in CH₂Cl₂ (1.8 mL). Purification of the crude residue by flash chromatography on silica gel using 10:90 ethyl acetate:hexanes as eluent yielded **30b** (41 mg, 0.14 mmol, 53% yield) as a colorless solid: *R*_f = 0.61 (30:70 diethyl ether:pentanes, stained with KMnO₄); ¹H NMR (500 MHz, CDCl₃) δ 5.21 (d, *J* = 2.1 Hz, 1H), 3.71 (s, 3H), 3.50 (s, 3H), 2.89–2.72 (m, 3H), 2.06 (br s, 1H), 1.60–1.21 (m, 10H), 0.89 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 178.0, 171.4, 106.0, 57.2, 52.2, 37.9, 37.0, 35.43, 35.38, 34.4, 26.1, 21.53, 21.47; IR (thin film) 2929, 2855, 1777, 1738 cm⁻¹; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₅H₂₄O₅Na 307.1521; Found 307.1523.

Preparation of Lactone 30c. Synthesized according to the general procedure from *N*-acyloxyphthalimide **29c** (30 mg, 0.07 mmol, 1.0 equiv), butenolide **28** (13 mg, 0.07 mmol, 1.0 equiv), *i*-Pr₂NEt (12 μL, 0.07 mmol, 1.0 equiv), [Ru(bpy)₃](PF₆)₂ (0.6 mg, 0.0007 mmol, 0.01 equiv) and Hantzsch ester (25 mg, 0.10 mmol, 1.5 equiv) in CH₂Cl₂ (0.45 mL). Purification of the crude residue by flash chromatography on silica gel using 10:90 acetone:hexanes as eluent yielded **30c** (16 mg, 0.036 mmol, 52% yield) as a clear oil: *R*_f = 0.40 (70:30 ethyl acetate:hexanes, stained with KMnO₄); ¹H NMR (600 MHz, CDCl₃) δ 5.20 (d, *J* = 2.3 Hz, 1H), 4.21 (d, *J* = 2.4 Hz, 2H), 3.76–3.45 (m, 20H), 2.89–2.74 (m, 3H), 2.43 (app t, *J* = 2.4 Hz, 1H), 2.06 (dd, *J* = 4.5, 2.3 Hz, 1H), 1.58 (app t, *J* = 7.0 Hz, 2H), 0.94 (s, 3H), 0.93 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 177.8, 171.3, 106.2, 79.8, 74.7, 70.8, 70.7, 70.6, 70.4, 69.2, 67.6, 58.6, 57.2, 55.8, 52.2, 39.1, 38.4, 36.9, 33.7, 24.6, 24.2; IR (thin film) 3267, 2932, 2873, 1775, 1739, 1440, 1364, 1111, 939 cm⁻¹; HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₂₂H₃₆O₉Na 467.2257; Found 467.2248.

General Procedure for the Reduction/Cyclization/Oxidation to Generate Dioxabicyclo[3.3.0]octanones 34a–c. A round-bottom flask was charged with the coupled product **30a–c** (1.0 equiv), toluene (0.1 M), and a magnetic stir bar under an atmosphere of argon. After cooling the solution to –78 °C, *i*-Bu₂AlH (2.1 equiv, 1 M in toluene) was added dropwise. The solution was maintained at –78 °C for 30 min. The reaction was quenched by the addition of saturated solution of Rochelle's salt (aq) (0.1 M) at –78 °C. The mixture was allowed to warm to rt and stirred at rt for 1 h. The biphasic mixture was transferred to a separatory funnel and extracted with CH₂Cl₂ (3 × 10

mL). The combined organic layers were dried over Na_2SO_4 and concentrated by use of a rotary evaporator to yield a crude mixture of bicyclic lactol epimers **33a–c**. A 1-dram scintillation vial was charged with this crude mixture of bicyclic lactol epimers **33a–c** (1.0 equiv), toluene (0.1 M), Ag_2CO_3 (50 wt % on Celite, 3.0 equiv), and a magnetic stir bar under an atmosphere of argon. The reaction vessel was capped and heated to 110 °C. After 1 h, the black suspension was allowed to cool to rt, filtered over Celite and concentrated by use of a rotary evaporator. The resulting residue was purified by flash column chromatography on silica gel to yield bicyclic lactones **34a–c**.

Preparation of Dioxabicyclo[3.3.0]octan-3-one 34a. Synthesized according to the general procedure described above from **30a** (50 mg, 0.21 mmol, 1.0 equiv) and 1 M solution of *i*-Bu₂AlH in toluene (430 μL , 0.43 mmol, 2.2 equiv) in toluene (2 mL, 0.1 M). Diagnostic data for lactol intermediate **33a**: ^1H NMR (500 MHz, CDCl_3) δ 5.68 (d, J = 12.1 Hz, 1H, OH confirmed by D_2O exchange); MS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{11}\text{H}_{20}\text{O}_4\text{Na}$ 239.1; Found 239.1. Conversion of lactol **33a** to lactone **34a** was achieved with Ag_2CO_3 (340 mg, 0.62 mmol, 3.0 equiv) in toluene (2 mL, 0.1 M). Purification of the crude residue by flash chromatography on silica gel using 20:80 ethyl acetate:hexanes as eluent yielded **34a** (28 mg, 0.13 mmol, 64% yield; 73% based on recovered starting material) as a colorless solid: R_f 0.45 (30:70 ethyl acetate:hexanes, stained with ceric ammonium molybdate); ^1H NMR (500 MHz, CDCl_3) δ 6.05 (d, J = 6.1 Hz, 1H), 5.04 (s, 1H), 3.36 (s, 3H), 2.97–2.93 (m, 1H), 2.86 (dd, J = 18.1, 11.2 Hz, 1H), 2.62 (dd, J = 18.1, 3.6 Hz, 1H), 1.95 (d, J = 1.4 Hz, 1H) 0.93 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ 175.6, 109.1, 108.7, 64.4, 55.0, 39.2, 36.6, 31.5, 27.5; IR (thin film) 1784 cm^{-1} ; HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{11}\text{H}_{18}\text{O}_4\text{Na}$ 237.1103; Found 237.1110.

Preparation of Dioxabicyclo[3.3.0]octan-3-one 34b. Synthesized according to the general procedure described above from **30b** (33 mg, 0.12 mmol, 1.0 equiv) and 1 M solution of *i*-Bu₂AlH in toluene (240 μL , 0.24 mmol, 2.2 equiv) in toluene (1.2 mL, 0.1 M). Diagnostic data for lactol intermediate **33b**: ^1H NMR (500 MHz, CDCl_3) δ 5.73 (d, J = 12.0 Hz, 1H, OH confirmed by D_2O exchange). Conversion of lactol **33b** to lactone **34b** was achieved with Ag_2CO_3 (190 mg, 0.35 mmol, 3.0 equiv) in toluene (1.2 mL, 0.1 M). Purification of the crude residue by flash chromatography on silica gel using 15:85 ethyl acetate:hexanes as eluent yielded **34b** (18 mg, 0.071 mmol, 64% yield) as a colorless solid: R_f 0.45 (30:70 ethyl acetate:hexanes, stained with ceric ammonium molybdate); ^1H NMR (500 MHz, CDCl_3) δ 6.04 (d, J = 6.1 Hz, 1H), 5.07 (s, 1H), 3.35 (s, 3H), 3.01–2.95 (m, 1H), 2.86 (dd, J = 18.1, 11.0 Hz, 1H), 2.61 (dd, J = 18.1, 3.5 Hz, 1H), 2.04 (br s, 1H), 1.56–1.19 (m, 10H), 0.84 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ 175.6, 109.2, 108.3, 55.0, 38.4, 36.8, 35.8, 35.7, 33.9, 26.2, 21.6, 20.8; IR (thin film) 1785 cm^{-1} ; HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{14}\text{H}_{22}\text{O}_4\text{Na}$ 277.1416; Found 277.1417.

Preparation of Dioxabicyclo[3.3.0]octan-3-one 34c. Synthesized according to the general procedure described above from **30c** (94 mg, 0.21 mmol, 1.0 equiv) and 1 M solution of *i*-Bu₂AlH in toluene (460 μL , 0.46 mmol, 2.2 equiv) in toluene (2.1 mL, 0.1 M). Conversion of lactol **33c** to lactone **34c** was achieved with Ag_2CO_3 (340 mg, 0.63 mmol, 3.0 equiv) in toluene (2.1 mL, 0.1 M). Purification of the crude residue by flash chromatography on silica gel using 30:70 ethyl acetate:hexanes as eluent yielded **34c** (35 mg, 0.08 mmol, 40% yield) as a colorless solid: R_f 0.30 (30:70 ethyl acetate:hexanes, stained with ceric ammonium molybdate); ^1H NMR (500 MHz, CDCl_3) δ 6.04 (d, J = 6.1 Hz, 1H), 5.03 (s, 1H), 4.21 (d, J = 2.3 Hz, 2H), 3.72–3.50 (m, 14H), 3.00–2.95 (m, 1H), 2.86 (dd, J = 18.3, 11.2 Hz, 1H), 2.63 (dd, J = 18.3, 3.8 Hz, 1H), 2.43 (t, J = 2.3 Hz, 1H), 2.10 (s, 1H), 1.65–1.48 (m, 2H), 0.95 (s, 3H), 0.87 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ 175.7, 109.0, 108.5, 79.8, 74.7, 70.7, 70.6, 70.5, 69.2, 67.8, 63.1, 58.6, 54.9, 39.9, 39.1, 36.6, 33.4, 25.2, 24.9; IR (thin film) 1781 cm^{-1} ; HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{21}\text{H}_{34}\text{O}_8\text{Na}$ 437.2151; Found 437.2149.

Preparation of Lactone 36. Synthesized according to the general fragment coupling procedure from *N*-acyloxypthalimide **29a** (150 mg, 0.6 mmol, 1.0 equiv), butenolide **35** (69 mg, 0.60 mmol, 1.0

equiv), *i*-Pr₂NEt (100 μL , 0.6 mmol, 1.0 equiv), $[\text{Ru}(\text{bpy})_3](\text{PF}_6)_2$ (5 mg, 0.006 mmol, 0.01 equiv) and Hantzsch ester (230 mg, 0.90 mmol, 1.5 equiv) in CH_2Cl_2 (4 mL). Purification of the crude residue by flash chromatography on silica gel using 5:95 ethyl acetate:hexanes $\rightarrow$ 10:90 ethyl acetate:hexanes as eluent yielded **36** (75 mg, 0.44 mmol, 73% yield) as a clear oil: R_f 0.27 (10:90 ethyl acetate:hexanes; stained with KMnO_4); ^1H NMR (500 MHz, CDCl_3) δ 5.19 (d, J = 2.5 Hz, 1H), 3.49 (s, 3H), 2.65 (dd, J = 18.5, 10.0 Hz, 1H), 2.35 (dd, J = 18.5, 5.5 Hz, 1H), 2.14 (ddd, J = 10.0, 5.5, 3.0 Hz, 1H), 0.92 (s, 9H); ^{13}C NMR (151 MHz, CDCl_3) δ 176.5, 107.4, 57.0, 51.7, 31.5, 30.0, 27.0; IR (thin film) 2965, 1779, 1175, 1115 cm^{-1} ; HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_9\text{H}_{16}\text{O}_3\text{Na}$ 195.0997; Found 195.0993.

Preparation of Ester 30a. A round-bottom flask was charged with *i*-Pr₂NH (110 μL , 0.75 mmol, 3.8 equiv), THF (4.6 mL, 0.1 M), and a magnetic stir bar under an atmosphere of argon. After cooling the solution to –78 °C, 1.93 M *n*-BuLi in hexanes (260 μL , 0.50 mmol, 2.5 equiv) was added dropwise. The resulting solution was then warmed to 0 °C and stirred for 30 min. Next, an aliquot of the LDA solution (3 mL, 0.3 mmol, 1.5 equiv) was added dropwise to a solution of **36** (34 mg, 0.20 mmol, 1.0 equiv) in THF (2 mL, 0.1 M) at –78 °C. After 1 h at –78 °C, a solution of methyl bromoacetate (45 mg, 0.30 mmol, 1.5 equiv) in THF (0.3 mL) was added dropwise. The reaction was allowed to warm to rt, H_2O (5 mL) was added, and the resulting biphasic mixture was extracted with CH_2Cl_2 (2 $\times$ 10 mL). The combined organic extracts were dried over Na_2SO_4 and concentrated by use of a rotary evaporator. The resulting residue was purified by flash chromatography on silica gel using 5:95 ethyl acetate:hexanes as eluent to yield **30a** (27 mg, 0.11 mmol, 56% yield) as a clear oil: R_f 0.45 (10:90 ethyl acetate:hexanes; stained with KMnO_4).

Preparation of Lactone 37 from 39. A 1-dram scintillation vial was charged with *N*-acyloxypthalimide **39** (25 mg, 0.066 mmol, 1.0 equiv), butenolide (*S*)-**35** (8 mg, 0.07 mmol, 1.0 equiv), Hantzsch ester (25 mg, 0.11 mmol, 1.5 equiv), $[\text{Ru}(\text{bpy})_3](\text{PF}_6)_2$ (0.6 mg, 0.0007 mmol, 0.01 equiv), *i*-Pr₂NEt-HBF₄ (29 mg, 0.13 mmol, 2.2 equiv), and a magnetic stir bar. The vial was sealed with a screw cap bearing a Teflon septum and CH_2Cl_2 (0.1 M, sparged with argon for 10 min) was added. Next, the vial was placed in the center of a 30 cm-loop of low-intensity blue LEDs. Heterogeneous reaction mixture was irradiated by low intensity blue LEDs and stirred vigorously at rt for 18 h. The mixture was filtered over silica gel using Et₂O as eluent and concentrated by use of a rotary evaporator. The resulting yellow residue was purified by flash chromatography on silica gel using 5:95 ethyl acetate:hexanes as eluent to yield **37** (12 mg, 0.039 mmol, 30% yield) as a clear oil: R_f 0.30 (10:90 ethyl acetate:hexanes; stained with KMnO_4); ^1H NMR (500 MHz, CDCl_3) δ 5.21 (d, J = 1.7 Hz, 1H), 4.83 (d, J = 1.7 Hz, 1H), 4.60 (br s, 1H), 3.50 (s, 3H), 2.75 (dd, J = 19.6, 11.2 Hz, 1H), 2.51 (d, J = 8.7 Hz, 1H), 2.46–2.47 (m, 1H), 2.43–2.41 (m, 1H), 2.35 (dd, J = 12.5, 5.2 Hz, 1H), 1.97–1.91 (m, 1H), 1.82–1.73 (m, 4H), 1.64 (dd, J = 13.9, 4.1 Hz, 1H), 1.61–1.56 (m, 2H), 1.45–1.34 (m, 1H), 1.28–1.25 (m, 1H), 0.98 (s, 3H), 0.94 (s, 3H), 0.83 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ 176.8, 153.5, 114.9, 107.6, 66.1, 57.0, 56.2, 54.3, 52.0, 47.1, 37.9, 37.7, 37.0, 36.4, 34.6, 28.9, 26.2, 25.9, 20.9, 15.5; IR (thin film) 1787 cm^{-1} ; $[\alpha]_D^{23}$ +90.8, $[\alpha]_D^{23}$ 377 + 93.2, $[\alpha]_D^{23}$ 346 + 106, $[\alpha]_D^{23}$ 435 + 181 (c = 1.0, CHCl_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{19}\text{H}_{30}\text{O}_3\text{Na}$ 329.2093; Found 329.2099.

Preparation of Lactone 37 from 40. A 1-dram scintillation vial was charged with **40** (47 mg, 0.2 mmol, 1.0 equiv), K_2HPO_4 (38 mg, 0.22 mmol, 1.1 equiv), $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$ (5 mg, 0.004 mmol, 0.02 equiv), 3:1 DMF: CH_3CN (0.5 mL, 0.4 M), H_2O (36 μL , 2.0 mmol, 10 equiv), (*S*)-**35** (23 mg, 0.2 mmol, 1.0 equiv), and a magnetic stir bar. The vial was then sealed with a screw cap bearing Teflon septum. The septum of the vial was pierced with a 21 gauge $\times$ 1.5" needle that was inserted just barely through the septum with the tip of the needle kept above the fluid level inside the vial. A separate 22 gauge $\times$ 3" needle attached to a flow of argon was also pierced through the septum, and the tip of the needle was pushed to the bottom of the vial and submersed in the fluid. The reaction mixture was degassed by sparging with argon for 15 min. Both needles were removed, and the sealed vial was then placed on a stir plate equipped with 2 $\times$ 34 W blue

LED lamps and a rack to hold the vial inside of a cardboard box to block light pollution from entering the lab. The vial was placed approximately 4 cm from the lamps and stirred vigorously. The sample was irradiated by the lamps for 18 h inside the closed box, allowing the temperature of the reaction mixture to rise to 60 °C and the air inside the box to 40–45 °C because of heat given off from the LEDs. The reaction was allowed to cool to rt and transferred to a separatory funnel and extracted with Et₂O (3 × 5 mL). The combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The residue was purified by flash column chromatography on silica gel using 5:95 ethyl acetate:hexanes as eluent to yield **37** (27 mg, 0.088 mmol, 44% yield) as a clear oil.

Preparation of Ester 41. A round-bottom flask was charged with *i*-Pr₂NH (280 μL, 2.0 mmol, 13 equiv), THF (3 mL, 0.05 M), and a magnetic stir bar under an atmosphere of argon. After cooling the solution to –78 °C, 2.3 M *n*-BuLi in hexanes (800 μL, 1.8 mmol, 12 equiv) was added dropwise. The resulting solution was then warmed to 0 °C and stirred for 30 min. Next, an aliquot of the LDA solution (550 μL, 0.22 mmol, 1.5 equiv) was added dropwise to a solution of **37** (46 mg, 0.15 mmol, 1.0 equiv) in THF (1.5 mL, 0.1 M) at –78 °C. After 1 h at –78 °C, a solution of methyl iodoacetate (60 mg, 0.3 mmol, 2.0 equiv) in THF (0.3 mL) was added dropwise. The reaction was allowed to warm to rt, H₂O (5 mL) was added, and the resulting biphasic mixture was extracted with CH₂Cl₂ (2 × 50 mL). The combined organic extracts were dried over Na₂SO₄ and concentrated by use of a rotary evaporator. The resulting residue was purified by flash chromatography on silica gel using 5:95 ethyl acetate:hexanes as eluent to yield **41** (26 mg, 0.069 mmol, 45% yield) as a clear oil: R_f 0.30 (10:90 ethyl acetate:hexanes; stained with KMnO₄); ¹H NMR (500 MHz, CDCl₃) δ 5.20 (s, 1H), 4.84 (d, *J* = 1.8 Hz, 1H), 4.70 (br s, 1H), 3.74 (s, 3H), 3.50 (s, 3H), 2.94–2.90 (m, 1H), 2.81–2.78 (m, 2H), 2.63 (d, *J* = 8.5 Hz, 1H), 2.34 (dd, *J* = 12.8, 5.7 Hz, 1H), 2.19 (br d, *J* = 2.1 Hz, 1H), 1.95–1.89 (m, 1H), 1.83–1.70 (m, 3H), 1.65 (dd, *J* = 13.9, 4.0 Hz, 1H), 1.61–1.58 (m, 2H), 1.43–1.35 (m, 1H), 1.27–1.25 (m, 2H), 1.00 (s, 3H), 0.94 (s, 3H), 0.82 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 178.5, 171.3, 153.4, 115.0, 107.2, 57.1, 57.0, 54.9, 54.1, 52.3, 47.4, 39.6, 37.82, 37.79, 37.0, 36.4, 34.5, 29.0, 26.1, 25.8, 21.2; IR (thin film) 1781, 1742 cm^{–1}; [α]_D²³ +73.0, [α]_D²³₅₇₇ +76.4, [α]_D²³₅₄₆ +86.1, [α]_D²³₄₃₅ +146 (*c* = 1.0, CHCl₃); HRMS (ESI-TOF) *m/z*: [M+Na]⁺ Calcd for C₂₂H₃₄O₅Na 401.2304; Found 401.2295.

Preparation of Dioxabicyclo[3.3.0]octan-3-one 42. A round-bottom flask was charged with **41** (4 mg, 0.01 mmol, 1.0 equiv), toluene (0.1 mL, 0.1 M), and a magnetic stir bar under an atmosphere of argon. After cooling the solution to –78 °C, 1 M solution of *i*-Bu₂AlH in toluene (22 μL, 0.022 mmol, 2.2 equiv) was added dropwise. The solution was maintained at –78 °C for 30 min. The reaction was quenched by the addition of saturated solution of Rochelle's salt (aq) (0.1 mL) at –78 °C. The mixture was allowed to warm to rt and stirred at rt for 1 h. Biphasic mixture was transferred to a separatory funnel and extracted with CH₂Cl₂ (3 × 10 mL). The combined organic layers were dried over Na₂SO₄ and concentrated by use of a rotary evaporator to yield a crude mixture of bicyclic lactol epimers **S3**. Diagnostic chemical shifts of the lactol intermediate **S3**: ¹H NMR (600 MHz, CDCl₃) δ 5.86 (d, *J* = 6.0 Hz, 1H), 5.75 (d, *J* = 12.0 Hz, 1H), 5.47 (dd, *J* = 12.0, 6.0 Hz, 1H), 4.96 (s, 1H), 4.85 (d, *J* = 2.4 Hz, 1H), 4.64 (d, *J* = 1.8 Hz, 1H), 3.50 (s, 3H).

A 1-dram scintillation vial was charged with crude mixture of bicyclic lactol epimers **S3**, toluene (0.1 mL, 0.1 M), Ag₂CO₃ (50 wt % on Celite, 17 mg, 0.03 mmol, 3.0 equiv), and a magnetic stir bar under an atmosphere of argon. The reaction vessel was capped and heated to 110 °C. After 1 h, the black suspension was allowed to cool to rt, filtered over Celite and concentrated by use of a rotary evaporator. The resulting residue was purified by flash column chromatography on silica gel using 10:90 ethyl acetate:hexanes as eluent to yield lactone **42** (3.5 mg, 0.01 mmol, 80% yield) as a clear oil: R_f 0.30 (10:90 ethyl acetate:hexanes; stained with KMnO₄); ¹H NMR (600 MHz, CDCl₃) δ 6.06 (d, *J* = 6.6 Hz, 1H), 4.98 (s, 1H), 4.84 (d, *J* = 1.8 Hz, 1H), 4.62 (br s, 1H), 3.36 (s, 3H), 3.07–3.05 (m, 1H), 2.90 (dd, *J* = 18.6, 11.4 Hz, 1H), 2.66 (dd, *J* = 18.6, 3.6 Hz, 1H), 2.53 (d, *J* = 9.0 Hz, 1H), 2.37–2.34 (m, 1H), 2.22 (s, 1H), 1.96–1.91 (m, 1H), 1.85–1.80 (m,

1H), 1.80–1.71 (m, 3H), 1.63 (dt, *J* = 13.8, 4.2 Hz, 1H), 1.59–1.56 (m, 2H), 1.44–1.38 (m, 1H), 1.31–1.23 (m, 1H), 1.00 (s, 3H), 0.95 (s, 3H), 0.80 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 175.5, 154.1, 114.6, 109.4, 109.3, 66.0, 56.6, 55.0, 54.5, 47.0, 39.9, 38.7, 37.8, 37.0, 36.8, 36.3, 34.6, 28.9, 26.4, 25.8, 21.2; IR (thin film) 2929, 2865, 1787, 1175, 1074, 1003, 932 cm^{–1}; [α]_D²³ +73.9, [α]_D²³₅₇₇ +77.2, [α]_D²³₅₄₆ +89.4, [α]_D²³₄₃₅ +153 (*c* = 1.0, CHCl₃); HRMS (ESI-TOF) *m/z*: [M+Na]⁺ Calcd for C₂₁H₃₂O₄Na 371.2198; Found 371.2193.

Preparation of (+)-Cheloviolene A (7). A 1-dram scintillation vial was charged with lactone **42** (13 mg, 0.036 mmol, 1.0 equiv), 1:1 N HCl (aq):THF (1.4 mL, 0.025 M), and a magnetic stir bar under ambient atmosphere. The resulting biphasic mixture was stirred vigorously at 40 °C for 12 h. The reaction mixture was transferred to a separatory funnel and extracted with CH₂Cl₂ (3 × 10 mL). The combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The residue was purified by flash column chromatography on silica gel using 20:80 ethyl acetate:hexanes as eluent to yield (+)-cheloviolene A (**7**) as a colorless solid (9 mg, 0.026 mmol, 70% yield): R_f 0.30 (20:80 ethyl acetate:hexanes, stained with KMnO₄). Recrystallization of the solid from hexanes:ethyl acetate afforded (+)-cheloviolene A (**6**) as a colorless crystalline solid. The ¹H and ¹³C NMR data in CDCl₃ and IR data were identical to that reported earlier for synthetic (+)-cheloviolene A.³¹ [α]_D²¹ +49.1, [α]_D²¹₅₇₇ +52.7, [α]_D²¹₅₄₆ +56.2, [α]_D²¹₄₃₅ +77.6 (*c* = 0.11, CHCl₃); mp: 155–156 °C (recrystallized from hexanes:ethyl acetate).

Preparation of Methyl Oxalate S4. A round-bottom flask was charged with **43** (146 mg, 0.694 mmol, 1.0 equiv), CH₂Cl₂ (7 mL, 0.1 M), DMAP (170 mg, 1.4 mmol, 2.0 equiv), Et₃N (210 mg, 2.1 mmol, 3.0 equiv), methyl chlorooxacetate (190 μL, 2.1 mmol, 3.0 equiv), and a magnetic stir bar under ambient atmosphere. The resulting solution was heated to 35 °C and maintained at that temperature for 30 min. After 30 min, the reaction was allowed to cool to rt, quenched with H₂O, transferred to a separatory funnel, and extracted with CH₂Cl₂ (3 × 10 mL). Combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The resulting residue was purified by flash column chromatography on silica gel using 10:90 ethyl acetate:hexanes as eluent to yield **S4** (181 mg, 0.61 mmol, 88% yield) as a clear oil: R_f 0.12 (10:90 ethyl acetate:hexanes; stained with *p*-anisaldehyde); ¹H NMR (500 MHz, CDCl₃) δ 3.84 (s, 3H), 2.79 (td, *J* = 12.2, 3.2 Hz, 1H), 2.48–2.32 (m, 3H), 2.18 (d, *J* = 12.5 Hz, 1H), 1.98 (ddd, *J* = 15.2, 10.0, 2.9 Hz, 1H), 1.88–1.72 (m, 2H), 1.75 (s, 3H), 1.61–1.43 (m, 3H), 1.44–1.30 (m, 1H), 0.97 (s, 3H), 0.77 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 212.5, 158.5, 156.3, 95.8, 63.2, 53.6, 49.1, 45.1, 44.1, 36.3, 34.3, 30.8, 25.2, 24.0, 20.8, 19.4; IR (thin film) 3021, 2964, 2873, 1742, 1646, 1216, 1161 cm^{–1}; [α]_D²³ –104, [α]_D²³₅₇₇ –113, [α]_D²³₅₄₆ –131, [α]_D²³₄₃₅ –247 (*c* = 0.4, CHCl₃); HRMS (ESI-TOF) *m/z*: [M+Na]⁺ Calcd for C₁₆H₂₄O₅Na 319.1521; Found 319.1522.

Preparation of Lithium Oxalate 44. A round-bottom flask was charged with **S4** (177 mg, 0.597 mmol, 1.0 equiv), 1:1 THF:H₂O (6 mL, 0.1 M), and a magnetic stir bar under ambient atmosphere. After cooling the biphasic mixture to 0 °C, 0.5 N LiOH (aq) (1.1 mL, 0.57 mmol, 0.95 equiv) was added dropwise. The mixture was then vigorously stirred at 0 °C for 5 min. Next, the homogeneous solution was concentrated by use of a rotary evaporator (50 °C, 12 Torr). The resulting colorless solid was washed with pentanes (3 × 5 mL) and dried further under high vacuum (rt, 0.5 Torr) to yield **44** as a colorless solid (158 mg, 0.547 mmol, 97% yield): ¹H NMR (600 MHz, CD₃OD) δ 2.92 (td, *J* = 12.1, 3.3 Hz, 1H), 2.60 (td, *J* = 12.0, 6.8 Hz, 1H), 2.42 (dt, *J* = 15.0, 8.7 Hz, 1H), 2.32 (ddd, *J* = 12.3, 6.9, 2.5 Hz, 1H), 2.14 (dd, *J* = 12.5, 1.9 Hz, 1H), 1.92 (ddd, *J* = 15.0, 10.2, 3.0 Hz, 1H), 1.88–1.78 (m, 2H), 1.70 (s, 3H), 1.63–1.41 (m, 3H), 1.40–1.30 (m, 1H), 0.99 (s, 3H), 0.77 (s, 3H); ¹³C NMR (151 MHz, CD₃OD) δ 215.7, 166.7, 166.1, 94.0, 65.2, 46.1, 45.1, 37.6, 35.1, 31.2, 26.2, 24.5, 22.1, 19.6; IR (thin film) 3513, 2900, 2819, 1694, 1448, 1420, 1050 cm^{–1}; [α]_D²¹ –95.8, [α]_D²¹₅₇₇ –101, [α]_D²¹₅₄₆ –116, [α]_D²¹₄₃₅ –199 (*c* = 0.6, CH₃OH); HRMS (ESI-TOF) *m/z*: [M][–] Calcd for C₁₅H₂₁O₅ 281.1389; Found 281.1388.

Preparation of Lactone 45. A 1-dram scintillation vial was charged with **44** (29 mg, 0.1 mmol, 1.0 equiv), Ir[dF(CF₃)-

ppy]₂(dtbbpy)PF₆ (2 mg, 0.002 mmol, 0.02 equiv), 3:1 DME:DMF (2 mL, 0.05 M), H₂O (18 μ L, 1.0 mmol, 10 equiv), (**R**)-**35** (11 mg, 0.1 mmol, 1.0 equiv), and a magnetic stir bar. The vial was then sealed with a screw cap bearing Teflon septum. The septum of the vial was pierced with a 21 gauge $\times$ 1.5" needle that was inserted just barely through the septum with the tip of the needle kept above the fluid level inside the vial. A separate 22 gauge $\times$ 3" needle attached to a flow of argon was also pierced through the septum, and the tip of the needle was pushed to the bottom of the vial and submersed in the fluid. The reaction mixture was degassed by sparging with argon for 15 min. Both needles were removed, and the sealed vial was then placed on a stir plate equipped with 2 $\times$ 34 W blue LED lamps and a rack to hold the vial inside of a cardboard box to block light pollution from entering the lab. The vial was placed approximately 4 cm from the lamps and stirred vigorously. The sample was irradiated by the lamps for 18 h inside the closed box, allowing the temperature of the reaction mixture to rise to 60 $^{\circ}$ C and the air inside the box to 40–45 $^{\circ}$ C because of heat given off from the LEDs. The reaction was allowed to cool to rt and transferred to a separatory funnel and extracted with Et₂O (3 $\times$ 5 mL). The combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The residue was purified by flash column chromatography on silica gel using 10:90 ethyl acetate:hexanes $\rightarrow$ 15:85 ethyl acetate:hexanes as eluent to yield **45** (17 mg, 0.071 mmol, 71% yield) as a colorless solid: R_f = 0.30 (20:80 ethyl acetate:hexanes; stained with *p*-anisaldehyde); ¹H NMR (600 MHz, CDCl₃) δ 2.13 (d, J = 11.9 Hz, 2H), 1.99–1.85 (m, 3H), 1.72 (dddd, J = 24.7, 12.7, 8.8, 5.6 Hz, 4H), 1.64–1.50 (m, 2H), 1.56 (s, 3H), 1.34 (qd, J = 12.2, 8.3 Hz, 1H), 1.25 (td, J = 13.4, 13.0, 2.6 Hz, 1H), 0.91 (s, 3H), 0.87 (s, 3H); ¹³C NMR (151 MHz, CDCl₃) δ 178.6, 93.1, 79.6, 56.5, 51.0, 46.5, 38.6, 37.5, 35.0, 31.6, 28.0, 27.7, 19.6, 19.0; IR (thin film) 3419, 1754, 1651, 1644, 1289, 1130 cm⁻¹; [α]_D²¹ + 55.6, [α]_D²³ + 52.2, [α]_D²³ + 57.5, [α]_D²³ + 85.6 (c = 0.2, CHCl₃); HRMS (ESI-TOF) m/z : [M+Na]⁺ Calcd for C₁₄H₂₂O₃Na 261.1467; Found 261.1466.

Preparation of Benzoate Ester S5. A 1-dram scintillation flask was charged with **45** (17 mg, 0.71 mmol, 1.0 equiv), CH₂Cl₂ (1.4 mL, 0.05 M), DMAP (9 mg, 0.07 mmol, 1.0 equiv), Et₃N (25 μ L, 0.18 mmol, 2.5 equiv), 4-nitrobenzoyl chloride (16 mg, 0.085 mmol, 1.2 equiv), and a magnetic stir bar under an atmosphere of argon at rt. The flask was capped and heated to 35 $^{\circ}$ C for 18 h. The reaction was quenched with addition of sat. NaHCO₃ (aq). The resulting biphasic mixture was transferred to a separatory funnel and extracted with Et₂O (3 $\times$ 5 mL). The combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The resulting residue was purified by flash column chromatography on silica gel using 10:90 ethyl acetate:hexanes as eluent to yield **S5** (22 mg, 0.057 mmol, 80% yield) as a colorless solid: R_f = 0.27 (10:90 ethyl acetate:hexanes; stained with *p*-anisaldehyde). Recrystallization of the solid from hot hexanes afforded a crystal suitable for single-crystal X-ray diffraction analysis. ¹H NMR (600 MHz, CDCl₃) δ 8.30 (d, J = 8.8 Hz, 2H), 8.17 (d, J = 8.8 Hz, 2H), 2.58 (d, J = 11.9 Hz, 1H), 2.47 (ddd, J = 15.8, 6.0, 3.1 Hz, 1H), 2.11–1.85 (m, 5H), 1.79 (dq, J = 11.2, 5.7, 5.3 Hz, 2H), 1.64–1.57 (m, 1H), 1.56 (s, 3H), 1.41–1.30 (m, 2H), 0.99 (s, 3H), 0.94 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 173.8, 162.7, 151.0, 135.4, 131.1, 123.9, 92.0, 87.2, 55.4, 52.4, 44.2, 40.0, 34.9, 34.3, 30.3, 27.1, 27.0, 21.1, 20.0; IR (thin film) 2958, 2934, 2869, 1771, 1731, 1530, 1348, 1281 cm⁻¹; [α]_D²² + 38.7, [α]_D²² + 40.5, [α]_D²² + 45.4, [α]_D²² + 80.8 (c = 0.8, CHCl₃); HRMS (ESI-TOF) m/z : [M+Na]⁺ Calcd for C₂₁H₂₅NO₆Na 410.1580; Found 410.1599; mp: 135–137 $^{\circ}$ C (recrystallized from hexanes).

Preparation of Hydrazone S6. A 1-dram vial was charged with **43** (10 mg, 0.05 mmol, 1.0 equiv), 1:1 THF:H₂O (0.5 mL, 0.1 M), *p*-toluenesulfonyl hydrazide (28 mg, 0.15 mmol, 3.0 equiv), CSA (3 mg, 0.02 mmol, 0.3 equiv), and a magnetic stir bar under ambient atmosphere at rt. The resulting biphasic mixture was stirred vigorously for 18 h at rt. The mixture was transferred to a separatory funnel, diluted with H₂O (5 mL), and extracted with EtOAc (3 $\times$ 10 mL). The combined organic extracts were dried over Na₂SO₄ and concentrated by use of a rotary evaporator. The resulting residue was purified by flash column chromatography on silica gel using 20:80

ethyl acetate:hexanes as eluent to yield **S6** (18 mg, 0.048 mmol, 95% yield) as a colorless solid: R_f = 0.2 (20:80 ethyl acetate:hexanes; stained with KMnO₄). Recrystallization of the solid from benzene and pentanes via a vapor diffusion method afforded a crystal suitable for single-crystal X-ray diffraction analysis. ¹H NMR (600 MHz, CDCl₃) δ 7.84 (d, J = 8.4 Hz, 2H), 7.34 (d, J = 7.9 Hz, 2H), 2.42 (s, 3H), 2.39–2.34 (m, 1H), 2.32 (d, J = 9.9 Hz, 1H), 2.19 (ddd, J = 18.4, 10.1, 6.1 Hz, 1H), 2.02–1.88 (m, 2H), 1.67–1.52 (m, 4H), 1.51–1.43 (m, 1H), 1.26 (ddd, J = 12.9, 11.4, 9.1 Hz, 1H), 1.13 (s, 3H), 1.07 (ddd, J = 13.7, 9.1, 4.7 Hz, 1H), 0.87 (s, 3H), 0.84 (s, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 165.5, 144.7, 135.1, 130.1, 128.4, 81.2, 52.5, 51.9, 44.6, 40.7, 35.0, 31.9, 31.6, 26.1, 25.3, 21.8, 18.9, 18.8; IR (thin film) 3426, 3224, 2964, 1633, 1598, 1455, 1336, 1166 cm⁻¹; [α]_D²³ – 79.2, [α]_D²³ – 81.1, [α]_D²³ – 105, [α]_D²³ – 183 (c = 1.3, CHCl₃); HRMS (ESI-TOF) m/z : [M+Na]⁺ Calcd for C₂₀H₃₀N₂O₃Na 401.1875; Found 401.1868; mp: 153–156 $^{\circ}$ C (recrystallized benzene and pentanes via a vapor diffusion method).

Preparation of Ester 66. A round-bottom flask was charged with lactone **64** (58 mg, 0.17 mmol, 1.0 equiv),³¹ tetrahydrofuran (1.7 mL, 0.1 M), and a magnetic stir bar under an argon atmosphere. The solution was cooled to –78 $^{\circ}$ C and a 1 M solution of LiHMDS in tetrahydrofuran (210 μ L, 0.21 mmol, 1.2 equiv) was added dropwise. After stirring the reaction mixture for 1 h at –78 $^{\circ}$ C, *tert*-butyl bromoacetate (38 μ L, 0.26 mmol, 1.5 equiv) was added dropwise. After 1 h at –78 $^{\circ}$ C, the reaction was quenched by addition of sat. NH₄Cl (aq) (5 mL). The resulting mixture was allowed to warm to rt and transferred to a separatory funnel and extracted with Et₂O (3 $\times$ 10 mL). The combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The residue was purified by flash column chromatography on silica gel using 5:95 ethyl acetate:hexanes as eluent to yield product **66** as a clear oil (67 mg, 0.15 mmol, 87% yield): R_f = 0.19 (5:95 ethyl acetate:hexanes, stained with *p*-anisaldehyde); ¹H NMR (600 MHz, CDCl₃) δ 5.53 (d, J = 1.9 Hz, 1H), 3.51 (td, J = 10.7, 4.1 Hz, 1H), 2.88 (dt, J = 9.0, 4.8 Hz, 1H), 2.67 (dd, J = 15.4, 5.4 Hz, 1H), 2.59 (dd, J = 15.4, 8.1 Hz, 1H), 2.12 (dtd, J = 22.4, 13.1, 11.4, 5.5 Hz, 2H), 1.99 (t, J = 2.9 Hz, 1H), 1.68–1.60 (m, 2H), 1.60–1.15 (m, 22H), 0.98 (dt, J = 10.4, 6.7 Hz, 1H), 0.92 (d, J = 6.6 Hz, 3H), 0.91–0.79 (m, 8H), 0.76 (d, J = 7.0 Hz, 3H); ¹³C NMR (151 MHz, CDCl₃) δ 178.6, 170.2, 101.0, 81.4, 77.2, 56.4, 47.9, 39.7, 38.9, 38.3, 35.5, 35.3, 34.6, 31.6, 28.2, 26.2, 25.4, 23.0, 22.5, 21.6, 21.5, 21.2, 20.3, 15.5; IR (thin film) 2953, 2929, 1770, 1727, 1369, 1151 cm⁻¹; [α]_D²¹ + 78.5, [α]_D²¹ + 82.1, [α]_D²¹ + 88.5, [α]_D²¹ + 138 (c = 0.27, CHCl₃); HRMS (ESI-TOF) m/z : [M+Na]⁺ Calcd for C₂₇H₄₆O₅Na 473.3243; Found 473.3250.

Preparation of Lactol 67. A round-bottom flask was charged with **66** (64 mg, 0.14 mmol, 1.0 equiv), THF (1.2 mL, 0.1 M), and a magnetic stir bar under an argon atmosphere. The solution was cooled to 78 $^{\circ}$ C and a 1 M solution of DIBALH in toluene (170 μ L, 0.17 mmol, 1.2 equiv) was added dropwise. After 30 min at –78 $^{\circ}$ C, TLC analysis of the reaction (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate) indicated presence of the starting material and additional 1 M solution of DIBALH in toluene (85 μ L, 0.085 mmol, 0.6 equiv) was added, followed by another portion of 1 M solution of DIBALH in toluene (85 μ L, 0.085 mmol, 0.6 equiv) after 30 min (total 1 M solution of DIBALH in toluene added: 340 μ L, 0.34 mmol, 2.4 equiv). After 30 min at –78 $^{\circ}$ C, TLC analysis of the reaction (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate) indicated complete consumption of the starting material and the reaction was quenched by addition of 100 μ L of MeOH, followed by the addition of saturated solution of Rochelle's salt (aq) (5 mL) at –78 $^{\circ}$ C. The reaction was allowed to warm to rt and stirred at rt for 30 min. Biphasic mixture was transferred to a separatory funnel and extracted with Et₂O (3 $\times$ 10 mL). The combined organic layers were dried over Na₂SO₄ and concentrated by use of a rotary evaporator. The residue was purified by flash column chromatography on silica gel using 10:90 ethyl acetate:hexanes as eluent to yield a 2.1:1 mixture of lactol epimers **67** as a colorless solid (46 mg, 0.10 mmol, 72% yield): R_f = 0.21 (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate); ¹H NMR (600 MHz, C₆D₆) δ 5.84 (t, J = 6.3 Hz, 1H), 5.53 (s, 2H), 5.37 (d, J = 2.3 Hz, 2H), 5.34 (d, J = 2.6 Hz,

1H), 3.65 (qd, $J = 10.9, 3.9$ Hz, 3H), 3.23 (d, $J = 5.5$ Hz, 2H), 2.97–2.80 (m, 2H), 2.69–2.62 (m, 1H), 2.62–2.46 (m, 8H), 2.41 (dt, $J = 9.9, 4.9$ Hz, 2H), 2.21 (d, $J = 12.3$ Hz, 1H), 2.16 (d, $J = 11.9$ Hz, 2H), 2.05–1.98 (m, 1H), 1.85–1.74 (m, 2H), 1.60–1.07 (m, 71H), 1.03 (d, $J = 6.9$ Hz, 6H), 1.01 (d, $J = 7.1$ Hz, 6H), 0.98–0.87 (m, 16H), 0.85 (d, $J = 6.5$ Hz, 6H), 0.83 (d, $J = 6.5$ Hz, 4H), 0.80–0.71 (m, 6H); ^{13}C NMR (151 MHz, C_6D_6) δ 172.8, 172.3, 104.7, 103.2, 102.0, 101.0, 80.9, 80.4, 75.6, 75.4, 61.1, 58.9, 49.2, 49.1, 44.8, 41.5, 40.9, 40.8, 40.5, 37.7, 37.6, 37.2, 36.8, 36.7, 35.4, 35.3, 32.2, 32.1, 28.7, 28.6, 27.1, 27.0, 25.8, 23.9, 23.6, 23.1, 22.52, 22.47, 22.45, 22.4, 22.03, 21.95, 21.4, 21.1, 16.54, 16.46; IR (thin film) 3463, 3438, 2924, 2867, 1704, 1636, 1367, 1152 cm^{-1} ; $[\alpha]_{\text{D}}^{21} + 101$, $[\alpha]_{\text{D}}^{21} + 101$, $[\alpha]_{\text{D}}^{21} + 112$, $[\alpha]_{\text{D}}^{21} + 159$ ($c = 0.15$, CHCl_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{27}\text{H}_{48}\text{O}_5\text{Na}$ 475.3399; Found 475.3395.

Preparation of Dioxabicyclo[3.3.0]octan-3-one 68. A 2-dram scintillation vial was charged with lactol **67** (34 mg, 0.075 mmol, 1.0 equiv), 1:1 4 M HCl (aq):THF (3 mL, 0.025 M), and a magnetic stir bar under ambient atmosphere. The resulting biphasic mixture was stirred vigorously at rt for 30 min. The reaction mixture was transferred to a separatory funnel and extracted with Et_2O (3×10 mL). The combined organic layers were dried over Na_2SO_4 and concentrated by use of a rotary evaporator. The residue was purified by flash column chromatography on silica gel using 10:90 ethyl acetate:hexanes $\rightarrow$ 25:75 ethyl acetate:hexanes as eluent to yield product **68** as a colorless solid (11 mg, 0.046 mmol, 61% yield): $R_f = 0.30$ (30:70 ethyl acetate:hexanes, stained with ceric ammonium molybdate). Spectral data were consistent with reported values.³¹

Preparation of Methyl Oxalate S7. A round-bottom flask was charged with **54** (560 mg, 2.7 mmol, 1.0 equiv), DMAP (33 mg, 0.27 mmol, 0.1 equiv), dichloromethane (13 mL, 0.20 M), and a stir bar under ambient atmosphere. Next, Et_3N (0.45 mL, 3.2 mmol, 1.2 equiv) and methyl chlorooxacetate (0.30 mL, 3.2 mmol, 1.2 equiv) were added sequentially. The resulting yellow solution was maintained at rt for 10 min, at which point TLC analysis (10:90 ethyl acetate:hexanes, stained with *p*-anisaldehyde) indicated complete consumption of the starting material. The reaction was quenched via addition of sat. NH_4Cl (aq) (15 mL). The resulting biphasic mixture was transferred to a separatory funnel and extracted with CH_2Cl_2 (3×25 mL). Combined organic layers were dried over MgSO_4 and concentrated by use of a rotary evaporator. The crude product was purified by flash column chromatography on silica gel using 5:95 ethyl acetate:hexanes as eluent to yield the desired product **S7** as a clear oil (720 mg, 2.45 mmol, 91% yield): $R_f = 0.43$ (10:90 ethyl acetate:hexanes, stained with *p*-anisaldehyde); ^1H NMR (500 MHz, CDCl_3) δ 4.92 (d, $J = 2.3$ Hz, 1H), 4.86 (d, $J = 2.3$ Hz, 1H), 3.88 (s, 3H), 3.20 (dd, $J = 8.1, 1.7$ Hz, 1H), 2.44 (dddd, $J = 12.0, 10.1, 3.4, 1.6$ Hz, 1H), 2.38–2.26 (m, 2H), 1.92–1.72 (m, 4H), 1.69–1.59 (m, 2H), 1.49 (s, 3H), 1.45–1.31 (m, 1H), 1.25 (dt, $J = 14.3, 3.5$ Hz, 1H), 0.98 (s, 3H), 0.91 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ 159.3, 157.1, 150.2, 117.2, 99.7, 57.7, 53.5, 50.9, 37.6, 37.3, 36.2, 35.9, 33.9, 28.7, 26.1, 24.3, 21.7; IR (thin film) 2953, 2937, 2867, 1766, 1739, 1154 cm^{-1} ; $[\alpha]_{\text{D}}^{23} + 11.5$, $[\alpha]_{\text{D}}^{23} + 13.6$, $[\alpha]_{\text{D}}^{23} + 14.4$, $[\alpha]_{\text{D}}^{23} + 29.1$ ($c = 1.0$, CHCl_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{17}\text{H}_{26}\text{O}_4\text{Na}$ 317.1729; Found 317.1732.

Preparation Lithium Oxalate 69. A round-bottom flask was charged with **S7** (700 mg, 2.38 mmol, 1.0 equiv), 1:1 THF: H_2O (4.8 mL, 0.5 M), and a stir bar under ambient atmosphere. The resulting biphasic mixture was cooled to 0 °C. Next, 0.5 N LiOH (aq) (4.8 mL, 1.0 equiv) was added dropwise. The mixture was then stirred vigorously at 0 °C for 5 min. The stir bar was removed and homogeneous solution was concentrated by use of a rotary evaporator with water bath warmed gradually from rt to 45 °C. The resulting colorless solid was washed with pentanes (3×5 mL) and dried further under high vacuum to yield product **69** as a colorless solid (680 mg, 2.38 mmol, 100% yield); ^1H NMR (500 MHz, CD_3OD) δ 4.91 (d, $J = 2.6$ Hz, 1H), 4.86 (d, $J = 2.6$ Hz, 1H), 2.51–2.37 (m, 2H), 2.35–2.27 (m, 1H), 1.95–1.84 (m, 1H), 1.84–1.67 (m, 5H), 1.46 (s, 3H), 1.43–1.34 (m, 1H), 1.31–1.15 (m, 1H), 1.00 (s, 3H), 0.93 (s, 3H); ^{13}C NMR (126 MHz, CD_3OD) δ 167.0, 166.3, 152.7, 116.8, 97.0, 58.5, 51.9, 38.6, 38.5, 37.0, 36.9, 34.2, 29.8, 26.4, 25.1, 22.2; IR (thin film)

2954, 2935, 2865, 1710, 1693, 1667, 1251, 1161 cm^{-1} ; $[\alpha]_{\text{D}}^{23} + 20.9$, $[\alpha]_{\text{D}}^{23} + 22.7$, $[\alpha]_{\text{D}}^{23} + 21.4$, $[\alpha]_{\text{D}}^{23} + 40.4$ ($c = 0.6$, MeOH); HRMS (ESI-TOF) m/z : $[\text{M}]^-$ Calcd for $\text{C}_{16}\text{H}_{23}\text{O}_4$ 279.1596; Found 279.1595.

Preparation of Potassium Oxalate 72. A round-bottom flask was charged with **S7** (81 mg, 0.28 mmol, 1.0 equiv), 1:1 THF: H_2O (1.4 mL, 0.2 M), and a stir bar under ambient atmosphere. The resulting biphasic mixture was cooled to 0 °C. Next, 0.85 N KOH (aq) (320 μL , 1.0 equiv) was added dropwise. The mixture was then stirred vigorously at 0 °C for 5 min. The stir bar was removed and homogeneous solution was concentrated by use of a rotary evaporator with water bath warmed gradually from rt to 45 °C. The resulting colorless solid was washed with pentanes (3×5 mL) and dried further under high vacuum to yield product **72** as a colorless solid (86 mg, 0.28 mmol, 99% yield); ^1H NMR (600 MHz, CD_3OD) 4.93–4.89 (m, 2H), 2.49–2.38 (m, 2H), 2.31 (d, $J = 12.9$ Hz, 1H), 1.93–1.84 (m, 1H), 1.84–1.69 (m, 5H), 1.48–1.44 (m, 3H), 1.41 (q, $J = 12.3, 11.4$ Hz, 1H), 1.25 (d, $J = 13.8$ Hz, 1H), 1.00 (t, $J = 2.5$ Hz, 3H), 0.95–0.91 (m, 3H); ^{13}C NMR (151 MHz, CD_3OD) δ 166.9, 166.5, 152.8, 117.0, 97.2, 58.7, 52.0, 38.7, 38.6, 37.2, 37.1, 34.3, 30.0, 26.5, 25.3, 22.4; IR (thin film) 3608, 1713, 1663, 1641, 1233, 893 cm^{-1} ; $[\alpha]_{\text{D}}^{23} + 14.6$, $[\alpha]_{\text{D}}^{23} + 14.9$, $[\alpha]_{\text{D}}^{23} + 16.7$, $[\alpha]_{\text{D}}^{23} + 31.2$ ($c = 1.7$, MeOH); HRMS (ESI-TOF) m/z calculated for $\text{C}_{16}\text{H}_{23}\text{O}_4$ $[\text{M}]^-$ 279.1596, observed 269.1598. HRMS (ESI-TOF) m/z : $[\text{M}]^-$ Calcd for $\text{C}_{16}\text{H}_{23}\text{O}_4$ 279.1596; Found 279.1598.

Preparation of Lactone 70 from 69. On the bench under ambient atmosphere, 1-dram scintillation vial was charged with **69** (86 mg, 0.30 mmol, 1.0 equiv), *ent*-**S9** (82 mg, 0.30 mmol, 1.0 equiv), $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$ (7 mg, 0.006 mmol, 0.02 equiv), tetrahydrofuran (500 μL , 0.6 M), H_2O (27 μL , 1.5 mmol, 5 equiv), and a stir bar. The vial was then sealed with screw cap bearing Teflon septum. The septum of the vial was pierced with a 21 gauge $\times 1.5$ " needle that was inserted just barely through the septum with the tip of the needle kept above the fluid level inside the vial. A separate 22 gauge $\times 3$ " needle attached to a flow of argon was also pierced through the septum, and the tip of the needle was pushed to the bottom of the vial and submersed in the fluid. The reaction mixture was degassed by sparging with argon for 15 min. Both needles were removed, and the sealed vial was then placed on a stir plate equipped with 2×34 W blue LED lamps and a rack to hold the vial inside of a cardboard box to block light pollution from entering the lab. The vial was placed approximately 4 cm from the lamps and stirred vigorously. The sample was irradiated by the lamps for 18 h inside the closed box, allowing the temperature of the reaction mixture to rise to 60 °C and the air inside the box to 40–45 °C because of heat given off from the LEDs. The reaction was allowed to cool to rt, diluted with Et_2O (1 mL) and filtered over MgSO_4 . The filtrate was concentrated by use of a rotary evaporator. The crude product was purified by flash column chromatography on silica gel using 2:98 ethyl acetate:hexanes as eluent to yield the desired product **70** as a thick colorless foam (102 mg, 0.22 mmol, 73% yield): $R_f = 0.33$ (5:95 ethyl acetate:hexanes, stained with *p*-anisaldehyde); ^1H NMR (600 MHz, CDCl_3) δ 5.59 (d, $J = 3.3$ Hz, 1H), 4.87 (s, 1H), 4.62 (s, 1H), 3.59 (td, $J = 10.7, 4.7$ Hz, 1H), 2.68–2.59 (m, 2H), 2.36 (dd, $J = 12.3, 5.4$ Hz, 1H), 2.26–2.17 (m, 1H), 2.14 (d, $J = 11.1$ Hz, 1H), 2.05 (q, $J = 9.8$ Hz, 1H), 1.83–1.73 (m, 5H), 1.73–1.57 (m, 5H), 1.45–1.34 (m, 2H), 1.28 (dd, $J = 12.4, 6.7$ Hz, 2H), 1.03–0.97 (m, 4H), 0.97–0.92 (m, 7H), 0.92–0.86 (m, 4H), 0.84 (s, 3H), 0.80 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (151 MHz, CDCl_3) δ 172.6, 153.4, 115.2, 101.9, 77.7, 60.9, 55.7, 53.7, 52.2, 48.1, 47.9, 40.0, 37.7, 37.4, 37.0, 36.5, 34.5, 31.6, 28.9, 25.8, 25.7, 25.4, 23.2, 22.5, 21.2, 21.1, 15.7; IR (thin film) 2953, 2922, 2868, 1789, 1364 cm^{-1} ; $[\alpha]_{\text{D}}^{23} - 52.0$, $[\alpha]_{\text{D}}^{23} - 54.3$, $[\alpha]_{\text{D}}^{23} - 65.8$, $[\alpha]_{\text{D}}^{23} - 104$ ($c = 0.6$, CHCl_3); HRMS (ESI-TOF) m/z : $[\text{M}+\text{Na}]^+$ Calcd for $\text{C}_{28}\text{H}_{45}\text{ClO}_3\text{Na}$ 487.2955; Found 487.2971.

Preparation of Lactone 70 from 72. On the bench under ambient atmosphere, 1-dram scintillation vial was charged with **72** (63 mg, 0.20 mmol, 1.0 equiv), *ent*-**S9** (55 mg, 0.20 mmol, 1.0 equiv), $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$ (5 mg, 0.004 mmol, 0.02 equiv), tetrahydrofuran (330 μL , 0.6 M), H_2O (18 μL , 1.0 mmol, 5 equiv), and a stir bar. The vial was then sealed with screw cap bearing Teflon

septum. The septum of the vial was pierced with a 21 gauge $\times$ 1.5" needle that was inserted just barely through the septum with the tip of the needle kept above the fluid level inside the vial. A separate 22 gauge $\times$ 3" needle attached to a flow of argon was also pierced through the septum, and the tip of the needle was pushed to the bottom of the vial and submersed in the fluid. The reaction mixture was degassed by sparging with argon for 15 min. Both needles were removed, and the sealed vial was then placed on a stir plate equipped with 2 $\times$ 34 W blue LED lamps and a rack to hold the vial inside of a cardboard box to block light pollution from entering the lab. The vial was placed approximately 4 cm from the lamps and stirred vigorously. The sample was irradiated by the lamps for 18 h inside the closed box, allowing the temperature of the reaction mixture to rise to 60 °C and the air inside the box to 40–45 °C because of heat given off from the LEDs. The reaction was allowed to cool to rt, diluted with Et₂O (1 mL) and filtered over MgSO₄. The filtrate was concentrated by use of a rotary evaporator. The crude product was purified by flash column chromatography on silica gel using 2:98 ethyl acetate:hexanes as eluent to yield the desired product **70** as a thick colorless foam (67 mg, 0.14 mmol, 72% yield): R_f = 0.33 (5:95 ethyl acetate:hexanes, stained with *p*-anisaldehyde). Spectral data were consistent with reported values.³¹

Preparation of Lactone 71 from 70. On the bench under ambient atmosphere, 1-dram scintillation vial was charged with **70** (86 mg, 0.18 mmol, 1.0 equiv), Bu₃N (44 μ L, 1.8 mmol, 10 equiv), Ir[dF(CF₃)ppy]₂(dtbbpy)PF₆ (4 mg, 0.004 mmol, 0.02 equiv), tetrahydrofuran (1.8 mL, 0.1 M), H₂O (33 μ L, 1.8 mmol, 10 equiv), and a stir bar. The vial was then sealed with screw cap bearing Teflon septum. The septum of the vial was pierced with a 21 gauge $\times$ 1.5" needle that was inserted just barely through the septum with the tip of the needle kept above the fluid level inside the vial. A separate 22 gauge $\times$ 3" needle attached to a flow of argon was also pierced through the septum, and the tip of the needle was pushed to the bottom of the vial and submersed in the fluid. The reaction mixture was degassed by sparging with argon for 15 min. Both needles were removed, and the sealed vial was then placed on a stir plate equipped with 2 $\times$ 34 W blue LED lamps and a rack to hold the vial inside of a cardboard box to block light pollution from entering the lab. The vial was placed approximately 4 cm from the lamps and stirred vigorously. The sample was irradiated by the lamps for 4 h inside the closed box, allowing the temperature of the reaction mixture to rise to 60 °C and the air inside the box to 40–45 °C because of heat given off from the LEDs. The reaction was allowed to cool to rt and transferred to a separatory funnel with Et₂O (10 mL). The solution was then sequentially washed with 1 N HCl (aq) (10 mL), H₂O (10 mL), and brine (10 mL). The organic layer was dried over MgSO₄ and concentrated by use of a rotary evaporator. The crude product was purified by flash column chromatography on silica gel using 2:98 ethyl acetate:hexanes as eluent to yield the desired product **71** as a thick colorless foam (79 mg, 0.18 mmol, 100% yield): R_f = 0.28 (5:95 ethyl acetate:hexanes, stained with *p*-anisaldehyde). Spectral data were consistent with reported values.³¹

Preparation of Lactone 71 from 69. On the bench under ambient atmosphere, 1-dram scintillation vial was charged with **69** (86 mg, 0.30 mmol, 1.0 equiv), *ent*-59 (82 mg, 0.30 mmol, 1.0 equiv), Ir[dF(CF₃)ppy]₂(dtbbpy)PF₆ (7 mg, 0.006 mmol, 0.02 equiv), tetrahydrofuran (500 μ L, 0.6 M), H₂O (27 μ L, 1.5 mmol, 5 equiv), and a stir bar. The vial was then sealed with screw cap bearing Teflon septum. The septum of the vial was pierced with a 21 gauge $\times$ 1.5" needle that was inserted just barely through the septum with the tip of the needle kept above the fluid level inside the vial. A separate 22 gauge $\times$ 3" needle attached to a flow of argon was also pierced through the septum, and the tip of the needle was pushed to the bottom of the vial and submersed in the fluid. The reaction mixture was degassed by sparging with argon for 15 min. Both needles were removed, and the sealed vial was then placed on a stir plate equipped with 2 $\times$ 34 W blue LED lamps and a rack to hold the vial inside of a cardboard box to block light pollution from entering the lab. The vial was placed approximately 4 cm from the lamps and stirred vigorously. The sample was irradiated by the lamps for 18 h inside the closed box, allowing the temperature of the reaction mixture to rise to 60 °C and the air inside the box to 40–45 °C because of heat given off from the LEDs. After

18 h, a degassed solution of Bu₃N (0.7 mL, 3 mmol, 10 equiv) in tetrahydrofuran (2.5 mL, 0.12 M) was added via syringe. The sample was irradiated by lamps for additional 4 h inside the closed box. The reaction was allowed to cool to rt and transferred to a separatory funnel with Et₂O (15 mL). The solution was then sequentially washed with 1 N HCl (aq) (10 mL), H₂O (10 mL), and brine (10 mL). The organic layer was dried over MgSO₄ and concentrated by use of a rotary evaporator. The crude product was purified by flash column chromatography on silica gel using 2:98 ethyl acetate:hexanes as eluent to yield the desired product **71** as a thick colorless foam (97 mg, 0.23 mmol, 75% yield): R_f = 0.28 (5:95 ethyl acetate:hexanes, stained with *p*-anisaldehyde). Spectral data were consistent with reported values.³¹

Preparation of Ester 73. A round-bottom flask was charged with **71** (420 mg, 0.98 mmol, 1.0 equiv), tetrahydrofuran (10 mL, 0.1 M), and a stir bar under an argon atmosphere. The solution was cooled to –78 °C. Next, a 1 M solution of LiHMDS in tetrahydrofuran (1.2 mL, 1.2 mmol, 1.2 equiv) was added dropwise. After stirring the reaction mixture for 15 min at –78 °C, *tert*-butyl bromoacetate (220 μ L, 1.5 mmol, 1.5 equiv) was added dropwise. After 1 h at –78 °C, the reaction was quenched via addition of sat. NH₄Cl (aq) (10 mL). The resulting mixture was allowed to warm to rt and transferred to a separatory funnel and extracted with Et₂O (3 $\times$ 25 mL). Combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The crude product was purified by flash column chromatography on silica gel using 2:98 ethyl acetate:hexanes as eluent to yield the desired product **73** as a clear oil (480 mg, 2.45 mmol, 91% yield): R_f = 0.31 (5:95 ethyl acetate:hexanes, stained with *p*-anisaldehyde); ¹H NMR (600 MHz, CDCl₃) δ 5.58 (d, *J* = 1.3 Hz, 1H), 4.85 (d, *J* = 2.2 Hz, 1H), 4.62 (d, *J* = 2.1 Hz, 1H), 3.55 (td, *J* = 10.7, 4.2 Hz, 1H), 2.86 (ddd, *J* = 8.6, 5.3, 2.8 Hz, 1H), 2.72 (dd, *J* = 15.3, 5.3 Hz, 1H), 2.62 (d, *J* = 8.8 Hz, 1H), 2.60 (d, *J* = 9.0 Hz, 1H), 2.37–2.30 (m, 1H), 2.18–2.08 (m, 3H), 2.02 (dt, *J* = 11.2, 8.0 Hz, 1H), 1.82–1.71 (m, 3H), 1.71–1.60 (m, 5H), 1.47 (s, 9H), 1.46–1.42 (m, 2H), 1.38 (dt, *J* = 13.7, 3.3 Hz, 2H), 1.25 (dd, *J* = 10.4, 3.3 Hz, 2H), 1.03–0.95 (m, 4H), 0.94 (d, *J* = 6.5 Hz, 3H), 0.92 (s, 3H), 0.90 (d, *J* = 7.0 Hz, 3H), 0.89–0.82 (m, 2H), 0.79 (s, 3H), 0.77 (d, *J* = 7.0 Hz, 3H); ¹³C NMR (151 MHz, CDCl₃) δ 179.1, 170.5, 154.2, 114.9, 102.4, 81.7, 57.0, 56.3, 53.8, 48.3, 48.1, 40.2, 40.0, 39.5, 38.0, 37.5, 37.2, 36.7, 34.74, 34.70, 31.8, 29.3, 28.49, 28.46, 26.00, 25.95, 25.8, 23.2, 22.7, 21.5, 21.3, 15.7; IR (thin film) 2953, 2924, 2869, 1780, 1732, 1456, 1367, 1152 cm^{–1}; [α]_D²³ –41.9, [α]_D²³₅₇₇ –45.2, [α]_D²³₅₄₆ –52.6, [α]_D²³₄₃₅ –83.9 (*c* = 0.9, CHCl₃); HRMS (ESI-TOF) *m/z*: [M + Na]⁺ Calcd for C₃₄H₅₆O₅Na 567.4025; Found 567.4030.

Preparation of (+)-Cheloviolene B (8). A round-bottom flask was charged with **73** (278 mg, 0.51 mmol, 1.0 equiv), toluene (5.1 mL, 0.1 M), and a stir bar under an argon atmosphere. The solution was cooled to –78 °C. Next, a 1 M solution of DIBALH in toluene (610 μ L, 0.61 mmol, 1.2 equiv) was added dropwise. After 1 h, TLC analysis of the reaction (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate) indicated presence of the starting material and additional 1 M solution of DIBALH in toluene (300 μ L, 0.3 mmol, 0.5 equiv) was added, followed by another portion of 1 M solution of DIBALH in toluene (300 μ L, 0.3 mmol, 0.5 equiv) after 30 min (total 1 M solution of DIBALH in toluene added: 1.2 mL, 1.2 mmol). After 30 min, TLC analysis indicated complete consumption of the starting material and the reaction was quenched via addition of 100 μ L of MeOH, followed by the addition of saturated solution of Rochelle's salt (aq) (10 mL) at –78 °C. The reaction was allowed to warm to rt and stirred at rt for 30 min. Biphasic mixture was transferred to a separatory funnel and extracted with Et₂O (3 $\times$ 25 mL). Combined organic layers were dried over MgSO₄ and concentrated by use of a rotary evaporator. The crude product was quickly passed through a pH 7 silica gel⁸³ column using 5:95 ethyl acetate:hexanes to elute crude lactol epimers **75** as a clear oil [crude mass: 155 mg; 1.1:1 dr; R_f minor = 0.45; R_f major = 0.3 (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate); diagnostic ¹H NMR shifts (600 MHz, CDCl₃): major: δ 5.26 (t, *J* = 4.6 Hz, 1H), 5.23 (d, *J* = 1.5 Hz, 1H), 4.82 (d, *J* = 2.3 Hz, 1H), 4.65 (d, *J* = 2.3 Hz, 1H); minor: δ 5.52 (dd, *J* = 7.9, 5.7 Hz, 1H), 5.20 (d, *J* = 2.3 Hz, 1H), 4.81 (d, *J* = 2.2 Hz, 1H), 4.60 (d, *J* = 2.3 Hz, 1H)] and 10:90 ethyl acetate:hexanes to elute

crude bicyclic lactol epimers **76** a clear oil [crude mass: 72 mg; 1.7:1 dr; R_f = 0.15 (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate); diagnostic ^1H NMR shifts (600 MHz, CDCl_3): major: δ 5.81 (d, J = 6.0 Hz, 1H), 5.46 (dd, J = 11.1, 5.5 Hz, 1H), 5.25 (s, 1H), 4.84 (s, 1H), 4.60 (s, 1H); minor: δ 5.89 (d, J = 5.8 Hz, 1H), 5.67–5.62 (m, 1H), 5.35 (s, 1H), 4.82 (s, 1H), 4.61 (s, 1H)].

A 2-dram scintillation vial was charged with crude bicyclic lactol epimers **76** (72 mg, 0.15 mmol, 1.0 equiv), dichloromethane (1.5 mL, 0.1 M), and a stir bar under ambient atmosphere. Next, PCC (65 mg, 0.30 mmol, 2.0 equiv) was added in one portion at rt. Heterogeneous reaction mixture was stirred vigorously at rt for 6 h, at which point TLC analysis (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate) indicated complete consumption of the starting material. The reaction mixture was filtered over pH 7 silica gel plug using 10:90 ethyl acetate:hexanes and concentrated by use of a rotary evaporator to yield crude bicyclic lactone **77** as a clear oil [crude mass: 70 mg; R_f = 0.24 (10:90 ethyl acetate:hexanes, stained with ceric ammonium molybdate); diagnostic ^1H NMR shifts (600 MHz, CDCl_3): δ 6.03 (d, J = 6.0 Hz, 1H), 5.42 (s, 1H), 4.84 (d, J = 2.1 Hz, 1H), 4.61 (d, J = 2.1 Hz, 1H)].

A 20 mL scintillation vial was charged with crude lactol epimers **75** (155 mg, 0.28 mmol, 1.0 equiv), crude bicyclic lactone **77** (70 mg, 0.15 mmol, 1.0 equiv), 1:1 4 N HCl (aq):THF (17 mL, 0.025 M), and a stir bar under ambient atmosphere. The resulting biphasic mixture was stirred vigorously at rt for 24 h. The reaction mixture was transferred to a separatory funnel and extracted with Et_2O (3×25 mL). Combined organic layers were dried over MgSO_4 and concentrated by use of a rotary evaporator. The crude product was purified by flash column chromatography on silica gel using 10:90 ethyl acetate:hexanes $\rightarrow$ 20:80 ethyl acetate:hexanes as eluent to yield (+)-cheloviolene **B** (**8**) as a colorless solid (90 mg, 0.27 mmol, 53% yield from **73**): R_f = 0.15 (20:80 ethyl acetate:hexanes, stained with ceric ammonium molybdate). Spectral data were consistent with reported values.³¹

Acid-Catalyzed Hydration of (+)-Dendrillolide C (11). A 1-dram scintillation vial was charged with **11** (9 mg, 0.028 mmol, 1 equiv), 1:1 4 M HCl (aq):THF (0.6 mL, 0.025 M), and a magnetic stir bar. The resulting heterogeneous mixture was stirred vigorously at 40 °C for 18 h. The reaction mixture was diluted with H_2O (1 mL) and extracted with Et_2O (3×2 mL). Organic layers were dried over MgSO_4 and concentrated by use of a rotary evaporator. Integration of ^1H NMR spectrum [CDCl_3 , 600 MHz: shifts corresponding to **8**: δ 6.09 (d, J = 6.0 Hz, 1H), 5.64 (s, 1H), 4.83 (d, J = 2.0 Hz, 1H), 4.61 (d, J = 2.0 Hz, 1H); shifts corresponding to **79**: δ 7.42 (d, J = 1.6 Hz, 1H), 7.21 (d, J = 1.7 Hz, 1H), 4.89 (d, J = 2.2 Hz, 1H), 4.71 (d, J = 2.1 Hz, 1H), 3.63 (s, 2H)] of the crude reaction mixture was used to establish the ratio of the two products formed, 1.2:1 (+)-cheloviolene **B** (**8**): furan **79**.³¹

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.7b02458.

Experimental procedures and characterization data (PDF)

X-ray crystallographic data of compound **7** (CIF)

X-ray crystallographic data of compound **S5** (CIF)

X-ray crystallographic data of compound **S6** (CIF)

X-ray crystallographic data of compound **ent-S9** (CIF)

X-ray crystallographic data of compound **8** (CIF)

■ AUTHOR INFORMATION

Corresponding Author

*leoverma@uci.edu

ORCID

Peng Zhao: 0000-0003-4581-406X

Young Ho Rhee: 0000-0002-2094-4426

Larry E. Overman: 0000-0001-9462-0195

Author Contributions

*M.R.G., Y.S., and C.R.J. contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support was provided by the NSF (CHE1265964) and by a graduate fellowship award to Y.S. (1F31GM113494). J.L.'s research at UC Irvine was supported by the National Research Foundation of Korea (NRF-2015R1A2A1A15056116). We thank Materia for a gift of Hoveyda-Grubbs second-generation olefin metathesis catalyst. NMR and mass spectra were obtained at UC Irvine using instrumentation acquired with the assistance of NSF and NIH Shared Instrumentation grants.

■ REFERENCES

- (1) For reviews, see: (a) Fürstner, A. *Chem. Rev.* **1999**, 99, 991–1046. (b) Nicolaou, K. C.; Snyder, S. A.; Montagnon, T.; Vassilikogiannakis, G. *Angew. Chem., Int. Ed.* **2002**, 41, 1668–1698. (c) Nicolaou, K. C.; Bulger, P. G.; Sarlah, D. *Angew. Chem., Int. Ed.* **2005**, 44, 4442–4489. (d) Nicolaou, K. C.; Bulger, P. G.; Sarlah, D. *Angew. Chem., Int. Ed.* **2005**, 44, 4490–4527. (e) Quasdorf, K. W.; Overman, L. E. *Nature* **2014**, 516, 181–191. (f) Gil, A.; Albericio, F.; Álvarez, M. *Chem. Rev.* **2017**, 117, 8420–8446.
- (2) For recent examples, see: (a) Wang, W.; Kishi, Y. *Org. Lett.* **1999**, 1, 1129–1132. (b) Lebsack, A. D.; Overman, L. E.; Valentekovich, R. J. *J. Am. Chem. Soc.* **2001**, 123, 4851–4852. (c) Lu, H.-H.; Martinez, M. D.; Shenvi, R. A. *Nat. Chem.* **2015**, 7, 604–607. (d) Lathrop, S. P.; Pompeo, M.; Chang, W.-T. T.; Movassaghi, M. *J. Am. Chem. Soc.* **2016**, 138, 7763–7769.
- (3) (a) Damm, W.; Giese, B.; Hartung, J.; Hasskerl, T.; Houk, K. N.; Hüter, O.; Zipse, H. *J. Am. Chem. Soc.* **1992**, 114, 4067–4079. (b) Arnaud, R.; Postlethwaite, H.; Barone, V. J. *Phys. Chem.* **1994**, 98, 5913–5919.
- (4) (a) Giese, B. *Angew. Chem., Int. Ed. Engl.* **1983**, 22, 753–764. (b) Fischer, H.; Radom, L. *Angew. Chem., Int. Ed.* **2001**, 40, 1340–1371. (c) Beckwith, A. L.; Poole, J. S. *J. Am. Chem. Soc.* **2002**, 124, 9489–9497.
- (5) Hayen, A.; Koch, R.; Saak, W.; Haase, D.; Metzger, J. O. *J. Am. Chem. Soc.* **2000**, 122, 12458–12468.
- (6) (a) Barton, D. H. R.; Sas, W. *Tetrahedron* **1990**, 46, 3419–3430. (b) Jasperse, C. P.; Curran, D. P.; Fevig, T. L. *Chem. Rev.* **1991**, 91, 1237–1286.
- (7) For reviews, see: (a) Tucker, J. W.; Stephenson, C. R. J. *J. Org. Chem.* **2012**, 77, 1617–1622. (b) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. *Chem. Rev.* **2013**, 113, 5322–5363. (c) Schultz, D. M.; Yoon, T. P. *Science* **2014**, 343, 1239176.
- (8) For recent examples of photoredox-mediated fragment couplings in total synthesis, see: (a) Furst, L.; Narayanam, J. M. R.; Stephenson, C. R. J. *Angew. Chem., Int. Ed.* **2011**, 50, 9655–9659. (b) Sun, Y.; Li, R.; Zhang, W.; Li, A. *Angew. Chem., Int. Ed.* **2013**, 52, 9201–9204. (c) Müller, D. S.; Untiedt, N. L.; Dieskau, A. P.; Lackner, G. L.; Overman, L. E. *J. Am. Chem. Soc.* **2015**, 137, 660–663. (d) Tao, D. J.; Slutskyy, Y.; Overman, L. E. *J. Am. Chem. Soc.* **2016**, 138, 2186–2189.
- (9) (a) Search of SciFinder on 6/14/2017. (b) Ring system **1** is also called *cis*-tetrahydro[2,3-*b*]furan-2(3H)one.
- (10) Karuso, P.; Bergquist, P. R.; Cambie, R. C.; Buckleton, J. S.; Clark, G. R.; Rickard, C. E. F. *Aust. J. Chem.* **1986**, 39, 1643–1653.
- (11) (a) Mayol, L.; Piccialli, V.; Sica, D. *J. Nat. Prod.* **1986**, 49, 823–828. (b) Mayol, L.; Piccialli, V.; Sica, D. *Tetrahedron* **1986**, 42, 5369–5376.

- (12) von Salm, J. L.; Witowski, C. G.; Fleeman, R. M.; McClintock, J. B.; Amsler, C. D.; Shaw, L. N.; Baker, B. J. *J. Org. Lett.* **2016**, *18*, 2596–2599.
- (13) Mayer, A.; Köpke, B.; Anke, H.; Sterner, O. *Phytochemistry* **1996**, *43*, 375–376.
- (14) For reviews of rearranged spongian diterpenoids, see: (a) Keyzers, R. A.; Northcote, P. T.; Davies-Coleman, M. T. *Nat. Prod. Rep.* **2006**, *23*, 321–334. (b) González, M. *Curr. Bioact. Compd.* **2007**, *3*, 1–36.
- (15) Hochlowski, J. E.; Faulkner, D. J.; Matsumoto, G. K.; Clardy, J. J. *Org. Chem.* **1983**, *48*, 1141–1142.
- (16) (a) Buckleton, J. S.; Cambie, R. C.; Clark, G. R. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1991**, *47*, 1438–1440. (b) Bergquist, P. R.; Bowden, B. F.; Cambie, R. C.; Craw, P. A.; Karuso, P.; Poiner, A.; Taylor, W. C. *Aust. J. Chem.* **1993**, *46*, 623–632. Cheloviolene A was isolated also by Faulkner and Bobzin and called chelonaplysin B.¹⁷
- (17) Bobzin, S. C.; Faulkner, D. J. *J. Nat. Prod.* **1991**, *54*, 225–232.
- (18) Molinski, T. F.; Faulkner, D. J.; He, C. H.; Van Duyne, G. D.; Clardy, J. J. *Org. Chem.* **1986**, *51*, 4564–4567.
- (19) All natural products having the spongian skeleton are reported to have the absolute configuration depicted in **12**.
- (20) (a) The absolute configuration of norrisolide was established by Theodorakis' inaugural total synthesis, see: Brady, T. P.; Kim, S. H.; Wen, K.; Theodorakis, E. A. *Angew. Chem., Int. Ed.* **2004**, *43*, 739–742. (b) For a second enantioselective total synthesis, see: Granger, K.; Snapper, M. L. *Eur. J. Org. Chem.* **2012**, *2012*, 2308–2311.
- (21) (a) Keyzers, R. A.; Northcote, P. T.; Zubkov, O. A. *Eur. J. Org. Chem.* **2004**, *2004*, 419–425. (b) Anketetty, S.; Amsler, C. D.; McClintock, J. B.; Baker, B. J. *J. Nat. Prod.* **2004**, *67*, 1172–1174.
- (22) Schnermann, M. J.; Beaudry, C. M.; Egorova, A. V.; Polishchuk, R. S.; Sütterlin, C.; Overman, L. E. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 6158–6163.
- (23) Klausner, R. D.; Donaldson, J. G.; Lippincott-Schwartz, J. J. *Cell Biol.* **1992**, *116*, 1071–1080.
- (24) Takizawa, P. A.; Yucel, J. K.; Veit, B.; Faulkner, D. J.; Deerinck, T.; Soto, G.; Ellisman, M.; Malhotra, V. *Cell* **1993**, *73*, 1079–1090.
- (25) (a) Guizzunti, G.; Brady, T. P.; Malhotra, V.; Theodorakis, E. A. *J. Am. Chem. Soc.* **2006**, *128*, 4190–4191. (b) Guizzunti, G.; Brady, T. P.; Fischer, D.; Malhotra, V.; Theodorakis, E. A. *Bioorg. Med. Chem.* **2010**, *18*, 2115–2122 and references therein.
- (26) Schnermann, M. J.; Beaudry, C. M.; Genung, N. E.; Canham, S. M.; Untiedt, N. L.; Karanikolas, B. D. W.; Sütterlin, C.; Overman, L. E. *J. Am. Chem. Soc.* **2011**, *133*, 17494–17503.
- (27) Jamison, C. R.; Overman, L. E. *Acc. Chem. Res.* **2016**, *49*, 1578–1586.
- (28) Lackner, G. L.; Quasdorf, K. W.; Overman, L. E. *J. Am. Chem. Soc.* **2013**, *135*, 15342–15345.
- (29) For discussions of the challenge in assembling chiral attached ring systems, see: (a) Overman, L. E.; Pennington, L. D. *Can. J. Chem.* **2000**, *78*, 732–738. (b) Overman, L. E.; Velthuisen, E. J. *J. Org. Chem.* **2006**, *71*, 1581–1587.
- (30) Schnermann, M. J.; Overman, L. E. *Angew. Chem., Int. Ed.* **2012**, *51*, 9576–9580.
- (31) For a preliminary report of a portion of these studies, see: Slutskyy, Y.; Jamison, C. R.; Zhao, P.; Lee, J.; Rhee, Y. H.; Overman, L. E. *J. Am. Chem. Soc.* **2017**, *139*, 7192–7195.
- (32) For other constructions of *cis*-2,8-dioxabicyclo[3.3.0]octan-3-ones containing carbon substituents at C-6, see: references **20a**, **b**, **25a**, and **26** (a) Nakata, T.; Nagao, S.; Oishi, T. *Tetrahedron Lett.* **1985**, *26*, 6465–6468. (b) Toyota, M.; Nishikawa, Y.; Fukumoto, K. *Heterocycles* **1998**, *47*, 675–678. (c) Toyota, M.; Hirota, M.; Nishikawa, Y.; Fukumoto, K.; Ihara, M. *J. Org. Chem.* **1998**, *63*, 5895–5902. (d) Weisser, R.; Yue, W.; Reiser, O. *Org. Lett.* **2005**, *7*, 5353–5356. (e) Brady, T. P.; Wallace, E. K.; Kim, S. H.; Guizzunti, G.; Malhotra, V.; Theodorakis, E. A. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 5035–5039. See also: (f) Corey, E. J.; Letavic, M. A. *J. Am. Chem. Soc.* **1995**, *117*, 9616–9617.
- (33) For examples of preparing butenolides in this way, see: (a) Hughes, G.; Kimura, M.; Buchwald, S. L. *J. Am. Chem. Soc.* **2003**, *125*, 11253–11258. (b) Conrad, J. C.; Parnas, H. H.; Snelgrove, J. L.; Fogg, D. E. *J. Am. Chem. Soc.* **2005**, *127*, 11882–11883. (c) Geurts, K.; Fletcher, S. P.; Feringa, B. L. *J. Am. Chem. Soc.* **2006**, *128*, 15572–15573.
- (34) (a) Ram, R. N.; Charles, I. *Tetrahedron* **1997**, *53*, 7335–7340. (b) Feray, L.; Bertrand, M. P. *Eur. J. Org. Chem.* **2008**, *2008*, 3164–3170.
- (35) Stewart, I. C.; Ung, T.; Pletnev, A. A.; Berlin, J. M.; Grubbs, R. H.; Schrodli, Y. *Org. Lett.* **2007**, *9*, 1589–1592.
- (36) Chapleo, C. B.; Svanholt, K. L.; Martin, R.; Dreiding, A. S. *Helv. Chim. Acta* **1976**, *59*, 100–107.
- (37) Okada, K.; Okamoto, K.; Morita, N.; Okubo, K.; Oda, M. *J. Am. Chem. Soc.* **1991**, *113*, 9401–9402.
- (38) (a) Giese, G.; Zwick, W. *Angew. Chem., Int. Ed. Engl.* **1978**, *17*, 66–67. (b) Srikanth, G. S. C.; Castle, S. L. *Tetrahedron* **2005**, *61*, 10377–10441.
- (39) Napolitano, J. E.; Gavín, J. A.; García, C.; Norte, M.; Fernández, J. J.; Daranas, A. H. *Chem. - Eur. J.* **2011**, *17*, 6338–6347.
- (40) See [Supporting Information](#) for details.
- (41) Pratsch, G.; Lackner, G. L.; Overman, L. E. *J. Org. Chem.* **2015**, *80*, 6025–6036.
- (42) (a) Takano, S.; Kudo, J.; Takahashi, M.; Ogasawara, K. *Tetrahedron Lett.* **1986**, *27*, 2405–2408. (b) Chen, W.; Hartwig, J. F. *J. Am. Chem. Soc.* **2012**, *134*, 15249–15252. (c) Xu, Z.-J.; Wu, Y. *Chem. - Eur. J.* **2017**, *23*, 2026–2030.
- (43) Product **31** epimerized completely to form **30a** upon exposure to DBU, which formed the basis of the stereochemical assignment for these epimers.
- (44) See, *inter alia* (a) Giese, B.; Farshchi, H.; Hartmanns, J.; Metzger, J. O. *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 600–601. (b) Krause, N. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 1764–1765. (c) Ischay, M. A.; Anzovino, M. E.; Du, J.; Yoon, T. P. *J. Am. Chem. Soc.* **2008**, *130*, 12886–12887. (d) Emmanuel, M. A.; Greenberg, N. R.; Oblinsky, D. G.; Hyster, T. K. *Nature* **2016**, *540*, 414–417.
- (45) Highly enantioenriched samples of both the (R)- and (S)-enantiomers 5-methoxy- and 5-menthoxybutenolides can be purchased from Accel Pharmtech or Proactive Molecular Research.
- (46) Qin, T.; Malins, L. R.; Edwards, J. T.; Merchant, R. R.; Novak, A. J. E.; Zhong, J. Z.; Mills, R. B.; Yan, M.; Yuan, C.; Eastgate, M. D.; Baran, P. S. *Angew. Chem., Int. Ed.* **2017**, *56*, 260–265.
- (47) We were able to reproduce the results of the coupling reaction between acrylonitrile and adamantane-derived *N*-acyloxypthalimide reported in reference **46**.
- (48) (a) Hedstrand, D. M.; Kruizinga, W. H.; Kellogg, R. M. *Tetrahedron Lett.* **1978**, *19*, 1255–1258. (b) Nakamura, K.; Fujii, M.; Mekata, H.; Oka, S.; Ohno, A. *Chem. Lett.* **1986**, *15*, 87–88. (c) Nguyen, J. D.; D'Amato, E. M.; Narayanam, J. M. R.; Stephenson, C. R. J. *Nat. Chem.* **2012**, *4*, 854–859. (d) Lackner, G. L.; Quasdorf, K. W.; Pratsch, G.; Overman, L. E. *J. Org. Chem.* **2015**, *80*, 6012–6024.
- (49) Chu, L.; Ohta, C.; Zuo, Z.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 10886–10889.
- (50) Hydrogen atom abstraction from solvent is likely responsible for premature reduction of the tertiary radical.
- (51) (a) Nawrat, C. C.; Jamison, C. R.; Slutskyy, Y.; MacMillan, D. W. C.; Overman, L. E. *J. Am. Chem. Soc.* **2015**, *137*, 11270–11273. (b) Jamison, C. R.; Slutskyy, Y.; Overman, L. E. *Org. Synth.* **2017**, *94*, 167–183.
- (52) Slutskyy, Y.; Jamison, C. R.; Lackner, G. L.; Müller, D. S.; Dieskau, A. P.; Untiedt, N. L.; Overman, L. E. *J. Org. Chem.* **2016**, *81*, 7029–7035.
- (53) Kozikowski, A. P.; Mugrage, B. B.; Wang, B. C.; Xu, Z. B. *Tetrahedron Lett.* **1983**, *24*, 3705–3708.
- (54) Curran, D. P. *J. Am. Chem. Soc.* **1983**, *105*, 5826–5833.
- (55) For similar observations, see: Tsantali, G. G.; Dimtsas, J.; Tsoleridis, C. A.; Takakis, I. M. *Eur. J. Org. Chem.* **2007**, *2007*, 258–265.
- (56) An alternative neutral reaction condition for conversion of isoxazoline **47** to the corresponding β -hydroxy ketone led to recovery of the starting material.⁵⁵

- (57) Kozikowski, A. P.; Adamczyk, M. *Tetrahedron Lett.* **1982**, 23, 3123–3126.
- (58) Kreiser, W.; Below, P. *Leibigs Ann. Chem.* **1985**, 1985, 203–209.
- (59) Stork, G.; Cama, L. D.; Coulson, D. R. *J. Am. Chem. Soc.* **1974**, 96, 5268–5270.
- (60) The efficiency of the reaction was highly dependent on the base used. Treatment of nitrile with KHMDs did not lead to the desired product. Deprotonation of nitrile with LDA formed the cyclized product in 60% yield.
- (61) For examples on the use of Stork epoxy-nitrile cyclization to form 3-, 4-, 5-, and 6-membered rings, see: Arseniayadis, S.; Kyler, K. S.; Watt, D. S. *Organic Reactions* **2005**, 31, 1–364.
- (62) (a) Cope, A. C.; Foster, T. T.; Towle, P. H. *J. Am. Chem. Soc.* **1949**, 71, 3929–3934. (b) DePuy, C. H.; King, R. W. *Chem. Rev.* **1960**, 60, 431–457.
- (63) The use of other halides, Br and I, as activating groups at the α -position of a 5-alkoxybutenolide led to lower yields of the desired coupling product because of the decomposition of the acceptors upon irradiation with blue LEDs.
- (64) The direct chlorination of 5-(*L*-menthyloxy)butenolide to form 3-chloro-5-(*L*-menthyloxy)butenolide in low yield is reported, see: Hui, H.; Qinghua, C. *Chin. Sci. Bull.* **2000**, 45, 711–715.
- (65) Lim, W.; Kim, J.; Rhee, Y. H. *J. Am. Chem. Soc.* **2014**, 136, 13618–13621.
- (66) Menthol-derived allene is not required for the high stereo-selectivity in the reaction, similar results were obtained with cyclohexanol-derived allene (98% yield, 95% *ee*).
- (67) Gatti, M.; Drinkel, E.; Wu, L.; Pusterla, I.; Gaggia, F.; Dorta, R. *J. Am. Chem. Soc.* **2010**, 132, 15179–15181.
- (68) Initial exploration of the ring-closing metathesis step with a diene precursor derived from cyclohexanol lacking the phenyl substituent gave low yields of the dihydrofuran product.
- (69) Fouteris, M. A.; Koutsourea, A. I.; Nikolaropoulos, S. S.; Riahi, A.; Muzart, J. J. *Mol. Catal. A: Chem.* **2006**, 250, 70–74.
- (70) For a summary of allylic oxidation conditions that were less successful, see [Supporting Information](#).
- (71) Partial epimerization of the allylic acetal stereocenter occurs under allylic oxidation reaction conditions (~10%). The menthol auxiliary is used to allow crystallization of butenolide **59** as a single diastereomer.
- (72) The expected absolute configuration of *ent*-**59** was confirmed by single-crystal X-ray diffraction analysis.⁴⁰
- (73) The formation of monoesters of oxalic acid from secondary alcohols by reaction with 2 equiv of oxalyl chloride in ether, followed by concentration to isolate the crude hemi oxalic acid and its subsequent coupling with arylhalides was reported by Zhang and MacMillan, see: Zhang, X.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2016**, 138, 13862–13865.
- (74) (a) Moradei, O. M.; Paquette, L. A. *Org. Synth.* **2003**, 80, 66–74. (b) Enantiopure samples of (*R*)- and (*S*)-enantiomers can be purchased from Proactive Molecular Research.
- (75) Narayanam, J. M. R.; Tucker, J. W.; Stephenson, C. R. J. *J. Am. Chem. Soc.* **2009**, 131, 8756–8757.
- (76) No coupling products were observed in other ethereal solvents, such as Et₂O; MTBE; 2-MeTHF; 1,4-dioxane.
- (77) Higher concentration likely promotes more efficient trapping of the tertiary radical by the acceptor *ent*-**59**, leading to an increase in yield of the reaction.
- (78) Sullivan, B.; Faulkner, D. J. *J. Org. Chem.* **1984**, 49, 3204–3206.
- (79) Houk, K. N.; Paddon-Row, M. N.; Rondan, N. G.; Wu, Y.-D.; Brown, F. K.; Spellmeyer, D. C.; Metz, J. T.; Li, Y.; Loncharich, R. J. *Science* **1986**, 231, 1108–1117.
- (80) Goswami, L. N.; Houston, Z. H.; Sarma, S. J.; Jalisatgi, S. S.; Hawthorne, M. F. *Org. Biomol. Chem.* **2013**, 11, 1116–1126. (b) Pawlak, M.; Mistlberger, G.; Bakker, E. *J. Mater. Chem.* **2012**, 22, 12796–12801.
- (81) Eey, S. T. C.; Lear, M. J. *Org. Lett.* **2010**, 12, 5510–5513.
- (82) Andrews, R. S.; Becker, J. J.; Gagné, M. R. *Angew. Chem., Int. Ed.* **2010**, 49, 7274–7276.
- (83) Coffey, D. S.; Overman, L. E.; Stappenbeck, F. J. *Am. Chem. Soc.* **2000**, 122, 4904–4914.