

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

The Environmental “Riskscape” and Social Inequality: Implications for Explaining Maternal and Child Health Disparities

### Permalink

<https://escholarship.org/uc/item/6vp0g328>

### Journal

Environmental Health Perspectives, 114(8)

### ISSN

1542-4359

### Authors

Morello-Frosch, Rachel  
Shenassa, Edmond D

### Publication Date

2006-08-01

### DOI

10.1289/ehp.8930

Peer reviewed

## The Environmental “Riskscape” and Social Inequality: Implications for Explaining Maternal and Child Health Disparities

Rachel Morello-Frosch<sup>1,2</sup> and Edmond D. Shenassa<sup>2,3</sup>

<sup>1</sup>Center for Environmental Studies; <sup>2</sup>Department of Community Health, School of Medicine; and <sup>3</sup>Centers for Behavioral and Preventive Medicine, Brown University, Providence, Rhode Island, USA

**BACKGROUND:** Research indicates that the double jeopardy of exposure to environmental hazards combined with place-based stressors is associated with maternal and child health (MCH) disparities.

**OBJECTIVE AND DISCUSSION:** Our aim is to present evidence that individual-level and place-based psychosocial stressors may compromise host resistance such that environmental pollutants would have adverse health effects at relatively lower doses, thus partially explaining MCH disparities, particularly poor birth outcomes. Allostatic load may be a physiologic mechanism behind the moderation of the toxic effect of environmental pollutants by social stressors. We propose a conceptual framework for holistic approaches to future MCH research that elucidates the interplay of psychosocial stressors and environmental hazards in order to better explain drivers of MCH disparities.

**CONCLUSION:** Given the complexity of the link between environmental factors and MCH disparities, a holistic approach to future MCH research that seeks to untangle the double jeopardy of chronic stressors and environmental hazard exposures could help elucidate how the interplay of these factors shapes persistent racial and economic disparities in MCH.

**KEY WORDS:** birth outcomes, environment, health disparities, stress. *Environ Health Perspect* 114:1150–1153 (2006). doi:10.1289/ehp.8930 available via <http://dx.doi.org/> [Online 6 April 2006]

A formidable challenge in the field of maternal and child health (MCH) has been explaining the persistent racial and socioeconomic health disparities, particularly in birth outcomes in the United States (Singh and Yu 1995). Despite declines in the overall infant mortality, there remains a significant disparity between black and white infant mortality rates. Nationally, black women are twice as likely as white women to give birth to a very low-birth-weight baby. For preterm births, although the gap between the two racial groups has narrowed recently, the disparity between the two groups remains large: a 6.7% difference (National Center for Health Statistics 2003).

Here we discuss the interplay of environmental hazards with place-based and individual-level psychosocial stressors and its implications for MCH research. Although a strong body of literature has shed much light on the individual-level factors (e.g., health behaviors, interpregnancy interval, and access to adequate health care) (Hessol et al. 1998; Hummer 1993; Rawlings et al. 1995; Starfield et al. 1991) and place-based drivers of MCH disparities (e.g., neighborhood poverty, relative income inequality, poor housing, and segregation) (Buka et al. 2002; Guest et al. 1998; Huynh et al. 2005; Laveist 1993; Matteson et al. 1998; Morenoff 2003; O’Campo et al. 1997; Shenassa et al. 2004), there has been little cross-pollination between this field and the research investigating links between environmental hazard exposures and birth outcomes (Parker et al. 2005; Ritz and Yu 1999; Ritz et al. 2000; Sadler et al. 1999; Whyatt et al. 2004; Wilhelm and Ritz 2003).

Place-based stressors are biologically relevant components of the human environment

and can function independently of individual-level stressors to determine health (Diez-Roux 1998; Shenassa 2001). These place-based factors can influence birth outcomes in three ways: *a*) by affecting birth outcomes directly (Huynh et al. 2005; Rich-Edwards and Grizzard 2005); *b*) by increasing exposures to environmental hazards, such as air pollutants (Parker et al. 2005; Woodruff et al. 2003); and *c*) by enhancing susceptibility to the toxic effects of contaminant exposures (Ponce et al. 2005). This third pathway concerning the interaction of place-based stressors with environmental hazards points toward the next generation of studies to understand the combined effects of environmental and psychosocial drivers of MCH disparities. We first discuss the confluence of place-based psychosocial stressors and environmental hazard exposures and its implications for future research on MCH disparities with a focus on birth outcomes. We then propose a possible physiologic link between place-based stressors and environmental hazards in ways that may enhance susceptibility to toxics. We conclude by outlining a conceptual framework for future MCH research.

### Social Inequality and Environmental Health Disparities

Wide-ranging political, socioeconomic, and discriminatory forces coupled with spatial patterns of industrialization and development have segregated people of color, particularly African Americans, into communities with some of the highest indices of urban poverty and material deprivation (Morello-Frosch and Jesdale 2006; Schultz et al. 2002; Williams

and Collins 2004). Researchers and policy makers concerned about environmental justice argue that communities of color and the poor face a higher frequency and magnitude of exposures to environmental as well as psychosocial stressors [Institute of Medicine (IOM) 1999; O’Neill et al. 2003]. Concern has centered on the limited science related to the cumulative impact of multiple exposures to environmental hazards and the potential vulnerability of poor communities to their toxic effects [National Environmental Justice Advisory Council (NEJAC) 2004]. This combination and potential interaction of elevated environmental hazard exposures, on the one hand, and socioeconomic stressors, on the other, have been described as a form of “double jeopardy” (IOM 1999).

Understanding the MCH implications of these “geographies of exposure and susceptibility” (Jerrett and Finkelstein 2005) or “riskscapes” (Morello-Frosch et al. 2001) requires consideration of the timing of exposure to psychosocial stressors as well as environmental hazards during the life course (e.g., during the prenatal years, infancy, adolescence, or adulthood) and socioeconomic, political, cultural, and gender dynamics. For example, the lack of child care for agricultural workers often forces families, mostly mothers, to take their children to the fields while they work, thereby increasing young children’s exposures to pesticides (Natural Resources Defense Council 1999). Many of these pesticides are known neurotoxicants and carcinogens, and the potential long-term effects of childhood and prenatal exposures are just being explored and understood (Berkowitz et al. 2004; Castorina et al. 2003; Eskenazi et al. 2004; Gladen et al. 2003; Perera et al. 2003; Torres-Arreola et al. 2003; Whyatt et al. 2004; Young et al. 2005). Similarly, neighborhood-level factors associated with

Address correspondence to R. Morello-Frosch, Center for Environmental Studies and Department of Community Health, School of Medicine, Brown University, 135 Angell St., Box 1943, Providence, RI 02912 USA. Telephone: (401) 863-9429. Fax: (401) 863-3503. E-mail: [rmf@brown.edu](mailto:rmf@brown.edu)

We thank S. Yu for her comments on an earlier version of this manuscript.

E.D.S. was supported by grant R40MC03600-01-00 from the Maternal and Child Health Bureau, Department of Health and Human Services.

The authors declare they have no competing financial interests.

Received 16 December 2005; accepted 6 April 2006.

racial residential segregation may affect health by influencing access to affordable markets with fresh fruits and vegetables and access to health services (Diez-Roux 1997; Morland et al. 2002). Women without access to adequate prenatal care, for example, are likely to have compromised nutritional status, which in turn can heighten the impact of lead exposure both *in utero* and during early childhood (Lee et al. 2005; Zierold 2004).

### State of the Evidence

Research on birth outcomes points to the validity of integrating social with environmental health risks in future MCH research (Gee and Payne-Sturges 2004; Morenoff 2003; O'Campo et al. 1997). Evidence shows a consistent relationship between residence in poverty-stricken (Collins et al. 1997; O'Campo et al. 1997; Papacek et al. 2002), segregated (Guest et al. 1998; Laveist 1993) neighborhoods and poor birth outcomes. Moreover, preliminary work suggests substantial racial and ethnic disparities in environmental hazards exposures (Centers for Disease Control and Prevention 2005; IOM 1999; Morello-Frosch 2002; NEJAC 2004), including during pregnancy (Woodruff et al. 2003), and studies have begun to link pollutant exposures and negative birth and developmental outcomes (Dejmek et al. 1999; Ritz and Yu 1999; Ritz et al. 2000, 2002). One recent study of individual factors, pollutant exposures, and neighborhood measures of socioeconomic hardship (Ponce et al. 2005) found that preterm birth risk was affected by the interaction of residential traffic-related air pollution exposure and measures of neighborhood economic hardship.

Distilling the results of this diverse body of MCH research reveals two critical paths for future inquiry. The first is the direct health effects of hazardous social and physical environments to which communities of color and the poor are disproportionately exposed. To date, MCH studies have emphasized this first line of inquiry by analyzing the effects of individual and place-based socioeconomic status (SES) stressors, on one hand, or by assessing the effect of individual factors and environmental hazards, on the other.

The second line of inquiry, as outlined below, examines all of these factors in an integrated fashion by exploring how the multi-level interplay and possible interaction of psychosocial stressors with environmental hazards may shape disparities in birth outcomes. For example, previous pollutant exposures may enhance susceptibility to the toxic effects of current pollutant exposures, particularly if the body's defense mechanisms and ability to recover or detoxify have been compromised through prior exposures to harmful agents. Similarly, exposure to place-based psychosocial stressors, such as persistent

poverty, material deprivation, and a lack of services, may lead to chronic stress, which can weaken the body's defense systems (Cohen 1999; McEwen 1998).

### Physiologic Mechanisms


The concept of allostasis provides a framework for measuring the physiologic manifestations of chronic psychosocial and environmental stressors. Allostasis refers to the ability of the body's stress-response systems to regulate internal physiology in response to psychosocial or physical stressors. The related concept of allostatic load refers to the cumulative physiologic degradation, over the life course, that can result from chronic stress exposure, and the accompanying long-term shift that occurs in the body's homeostatic functions, with harmful consequences (Geronimus 1992; McEwen 1998; Seeman et al. 1997). The physiologic effect of prolonged stressors can exact a toll on the body that is caused by chronic activation of biologic systems, such as the hypothalamic-pituitary-adrenal axis, which releases hormones (e.g., glucocorticoids) that can have several metabolic and psychological effects, including the mobilization of energy reserves and suppression of the immune system (Seeman et al. 1997; Sterling and Eyer 1988). Chronic activation of this system can lead to "wear and tear" on major organ systems (McEwen 1998).

The mechanism of allostatic load provides a potential pathway by which place-based

stressors can modify the toxic effect of environmental hazard exposures to produce disparate patterns of birth outcomes between and within populations. This is also in line with the concept of "weathering" proposed by Geronimus and others, suggesting that chronic stress associated with the combined effects of poverty, racial discrimination, and material deprivation causes the health of African-American mothers to deteriorate particularly rapidly, leading to poorer birth outcomes with increased age (Collins and Williams 1999; Geronimus 1992). The biomechanics of stress and allostasis can be considered a possible mediator of the heightened susceptibility to the adverse effects of pollution exposures observed among people with low SES.

### A Framework for Future Research

The framework in Figure 1 suggests how area-level and individual-level stressors and buffers may combine to shape environmental hazard exposures, affect individual allostatic load, and in turn enhance susceptibility to the toxic effects of pollution exposures. At the bottom of Figure 1 is a variation of the exposure-health outcome continuum that outlines how environmental toxins might cause disease (National Research Council 1991). Traditionally, the exposure-health outcome continuum includes the emission of a contaminant from an indoor source such as smoking or an outdoor source such as an industrial facility, through human



**Figure 1.** The interplay of community and individual stressors/buffers that shape exposures and susceptibility to environmental hazards. Thick arrows indicate relationships that have been studied in the epidemiologic and sociology literature; dashed arrows indicate relationships that have not been extensively explored.

exposure via various media (e.g., air), and the occurrence of an adverse health outcome (e.g., low birth weight). The framework depicted in Figure 1 implies that the presence of an environmental contaminant must first lead to human exposure and then overcome the body's defense systems to have an adverse health effect. The internal dose may not have an adverse health effect until it achieves a biologically effective dose that depends on rates of bioaccumulation, biotransformation, elimination, and, most relevant to our discussion, an individual's susceptibility.

Animal studies suggest that stress can moderate a response to environmental toxins and other agents. For example, chronic stress may increase the absorption of environmental toxicants into the body through increased respiration, consumption, or perspiration (Gordon 2003). Similarly, allostatic load may amplify susceptibility to the toxic effects of pollutants, leading to adverse birth outcomes. Indeed, stress may alter physiologic functioning, through stress-dependent hormones that can affect *in utero* development and shift the threshold for toxicity, thereby leading to adverse birth outcomes at lower exposures (Paarlberg et al. 1995). Moreover, the body's biotransformation or detoxification systems can remove or nullify toxins, but under conditions of chronic stress, the body's defense system may be impaired, resulting in compromised organ resistance. Finally, illness caused by chronic stress may compromise a sick individual's capacity to cope with environmental hazard exposures (Rios et al. 1993).

## Conclusion

Allostatic load may be a critical physiologic mechanism that explains the excess burden of adverse birth outcomes related to certain pollutants observed among low-SES populations and some communities of color. Maternal immune systems that are shaped by chronic stressors before conception and during pregnancy may enhance particular vulnerabilities to adverse pregnancy outcomes. This is compounded by race- and class-based disparities in exposures to environmental hazards that are driven by the distribution of power, privilege, and economic resources (Morello-Frosch 2002). These environmental health disparities are likely to be moderated by the degree and magnitude of chronic community and individual-level stressors that may be reflected in individuals' allostatic load. Therefore, a holistic approach to future MCH research that seeks to untangle the double jeopardy of chronic stressors and environmental hazard exposures could help elucidate how the interplay of these individual- and community-level factors shape persistent racial and economic disparities in birth outcomes. Most important, for researchers and practitioners concerned

about environmental justice, this line of inquiry could suggest new strategies for alleviating systemic drivers of racial and socioeconomic disparities in birth outcomes.

## REFERENCES

- Berkowitz GS, Wetmur JG, Birman-Deych E, Obel J, Lapinski RH, Godbold JH, et al. 2004. *In utero* pesticide exposure, maternal paraoxonase activity, and head circumference. *Environ Health Perspect* 112:388–391.
- Buka S, Brennan RT, Rich-Edwards JW, Raudenbush SW, Earls F. 2002. Neighborhood support and the birthweight of urban infants. *Am J Epidemiol* 157:1–8.
- Castorina R, Bradman A, McKone TE, Barr DB, Harnly ME, Eskenazi B. 2003. Cumulative organophosphate pesticide exposure and risk assessment among pregnant women living in an agricultural community: a case study from the CHAMACOS cohort. *Environ Health Perspect* 111:1640–1648.
- Centers for Disease Control and Prevention. 2005. Third National Report on Human Exposure to Environmental Chemicals. NCEH Pub. No. 05-0570. Atlanta, GA:Centers for Disease Control and Prevention, National Center for Environmental Health.
- Cohen S. 1999. Social status and susceptibility to respiratory infections. *Ann NY Acad Sci* 896:246–253.
- Collins C, Williams D. 1999. Segregation and mortality: the deadly effects of racism. *Social Forum* 14:495–523.
- Collins JW, Herman AA, David RJ. 1997. Very-low-birthweight infants and income incongruity among African American and white parents in Chicago. *Am J Public Health* 87:414–417.
- Dejmek J, Selevan S, Beneš I, Solansky I, Šrám R. 1999. Fetal growth and maternal exposure to particulate matter during pregnancy. *Environ Health Perspect* 107:475–480.
- Diez-Roux A. 1997. Neighborhood environments and coronary heart disease: a multilevel analysis. *Am J Epidemiol* 146:48–63.
- Diez-Roux A. 1998. Bringing context back into epidemiology: variables and fallacies in multilevel analysis. *Am J Public Health* 88:216–222.
- Eskenazi B, Harley K, Bradman A, Weltzien E, Jewell NP, Barr DB, et al. 2004. Association of *in utero* organophosphate pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect* 112:1116–1124.
- Gee G, Payne-Sturges D. 2004. Environmental health disparities: a framework integrating psychosocial and environmental concepts. *Environ Health Perspect* 112:1645–1653.
- Geronimus AT. 1992. The weathering hypothesis and the health of African-American women and infants: evidence and speculations. *Ethn Dis* 2:207–221.
- Gladen BC, Shkiryak-Nyzhnyk ZA, Chyslovska N, Zadorozhnaja TD, Little RE. 2003. Persistent organochlorine compounds and birth weight. *Ann Epidemiol* 13:151–157.
- Gordon C. 2003. Role of environmental stress in the physiological response to chemical toxins. *Environ Res* 92:1–7.
- Guest AM, Almgren G, Hussey JM. 1998. The ecology of race and socioeconomic distress: infant and working-age mortality in Chicago. *Demography* 35:23–34.
- Hessol NA, Fuentes-Afflick E, Bacchetti P. 1998. Risk of low birth weight infants among black and white parents. *Obstet Gynecol* 92:814–822.
- Hummer RA. 1993. Racial differences in infant mortality in the U.S.: an examination of social and health determinants. *Soc Forces* 72:529–554.
- Huynh M, Parker J, Harper S, Pamuk E, Schoendorf K. 2005. Contextual effect of income inequality on birth outcomes. *Int J Epidemiol* 34:888–895.
- IOM. 1999. *Toward Environmental Justice: Research, Education, and Health Policy Needs*. Washington, DC:Institute of Medicine, Committee on Environmental Justice, Health Sciences Policy Program, Health Sciences Section.
- Jerrett M, Finkelstein M. 2005. Geographies of risk in studies linking chronic air pollution exposure to health outcomes. *J Toxicol Environ Health* 68:1207–1242.
- Laveist TA. 1993. Segregation, poverty, and empowerment: health consequences for African Americans. *Milbank Q* 71:41–64.
- Lee M, Chun O, Song W. 2005. Determinants of the blood lead levels of US women of reproductive age. *J Am Coll Nutr* 24:1–9.
- Matteson DW, Burr JA, Marshall JR. 1998. Infant mortality: a multi-level analysis of individual and community risk factors. *Soc Sci Med* 47:1841–1854.
- McEwen BS. 1998. Protective and damaging effects of stress mediators. *N Engl J Med* 338:171–179.
- Morello-Frosch R. 2002. The political economy of environmental discrimination. *Environ Plann C Gov Policy* 20:477–496.
- Morello-Frosch R, Jesdale BM. 2006. Separate and unequal: residential segregation and estimated cancer risks associated with ambient air toxics in U.S. metropolitan areas. *Environ Health Perspect* 114:386–393.
- Morello-Frosch R, Pastor M, Sadd J. 2001. Environmental justice and southern California's "riskscape": the distribution of air toxics exposures and health risks among diverse communities. *Urban Affairs Rev* 36:551–578.
- Morenoff J. 2003. Neighborhood mechanisms and the spatial dynamics of birth weight. *Am J Sociol* 108:976–1017.
- Morland K, Wing S, Diez Roux A, Poole C. 2002. Neighborhood characteristics associated with the location of food stores and food service places. *Am J Prev Med* 22:23–29.
- National Center for Health Statistics. 2003. *National Vital Statistics Report*. Washington, DC:National Center for Health Statistics.
- National Research Council. 1991. *Human Exposure Assessment for Airborne Pollutants: Advances and Opportunities*. Washington, DC:National Academy Press.
- Natural Resources Defense Council. 1999. *Trouble on the Farm: Growing Up with Pesticides in Agricultural Communities*. San Francisco:Natural Resources Defense Council.
- NEJAC. 2004. *Ensuring Risk Reduction in Communities with Multiple Stressors: Environmental Justice and Cumulative Risks/Impacts—Draft Report*. New Orleans:National Environmental Justice Advisory Council.
- O'Campo P, Xue X, Wang M-C, Caughy MOB. 1997. Neighborhood risk factors for low birthweight in Baltimore: a multilevel analysis. *Am J Epidemiol* 87:1113–1118.
- O'Neill MS, Jerrett M, Kawachi I, Levy JI, Cohen AJ, Gouveia N, et al. 2003. Health, wealth, and air pollution: advancing theory and methods. *Environ Health Perspect* 111:1861–1870.
- Paarlberg KM, Vingerhoets AJ, Passchier J, Dekker GA, Van Geijn HP. 1995. Psychosocial factors and pregnancy outcome: a review with emphasis on methodological issues. *J Psychosom Res* 39:563–595.
- Papacek EM, Collins JW, Schulte NF, Goergen C, Drolet A. 2002. Differing postneonatal mortality rates of African-American and white infants in Chicago: an ecologic study. *Matern Child Health J* 6:99–105.
- Parker J, Woodruff T, Basu R, Schoendorf K. 2005. Air pollution and birth weight among term infants in California. *Pediatrics* 115:121–128.
- Perera FP, Rauh V, Tsai WY, Kinney P, Camann D, Barr D, et al. 2003. Effects of transplacental exposure to environmental pollutants on birth outcomes in a multiethnic population. *Environ Health Perspect* 111:201–205.
- Ponce N, Hoggatt K, Wilhelm M, Ritz B. 2005. Preterm birth: the interaction of traffic-related air pollution with economic hardship in Los Angeles neighborhoods. *Am J Epidemiol* 162:140–148.
- Rawlings J, Rawlings V, Read J. 1995. Prevalence of low birth weight and preterm delivery in relation to the interval between pregnancies among white and black women. *N Engl J Med* 332:69–74.
- Rich-Edwards JW, Grizzard T. 2005. Psychosocial stress and neuroendocrine mechanisms in preterm delivery. *Am J Obstet Gynecol* 192:S30–S35.
- Rios R, Poje G, Detels R. 1993. Susceptibility to environmental pollutants among minorities. *Toxicol Ind Health* 9:797–820.
- Ritz B, Yu F. 1999. The effect of ambient carbon monoxide on low birth weight among children born in southern California between 1989 and 1993. *Environ Health Perspect* 107:17–25.
- Ritz B, Yu F, Chapa G, Fruin S. 2000. Effect of air pollution on preterm birth among children born in southern California between 1989 and 1993. *Epidemiology* 11:502–511.
- Ritz B, Yu F, Chapa G, Shaw G, Harris J. 2002. Ambient air pollution and risk of birth defects in southern California. *Am J Epidemiol* 155:17–25.
- Sadler L, Belanger K, Saftlas A, Leaderer B, Hellenbrand K, McSharry J, et al. 1999. Environmental tobacco smoke exposure and small-for-gestational-age birth. *Am J Epidemiol* 150:695–705.
- Schultz A, Williams D, Israel B, Lempert L. 2002. Racial and spatial relations as fundamental determinants of health in Detroit. *Milbank Q* 80:677–707.
- Seeman TE, Singer BH, Rowe JW, Horwitz RI, McEwen BS. 1997. Price of adaptation—allostatic load and its health

- consequences. *MacArthur studies of successful aging. Arch Intern Med* 157:2259–2268.
- Shenassa E. 2001. Society, physical health and modern epidemiology. *Epidemiology* 12:467–470.
- Shenassa ED, Stubbendick A, Brown MJ. 2004. Social disparities in housing and related pediatric injury: a multilevel study. *Am J Public Health* 94:633–639.
- Singh GK, Yu SM. 1995. Infant mortality in the United States: trends, differentials, and projections, 1950 through 2010. *Am J Public Health* 85:957–964.
- Starfield B, Shapiro S, Weiss J, Liang K-Y, Ra K, Paige D, et al. 1991. Race, family income, and low birth weight. *Am J Epidemiol* 134:1167–1174.
- Sterling P, Eyer J. 1988. Allostasis: a new paradigm to explain arousal pathology. In: *Handbook of Life Stress, Cognition, and Health* (Fisher S, Reason JT, eds). Chichester, NY:Wiley, 750.
- Torres-Arreola L, Berkowitz G, Torres-Sanchez L, Lopez-Cervantes M, Cebrian ME, Uribe M, et al. 2003. Preterm birth in relation to maternal organochlorine serum levels. *Ann Epidemiol* 13:158–162.
- Whyatt RM, Rauh V, Barr DB, Camann DE, Andrews HF, Garfinkel R, et al. 2004. Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ Health Perspect* 112:1125–1132.
- Wilhelm M, Ritz B. 2003. Residential proximity to traffic and adverse birth outcomes in Los Angeles County, California, 1994–1996. *Environ Health Perspect* 111:207–216.
- Williams D, Collins C. 2004. Reparations: a viable strategy to address the enigma of African American health. *Am Behav Sci* 47:977–1000.
- Woodruff T, Parker J, Kyle A, Schoendorf K. 2003. Disparities in exposure to air pollution during pregnancy. *Environ Health Perspect* 111:942–946.
- Young JG, Eskenazi B, Gladstone EA, Bradman A, Pedersen L, Johnson C, et al. 2005. Association between in utero organophosphate pesticide exposure and abnormal reflexes in neonates. *Neurotoxicology* 26:199–209.
- Zierold K. 2004. Trends in blood lead levels among children enrolled in the special supplemental nutrition program for women, infants, and children from 1996 to 2000. *Am J Public Health* 94:1513–1515.