

UCLA

Proceedings of UCLA Health

Title

Free Wall Rupture: A Rare Complication of Myocardial Infarction

Permalink

<https://escholarship.org/uc/item/6vv78887>

Journal

Proceedings of UCLA Health, 25(1)

Author

Shah, Janki

Publication Date

2021-03-17

CLINICAL VIGNETTE

Free Wall Rupture: A Rare Complication of Myocardial Infarction

Janki Shah, MD

Clinical Presentation

An 81-year-old female was brought to the ED by paramedics for chest pain with nausea and vomiting. She was clammy and diaphoretic in the field and EKG showed ST elevations. She was given aspirin and sublingual nitroglycerin which improved her chest discomfort. Her past medical history includes hypertension, hyperlipidemia and mildly elevated coronary artery calcification score. She also had elevated A1C and mild to moderate bilateral carotid stenosis.

Initial ED vital signs included blood pressure of 96/64, pulse of 77, and O2 saturation of 99% on room air. She was alert and in no acute distress. Cardiac exam showed normal heart sounds without murmurs. Chest was clear and her skin was warm and dry. Initial labs included: WBC 13.8, creatinine of 1.3, glucose 161, BNP 325 and troponin 22.5ng/mL. EKG showed normal sinus rhythm with less than 1 mm ST elevations in the anterior leads, lateral 0.5 mm ST depressions with T wave inversions. She was taken to the cardiac catheterization laboratory, where she became unresponsive. Code Blue was called for pulseless

asystolic cardiac arrest. CPR was initiated and ACLS protocols were followed. Bedside echocardiogram showed a large pericardial effusion. She underwent emergent pericardiocentesis and 500 mL of fresh blood was removed. Despite prolonged efforts, she could not be resuscitated and was pronounced about 1 hour after code blue initiation.

Discussion

This patient had acute myocardial infarction complicated by left ventricular free wall rupture. Free wall rupture, rupture of the interventricular septum, and acute mitral regurgitation due to papillary muscle rupture are three mechanical complications of acute myocardial infarctions. Interventricular septal rupture is the most common mechanical complication in patients with ST elevation myocardial infarction (STEMI) and free wall rupture is the least common. Table 1 presents the rates of the three types of rupture in STEMI and non-STEMI (NSTEMI) patients, from the National Inpatient Sample database of 9,000,000 MI hospitalizations between 2003-2015.¹

Table 1

COMPLICATION	RATE IN STEMI	RATE IN NSTEMI
Ventricular septal rupture	0.21%	0.04%
Papillary muscle rupture	0.05%	0.01%
Free wall rupture	0.01%	0.01%

These complications can be fatal or lead to cardiogenic shock. They are caused by rupture or tearing of necrotic myocardial tissue. The incidence of these mechanical complications is approximately 3 per 1000 patients in the setting of an acute myocardial infarction in the reperfusion era, with most occurring in STEMI patients.² Prior to the widespread use of reperfusion, rupture of the free wall occurred in approximately 2 percent of acute MIs.³

Rupture of the myocardium involves the left ventricle more often than the right ventricle. Patients with had early, less than 72 hours free wall rupture, more commonly had anterior infarction sites with an abrupt tear in the infarcted myocardial

wall. Patients with late rupture, greater than 4 days from MI had infarct expansion.⁴

Risk factors for myocardial rupture, in studies from the fibrinolytic era, include: no prior history of angina or MI; ST elevation on initial EKG; and peak CK-MB level > 150 IU/L. Patients with all 3 of these risk factors had 9.2 times higher risk of myocardial rupture.⁵ Other risk factors include first MI, large transmural MIs, anterior MI, female gender, and age >70 years. Protective factors include left ventricular hypertrophy and the presence of extensive collaterals.⁶ Over 50% of free wall ruptures occur within the first 5 days after a myocardial infarction and over 90% occur within the first 14 days.⁷

Clinical presentation can be variable. Patients with a silent MI who have cardiac rupture present with sudden death. Patients with known MI can have complete rupture or incomplete/subacute rupture, which occurs when organized clot and pericardial tissue seal the perforation site.⁸ Complete rupture leads to pericardial tamponade from hemopericardium with sudden right sided congestive heart failure and shock with progression to pulseless electrical activity and death.⁹ In contrast, incomplete rupture may be present as pericardial chest pain, nausea, restlessness, acute hypotension, with or without EKG features of pericarditis.

Diagnosis is made by echocardiography which shows a large pericardial effusion and signs of tamponade. Initial treatment includes emergency pericardiocentesis, fluids, inotropic support, and vasopressor support if indicated.

REFERENCES

1. **Elbadawi A, Elgendy IY, Mahmoud K, Barakat AF, Mentias A, Mohamed AH, Ogunbayo GO, Megaly M, Saad M, Omer MA, Paniagua D, Abbott JD, Jneid H.** Temporal Trends and Outcomes of Mechanical Complications in Patients With Acute Myocardial Infarction. *JACC Cardiovasc Interv.* 2019 Sep 23;12(18):1825-1836. doi: 10.1016/j.jcin.2019.04.039. PMID: 31537282.
2. **Goldswieg AM, Wang Y, Forrest JK, Cleman MW, Minges KE, Mangi AA, Aronow HD, Krumholz HM, Curtis JP.** Ventricular septal rupture complicating acute myocardial infarction: Incidence, treatment, and outcomes among medicare beneficiaries 1999-2014. *Catheter Cardiovasc Interv.* 2018 Nov 15;92(6):1104-1115. doi: 10.1002/ccd.27576. Epub 2018 Mar 7. PMID: 29513365.
3. **Figueras J, Alcalde O, Barrabés JA, Serra V, Alguersuari J, Cortadellas J, Lidón RM.** Changes in hospital mortality rates in 425 patients with acute ST-elevation myocardial infarction and cardiac rupture over a 30-year period. *Circulation.* 2008 Dec 16;118(25):2783-9. doi: 10.1161/CIRCULATIONAHA.108.776690. Epub 2008 Dec 8. PMID: 19064683.
4. **Nakatsuchi Y, Minamino T, Fujii K, Negoro S.** Clinicopathological characterization of cardiac free wall rupture in patients with acute myocardial infarction: difference between early and late phase rupture. *Int J Cardiol.* 1994 Dec;47(1 Suppl):S33-8. doi: 10.1016/0167-5273(94)90324-7. PMID: 7737750.
5. **Pohjola-Sintonen S, Muller JE, Stone PH, Willich SN, Antman EM, Davis VG, Parker CB, Braunwald E.** Ventricular septal and free wall rupture complicating acute myocardial infarction: experience in the Multicenter Investigation of Limitation of Infarct Size. *Am Heart J.* 1989 Apr;117(4):809-18. doi: 10.1016/0002-8703(89)90617-0. PMID: 2648779.
6. **Becker RC, Hochman JS, Cannon CP, Spencer FA, Ball SP, Rizzo MJ, Antman EM.** Fatal cardiac rupture among patients treated with thrombolytic agents and adjunctive thrombin antagonists: observations from the Thrombolysis and Thrombin Inhibition in Myocardial Infarction 9 Study. *J Am Coll Cardiol.* 1999 Feb;33(2):479-87. doi: 10.1016/s0735-1097(98)00582-8. PMID: 9973029.
7. **Batts KP, Ackermann DM, Edwards WD.** Postinfarction rupture of the left ventricular free wall: clinicopathologic correlates in 100 consecutive autopsy cases. *Hum Pathol.* 1990 May;21(5):530-5. doi: 10.1016/0046-8177(90)90010-3. PMID: 2338333.
8. **McMullan MH, Maples MD, Kilgore TL Jr, Hindman SH.** Surgical experience with left ventricular free wall rupture. *Ann Thorac Surg.* 2001 Jun;71(6):1894-8; discussion 1898-9. doi: 10.1016/s0003-4975(01)02625-x. PMID: 11426765.
9. **Holger Thiele, MD, FESCJ Dawn Abbott, MD, FACC.** Acute myocardial infarction: Mechanical complications. In: *UpToDate*, Post TW (Ed), UpToDate, Waltham, MA. (Accessed November 2021).