

# UCSF

## UC San Francisco Previously Published Works

### Title

Systematic Review of the Influence of Physical Work Environment on Office Workers' Physical Activity Behavior

### Permalink

<https://escholarship.org/uc/item/6wp4620b>

### Journal

Workplace Health & Safety, 70(2)

### ISSN

2165-0799

### Authors

Michalchuk, Victoria F  
Lee, Soo-Jeong  
Waters, Catherine M  
[et al.](#)

### Publication Date

2022-02-01

### DOI

10.1177/21650799211039439

Peer reviewed



Published in final edited form as:

*Workplace Health Saf.* 2022 February ; 70(2): 97–119. doi:10.1177/21650799211039439.

## Systematic Review of the Influence of Physical Work Environment on Office Workers' Physical Activity Behavior

Victoria F. Michalchuk, RN, PhD<sup>1</sup>, Soo-Jeong Lee, RN, PhD, FAAOHN<sup>1</sup>, Catherine M. Waters, RN, PhD, FAAN<sup>1</sup>, Oi Saeng Hong, RN, PhD, FAAOHN<sup>1</sup>, Yoshimi Fukuoka, RN, PhD, FAAN<sup>1</sup>

<sup>1</sup>University of California San Francisco.

### Abstract

**Background:** Many American workers spend over 7 hours a day at work in primarily sedentary office work. Physical activity is a key aspect of optimizing health and preventing disease; yet, 80% of American adults do not meet the recommended guidelines for physical activity. In this systematic review, the relationship between physical work environment and physical activity among office workers was explored.

**Methods:** Of the 321 studies screened, 26 studies met the eligibility criteria and were included for evaluation in this systematic review.

**Results:** Of the 26 studies, four were cross-sectional studies, 14 were quasi-experimental studies, and eight were randomized control trials. Physical activity during the workday was measured using self-report surveys and electromechanical devices such as accelerometers. Physical work environments examined by the studies included different types of desks ( $n = 16$ ), office arrangements ( $n = 5$ ), and building design ( $n = 5$ ). In nine studies, office environments and building work environments designed to promote activity using active design principles such as stairs and flexible workspaces were associated with increased physical activity. Sit–stand desks reduced overall sitting time, but had a minimal effect on physical activity.

**Conclusion/Application to practice:** Offices and buildings designed for activity had the largest impact on physical activity among office workers. To increase physical activity in office workers, focus should be placed on opportunities to increase incidental movement that can

---

Address correspondence to: Victoria F. Michalchuk, RN, PhD, Department of Community Health Systems, School of Nursing, University of California San Francisco, 2 Koret Way, N505, San Francisco, CA 94143, USA; Victoria.Michalchuk@ucsf.edu.

#### Author Contributions

Dr. Victoria F. Michalchuk conceptualized and designed the study, conducted the data analysis, and wrote the manuscript. Drs. Soo-Jeong Lee, Catherine M. Waters, and Oi Saeng Hong contributed to critical revision of the manuscript. Dr. Yoshimi Fukuoka contributed to data analysis and critical revision of the manuscript. All authors approved the final version to be published and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

#### Authors' Note

The contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIOSH (National Institute for Occupational Safety & Health).

#### Conflict of Interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

#### Supplemental Material

Supplemental material for this article is available online.

increase physical activity throughout the workday. Occupational health nurses should advocate workspace designs that can increase physical activity in workers.

### Keywords

workplace physical activity; leisure-time physical activity; work environment; office environment; office workers

---

### Background

Regular physical activity is important in decreasing the risk of disease, optimizing health, and preventing chronic diseases such as type 2 diabetes, hypertension, osteoporosis, high blood cholesterol, coronary heart disease, stroke, and excess weight gain (Löllgen et al., 2009; 2018 Physical Activity Guidelines Advisory Committee, 2018; Warburton & Bredin, 2017). The physical activity guideline for Americans recommends adults perform at least 150 minutes of moderate-intensity aerobic activity per week or 75 minutes of vigorous-intensity aerobic activity a week, or a combination of the two (Piercy et al., 2018). Also recommended are muscle-strengthening activities involving all the major muscle groups at least 2 days a week. Despite the known benefits of physical activity, 80% of adults in the United States do not meet the physical activity guidelines. In the United States, estimates are that nearly US\$117 billion in annual health care costs and 10% of all premature mortality are associated with failure to meet recommended physical activity levels (Carlson et al., 2015). Furthermore, recent systematic reviews suggest that engaging in excessive sedentary behaviour increases the risk of morbidity and mortality, independent of physical activity (Ekelund et al., 2019; Ku et al., 2018).

Sedentary and light activity jobs have steadily increased over the past 60 years as the number of workers employed in service occupations that mostly entail sitting work has increased (Church et al., 2011). As of 2016, 80% of civilian jobs in the United States were considered sedentary or light work (U.S. Bureau of Labor Statistics [USBLS], 2017). In 2019, there were over 129 million full-time employed adults in the U.S. workforce (USBLS, 2019), and, on average, workers in the United States spend over 7 hours a day at their place of employment (Centers for Disease Control and Prevention, 2017). Given the rapidly increasing number of office workers who engage in longer periods of sedentary behavior, the workplace will play an important role in promoting health and preventing chronic illnesses.

Over time, if targeted attention is not given to physical activity behavior in the workplace, sitting time is estimated to increase about 2% per year, and both leisure and work time physical activity will decrease (Lindsay et al., 2016). The decrease in labor-intensive jobs paired with the decrease in leisure-time physical activity heightens the importance of understanding the influence of the physical work environment on the physical activity behavior of working adults, particularly workers with sedentary or light activity jobs. There is strong evidence that physical activity is a key aspect of optimizing health and preventing disease. For example, systematic reviews of work environments have considered the relationship between desk type and heart rate, blood pressure, cardiometabolic risk factors, body mass index (BMI), work productivity, and mood (MacEwen et al., 2015), or

open-plan workspaces work environments and health components such as sickness absences, job satisfaction, job concentration, work fatigue, or musculoskeletal disorders (De Croon et al., 2005; Richardson et al., 2017). Numerous studies have considered the work environment and reducing sedentary behavior at work (Becker et al., 2019; Chau et al., 2016; MacEwen et al., 2015; Neuhaus et al., 2014; Tew et al., 2015). However, no systematic reviews have focused on the relationship between physical activity and physical or structural office design components (desk, office layouts, floor plans, and building design) of the work environment. Therefore, the purpose of this systematic review was to determine the relationship between the physical work environment and overall physical activity, work-related physical activity, and leisure-time physical activity in office workers.

## Methods

This systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement and checklist guidelines (Moher et al., 2009).

Inclusion criteria for the systematic review were as follows: (a) studies that included office-based adult workers in the sample, (b) the setting was office-based, (c) physical work environment or office design was an independent variable, and (d) physical activity was an outcome. The physical activity measurement could be of any type and intensity of physical activity, including steps, stepping, or walking time, as long as it was assessed in an office-based setting among adult workers. The following research designs were considered for the systematic review: cross-sectional, case-control, cohort, quasi-experimental, or randomized control. Qualitative studies were excluded from this review. Publications had to be in English-language peer-reviewed journals. No time period for publication was set, which allowed for a broader scan of the literature in an understudied area of research.

## Information Sources and Search

A pre-planned systematic search strategy was developed in collaboration with a medical librarian for use with three electronic databases: PubMed, Embase, and Web of Science. In addition, a hand search and reference list review were conducted. The last search date for each database was May 1, 2019. Appendix A (Supplemental Material) shows detailed search terms for each database. The phrasing differed slightly for each database to account for official keywords, such as MeSH terms, used in each database. In summary, the MeSH and keyword search terms used for PubMed included exercise (MeSH), physical activity, sedentary behavior, workplace (MeSH), work environment, interior design and furnishings, office design (MeSH), workplace design, and sit–stand. In the PubMed search, exercise was used in addition to physical activity because it is defined as a MeSH term within PubMed and yielded a higher quantity of relevant articles. Search terms used for Embase and Web of Science included exercise, physical activity, sitting, standing, sedentary time, workplace, work environment, office worker, workstation, office, interior design and furnishings, office design, and workplace design. To increase the sensitivity of the search, both physical activity and sedentary behavior were included in the search terms.

## Selection Process

The retrieved articles were imported into EndNote reference management software (The EndNote Team, 2013), duplicates were removed, and then the remaining articles were uploaded into Covidence systematic review software, which is recommended by Cochrane (Veritas Health Innovation, 2013). In the first phase of screening, the first researcher (V.F.M.) assessed study titles and abstracts against the eligibility criteria. For the titles and abstracts that did not contain information about the specific study population (i.e., office workers) and the phenomenon of interest (i.e., physical activity), the researcher (V.F.M.) reviewed full text to determine its eligibility. The selected studies were independently reviewed by the second researcher (Y.F.). Discrepancies were resolved through discussion between the two researchers (V.F.M. and Y.F.).

## Data Collection Process, Data Items, and Synthesis of Results

The following information was extracted from each of the studies included in the systematic review: study design, aims, sample characteristics (including location, sample size, age, and gender), intervention (if any), overall physical activity (physical activity measured all day), physical activity at work, and study limitations. Natural experiment studies and intervention studies that involved non-randomized pre- and post-comparisons without a control group were considered quasi-experimental studies. Physical activity measures included self-report surveys and electromechanical devices such as questionnaires and accelerometers. Because of the heterogeneity in types of study outcomes, outcome measures, and study designs, we qualitatively evaluated and synthesized the results of the studies. We did not conduct a meta-analysis and did not assess publication bias.

## Risk of Bias Within Studies

The risk of bias for each study included in the systematic review was assessed using the Joanna Briggs Appraisal Tool (Moola et al., 2020; Tufanaru et al., 2017). This assessment tool was designed to identify potential risk of bias within studies. A study was classified as “minimal risk” if there were “yes” answers to 90% or greater to the tool’s questions. A study was classified as “moderate risk” if there were “yes” answers to 50% to 89% of the tool’s questions

## Results

The study selection process is presented in Figure 1. The initial search yielded 493 records. After removing duplicates, the titles and abstracts of 321 records were reviewed; 285 records did not meet at least one of the eligibility criteria, yielding 36 records eligible for the next screening stage. Of these, 10 studies that did not meet all the eligibility criteria were excluded, yielding a total of 26 studies for the systematic review. Studies were excluded for not specifying the type of work environment examined in the study ( $n = 1$ ); only describing the study protocol, but no results ( $n = 3$ ); not including physical activity as a study variable ( $n = 3$ ); and not including work environment as a study variable ( $n = 3$ ).

## Study Characteristics

The 26 studies included in the systematic review were published between 2012 and 2019. Eight studies were randomized control trials, 14 were quasi-experimental studies, and four were cross-sectional studies. See Table 1 for sample characteristics, study methods, and description of interventions (when conducted) of the studies. The studies were conducted in several countries: the United States ( $n = 9$ ), Australia ( $n = 8$ ), Europe ( $n = 6$ ), Japan ( $n = 1$ ), Canada ( $n = 1$ ), and New Zealand ( $n = 1$ ). Sample sizes ranged from 11 to 1,098; 69% ( $n = 18$ ) had a sample size between 11 and 49, 12% ( $n = 3$ ) had a sample size between 50 and 99, and 19% ( $n = 5$ ) had a sample size of 100 or greater. Study participants were mostly in middle adulthood; the mean age ranged from 32 to 51 years. A majority of the samples consisted of participants who were female and had a university education or higher. The studies took place at a variety of workplaces, with college/university being the most common work setting ( $n = 7$ ). The most common type of physical work environment design at baseline was seated desks ( $n = 17$ ), followed by sit–stand desk ( $n = 3$ ) and unassigned or open desk ( $n = 3$ ).

Of the 26 studies, 22 were intervention studies: eight randomized control trials and 14 quasi-experimental studies. The three types of physical work environment designs identified in the 22 intervention studies were desk type ( $n = 14$ ), office type ( $n = 4$ ), and building design ( $n = 4$ ). The 14 desk-type intervention studies examined sit–stand desk ( $n = 8$ ), set height standing desk ( $n = 1$ ), treadmill desk ( $n = 4$ ), and both sit–stand and treadmill desks ( $n = 1$ ).

The four office-type intervention studies examined office arrangement and office layout configurations (Candido et al., 2019; Maylor et al., 2018; Wahlstrom et al., 2019; Wallmann-Sperlich et al., 2019). These studies reviewed spatial design characteristics such as how workstations were placed within the office space (i.e., assigned vs. unassigned workstations), and how supplies such as trash cans and printers were arranged throughout the office. The setting for these studies included private offices, cubicles, and open neighborhood. The four building design intervention studies compared the influence of office building design (active building vs. traditional) on office workers' physical activity and sedentary behavior (Engelen et al., 2016; Eyler et al., 2018; Gorman et al., 2013; Jancey et al., 2016).

The duration of the 22 intervention studies ranged from 5 days to 18 months, but the vast majority of studies were between 1 and 6 months. The majority of these studies ( $n = 19$ ) implemented interventions during all workdays. In one study on treadmill desk (Schuna et al., 2014), the frequency of the intervention was twice daily. In two studies on using a standing desk intervention (Miyachi et al., 2015) and treadmill desk intervention (Malaeb et al., 2019), participants utilized the intervention during the workday at their discretion. A majority of the desk intervention studies used a sitting desk for the comparison group. In one study, sit–stand desk was the control activity that was compared with the treadmill desk intervention (Bergman et al., 2018). The office arrangement and building design intervention studies used previous work setting conditions or “traditional offices” as control activities.

Four studies were non-intervention studies using a descriptive, correlational cross-sectional research design (Carr et al., 2016; Lindberg et al., 2018; McGann et al., 2015; Renaud et al., 2018). These studies focused on three types of physical work environment designs: desk type, office type, and building design. In these studies, the association with workers' physical activity level was examined on the type of sit–stand desk (Carr et al., 2016), sit–stand desk usage (Renaud et al., 2018), different corridor and staircase designs (McGann et al., 2015), and office arrangements (Lindberg et al., 2018). In their analyses, three studies did not control for any confounding factors and only Lindberg et al. (2018) controlled for gender and work type, defined as self-reported computer dominant job or non-computer dominant job.

### Physical Activity and Sedentary Behavior Measurements

Table 2 lists the physical activity measurement tools used in the 26 studies included in this systematic review. Three studies used only subjective self-report measures (Engelen et al., 2016; Renaud et al., 2018; Wallmann-Sperlich et al., 2019), and 13 studies used only objective measures (Candido et al., 2019; Carr et al., 2016; Gilson et al., 2012; Gorman et al., 2013; Koepp et al., 2013; Lindberg et al., 2018; Mansoubi et al., 2016; Maylor et al., 2018; Miyachi et al., 2015; Schuna et al., 2014; Tobin et al., 2016; Wahlstrom et al., 2019; Zhu et al., 2018). Both subjective and objective physical activity measures were used simultaneously in 10 studies (Bergman et al., 2018; Chau et al., 2014, 2016; Dutta et al., 2014, 2019; Eyler et al., 2018; Jancey et al., 2016; Malaeb et al., 2019; McGann et al., 2015; Pierce et al., 2019). Subjective physical activity was measured using the Occupational Sitting and Physical Activity Questionnaire (OSPAQ) in five studies and the International Physical Activity Questionnaire (IPAQ) in two studies. Other self-reported physical activity measures included the Active Australia Questionnaire, the Baecke Questionnaire for Habitual Physical Activity, the Workforce Sitting Questionnaire (WSQ), and the Marshall Sitting Questionnaire. A single item question was also used to measure physical activity.

Objective physical activity was measured using one or more electromechanical devices. Five studies used only the ActivPAL accelerometer (Chau et al., 2014; Eyler et al., 2018; Gorman et al., 2013; Maylor et al., 2018; Tobin et al., 2016), three studies used only the ActiGraph accelerometer to measure physical activity (Jancey et al., 2016; McGann et al., 2015; Schuna et al., 2014), and four studies used a combination of the two electromechanical devices to measure physical activity (Bergman et al., 2018; Chau et al., 2016; Mansoubi et al., 2016; Wahlstrom et al., 2019). Nine studies used other types of electromechanical devices: Fitbit Charge 2 accelerometer, Modular Signal Recorder accelerometer, Gruve accelerometer, Armband accelerometer by SenseWear, Actical accelerometer, EcgMove3 accelerometer, Actimarker accelerometer, or a Keep Walking–Stay Fit pedometer. Of the 26 studies, 24 studies measured sedentary behavior in addition to physical activity behavior.

### Randomized Control Intervention Study Findings

Among the eight randomized control trials, findings were reported on overall physical activity in two studies (Bergman et al., 2018; Miyachi et al., 2015), physical activity at work in two studies (Chau et al., 2014; Tobin et al., 2016), and both overall and at work physical

activity in four studies (Dutta et al., 2014; Maylor et al., 2018; Pierce et al., 2019; Schuna et al., 2014). Four sit–stand desk intervention studies found that providing sit–stand desks had a little effect on workers’ overall or work-related physical activity when compared with traditional-sitting desks (Chau et al., 2014; Dutta et al., 2014; Pierce et al., 2018; Tobin et al., 2016). Results ranged from a 13-minute/day increase in stepping time at work (Chau et al., 2014) to a 2.1-minute/8-hour workday increase in stepping time (Tobin et al., 2016). The sit–stand desk interventions did significantly decreased workers’ overall sitting time. Compared with the traditional desk groups, the net reduction in sitting time during the workday ranged from 4.8 minutes/hour (Dutta et al., 2014) to 99.9 minutes/day (Tobin et al., 2016). In Dutta et al.’s (2014) study. Miyachi and colleagues (2015) found a significant increase in overall time spent in light physical activity in the standing desk intervention group compared with the traditional-sitting desk group.

Workers using a treadmill desk as the intervention in two randomized control trials resulted in statistically significant increases in light physical activity (Bergman et al., 2018; Schuna et al., 2014). Compared with workers in the sit–stand desk group, workers in the treadmill desk intervention group engaged in walking for additional 22 minutes/day (Bergman et al., 2018). In treadmill desk users compared with sitting desk users, the net significant increase in overall light physical activity at work was 1.6 and 2.9 minutes/hour (Schuna et al., 2014). However, there were no significant changes in moderate- or vigorous-intensity physical activity among workers using a treadmill desk in any of the randomized control studies. As compared with the sitting desk group, workers in a multicomponent intervention that incorporated environmental changes to the office layout significantly increased their stepping time at work by 12 minutes/day (Maylor et al., 2018). No significant changes, however, were found in overall stepping time, overall physical activity, overall sitting time, or sitting time at work between control and intervention group participants.

### Quasi-Experimental Intervention Study Findings

Among the 14 quasi-experimental studies included in the systematic review, overall physical activity was reported in one study (Malaeb et al., 2019), work-related physical activity was reported in 11 studies (Candido et al., 2019; Chau et al., 2016; Dutta et al., 2019; Eyler et al., 2018; Gilson et al., 2012; Gorman et al., 2013; Jancey et al., 2016; Mansoubi et al., 2016; Wahlstrom et al., 2019; Wallmann-Sperlich et al., 2019; Zhu et al., 2018), and both overall and work-related physical activity were reported in two studies (Engelen et al., 2016; Koeppe et al., 2013). Sit–stand desk intervention studies found no significant effect on office workers’ stepping time, light physical activity, or moderate to vigorous levels of overall or work-related physical activity (Chau et al., 2016; Gilson et al., 2012; Mansoubi et al., 2016; Zhu et al., 2018). In one study, there was a significant decrease in sitting time at work (Mansoubi et al., 2016).

Treadmill desk interventions were found to significantly increase office workers’ walking time at work and decrease sedentary behaviors in the short and long term. Koeppe et al. (2013) found that at 12-month follow-up, workers in the treadmill desk intervention group increased the average walking time at work from 70 to 109 minutes/workday and decreased the average daily sedentary time by 43 minutes/workday. In another study using a treadmill



desk intervention, the intervention increased the overall step count among office workers from the baseline assessment (Malaeb et al., 2019), but the researchers did not report the  $p$  value. Zhu et al. (2018) found that a treadmill desk intervention decreased workers' average sitting time by 53 minutes/workday at 18 months post-intervention.

Among the three studies that used office design modifications as interventions, two studies found significant effects on workers' physical activity. In Wahlstrom et al.'s (2019) study, workers in flex offices significantly increased their walking time at work from 39 minutes/workday at baseline to 47 minutes/workday as well as moderate- to vigorous-intensity physical activity at work from 19 minutes/workday at baseline to 27 minutes/day at 18 months postintervention. Wallmann-Sperlich et al. (2019) also found a significant decrease in average sitting time at work after 7 months of workers participating in an office design modification intervention that included adding sit-stand desk, 26 treadmill desks, sit-stand meeting space, shared trash bins, and sit-stand break tables.

The four quasi-experimental studies that used building design interventions showed a significant increase in workers' light physical activity (Engelen et al., 2016; Eyler et al., 2018; Gorman et al., 2013; Jancey et al., 2016). The average minutes spent in light activities at work increased from 35 minutes/workday at baseline to 57 minutes/workday post-intervention in the study by Jancey et al. (2016). On the contrary, none of the studies found a significant change in time steps at work or in overall time spent engaging in moderate- to vigorous-intensity physical activity. There, however, were increasing trends in stepping time at work, time spent engaging in moderate-or vigorous intensity physical activity at work, and total average steps per day for office workers in the intervention group (Eyler et al., 2018; Gorman et al., 2013; Jancey et al., 2016).

### Non-Intervention Study Findings

Among the four cross-sectional studies, one study reported findings on overall physical activity and work-related physical activity (Renaud et al., 2018), and three studies reported findings on only work-related physical activity (Carr et al., 2016; Lindberg et al., 2018; McGann et al., 2015). Two studies examined the relationship between having sit-stand desks and workers' physical activity behavior. In a study by Carr and colleagues (2016), using a sit-stand desk, compared with sitting desks, was significantly associated with increased standing time at work and decreased sitting time at work, but not associated with walking time at work.

Renaud et al. (2018) found that walking time at work was greater in employees who used their sit-stand desk more often (less than once per week, but at least once a month; once or twice per week; 3-4 times per week; once or twice per day; 3 or more times per day) than those who did not utilize the sit-stand desk features. Sit-stand desk users also met the physical activity guidelines (moderate- to vigorous-physical activity 150 minutes per week) more often than workers who did not use the sit-stand desk features. In the studies that compared the office or building design floor plan, physical activity time at work was greater among employees working in buildings with accessible stairwells compared with buildings without accessible stairwells (McGann et al., 2015) and also greater in flex office spaces compared with private or cubical cell offices (Lindberg et al., 2018).

## Risk of Bias

Tables 3, 4, and 5 present the summary of the risk of bias organized by study design type based on the Joanna Briggs Critical Appraisal Tool for Bias (Moola et al., 2020; Tufanaru et al., 2017). The eight randomized control studies had a moderate risk of bias because allocation to the intervention groups could not be concealed nor could participants be blinded to their intervention assignment. Tobin et al. (2016) did not describe the study's randomization procedure determining how participants were chosen for the study, nor did they describe how study participants were assigned the sit–stand desk intervention. Among the 14 quasi-experimental studies, 12 studies had a minimal risk of bias and two studies had a moderate risk of bias. Eleven of the 14 quasi-experimental studies conducted pre–post comparisons and did not have a control group. Engelen et al. (2016) and Eyler et al. (2018) did not compare the baseline buildings for similarities or differences in desk type, square footage, stairwells, or amenities in the pre-phase before group moved to their new work environments (post-phase). Among the four cross-sectional studies, three had a minimal risk of bias and one study that did not measure how long employees used a sit–stand desk had a moderate risk of bias (Renaud et al., 2018).

## Discussion

This systematic review aimed to identify the relationship between the physical work environment and overall physical activity, work-related physical activity, or leisure-time physical activity in office workers. After reviewing 26 studies, this systematic review found that work environments built with active design principles are the most likely to result in increasing workers' physical activity at work. Participants in work environments with flexible space and open floor plans with active design building principles spent more time walking and engaging in light physical activity at work than those in traditional spaces (Candido et al., 2019; Eyler et al., 2018; Gorman et al., 2013; Jancey et al., 2016; Wahlstrom et al., 2019). Office workers in these environments were consistently the most physically active at work even after 12 months or longer follow-up periods (Eyler et al., 2018; Wahlstrom et al., 2019).

“Active design” is a newer building design concept that includes environmental and structural design, policy, and workplace culture to create an environment that promotes physical activity, promotes active living, and improves the quality of life of building occupants (Center for Active Design, 2010). The building design encourages movement by including features such as central staircases; shared and centralized facilities such as break rooms, bathrooms, printers, and trash cans; and shared and diverse workspaces for sitting and standing work. Our review results align with a previous review showing that programs promoting incidental physical activity within and around the workplace had the strongest potential to increase physical activity of workers (Marshall, 2004). A recent systematic review of workplace physical activity interventions in working adults found that lifestyle-based interventions to increase physical activity had lack of compliance and low participation (Mulchandani et al., 2019). Unlike sit–stand or treadmill desk-based interventions that require participant adherence, office arrangement and building designs

with active design guidelines focus on providing more opportunities for incidental activity, and therefore encourage more movement and less sitting (Center for Active Design, 2010).

Another noteworthy finding of this systematic review is the distinction between physical activity and sedentary behavior. This systematic review was focused on changes in physical activity; however, 23 of the 26 studies assessed sedentary behaviors among office workers. This pattern aligns with the literature; many previous intervention studies measured sedentary time or sedentary behavior as a primary outcome and physical activity as a secondary outcome (MacDonald et al., 2018; Prince et al., 2019). However, sedentary behavior and physical activity are two independent concepts that are related but not interchangeable (Thivel et al., 2018). The results of this systematic review showed that a desk-type intervention had the greatest impact on decreasing sitting time, which aligned with previous systematic reviews of adjustable workstation desk (Chau et al., 2016; Neuhaus et al., 2014; Tew et al., 2015). However, this systematic review result found that desk had little effect on increasing physical activity. Although changing a worker's desk can reduce sitting time, changes made to desk and workstations alone may not simultaneously change physical activity behavior.

The findings of this systematic review highlight a wide range of physical activity measures used and variations in reporting of these outcomes across the 26 studies. Physical activity was measured using 17 unique methods in the studies included in this systematic review. More than half of the studies ( $n = 15$ ) only measured work-related physical activity. In addition, the data analysis methods varied across the studies: Some studies reported the percentage of time in work-related, leisure, or overall physical activity; others reported minutes per day or minutes per workday of work-related, leisure, or overall physical activity, while others reported stepping time or step counts. In addition, some studies used physical activity intensity categories such as light physical activity or moderate physical activity, while others used walking. These variations in physical activity measurement and physical activity reporting make it difficult to compare study results.

### Strengths and Limitations

This systematic review examined the physical workplace as the phenomenon of interest in relation to workers' physical activity. The strengths included a comprehensive search strategy developed with a research librarian as well as the inclusion of work-related, leisure-time, and overall physical activities to examine a more holistic understanding of physical activity in office workers. Despite the strengths of this systematic review, several limitations need to be acknowledged. First, this systematic review only searched three databases, gray literature was not searched, and non-English studies were excluded. Thus, there may be additional studies that were not included in this review, specifically white papers that exist in the building industry. Second, only eight of the 26 studies reviewed were randomized control trials. Although the overall quality of studies in this review was strong, based on the level of evidence the authors cannot confirm causality between physical work environment and office workers' physical activity behavior. After critical appraisal of all studies in this review, the overall quality of the evidence is strong. Given the nature of desk and office

design physical environment intervention research, blinding researchers or participants is not feasible and quasi-experimental studies are more common and practical.

### **Implications for Occupational Health**

Occupational health nurses and program managers have the opportunity to positively influence the work environment to promote regular physical activity of workers and prevent chronic diseases. Occupational health nurses should be aware of the important role of the physical work environment in physical activity behavior among workers. To increase physical activity in office workers or low activity occupations, the focus must shift from limiting sedentary behavior to increasing activity throughout the day. The findings from this review suggest that workplace wellness programs should target how the office space is built as well as encouraging individual physical activity behaviors to be the most effective. To achieve this level of health promotion, occupational health nurses must engage with organization leaders to gain business support and company-level policy change.

### **Conclusion**

The results of this systematic review indicate that physical work environments built with active design principles are the most effective in increasing workers' physical activity. This review also identified that many studies did not assess physical activity outside of work time, and thus, the relationship between the physical work environment and workers' overall physical activity level is unclear in the current literature. Future research is needed to determine the effect of activity design office environments on overall total physical activity in office workers. The findings from this systematic review will help shape evidence-based solutions that can increase physical activity while reducing sedentary time in office workers.

### **Supplementary Material**

Refer to Web version on PubMed Central for supplementary material.

### **Acknowledgment**

The authors acknowledge Min-Lin Fang, MLIS, Research Librarian, at Academic Health Sciences Library, University of California, San Francisco, for her assistance in developing search strategies and literature retrieval.

### **Funding**

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Victoria F. Michalchuk was supported in part by the NIOSH's (National Institute for Occupational Safety & Health) Traineeship (Grant Number T42 OH008429).

### **Author Biographies**

Victoria F. Michalchuk, RN, PhD, is a doctoral graduate of University of California, San Francisco, School of Nursing. Her research focuses on the effects of workplace characteristics and physical activity in workers.

Soo-Jeong Lee, RN, PhD, FAAOHN, is an associate professor in the School of Nursing at University of California, San Francisco.

Catherine M. Waters, RN, PhD, FAAN, is a professor and associate dean in the School of Nursing at University of California, San Francisco.

Oi Saeng Hong, RN, PhD, FAAOHN, is a professor and director of Occupational and Environmental Health Nursing Program, Northern California NIOSH Education and Research Center.

Yoshimi Fukuoka, RN, PhD, FAAN, is a professor in the School of Nursing at University of California, San Francisco.

## References


- Becker I, Wallmann-Sperlich B, Rupp R, & Bucksch J. (2019). [Workplace interventions to reduce sedentary behavior: A systematic review]. *Gesundheitswesen*, 81(08–09), 606–614. 10.1055/s-0043-112746 [PubMed: 29108081]
- Bergman F, Wahlstrom V, Stomby A, Otten J, Lanthen E, Renklint R, Waling M, Sörlin A, Boraxbekk C-J, Wennberg P, Öhberg F, Levine JA, & Olsson T. (2018). Treadmill workstations in office workers who are overweight or obese: A randomised controlled trial. *Lancet Public Health*, 3(11), e523–e535. 10.1016/s2468-2667(18)30163-4 [PubMed: 30322782]
- Candido C, Thomas L, Haddad S, Zhang F, Mackey M, & Ye W. (2019). Designing activity-based workspaces: Satisfaction, productivity and physical activity. *Building Research and Information*, 47(3), 275–289. 10.1080/09613218.2018.1476372
- Carlson SA, Fulton JE, Pratt M, Yang Z, & Adams EK (2015). Inadequate physical activity and health care expenditures in the United States. *Progress in Cardiovascular Diseases*, 57(4), 315–323. 10.1016/j.pcad.2014.08.002 [PubMed: 25559060]
- Carr LJ, Swift M, Ferrer A, & Benzo R. (2016). Cross-sectional examination of long-term access to sit-stand desks in a professional office setting. *American Journal of Preventive Medicine*, 50(1), 96–100. 10.1016/j.amepre.2015.07.013 [PubMed: 26437867]
- Center for Active Design. (2010). What is active design? <https://centerforactivedesign.org/WhatsActiveDesign/>
- Centers for Disease Control and Prevention. (2017). Worksite physical activity. <https://www.cdc.gov/physicalactivity/worksite-pa/index.htm>
- Chau JY, Daley M, Dunn S, Srinivasan A, Do A, Bauman AE, & Van der Ploeg HP (2014). The effectiveness of sit-stand workstations for changing office workers' sitting time: Results from the Stand@Work randomized controlled trial pilot. *International Journal of Behavioral Nutrition and Physical Activity*, 11, Article 127. 10.1186/s12966-014-0127-7 [PubMed: 25291960]
- Chau JY, Sukala W, Fedel K, Do A, Engelen L, Kingham M, Sainsbury A, & Bauman AE (2016). More standing and just as productive: Effects of a sit-stand desk intervention on call center workers' sitting, standing, and productivity at work in the Opt to Stand pilot study. *Preventive Medicine Reports*, 3, 68–74. 10.1016/j.pmedr.2015.12.003 [PubMed: 26844191]
- Church TS, Thomas DM, Tudor-Locke C, Katzmarzyk PT, Earnest CP, Rodarte RQ, Martin CK, Blair SN, & Bouchard C. (2011). Trends over 5 decades in U.S. occupation-related physical activity and their associations with obesity. *PLOS ONE*, 6(5), Article e19657. 10.1371/journal.pone.0019657
- De Croon EM, Sluiter JK, Kuijer PP, & Frings-Dresen MH (2005). The effect of office concepts on worker health and performance: A systematic review of the literature. *Ergonomics*, 48(2), 119–134. 10.1080/00140130512331319409 [PubMed: 15764312]
- Dutta N, Koepp GA, Stovitz SD, Levine JA, & Pereira MA (2014). Using sit-stand workstations to decrease sedentary time in office workers: A randomized crossover trial. *International Journal of Environmental Research and Public Health*, 11(7), 6653–6665. 10.3390/ijerph110706653 [PubMed: 24968210]
- Dutta N, Walton T, & Pereira MA (2019). One-year follow-up of a sit-stand workstation intervention to decrease sedentary time in office workers. *Preventive Medicine Reports*, 13, 277–280. 10.1016/j.pmedr.2019.01.008 [PubMed: 30723663]

- Ekelund U, Tarp J, Steene-Johannessen J, Hansen BH, Jefferis B, Fagerland MW, Whincup P, Diaz KM, Hooker SP, Chernofsky A, Larson MG, Spartano N, Vasani RS, Dohrn IM, Hagströmer M, Edwardson C, Yates T, Shiroma E, Anderssen SA, & Lee I-M (2019). Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: Systematic review and harmonised meta-analysis. *British Medical Journal*, 366, Article 14570. 10.1136/bmj.14570
- Engelen L, Dhillon HM, Chau JY, Hespe D, & Bauman AE (2016). Do active design buildings change health behaviour and workplace perceptions? *Occupational Medicine*, 66(5), 408–411. 10.1093/occmed/kqv213 [PubMed: 26769894]
- Eyler AA, Hipp A, Valko CA, Ramadas R, & Zwald M. (2018). Can building design impact physical activity? A natural experiment. *Journal of Physical Activity & Health*, 15(5), 355–360. 10.1123/jpah.2017-0319 [PubMed: 29493373]
- Gilson N, Straker L, & Parry S. (2012). Occupational sitting: Practitioner perceptions of health risks, intervention strategies and influences. *Health Promotion Journal of Australia*, 23(3), 208–212. [PubMed: 23540321]
- Gorman E, Ashe MC, Dunstan DW, Hanson HM, Madden K, Winkler EA, McKay HA, & Healy GN (2013). Does an “activity-permissive” workplace change office workers’ sitting and activity time? *PLOS ONE*, 8(10), Article e76723. 10.1371/journal.pone.0076723
- Jancey JM, McGann S, Creagh R, Blackford KD, Howat P, & Tye M. (2016). Workplace building design and office-based workers’ activity: A study of a natural experiment. *Australian and New Zealand Journal of Public Health*, 40(1), 78–82. 10.1111/1753-6405.12464 [PubMed: 26455349]
- Koepp GA, Manohar CU, McCrady-Spitzer SK, Ben-Ner A, Hamann DJ, Runge CF, & Levine JA (2013). Treadmill desks: A 1-year prospective trial. *Obesity (Silver Spring)*, 21(4), 705–711. 10.1002/oby.20121 [PubMed: 23417995]
- Ku PW, Steptoe A, Liao Y, Hsueh MC, & Chen LJ (2018). A cut-off of daily sedentary time and all-cause mortality in adults: A meta-regression analysis involving more than 1 million participants. *BMC Medicine*, 16(1), Article 74. 10.1186/s12916-018-1062-2 [PubMed: 29793552]
- Lindberg CM, Srinivasan K, Gilligan B, Razjouyan J, Lee H, Najafi B, Canada KJ, Mehl MR, Currim F, Ram S, Lunden M, Heerwagen JH, Kampschroer K, & Sternberg EM (2018). Effects of office workstation type on physical activity and stress. *Occupational & Environmental Medicine*, 75(10), 689–695. 10.1136/oemed-2018-105077 [PubMed: 30126872]
- Lindsay DB., Devine S., Sealey RM., & Leicht AS. (2016). Time kinetics of physical activity, sitting, and quality of life measures within a regional workplace: A cross-sectional analysis. *BMC Public Health*, 16, Article 786. 10.1186/s12889-016-3487-x [PubMed: 27527583]
- Löllgen H, Böckenhoff A, & Knapp G. (2009). Physical activity and all-cause mortality: An updated meta-analysis with different intensity categories. *International Journal of Sports Medicine*, 30(3), 213–224. 10.1055/s-0028-1128150 [PubMed: 19199202]
- MacDonald B, Janssen X, Kirk A, Patience M, & Gibson AM (2018). An integrative, systematic review exploring the research, effectiveness, adoption, implementation, and maintenance of interventions to reduce sedentary behaviour in office workers. *International Journal of Environmental Research and Public Health*, 15(12), Article 2876. 10.3390/ijerph15122876 [PubMed: 30558258]
- MacEwen BT, MacDonald DJ, & Burr JF (2015). A systematic review of standing and treadmill desks in the workplace. *Preventive Medicine*, 70, 50–58. 10.1016/j.ypmed.2014.11.011 [PubMed: 25448843]
- Malaeb S, Perez-Leighton CE, Noble EE, & Billington C. (2019). A “NEAT” approach to obesity prevention in the modern work environment. *Workplace Health & Safety*, 67(3), 102–110. 10.1177/2165079918790980 [PubMed: 30370831]
- Mansoubi M, Pearson N, Biddle SJ, & Clemes SA (2016). Using sit-to-stand workstations in offices: Is there a compensation effect? *Medicine & Science in Sports & Exercise*, 48(4), 720–725. 10.1249/MSS.0000000000000802 [PubMed: 26496419]
- Marshall AL (2004). Challenges and opportunities for promoting physical activity in the workplace. *Journal of Science and Medicine in Sport*, 7(1 Suppl.), 60–66. 10.1016/s1440-2440(04)80279-2 [PubMed: 15214603]

- Maylor BD, Edwardson CL, Zakrzewski-Fruer JK, Champion RB, & Bailey DP (2018). Efficacy of a multicomponent intervention to reduce workplace sitting time in office workers a cluster randomized controlled trial. *Journal of Occupational and Environmental Medicine*, 60(9), 787–795. 10.1097/jom.0000000000001366 [PubMed: 29851741]
- McGann S, Creagh R, Tye M, Jancey J, Pages-Oliver R, & James H. (2015). Stairway to health: An analysis for workplace stairs design and use. University of Melbourne.
- Miyachi M, Kurita S, Tripette J, Takahara R, Yagi Y, & Murakami H. (2015). Installation of a stationary high desk in the workplace: Effect of a 6-week intervention on physical activity. *BMC Public Health*, 15, Article 368. 10.1186/s12889-015-1724-3 [PubMed: 25886894]
- Moher D, Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Loannidis JP, Clarke M, Devereaux PJ, & Kleijnen J. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: Explanation and elaboration. *British Medical Journal*, 339, Article b2700.
- Moola S, Munn Z, Tufanaru C, Aromataris E, Sears K, Sfetcu R, Currie M, Qureshi R, Mattis P, Lisy K, & Mu P-F. (2020). Chapter 7: Systematic reviews of etiology and risk. In Aromataris E. & Munn Z. (Eds.), *JBIManual for Evidence Synthesis*. JBI, 2020. <https://synthesismanual.jbi.global>
- Mulchandani R, Chandrasekaran AM, Shivashankar R, Kondal D, Agrawal A, Panniyammakal J, Tandon N, Prabhakaran D, Sharma M, & Goenka S. (2019). Effect of workplace physical activity interventions on the cardio-metabolic health of working adults: Systematic review and meta-analysis. *International Journal of Behavioral Nutrition and Physical Activity*, 16(1), Article 134. 10.1186/s12966-019-0896-0 [PubMed: 31856826]
- Neuhaus M, Eakin EG, Straker L, Owen N, Dunstan DW, Reid N, & Healy GN (2014). Reducing occupational sedentary time: A systematic review and meta-analysis of evidence on activity-permissive workstations. *Obesity Reviews*, 15(10), 822–838. 10.1111/obr.12201 [PubMed: 25040784]
- Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, Galuska DA, George SM, & Olson RD (2018). The Physical Activity Guidelines for Americans. *JAMA*, 320, 2020–2028. 10.1001/jama.2018.14854 [PubMed: 30418471]
- Pierce J, Legg S, Godfrey JR, & Kawabata E. (2019). The effects of introducing electric adjustable height desks in an office setting on workplace physical activity levels: A randomised control field trial. *Work: A Journal of Prevention Assessment & Rehabilitation*, 62(1), 139–150. 10.3233/wor-182849 [PubMed: 30689597]
- Prince SA, Elliott CG, Scott K, Visintini S, & Reed JL (2019). Device-measured physical activity, sedentary behaviour and cardiometabolic health and fitness across occupational groups: A systematic review and meta-analysis. *International Journal of Behavioral Nutrition and Physical Activity*, 16(1), Article 30. 10.1186/s12966-019-0790-9 [PubMed: 30940176]
- Renaud LR, Huysmans MA, Van der Ploeg HP, Speklé EM, & Van der Beek AJ (2018). Long-term access to sit-stand workstations in a large office population: User profiles reveal differences in sitting time and perceptions. *International Journal of Environmental Research and Public Health*, 15(9). 10.3390/ijerph15092019
- Richardson A, Potter J, Paterson M, Harding T, Tyler-Merrick G, Kirk R, Reid K, & McChesney J. (2017). Office design and health: A systematic review. *The New Zealand Medical Journal*, 130(1467), 39–49.
- Schuna JM Jr., Swift DL, Hendrick CA, Duet MT, Johnson WD, Martin CK, Church TS, & Tudor-Locke C. (2014). Evaluation of a workplace treadmill desk intervention: A randomized controlled trial. *Journal of Occupational and Environmental Medicine*, 56(12), 1266–1276. 10.1097/jom.0000000000000336 [PubMed: 25479296]
- Tew GA, Posso MC, Arundel CE, & McDaid CM (2015). Systematic review: Height-adjustable workstations to reduce sedentary behaviour in office-based workers. *Occupational Medicine*, 65(5), 357–366. 10.1093/occmed/kqv044 [PubMed: 25934982]
- The EndNote Team. (2013). EndNote. In (Version EndNote X9) Clarivate.
- Thivel D, Tremblay A, Genin PM, Panahi S, Rivière D, & Duclos M. (2018). Physical Activity, inactivity, and sedentary behaviors: Definitions and implications in occupational health. *Frontiers in Public Health*, 6, Article 288. 10.3389/fpubh.2018.00288 [PubMed: 30345266]

- Tobin R, Leavy J, & Jancey J. (2016). Uprising: An examination of sit-stand workstations, mental health and work ability in sedentary office workers in Western Australia. *Work*, 55(2), 359–371. 10.3233/wor-162410 [PubMed: 27689593]
- Tufanaru C, Munn Z, Aromataris E, Campbell J, & Hopp L. (2020). Chapter 3: Systematic reviews of effectiveness. In Aromataris E. & Munn Z. (Eds.), *Joanna Briggs Institute reviewer's manual for Evidence Synthesis*. JBI. <https://synthesismanual.jbi.global>
- 2018 Physical Activity Guidelines Advisory Committee. 2018 Physical Activity Guidelines Advisory Committee Scientific Report. Washington, DC: U.S. Department of Health and Human Services, 2018.
- U.S. Bureau of Labor Statistics. (2017). The Economics Daily: Physical strength required for jobs in different occupations in 2016. <http://www.bls.gov/opub/ted/2017/physical-strength-required-for-jobs-in-different-occupations-in-2016.htm>
- U.S. Bureau of Labor Statistics. (2019). Labor forces statistics from the current population survey. <https://www.bls.gov/cps/cpsaat09.htm>
- Veritas Health Innovation. (2013). Covidence systematic review software. <http://www.covidence.org>
- Wahlstrom V., Bergman F., Ohberg F., Eskilsson T., Olsson T., & Jarvholm LS. (2019). Effects of a multicomponent physical activity promoting program on sedentary behavior, physical activity and body measures: A longitudinal study in different office types. *Scandinavian Journal of Work, Environment & Health*, 45, 493–504. 10.5271/sjweh.3808
- Wallmann-Sperlich B, Hoffmann S, Salditt A, Bipp T, & Froboese I. (2019). Moving to an “active” biophilic designed office workplace: A pilot study about the effects on sitting time and sitting habits of office-based workers. *International Journal of Environmental Research and Public Health*, 16(9), Article 1559. 10.3390/ijerph16091559 [PubMed: 31060203]
- Warburton DER, & Bredin SSD (2017). Health benefits of physical activity: A systematic review of current systematic reviews. *Current Opinion in Cardiology*, 32(5), 541–556. 10.1097/HCO.0000000000000437 [PubMed: 28708630]
- Zhu W, Gutierrez M, Toledo MJ, Mullane S, Stella AP, Diemar R, Buman KF, & Buman MP (2018). Long-term effects of sit-stand workstations on workplace sitting: A natural experiment. *Journal of Science and Medicine in Sport*, 21(8), 811–816. 10.1016/j.jsams.2017.12.005 [PubMed: 29289496]





**Figure 1.** Study selection by Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

Table 1.

Study Characteristic and Methods

No.	Sample characteristics						Study methods			PA measure	
	Author (publishing year)	City, state, country	Sample size (n)	M age (SD)	Female (%)	Study design	Study focus area	Study frequency and duration	Self-report	Objective	
1	Chau et al. (2014)	New South Wales, Australia	42	38 (11)	86	RCT	Sit-stand desk versus sitting desk	4 weeks, all workdays	OSPAC <sup>a</sup>	ActivPAL 3 accelerometer	
2	Dutta et al. (2014)	Minnesota, USA	28	40 (NR)	68	RCT	Sit-stand desk versus sitting desk	4 weeks, all workdays	OSPAC <sup>a</sup> end of each week	(a) MSR accelerometer (b) Grave accelerometer	
3	Schuna et al. (2014)	Baton Rouge, Louisiana, USA	41	40 (10)	98	RCT	Shared treadmill desk versus sitting desk	3 months, twice daily for 45-minute sessions	None	ActiGraph	
4	Miyachi et al. (2015)	Tokyo, Japan	32	44 (10.2)	31	RCT	Shared standing desk in open-space office versus sitting desk in open-space office	6 weeks, 10 hours/week	None	Acti marker accelerometer	
5	Tobin et al. (2016)	Perth, Australia	37	35 (10.5)	86	RCT	Sit-stand desk versus sitting desk	4 weeks, all workdays	None	ActivPAL accelerometer	
6	Bergman et al. (2018)	Umeå, Sweden	80	51 (6.8)	55	RCT	Treadmill desk versus sit-stand desk	13 months, all workdays	Self-reported question	(a) ActivPAL accelerometer (b) ActiGraph accelerometer	
7	Mayloret et al. (2018)	Bedfordshire, UK	896	43 (2.5)	57	RCT	Multicomponent <sup>a</sup> policy and environmental changes versus usual work practices at sitting desk	8 weeks, all workdays	None	ActivPAL accelerometer	
8	Pierce et al. (2019)	Hastings, New Zealand	24	39 (9.5)	52	RCT	Sit-stand desk versus sitting desk	6 weeks, all workdays	Baecke Questionnaire for Habitual Physical Activity	Keep Walking-Stay Fit Pedometer	
9	Gilson et al. (2012)	Sydney, Australia	11	47 (9.8)	64	Quasi-Exp (pre/post)	Move from sitting desk in open workspace to shared sit-stand desk in open workspace	5 days, all workdays	None	Armband accelerometer (SenseWear)	
10	Gorman et al. (2013)	Vancouver, Canada	24	35 (8.1)	75	Quasi-Exp (pre/post)	Building design—Move from conventional sitting desk workplace to activity-permissive workplace environment (sit-stand desk, staircases, cafe-style meeting rooms)	3–6 months in new environment, all workdays	None	ActivPAL accelerometer	

No.	Sample characteristics						Study methods			PA measure	
	Author (publishing year)	City, state, country	Sample size (n)	M age (SD)	Female (%)	Study design	Study focus area	Study frequency and duration	Self-report	Objective	
11	Koepp et al. (2013)	Minnesota, USA	36	42 (9.9)	69	Quasi-Exp (pre/post)	Move from sitting desk to treadmill desk	12 months, all workdays	None	Actical accelerometer	
12	Jancey et al. (2016)	Perth, Australia	42	40 (11.93)	64	Quasi-Exp (pre/post)	Building design—Move from traditional office space to active design building (centralized break room and bathroom, open accessible staircase)	4 months, all workdays	Online survey stair use	ActiGraph accelerometer	
13	Chau et al. (2016)	Sydney, Australia	31	33 (10.8)	45	Quasi-Exp (pre/post)	Move from sitting desk to sit-stand desk	19 weeks, all workdays	(a) Active Australia Questionnaire after 19 weeks (b) OSPAQ after 19 weeks	(a) ActiPAL 2 accelerometer (b) ActiGraph accelerometer	
14	Engelen et al. (2016)	Sydney, Australia	34	NR	74	Quasi-Exp (pre/post)	Building design—Old office building to new health-promoting active design building (accessible stairs and centralized break rooms and bathrooms)	2 months, all workdays	(a) OSPAQ (b) "How many day 30 mins MVPA" in past week?" completed at 2 months before and 2 months post-move	NA	
15	Mansoubi et al. (2016)	UK	40	32 (8.6)	55	Quasi-Exp (pre/post)	Move from sitting desk to sit-stand desk	3 months, all workdays	None	(a) ActiPAL 3 accelerometer (b) ActiGraph accelerometer	
16	Eyler et al. (2018)	Missouri, USA	143	NR	NR	Quasi-Exp (pre/post with control group)	Building design—Pre- or post-move to new active design building (stairwell, natural light, sit-stand desk, common printing and trash bin areas, and commuter showers) versus non-movers in adjacent building	12 months, all workdays	Adapted International Physical Activity Questionnaire for workday	ActiPAL accelerometer	
17	Zhu et al. (2018)	Arizona, USA	36	39 (11.3)	75	Quasi-Exp (pre/post with control group)	Pre/post sit-stand desk and shared treadmill desk versus staff in sitting desk in a different workspace	18 months, all workdays	None	ActiPAL accelerometer	
18	Candido et al. (2019)	Sydney, Australia	20	NR	NR	Quasi-Exp (pre/post)	Office arrangement design—Move from assigned seating in open-plan configuration to unassigned activity-based working space (sit-stand desk, multipurpose spaces, biophilia)	Average 82 days, all workdays	None	Fitbit Charge 2 accelerometer	

No.	Sample characteristics						Study methods			PA measure	
	Author (publishing year)	City, state, country	Sample size (n)	M age (SD)	Female (%)	Study design	Study focus area	Study frequency and duration	Self-report	Objective	
19	Dutta et al. (2019)	Minnesota, USA	15	39 (9.7)	73	Quasi-Exp (pre/post)	Move from sitting desk to sit-stand desk	12 months, all workdays	OSPAQ <sup>a</sup> at the end of each week for 2 weeks	Gruve accelerometer	
20	Malaeb et al. (2019)	Minnesota, USA	25	47 (NR)	92	Quasi-Exp (pre/post)	Move from sitting desk to shared treadmill desk	2 weeks, 2.5 hours/workday	International Physical Activity Questionnaire Short Version weekly	Actical accelerometer	
21	Wahlstrom et al. (2019)	Sweden	86	48 (10.3)	86	Quasi-Exp (pre/post)	Office arrangement design— Move from assigned sitting desk to (a) cell office with PA promotion program or (b) flex office in active design office space (sit-stand meeting space, 26 treadmill desks, shared trash bins, and sitting and standing break tables) with PA promotion program	18 months, all workdays	None	(a) ActivPAL accelerometer worn 24 hours/day for 7 consecutive days (b) ActiGraph accelerometer worn during waking hours for 7 consecutive days	
22	Wallmann-Sperlich et al. (2019)	Krefeld, Germany	23	39 (10)	75	Quasi-Exp (pre/post)	Office arrangement design— Move from sitting desk open space to new active office space (sit-stand desk; natural lighting; ventilation; significant numbers of plants, views, and recycled and non-synthetic materials; possibilities for sitting and standing; and features permitting PA)	months, all workdays	(a) Marshall Sitting Questionnaire 1 month before relocation, and 3 and 7 months after relocation (b) OSPAQ <sup>a1</sup> 1 month before relocation, and 3 and 7 months after relocation	None	
23	McGann et al. (2015)	Australia	111	45 (NR)	77	Cross-sectional	Building design—Compared three building styles (placement of shared spaces, staircases, and floor plan arrangement)	NA	OSPAQ <sup>a</sup>	ActiGraph accelerometer	
24	Carr et al. (2016)	Midwest, USA	69	44 (10.7)	74	Cross-sectional	Compared sit-stand desk users for 6 months with sitting desk users	NA	None	ActivPAL 3 accelerometer	
25	Lindberg et al. (2018)	USA	231	44 (12.2)	50	Cross-sectional	Office arrangement design— Compared desk types (private, cubicle, open bench)	NA	None	EggMove3 accelerometer	
26	Renaud et al. (2018)	Europe	1,098	47 (7.8)	35	Cross-sectional	Compared PA in sit-stand desk users with sitting desk users	NA	(a) Workforce Sitting Questionnaire (b) OSPAQ <sup>a</sup>	None	

No.	Sample characteristics					Study methods			PA measure	
	Author (publishing year)	City, state, country	Sample size (n)	M age (SD)	Female (%)	Study design	Study focus area	Study frequency and duration	Self-report	Objective
									(c) Single Physical Activity Guidelines question	

Note. PA = physical activity; RCT= randomized control trial; OSPAQ = Occupational Sitting and Physical Activity Questionnaire; NR = not reported; MSR = Modular Signal Recorder; Quasi-Exp = quasi-experimental study; MVPA = moderate- to vigorous-physical activity; NA = not available.

<sup>a</sup>Multicomponent intervention included education presentation, brainstorming session, step challenge, individual health check meetings software prompts to move, weekly telephone support, and changes to work environment such as relocation of trash bins and printers

Table 2.

PA Results, Main Findings, and Limitations

No.	Author (publishing year)	PA results		Main findings	Study limitations
		Overall	At work		
1	Chaulet. (2014)	NR	(a) Net difference in stepping time (NR, $p = .433$ ) minutes/day at work from baseline to post-intervention: 13 minutes/day ( $p = .127$ ) change in intervention and 2 minutes/day ( $p = .823$ ) change in control group (OSPAQ) (b) Net difference in stepping time (NR, $p = .453$ ) minutes/day at work from baseline to post-intervention: 11 minutes/day ( $p = .081$ ) change in intervention and 3 minutes/day ( $p = .596$ ) change in control (ActivPAL)	Sit-stand desk reduces overall sitting time but has no effect on step time or PA at work	<ul style="list-style-type: none"> <li>• Short-term follow-up</li> <li>• Convenience sample</li> <li>• No blinding</li> <li>• No objective measure for frequency or duration of activity</li> </ul>
2	Dutta et al. (2014)	(a) Increase in AUs from baseline to post-intervention in total activity: 237,729 AU/hour ( $p > .05$ ) in intervention group versus 236,445 AU/hour ( $p > .05$ ) in control group (Gruve)	(a) Increase in AUs in work activity from baseline to post-intervention: 229,156 AU/hour ( $p < .05$ ) in intervention group versus 210,245 AU/hour ( $p < .05$ ) in control group (Gruve)	Sit-stand desk used over 4 months significantly reduced sitting time and increased standing and light activity during work hours	<ul style="list-style-type: none"> <li>• No blinding</li> <li>• Short-term follow-up</li> <li>• Small sample</li> </ul>
3	Schuna et al. (2014)	(a) 1.6 minutes/hour (95% CI = [0.5, 2.8]) increase in light PA from baseline to post-intervention between the control group and the intervention group (b) -0.1 minute/hour (95% CI = [-0.5, 0.4]) decrease in MVPA from baseline to post-intervention between the control group and the intervention group	(a) 2.9 minutes/hour (95% CI = [0.9, 5.0]) increase in light PA from baseline to post-intervention between the control group and the intervention group (b) -0.4 minutes/hour (95% CI = [-0.9, 0.1]) decrease in MVPA from baseline to post-intervention between the control group and the intervention group	Shared treadmill workstations decreased sedentary time and improved low-intensity and light PA behavior but not MVPA activity during the workday and overall in overweight/obese office workers	<ul style="list-style-type: none"> <li>• Only used overweight/obese workers</li> <li>• Short-term follow-up</li> </ul>
4	Miyachi et al. (2015)	(a) Change in total time spent on total PA ( $p = \text{NR}$ ): 544.6 ( $\pm 117.5$ ) minutes/day after intervention versus 536.1 ( $\pm 117.0$ ) minutes/day in control group (b) Change in total time spent on light PA ( $p = .019$ ) minutes/day daily from baseline to post-intervention: 481.9 ( $\pm 116.0$ ) minutes/day after intervention versus 479.1 ( $\pm 113.5$ ) minutes/day in control group (c) Change in total time spent on moderate PA ( $p = \text{NR}$ ) minutes/day daily from baseline to post-intervention:	NR	Installation of sit-stand desk increases time spent on overall PA, especially PA during weekdays	<ul style="list-style-type: none"> <li>• Minimal control over daily workload</li> <li>• Objective measure unable to determine duration of standing up</li> <li>• Small sample size</li> </ul>

No.	Author (publishing year)	PA results			Main findings	Study limitations
		Overall	At work			
5	Tobin et al. (2016)	58.2 (±20.7) minutes/day after intervention versus 53.4 (±17.0) minutes/day in control group (d) Change in total time spent on vigorous PA ( $p = \text{NR}$ ) minutes/day daily from baseline to post-intervention: (±11.1) minutes/day after intervention versus (±11.6) minutes/day in control group	(a) Net difference in stepping time by 2.1 minutes/8-hour workday ( $p = .761$ ) in intervention group from baseline to post-intervention relative to control group	NR	A significant reduction in sitting time, but no change in the amount of time participants spent stepping at work	<ul style="list-style-type: none"> <li>• Short-term study period</li> <li>• Small sample size</li> <li>• Unmeasured confounders such as work or life stress</li> <li>• Outcomes measured during working hours only</li> </ul>
6	Bergman et al. (2018)	(a) Increase in daily walking time (22 minutes/weekday, $p = .00045$ ) from baseline to 13 months (b) Increase in daily light activity PA time (3 minutes/weekday, $p = .005$ ) from baseline to 13 months (c) Change in daily MVPA activity time (2 minutes/weekday, $p = .23$ ) from baseline to 13 months	NR	NR	Treadmill workstations increased daily walking time among overweight or obese office workers compared with those with a sit-stand desk. However, MVPA decreased over the study period with the largest decrease in the intervention group after 13 months	<ul style="list-style-type: none"> <li>• Different companies had various health promotion programs during intervention periods</li> <li>• Lack of blinding</li> <li>• Only used overweight/obese workers</li> <li>• Different companies had various health promotion programs during intervention periods</li> </ul>
7	Maylor et al. (2018)	(a) Net increase in stepping time (1.0 minute/day, $p = .770$ ) from baseline to 8 weeks	(a) Net increase in stepping time at work (12 minutes/workday, $p < .001$ ) from baseline to 8 weeks		Decrease in prolonged sitting time at workout and increase in steps per day using an individual, organization, and environmental intervention approach without the use of a sit-stand desk	<ul style="list-style-type: none"> <li>• Only one worksite used</li> <li>• Unable to measure effect of individual interventions</li> <li>• Outcomes measured during working hours only</li> </ul>
8	Pierce et al. (2019)	(a) No change in leisure PA associated with intervention (NR, $p = .039$ )	(a) Increase in steps taken at work (NR, $p < .001$ ) from baseline to 8 weeks associated with intervention		Adding electronic adjustable height desk to the workplace was associated with increase steps at work, decrease in sitting time at work, and no change in leisure-time PA	<ul style="list-style-type: none"> <li>• Small sample size</li> <li>• Pedometers unable to detect postural changes</li> <li>• Pedometers unable to capture intensity level of PA</li> <li>• Shorten version of the Baecke Questionnaire has not been previously assessed</li> </ul>
9	Gilson et al. (2012)	NR	(a) Net difference in percentage of time at work spent in light activity PA (0.8%, 95% CI = [-6.8, 7.9]) from baseline to post-intervention (b) Net difference in percentage of time at work spent in		Desk had no overall effect on sedentary time or PA time at work	<ul style="list-style-type: none"> <li>• Armband accelerometer did not capture posture changes</li> <li>• Small sample size</li> <li>• Short measurement period</li> </ul>

No.	Author (publishing year)	PA results		Main findings	Study limitations
		Overall	At work		
10	Gorman et al. (2013)	NR	MVPA (-0.7%, 95% CI = [-1.8, 2.3]) from baseline to post-intervention  (a) Net increase in stepping time 1.2 minutes/workday ( <i>p</i> = .748) at work from baseline to post-intervention	Post-move to an activity-permissive workspace office worker had less time sitting and more time standing, but no change in stepping time or PA during work time	<ul style="list-style-type: none"> <li>Participants shared the intervention desk</li> <li>Change to PA outside work hours was not accessed</li> <li>Bias within sample population due to nature of their industry</li> <li>Timing of data collection premove was not described</li> </ul>
11	Koepp et al. (2013)	(a) Change in average daily PA (NR, <i>p</i> = NR) from baseline to post-intervention: 4,205 AU/day ( <i>p</i> .001) at 12 months of intervention, 4,460 AU/day ( <i>p</i> .001) at 6 months of intervention, and 3,353 AU/day ( <i>p</i> .001) at baseline	(a) Change in walking time (NR, <i>p</i> = NR) at work from baseline to post-intervention: 70 minutes/workday ( <i>p</i> .001) at baseline, 128 minutes/workday ( <i>p</i> .001) at 6 months of intervention, and 109 minutes/workday ( <i>p</i> .001) at 12 months of intervention	Overall PA and walking time at work increased over the course of a year in employees with access to a walking treadmill desk and had no significant impact on work performance	<ul style="list-style-type: none"> <li>Treadmill desk scatter throughout the office did not allow for community support among users</li> <li>Small sample size</li> <li>Unlike similar studies that report PA in METs or intensity categories, PA is only reported in AUs</li> </ul>
12	Jancey et al. (2016)	NR	(a) Change in the average minutes/workday time spent doing light activity at work ( <i>p</i> < .001) from baseline to post-intervention: 57.16 minutes/workday (95% CI = [52, 63]) after intervention and 35.13 minutes/workday (95% CI = [32, 39]) at baseline  (b) Change in the average minutes/workday time spent doing moderate activity at work ( <i>p</i> = .109) from baseline to post-intervention: 39.72 minutes/workday (95% CI = [35, 45]) after intervention and 36.13 minutes/workday (95% CI = [33, 40]) at baseline  (c) Change in the average minutes/workday time spent doing vigorous activity at work ( <i>p</i> = .658) from baseline to post-intervention: 0.29 minutes/workday (95% CI = [0.11, 0.53]) after intervention and 0.33 minutes/workday (95% CI = [0.16, 0.54]) at baseline	Average time spent doing light activity at work increased, time spent standing at work increased, and sitting time at work decreased	<ul style="list-style-type: none"> <li>No control group</li> <li>Large loss to follow-up</li> <li>Did not measure PA outside of work</li> </ul>
13	Chaunet al. (2016)	NR	(a) Net difference in walking time (-8 minutes/workday, <i>p</i> = .679) at work from baseline to post-intervention: -21 minutes/day ( <i>p</i> = .144) change in intervention versus -13 minutes/day ( <i>p</i> = .287) change in control (OSPAQ)  (b) Net difference in heavy labor time (14 minutes/workday, <i>p</i> = .125) at work from baseline to post-intervention: 11 minutes/day ( <i>p</i> = .396) change in intervention versus -3 minutes/day ( <i>p</i> = .749) change in control (OSPAQ)	No changes in walking time or PA, but sit-stand desk did reduce sitting and increase standing time	<ul style="list-style-type: none"> <li>Small convenience sample</li> <li>Data loss from accelerometer device malfunction</li> <li>Non-adherence to wearing accelerometer device</li> </ul>
14	Engelen et al. (2016)	(a) Decrease in number of days spent doing MVPA per week ( <i>p</i> > .05) at work from baseline to post-intervention: 4.6 days/week in intervention versus 4.9 days/week at baseline	(a) Change in percentage of time spent walking ( <i>p</i> > .05) at work from baseline to post-intervention: 11% of workday at baseline versus 10% of workday in intervention (OSPAQ)  (b) No change in percentage of time spent on heavy labor at work from baseline to post-intervention: 0% of	Workers sat less and new building provided more opportunities for incidental activity	<ul style="list-style-type: none"> <li>No objective measure of PA</li> <li>Small sample size</li> <li>Large loss to follow-up</li> <li>Study did not compare the four baseline sites for differences at baseline</li> </ul>



No.	Author (publishing year)	PA results		Main findings	Study limitations
		Overall	At work		
15	Mansoubi et al. (2016)	NR	workday in intervention and 0% of workday at baseline (OSPAQ) (a) No change in stepping time at work from baseline to post-intervention (ActiPAL) (b) No change in overall proportion of time spent in light activity on workdays from baseline to post-intervention (ActiGraph) (c) No change in overall proportion of time spent in MVPA on workdays from baseline to post-intervention (ActiGraph)	The use of sit-to-stand desk decreased sedentary time at work, increased light activity time at work, and had no effect on leisure-time moderate PA	<ul style="list-style-type: none"> <li>Short-term study period</li> <li>Small sample size</li> <li>Convenience sample</li> </ul>
16	Eyler et al. (2018)	NR	(a) Increase in average steps/day (NR, $p = .99$ ) from baseline to post-intervention in all groups: 1,591 steps/day ( $p < .001$ ) change in movers, 928 steps/day ( $p = .09$ ) change in non-movers, and 1,756 steps/day ( $p = .008$ ) change in control	PA increased in all study groups; it is unclear whether building had an effect on PA	<ul style="list-style-type: none"> <li>Wellness challenge to track PA for cash incentives started at time of post-data collection</li> <li>Self-selection bias in those who volunteered to wear accelerometer</li> <li>Unable to capture sample demographics</li> <li>Did not compare sitting and activity patterns at baseline for differences between groups</li> </ul>
17	Zhu et al. (2018)	NR	(a) No change to light PA or MVPA at the workplace	Sit-stand desk paired with motivational support decreased sitting time, increased standing time, and increased low-intensity PA in the workplace and is sustainable for 18 months	<ul style="list-style-type: none"> <li>Small sample</li> <li>High attrition at 18 months</li> </ul>
18	Candido et al. (2019)	NR	(a) Increase in average steps/day: 300 steps/day ( $p = NR$ ) after relocation to new office environment	The average steps per day increased after moving to a active building design environment	<ul style="list-style-type: none"> <li>Selection bias in those who volunteered to wear Fitbit</li> <li>PA data only collected from 20 volunteers</li> </ul>
19	Dutta et al. (2019)	NR	(a) 24,748 AU/hour (95% CI = [7,150, 42,347]) increase at work after 1 year follow-up from baseline	Overall PA during the workday remained about 12% higher, and sitting time remained reduced after 1 year of sit-stand desk relative to baseline	<ul style="list-style-type: none"> <li>Small sample size</li> <li>Blinding not achievable</li> <li>Low retention from original study</li> </ul>
20	Malaeb et al. (2019)	(a) Increase in total PA step count in intervention group compared with baseline ( $p < .01$ )	NR	Treadmill desk usage over 2 weeks increased overall step count and had positive body composition results	<ul style="list-style-type: none"> <li>Small sample size</li> <li>All participants had BMI &gt;25</li> <li>Short intervention period</li> <li>Accelerometer counts not converted to METs or minutes of activity</li> </ul>
21	Wahlstrom et al. (2019)	NR	(a) Change in walking time (NR, $p = .001$ ) minutes/workday from baseline to 18-month follow-up: 39 minutes/workday (95% CI = [35, 43]) at baseline and 47 minutes/workday (95% CI = [44, 52]) at 18-month follow-up in flex office; 42 minutes/workday (95% CI = [38, 46]) at baseline and 41 minutes/workday (95% CI = [40, 46]) at 18-month follow-up in cell office	Greatest increase in walking time, number of steps, and MVPA time during the workday compared with baseline occurred in flex offices. No changes in sitting time occurred	<ul style="list-style-type: none"> <li>All employees had sit-stand Desk before and after move regardless of office type</li> <li>Unbalanced distribution of gender and managers between the two groups</li> <li>Utilization of provided health and wellness hour not measured</li> </ul>

No.	Author (publishing year)	PA results		Study limitations
		Overall	At work	
22	Wallmann-Sperlich et al. (2019)	NR	(b) Change in MVPA time (NR, $p < .001$ ) minutes/workday from baseline to 18-month follow-up: 19 minutes/workday (95% CI = [15, 22]) at baseline and 27 minutes/workday (95% CI = [23, 30]) at 18-month follow-up in flex office; 16 minutes/workday (95% CI = [13, 19]) at baseline and 19 minutes/workday (95% CI = [15, 22]) at 18-month follow-up in cell office (a) Change in minutes/workday walking ( $p = .33$ ) from baseline to 7 months post-intervention: 70.2 minutes/workday (95% CI = [30.7, 109.7]) at baseline and 84 minutes/workday (95% CI = [50.1, 117.9]) at post-intervention measure	<ul style="list-style-type: none"> <li>Seasonal differences in moves were not accounted for</li> <li>Small sample size</li> <li>High attrition</li> <li>No objective PA measurement</li> <li>PA intensity levels not measured</li> </ul>
23	McGann et al. (2015)	NR	(a) Light-intensity PA was 4.6% in Building 1, 2.6% in Building 2, and 3.3% in Building 3. Building 1 has best quality staircases and corridors	<ul style="list-style-type: none"> <li>Did not discuss the participants workstations or desk</li> <li>Did not provide PA measures per building</li> </ul>
24	Carr et al. (2016)	NR	(a) Difference in average time walking hours/day at work in employees with sit-stand desk versus walking desk: 0.7 hours/day in sit desk and 0.9 hours/day in sit-stand desk ( $p = .22$ )	<ul style="list-style-type: none"> <li>Small sample size</li> <li>Limited generalizability</li> <li>PA intensity levels not measured</li> </ul>
25	Lindberg et al. (2018)	NR	(a) Workers in open bench seating exhibited 31.83% higher rates of PA than those in private offices (225.51 mG; 95% CI = [137, 314]) (b) Workers in open bench seating exhibited 20.16% higher rates of PA than those in cubicles (185 mG; 95% CI = [67, 304])	<ul style="list-style-type: none"> <li>Cannot confirm causal relationship between office type and PA</li> </ul>
26	Renaud et al. (2018)	(a) Meeting the guidelines for PA showed a positive trend with sit-stand desk users: 30% in non-users, 30% in monthly to weekly users, and 35% in daily users	(a) Walking hours/week at work showed a positive trend with sit-stand desk users: 2.3 hours/week in non-users, 2.2 hours/week in monthly to weekly users, and 3.2 hours/week in daily users	<ul style="list-style-type: none"> <li>Recall bias</li> <li>Social desirability</li> <li>Did not measure how long employees had used a sit-stand desk</li> <li>Limited PA measurement</li> </ul>

Note. PA = physical activity; NR = not reported; OSPAQ = Occupational Sitting and Physical Activity Questionnaire; AUs = activity units; CI = confidence interval; MVPA = moderate- to vigorous-physical activity; BMI = body mass index; mG = milli-Gs (g-force); METs = Metabolic equivalent of task.

**Table 3.** Joanna Briggs Critical Appraisal Tool for Bias in Randomized Control Trial Studies

No.	Author (publishing year)	(a) Was true randomization used for assignment of participants to treatment groups?	(b) Was allocation to treatment groups concealed?	(c) Were treatment groups similar at baseline?	(d) Were participants blind to treatment assignment?	(e) Were those delivering treatment blind to treatment assignment?	(f) Were outcome assessors blind to treatment assignment?	(g) Were treatment groups treated identically other than the intervention of interest?	(h) Was follow-up complete and if not, were differences between groups in terms of their follow-up adequately described and analyzed?	(i) Were participants analyzed in the groups to which they were randomized?	(j) Were outcomes measured in the same way for treatment groups?	(k) Were outcomes measured in a reliable way?	(l) Was appropriate statistical analysis used?	(m) Was the trial design appropriate, and any deviations from the standard RCT design (individual randomization, parallel groups) accounted for in the conduct in the analysis of the trial?
1	Chau et al. (2014)	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
2	Dutta et al. (2014)	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
3	Schunget al. (2014)	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
4	Miyachet al. (2015)	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
5	Tobin et al. (2016)	Unclear	Unclear	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
6	Bergmo et al. (2018)	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
7	Maylofet al. (2018)	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes
8	Pierce et al. (2019)	Yes	No	Yes	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes

Note: RCT= randomized control trial. Yes responses shaded In green.

**Table 4.**

Joanna Briggs Critical Appraisal Tool for Bias in Quasi-Experimental Studies

No.	Author (publishing year)	(a) it is clear in the study what is the cause and what is the effect?	(b) Were the participants included in any comparisons similar?	(c) Were the participants included in any comparisons receiving similar treatment/care, exposure or intervention of interest?	(d) Was there a control group?	(e) Were there multiple measurements of the outcome both pre and post the intervention/exposure?	(f) Was follow-up complete and if not, were differences between groups in terms of their follow-up adequately described and analyzed?	(g) Were the participants included in any comparisons measured in the same way?	(h) Were outcomes measured in a reliable way?	(i) Was appropriate statistical analysis used?
9	Gilson et al. (2012)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
10	Gorman et al. (2013)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
11	Koopp et al. (2013)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
12	Jancey et al. (2016)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
13	Chau et al. (2016)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
14	Engelen et al. (2016)	Yes	Unclear	Yes	No	No	Yes	Yes	Yes	Yes
15	Mansoubi et al. (2016)	Yes	Yes	No	No	Yes	Yes	Yes	Yes	Yes
16	Eyler et al. (2018)	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	Yes	Yes
17	Zhu et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
18	Candido et al. (2019)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
19	Dutta et al. (2019)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
20	Malaeb et al. (2019)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
21	Wahlstrom et al. (2019)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
22	Wallmann-Sperlich et al. (2019)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes

Note. Yes responses shaded in green.

**Table 5.**

Joanna Briggs Critical Appraisal Tool for Bias in Cross-Sectional Studies

No.	Author (publishing year)	(a) Were the criteria for inclusion in the sample clearly defined?	(b) Were the study subjects and the setting described in detail?	(c) Was the exposure measured in a valid and reliable way?	(d) Were objective, standard criteria used for measurement of the condition?	(e) Were confounding factors identified?	(f) Were strategies to deal with confounding factors stated?	(g) Were the outcomes measured in a valid and reliable way?	(h) Was appropriate analysis used?
23	McGann et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
24	Carr et al. (2016)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
25	Lindberg et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
26	Renaud et al. (2018)	Unclear	Yes	Yes	Unclear	Yes	Yes	Yes	Yes

*Note.* Yes responses shaded in green.