

UC Davis

Dermatology Online Journal

Title

Photodamage spares the distal digits owing to primate-like hand flexion

Permalink

<https://escholarship.org/uc/item/6x05n0bz>

Journal

Dermatology Online Journal, 25(1)

Authors

Dorrell, Deborah N
Feldman, Steven R

Publication Date

2019

DOI

10.5070/D3251042623

Copyright Information

Copyright 2019 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Photodamage spares the distal digits owing to primate-like hand flexion

Deborah N Dorrell¹ BA, Steven R Feldman^{1,2,3} MD PhD

Affiliations: ¹Center for Dermatology Research, Department of Dermatology, Wake Forest School of Medicine, Winston-Salem, North Carolina, USA, ²Department of Pathology, Wake Forest School of Medicine, Winston-Salem, North Carolina, USA, ³Department of Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, North Carolina

Corresponding Author: Steven R. Feldman MD, PhD, Department of Dermatology, Wake Forest School of Medicine, Medical Center Boulevard, Winston-Salem, NC 27157-1071, Tel: 336-716-7740, Fax: 336-716-7732, Email: sfeldman@wakehealth.edu

Abstract

Sun-exposed areas of the body — including the face, neck, and extensor forearms — are chronically exposed to UV light and display signs of photoaging. The skin distal to the proximal interphalangeal (PIP) joint is, however, typically spared owing to the natural rest position of the hand. The inward curvature of the fingers, termed the finger flexion cascade, orients the skin distal to the PIP joint toward the ground as in knuckle-walking primates. The near constant protection of this area of skin from sun may reflect our primate ancestry, the daily activities of which included swinging through trees and knuckle-walking. Primates have elongated, inward curved fingers that were once advantageous for gripping branches during arboreal locomotion. Although natural selection has favored shorter, straighter fingers, the human hand continues to assume a natural shape of flexion at rest. The UV untouched, healthy skin at the ends of the fingers is a reminder of our primate heritage. This finding may provide a colorful, memorable means to reinforce to patients how photoaging is a manifestation of sun exposure rather than age.

Keywords: photodamage, photoaging, finger flexion cascade

Discussion

Sun-exposed areas of the body — including the face, neck, and extensor forearms — are chronically exposed to UV light and display signs of photoaging, including solar purpura and hemosiderin deposition, solar lentigines and ephelides, and actinic keratoses

[1, 2]. The extensor forearms of many patients, especially older, light-skinned individuals, are often extensively involved. The skin distal to the proximal interphalangeal (PIP) joint is, however, typically spared (**Figure 1**) owing to the natural rest position of the hand (**Figure 2A**). The inward curvature of the fingers orients the skin distal to the PIP joint toward



Figure 1. This human hand shows considerable photodamage, but the skin of and distal to the proximal interphalangeal joint is spared from photodamage.



Figure 2. A) The natural finger flexion cascade of a human hand at rest, which is similar to the distal digit flexion of the black gorilla paw, **(B)**.

the ground as in knuckle-walking primates (**Figure 2B**).

The alignment of human fingers at rest is termed the finger flexion cascade or cascade sign and is caused by the resting tone and flexibility of the soft tissues of the hand. Each finger from the index to the pinky is sequentially flexed to a greater degree [3]. The finger flexion cascade provides a ready explanation for the apparent resistance to photodamage of skin distal to the PIP joints. The near constant protection of this area of skin from sun may reflect our primate ancestry, the daily activities of which included swinging through trees and knuckle-walking.

References

1. Lim HW, Hawk JLM, Rosen CF. Photodermatologic Disorders. In: Dermatology. Bologna J, Schaffer J, Cerroni L, editors. 4th ed. Elsevier Limited: 2018. p. 2310-7.
2. DeLeo VA. Sunscreens and Photoprotection. In: Dermatology. Bologna J, Schaffer J, Cerroni L, editors. 4th ed. Elsevier Limited: 2018. p. 1536-47.
3. Asha K, Varghese JG, Sriniviasan V, Bhojan K. An Analytical Study on Finger Flexion Cascade in General Population of Various Occupations. *Orthop Rheumatol Open Access.J* 2015;1(1). doi: 10.19080/OROAJ.2015.01.555554.
4. Kivell TL, Schmitt D, Walker A. Independent Evolution of Knuckle-Walking in African Apes Shows That Humans Did Not Evolve from a Knuckle-Walking Ancestor. *Proc Natl Acad Sci.* 2009;106:14241-6. [PMID: 19667206].
5. Young RW. Evolution of the human hand: the role of throwing and clubbing. *J Anat.* 2003;202:165-74. [PMID: 12587931].

There is debate among evolutionary biologists as to whether humans are most closely descended from terrestrial, knuckle-walking primates like gorillas or from arboreal apes like chimpanzees and bonobos. However, the main difference between the hand structures of these two groups of primates is in the wrist rather than the fingers [4]. Both groups have elongated, inward curved fingers that were once advantageous for gripping branches during arboreal locomotion. Fossils from early hominids also show curved fingers, but over time natural selection favored fingers better suited for throwing, clubbing, and using primitive tools. Therefore, human fingers have become shorter and straighter and thumbs have become longer and stronger. Our hands now have increased dexterity and the ability to form specialized power and precision grips [5]. But at rest, the hand continues to assume its age-old flexion cascade inherited from evolutionary ancestors. The UV untouched, healthy skin at the ends of the fingers is a reminder of this heritage. This finding may provide a colorful, memorable means to reinforce to patients how photoaging is a manifestation of sun exposure rather than age.