

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

HOMOGENEOUS CATALYTIC HYDROGENATIONS, 1: REDUCTIONS OF POLYNUCLEAR AROMATIC AND POLYNUCLEAR HETEROAROMATIC NITROGEN COMPOUNDS UTILIZING CARBON MONOXIDE WITH WATER OR HYDROGEN

### Permalink

<https://escholarship.org/uc/item/6xf0j5n5>

### Authors

Fish, R.H.

Cremer, G.A.

### Publication Date

1982



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## ENERGY & ENVIRONMENT DIVISION

Submitted to the Journal of The American  
Chemical Society

HOMOGENEOUS CATALYTIC HYDROGENATIONS, 1:  
REDUCTIONS OF POLYNUCLEAR AROMATIC AND  
POLYNUCLEAR HETEROAROMATIC NITROGEN  
COMPOUNDS UTILIZING CARBON MONOXIDE WITH  
WATER OR HYDROGEN

Richard H. Fish and Gregg A. Cremer

January 1982


RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

JAN 8 1982

LIBRARY AND  
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 6782



LBL-13734  
c.d.

## DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

HOMOGENEOUS CATALYTIC HYDROGENATIONS, 1:  
Reductions of Polynuclear Aromatic and  
Polynuclear Heteroaromatic Nitrogen Compounds  
Utilizing Carbon Monoxide with Water or Hydrogen

Richard H. Fish and Gregg A. Cremer

Energy and Environment Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, CA 94720


January 1982

This study was jointly supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division, and the Assistant Secretary of Fossil Energy, Office of Coal Research, Liquefaction Division of the U.S. Department of Energy through the Pittsburgh Energy Technology Center under Contract W-7405-ENG-48.

Homogeneous Catalytic Hydrogenations, 1.  
Reductions of Polynuclear Aromatic and Polynuclear  
Heteroaromatic Nitrogen Compounds Utilizing Carbon  
Monoxide with Water or Hydrogen

Richard H. Fish\* and Gregg A. Cremer  
Energy and Environment Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

Pettit and his co-workers recently demonstrated, in an elegant manner, the use of carbon monoxide and water as a reducing agent for the hydroformylation of olefins and of carbon monoxide, water and hydrogen in the reduction of nitroarenes<sup>1-3</sup>. The compounds responsible for these catalytic reactions, in Pettit's studies, were transition metal carbonyl hydrides that had been generated catalytically under the reaction conditions (eq. 1).




These results prompted us to study the utilization of these reagents for the reduction of polynuclear aromatic and polynuclear heteroaromatic nitrogen compounds. The importance of studying these types of polynuclear compounds and reducing agents stems from the need to learn more about the basic aspects of coal liquefaction and the hydroprocessing of coal liquids and shale oils<sup>4</sup>. Additionally, it is well known that homogeneous catalytic reductions proceed at lower temperatures and pressures and give higher regioselectivities when compared to their heterogeneous counterparts<sup>5a,b</sup>.

We wish to demonstrate, in this communication, the potential of carbon monoxide and water or carbon monoxide and hydrogen as reducing agents for polynuclear aromatic and polynuclear heteroaromatic nitrogen compounds, 1-8, (Chart 1) using transition metal carbonyl compounds as homogeneous catalysts.

We reacted a wide variety of transition metal carbonyl compounds with anthracene, 1, under water gas shift (wgs) conditions (CO, H<sub>2</sub>O, base)<sup>6</sup>, and found that only Mn<sub>2</sub>(CO)<sub>8</sub>(Bu<sub>3</sub>P)<sub>2</sub>, Fe(CO)<sub>4</sub>Bu<sub>3</sub>P and Co<sub>2</sub>(CO)<sub>6</sub>(φ<sub>3</sub>P)<sub>2</sub> produced the reduction product, 9,10-dihydroanthracene, 9, in yields of 13%, 8% and 3% respectively. The carbonyls of Ru, Rh, W, Os, Re and Mo all produced hydrogen and carbon dioxide (wgs reaction), but no compound 9.

These results can partially be rationalized by the fact that the latter transition metal carbonyl hydrides are unstable under the reactions conditions and undergo reductive elimination of hydrogen rather than hydrogen transfer to the polynuclear aromatic substrate, 1, i.e.,  $k_2 > k_1$  (eq. 2).



In view of these discoveries, we decided to learn more about the parameters of this reaction using Mn<sub>2</sub>(CO)<sub>8</sub>(Bu<sub>3</sub>P)<sub>2</sub> as the catalyst. The reaction of carbon monoxide and deuterium oxide (D<sub>2</sub>O) with 1 provided only 9,10-dideu--tetraanthracene, 10<sup>7</sup>, and this result strongly indicates that the hydrogen comes exclusively from water. In addition, the presence of added deuterium gas (D<sub>2</sub>) provided no deuterium incorporation; thereby eliminating a mechanism which includes oxidative addition of H<sub>2</sub> to any coordinatively unsaturated manganese species that might be formed<sup>3</sup>, and supports the facile reaction (eq. 1) of hydroxide ion on coordinated CO as the major reaction pathway to the carbonyl hydrides<sup>1</sup>. Several control experiments verified that the reduction

product, 9, did not undergo exchange of hydrogen for deuterium under the reaction conditions.

In an experiment to elucidate the stereochemistry of the anthracene reaction, we reacted an analog of 1, 9,10-dimethylantracene, 11, with  $\text{Mn}_2(\text{CO})_8(\text{Bu}_3\text{P})_2$  under water gas shift conditions ( $180^\circ$ , 5 hrs, 350 psi CO, 0.2M KOH in THF, sub/cat = 20), to provide a 30% yield of cis and trans-9,10-dihydro-9,10-dimethylantracene, 12 and 13, in a ratio of 53%:47% <sup>8</sup>. The mechanistic implication of this result strongly suggests a free radical process for this reaction. A similar result was recently reported by Halpern et al. <sup>9a</sup> for the reaction of 10 and  $\text{Mn}_2(\text{CO})_{12}$  and Taylor and Orchin <sup>9b</sup> with  $\text{Co}_2(\text{CO})_8$  under syn gas conditions ( $\text{CO}_2$ ,  $\text{H}_2$  1:1,  $\sim 200^\circ$ ).

We also found that in the absence of aqueous base and in the presence of hydrogen, i.e., synthesis gas (sg) conditions ( $\text{CO}/\text{H}_2 = 1$ ), that better yields of reduced polynuclear aromatic products could be obtained using  $\text{Mn}_2(\text{CO})_8(\text{Bu}_3\text{P})_2$  as the catalyst and compounds 1 - 3 as substrates. Thus, under sg conditions <sup>10</sup> reaction of 1 with  $\text{Mn}_2(\text{CO})_8(\text{Bu}_3\text{P})_2$  provided 9 in 30% yield, while reaction with 2 gave no product and 3 produced 4,5-dihdropyrene, 14, in 3% yield.


From these results, it is evident that bent polynuclear aromatic compounds are extremely unreactive under either wgs or sg conditions, when compared to 1, a linear polynuclear aromatic compound <sup>11</sup>.

The polynuclear heteroaromatic compounds, specifically with nitrogen as the heteroatom, are important to study, since they are highly prevalent in all coal and oil shale products <sup>12</sup>. We found a dramatic increase in reactivity for the polynuclear heterocyclic nitrogen compounds under either wgs or sg conditions compared to their carbon analogs, e.g., acridine more

reactive than anthracene. Table 1 summarizes our results with polynuclear heteroaromatic nitrogen compounds  $\underset{z}{4}$ - $\underset{z}{8}$ .

It is important to note the high regioselectivity observed in these reactions by the ease with which the nitrogen heterocyclic ring gets hydrogenated in preference to a benzene ring. This result might be explained by the lowered aromaticity of the nitrogen heterocyclic ring versus their carbon analog, which would provide a lower activation energy for hydrogenation<sup>13</sup>. The order of reactivity for compounds  $\underset{z}{4}$ - $\underset{z}{8}$  either under wgs or sg conditions is:<sup>14</sup>  $\underset{z}{4} \gg \underset{z}{6} > \underset{z}{8} > \underset{z}{5} \gg \underset{z}{7}$ .

The overall observed relative reactivities of compounds  $\underset{z}{1}$ - $\underset{z}{8}$ , either under wgs or sg conditions, is as follows:



This order, while qualitative, is still significant, since these polynuclear compounds are present in coal, coal liquids and shale oil and represents the first logical explanation for compound reactivity under catalytic hydrogenation conditions in these complex matrices.

Finally, the fact that Fe, Mn and Co carbonyls give reduced products, while other transition carbonyls were unreactive imply important mechanistic differences. Pertinently, we found in these studies, in addition to the instability factor for several of the transition carbonyl hydrides, a pronounced carbon monoxide inhibition on the catalytic hydrogenation reactions especially with the ruthenium carbonyls and report these results in the following paper<sup>15</sup>.

We are continuing to pursue this important area of fossil energy research and will report further details in forthcoming publications<sup>16</sup>.



### Acknowledgements

We wish to thank Professors R. Bergman, K.P.C. Vollhardt and T. Vermeulen as well as Dr. H. Heinemann for helpful discussions concerning the preparation of this manuscript. This study was jointly supported by the Director, Office of Energy Research, Office of Basic Energy Science, Chemical Sciences Division, and the Assistant Secretary of Fossil Energy, Office of Coal Research, Liquefaction Division, of the U.S. Department of Energy through the Pittsburgh Energy Technology Center under contract W-7405-ENG-48.

### References and Notes

- (1) Kang, H. G.; Mauldin, C. H.; Cole, T.; Slegeir, W.; Cann, K. and Pettit, R.; J. Amer. Chem. Soc., 1977, 99, 8323 and reference therein.
- (2) Cann, K.; Cole, T.; Slegeir, W.; Pettit, R.; ibid. 1978, 100, 3969.
- (3) Cole, T.; Ramage, R.; Cann, K., Pettit, R.; ibid. 1980, 102, 6182.
- (4) Coal Liquefaction Fundamentals ACS Syn. Series, 139, Whitehurst, D. D., Editor, (1980).
- (5a) James, B. J. "Homogeneous Hydrogenations"; John Wiley and Son: New York, N.Y. 1973.
- (5b) Parshall, G. W., Science 1980, 208, 1221.
- (6) The reaction conditions were as follows for the following carbonyls:  $\text{Re}(\text{CO})_4\text{Bu}_3\text{P}$ ,  $\text{Mn}_2(\text{CO})_8(\text{Bu}_3\text{P})_2$ ,  $\text{Co}_2(\text{CO})_6(\phi_3\text{P})_2$ ,  $\text{W}(\text{CO})_6$ ,  $\text{Re}(\text{CO})_{10}$ ,  $\text{RuCl}_2(\text{CO})_2(\phi_3\text{P})_2$ ,  $\text{Rh}_6(\text{CO})_{16}$ ,  $\text{Mo}(\text{CO})_5\text{Bu}_3\text{P}$ . A 45 ml Parr mini-reactor was used with tetrahydrofuran (12 ml) as solvent, 0.2 M KOH (3 ml),  $P_{\text{CO}}$  350 psi, 5 hr reaction time, 1 mmole of 1, 0.1 mmole of transition metal carbonyl catalyst (10:1 substrate to catalyst ratio).

The product percentage was found by area ratio of starting material and product using a HP 5880A capillary gas chromatograph with digital integration (12 m x 0.1 mm OV 101 at 60° to 200 (10°/min) and flame ionization detection. The reduction product, 9, was isolated by column chromatography on florisil (benzene) and identified by GC-EIMS and nmr spectroscopy ( $^1\text{H}$ , 250 MHz).


- (7) GC-EI mass spectral analysis showed m/e 182 (100%) consistent with two deuteriums being incorporated into compound 1 (m/e 178).

- (8) The cis and trans isomers of 9,10-dihydro-9,10-dimethylantracene, 12, 13 could be cleanly separated on a 12 m x 0.1 mm i.d., fused silica capillary column (OV 101) using a HP 5880A instrument with flame ionization detection (temperature programming 60°-200°, 10°/min).
- (9a) Sweany, R.; Buttler, S. C; and Halpern, J. Organometal Chem., 1981, 213, 487.
- (9b) Taylor, P. D; and Orchin. M. J., Org. Chem., 1972, 37, 3913.
- (10) The reactions of 1-3 with  $Mn_2(CO)_8(Bu_3P)_2$  as catalyst were run in a 45 ml Parr mini-reactor at 200°C, 350 psi H<sub>2</sub>, 350 psi CO in THF for 2 hr with a 10:1 substrate to catalyst ratio. Other carbonyls such as Fe(CO)<sub>4</sub> Bu<sub>3</sub>P, Cr(CO)<sub>6</sub>, Mo(CO)<sub>6</sub>, W(CO)<sub>6</sub>, Re<sub>2</sub>(CO)<sub>10</sub> and Ru<sub>3</sub>(CO)<sub>12</sub> produced no reduced product with 1-3 as substrates.
- (11) Friedman, S.; Meltin, S.; Svedi, A.; and Wender, I. J., Org. Chem. 1959, 24, 1287.
- (12) Larsen, J. W., Editor. Organic Chemistry of Coal, ACS Symposium Series 71, 1978.
- (13) Dewar, M.J.S. "The Molecular Orbital Theory of Organic Chemistry" McGraw-Hill, New York, N.Y., 1969.
- (14) The yields we report in Table 1 were mainly for comparison purposes, i.e., times of 2 hrs used for each reaction. However, longer reaction times provided, in most cases, quantitative conversions and thus the preparative aspects could be of potential interest to synthetic chemists.
- (15) See the following paper: Fish, R. H.; and Cremer, G. A., J. Amer. Chem. Soc., 1982, 104, 0000.


(16) Preliminary accounts of this work were presented at the 182nd National ACS Meeting, Aug. 23-28, 1981, New York, N.Y., Abstract INOR 49 and the Pacific Conference on Chemistry and Spectroscopy, Anaheim, CA, Oct. 19-21, 1981, Abstract 188.

Chart 1


Polynuclear Aromatic and Polynuclear Heteroaromatic  
Nitrogen Compounds Reacted under Water Gas Shift or  
Synthesis Gas Conditions




1




2




3




4




5



6



7



8

Table 1

Reductions of Polynuclear Heteroaromatic Nitrogen Compounds  
under Water Gas Shift <sup>a</sup> (WGS) and Syn Gas (SG) Conditions <sup>b</sup>  
with Transition Metal Carbonyls as Catalysts.

Substrate	Catalyst <sup>c</sup>	Sub/Cat	Temp.(°C)	Time(hr)	Conditions	Products (%) <sup>d</sup>
<u>4</u>	Fe(CO) <sub>5</sub>	10	180	2	WGS	9,10-dihydro-acridine (100)
<u>4</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	10	200	2	WGS	" (38)
<u>4</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	2	SG	" (100)
<u>4</u>	Co <sub>2</sub> (CO) <sub>6</sub> (φ <sub>3</sub> P) <sub>2</sub>	20	200	2	SG	" (100)
<u>5</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	2	SG	1,2,3,4, tetrahydro- 5,6-benzo- quinoline (7)
<u>5</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	5	WGS	" (4)
<u>5</u>	Fe(CO) <sub>4</sub> (Bu <sub>3</sub> P)	10	180	5	WGS	" (1)
<u>5</u>	Co <sub>2</sub> (CO) <sub>6</sub> (φ <sub>3</sub> P) <sub>2</sub>	20	200	2	SG	" (8)
<u>6</u>	Fe(CO) <sub>5</sub>	10	180	2	WGS	1,2,3,4- Tetrahydro- quinoline (0)
<u>6</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	5	WGS	" (4)
<u>6</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	5	SG	" (33)
<u>6</u>	Co <sub>2</sub> (CO) <sub>6</sub> (φ <sub>3</sub> P) <sub>2</sub>	20	200	1	SG	" (70)
<u>7</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	2	SG	1,2,3,4 tetrahydro- 7,8-benzo- quinoline (2)
<u>7</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	2	WGS	No product
<u>8</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	2	WGS	9,10-dihydro- phenanthridine (1)
<u>8</u>	Mn <sub>2</sub> (CO) <sub>8</sub> (Bu <sub>3</sub> P) <sub>2</sub>	20	200	2	SG	" (11)
<u>8</u>	Co <sub>2</sub> (CO) <sub>6</sub> (φ <sub>3</sub> P) <sub>2</sub>	20	200	2	SG	" (21)

a. Reaction run in THF (12 ml) with 0.2 M KOH (3 ml), 350 psi CO.

b. 350 psi H<sub>2</sub> and 350 psi CO in THF (15 ml).

c. 800 psi CO.

d. Determined by capillary GC using a digital integrator (HP 5880A). Isolated by column chromatography (florisil) and identified by GC-MS and nmr spectroscopy (250 MHz, <sup>1</sup>H).

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720