

UCSF

UC San Francisco Previously Published Works

Title

Cardiac Alpha1-Adrenergic Receptors: Novel Aspects of Expression, Signaling Mechanisms, Physiologic Function, and Clinical Importance

Permalink

<https://escholarship.org/uc/item/6xx665vs>

Journal

Pharmacological Reviews, 66(1)

ISSN

0031-6997

Authors

O'Connell, Timothy D
Jensen, Brian C
Baker, Anthony J
et al.

Publication Date

2014

DOI

10.1124/pr.112.007203

Peer reviewed

ASSOCIATE EDITOR: PAUL A. INSEL

Cardiac α_1 -Adrenergic Receptors: Novel Aspects of Expression, Signaling Mechanisms, Physiologic Function, and Clinical Importance

Timothy D. O'Connell, Brian C. Jensen, Anthony J. Baker, and Paul C. Simpson

Department of Integrative Biology and Physiology, University of Minnesota, Minneapolis, Minnesota (T.D.O.); Cardiology Division, University of North Carolina, Chapel Hill, North Carolina (B.C.J.); and Cardiology Division, VA Medical Center and Cardiovascular Research Institute, University of California, San Francisco, San Francisco, California (A.J.B., P.C.S.)

Abstract.....	309
I. Introduction.....	310
II. α_1 -Adrenergic Receptor Expression in the Heart	310
A. α_1 -Adrenergic Receptor Expression in the Heart in Animal Models	310
B. Unique Aspects of α_1 -Adrenergic Receptor Expression Profiles in Cardiac Myocytes	311
C. α_1 -Adrenergic Receptor Expression in Human Heart.....	311
D. α_1 -Adrenergic Receptor Levels Increase Proportionately in Human Heart Failure	311
E. Conclusions on α_1 -Adrenergic Receptor Heart Expression.....	311
III. α_1 -AR Signaling in Cardiac Myocytes	312
A. Conventional Models of α_1 -Adrenergic Receptor Signaling	312
B. New Model for General G-Protein-Coupled Receptor Signaling: G-Protein-Coupled Receptors at the Nucleus	312
C. α_1 -Adrenergic Receptors in the Nuclei in Cardiac Myocytes.....	312
1. Nuclear Localization of α_1 -Adrenergic Receptors in Cardiac Myocytes	313
2. Mechanism of α_1 -Adrenergic Receptor Nuclear Localization	314
3. Receptor Orientation in the Inner Nuclear Membrane	314
4. Catecholamine Uptake in Cardiac Myocytes	314
5. Functional Evidence for Nuclear α_1 -Adrenergic Receptor Signaling.....	315
6. Localization of Signaling Partners with α_1 -Adrenergic Receptors in Cardiac Myocyte Nuclei	316
7. Nuclear α_1 -Adrenergic Receptor Localization in Vivo.....	316
8. Pathophysiologic Implications of Nuclear α_1 -Adrenergic Receptor Signaling	316
9. Summary of α_1 -Adrenergic Receptor Nuclear Localization.....	316
IV. α_1 -Adrenergic Receptor Physiologic Function in the Heart.....	317
A. α_1 -Adrenergic Receptors Activate Physiologic or Adaptive Hypertrophy	317
1. α_1 -Adrenergic Receptor-Mediated Hypertrophy in Cell Culture Models	317
2. α_1 -Adrenergic Receptor-Mediated Hypertrophy in Animal Models	318
3. Summary of α_1 -Adrenergic Receptor in Hypertrophy.....	319
B. α_1 -Adrenergic Receptors Prevent Cardiac Myocyte Death	319
1. α_1 -Adrenergic Receptor-Mediated Myocyte Survival Signaling in Cell Culture Models.....	319
2. α_1 -Adrenergic Receptor-Mediated Myocyte Survival Signaling in Animal Models.....	320
3. Summary of α_1 -Adrenergic Receptor-Mediated Myocyte Survival Signaling	320

This work was supported by the National Institutes of Health National Institute of General Medical Sciences [Grants P20-RR017662 (to T.D.O.)]; National Institutes of Health National Heart, Lung, and Blood Institute [Grants K08-HL096836 (to B.C.J.) and R01-HL31113 (to P.C.S.)]; the Department of Veteran's Affairs [Grants IO1-BX 001078 and 001970 (to P.C.S.), IO1 BX000740 (to A.J.B.)]; the American Heart Association, Western States Affiliate (to P.C.S., A.J.B.); the American Heart Association, Greater Midwest Affiliate (to T.D.O.); the GlaxoSmithKline Research and Education Foundation for Cardiovascular Disease (to B.C.J.); and the University of California, San Francisco, Foundation for Cardiac Research (to B.C.J.).

Address correspondence to: Dr. Paul C. Simpson, VA Medical Center (111-C-8), 4150 Clement St., San Francisco, CA 94121. E-mail: paul.simpson@ucsf.edu; or Dr. Timothy D. O'Connell, E-mail: tdoconne@umn.edu.
dx.doi.org/10.1124/pr.112.007203.

C.	α_1 -Adrenergic Receptors Augment Contractile Function	321
1.	α_1 -Adrenergic Receptor Activation of Contraction in In Vitro Models	321
2.	α_1 -Adrenergic Receptor-Mediated Contraction in Transgenic and Gene-Deletion Mouse Models	321
3.	α_1 -Adrenergic Receptor Activation of Contraction in Humans	322
4.	Summary of α_1 -Adrenergic Receptor Activation of Contraction	322
D.	α_1 -Adrenergic Receptors Induce Ischemic Preconditioning	322
1.	α_1 -Adrenergic Receptor-Mediated Preconditioning in Animal Models	322
2.	α_1 -Adrenergic Receptor-Mediated Preconditioning in Humans	322
3.	Summary of α_1 -Adrenergic Receptor-Mediated Preconditioning	323
E.	Conclusions: α_1 -Adrenergic Receptors Are Cardioprotective	323
1.	α_1 -Adrenergic Receptors are Cardioprotective and Prevent Pathologic Remodeling in Heart Failure Unlike Other G_q -Coupled Receptors	323
2.	α_1 -Adrenergic Receptor-Mediated Cardioprotective Signaling Can Explain the Worsening of Heart Failure with α_1 -Blockers Observed in Clinical Trials	323
3.	α_1 -Agonist Therapies Might Improve Heart Failure Outcomes	323
V.	α_1 -Adrenergic Receptors in Human Heart Disease	324
A.	Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial: An α_1 -Adrenergic Receptor Antagonist in Hypertension Increases the Risk of Heart Failure ..	324
B.	Vasodilator-Heart Failure Trial: An α_1 -Adrenergic Receptor Antagonist Does Not Improve Survival in Heart Failure	326
C.	α_1 -Adrenergic Receptor Antagonist Therapy in Benign Prostatic Hyperplasia Might Exacerbate Heart Failure	326
D.	Carvedilol: A Nonselective $\beta_{1/2}$ -Adrenergic Receptor/ α_1 -Adrenergic Receptor Antagonist for Heart Failure	326
E.	Sympatholytics: Reducing Norepinephrine Levels Does Not Improve Heart Failure	327
F.	Conclusions and Implications: Are Myocardial α_1 -Adrenergic Receptors Cardioprotective in Humans?	327
VI.	Final Summary	327
	References	328

Abstract—Adrenergic receptors (AR) are G-protein-coupled receptors (GPCRs) that have a crucial role in cardiac physiology in health and disease. Alpha₁-ARs signal through G_{α_q} , and signaling through G_q , for example, by endothelin and angiotensin receptors, is thought to be detrimental to the heart. In contrast, cardiac alpha₁-ARs mediate important protective and adaptive functions in the heart, although alpha₁-ARs are only a minor fraction of total cardiac ARs. Cardiac alpha₁-ARs activate pleiotropic downstream signaling to prevent pathologic remodeling in heart failure. Mechanisms defined in animal and cell models include activation of adaptive hypertrophy, prevention of cardiac myocyte death, augmentation of contractility, and induction of ischemic preconditioning. Surprisingly,

at the molecular level, alpha₁-ARs localize to and signal at the nucleus in cardiac myocytes, and, unlike most GPCRs, activate “inside-out” signaling to cause cardioprotection. Contrary to past opinion, human cardiac alpha₁-AR expression is similar to that in the mouse, where alpha₁-AR effects are seen most convincingly in knockout models. Human clinical studies show that alpha₁-blockade worsens heart failure in hypertension and does not improve outcomes in heart failure, implying a cardioprotective role for human alpha₁-ARs. In summary, these findings identify novel functional and mechanistic aspects of cardiac alpha₁-AR function and suggest that activation of cardiac alpha₁-AR might be a viable therapeutic strategy in heart failure.

ABBREVIATIONS: α_1 -AR, α_1 -adrenergic receptor; β -AR, β -adrenergic receptor; α_1A -subtype, α_1A -adrenergic receptor; α_1B -subtype, α_1B -adrenergic receptor; α_1 -blocker, α_1 -adrenergic receptor antagonist; α_1AKO , α_1A -adrenergic receptor knockout; α_1BKO , α_1B -adrenergic receptor knockout; α_1ABKO , α_1AB -adrenergic receptor double knockout; α SkAct, α -skeletal actin; AR, adrenergic receptor; ATR, angiotensin receptor; ALLHAT, Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial; BPH, benign prostatic hyperplasia; BEST, β -Blocker Evaluation of Survival Trial; cav-3, caveolin-3; CAM, constitutively active mutant; DAG, diacylglycerol; ETR, endothelin receptor; EMT, extraneuronal monoamine transporter; ERK, extracellular signal-regulated kinase; GFP, green fluorescent protein; GPCR, G-protein-coupled receptor; HEK, human embryonic kidney; HW, heart weight; IP, inositol phosphate; IP₃, inositol 1,4,5-trisphosphate; KO, knockout; MEK, mitogen-activated protein kinase kinase; MOXCON, Moxonidine Congestive Heart Failure Trial; MOXSE, Moxonidine Safety and Efficacy Trial; MyHC, myosin heavy chain; NE, norepinephrine; NFAT, nuclear factor of activated T cells; NRVM, neonatal rat ventricular myocyte; NYHA, New York Heart Association; OCT3, organic cation transporter 3; PLC β_1 , phospholipase C β_1 ; PKC, protein kinase C; PKD, protein kinase D; V-HeFT, Vasodilator-Heart Failure Trial; WT, wild type.

I. Introduction

Adrenergic receptors (ARs) bind to and are activated by the endogenous catecholamine hormones epinephrine and norepinephrine (NE). Epinephrine is primarily produced in and released to the circulation from the adrenal gland, whereas NE is synthesized in and released by sympathetic nerve terminals in the peripheral nervous system and brain. In the heart, the two main ARs are the β -ARs, which comprise roughly 90% of the total cardiac ARs, and α_1 -ARs, which account for approximately 10% (see section II).

In general, acute activation of cardiac β_1 -ARs, the predominant β -AR subtype (80% or more of total β -ARs in heart), induces positive inotropic and chronotropic responses, although in heart failure, where sympathetic activation and catecholamine levels are increased, long-term activation of β_1 -ARs exacerbates pathologic remodeling (Bristow, 2000; Naga Prasad et al., 2001; Lohse et al., 2003).

Less is known about cardiac α_1 -ARs, but studies from the last thirty years indicate that long-term activation of cardiac α_1 -ARs activates beneficial trophic signaling in the developing heart and that these α_1 -AR-mediated trophic effects in the adult, in many ways, counteract the negative effects of overstimulation of β_1 -ARs in heart failure. This review will focus on these trophic effects of cardiac α_1 -ARs and how activation of α_1 -ARs might be beneficial in heart failure.

There are three α_1 -AR subtypes, the α_1A , α_1B , and α_1D , and all three are expressed in the heart in a cell-type specific manner (section II). All three α_1 -ARs are G-protein-coupled receptors (GPCR), and classic α_1 -AR signaling mechanisms involve coupling to the $G_q/11$ (G_{α_q}) family of G-proteins and activation of phospholipase $C\beta_1$ ($PLC\beta_1$) at the plasma membrane. Activation of $PLC\beta_1$ cleaves phosphatidylinositol (PI), increasing inositol triphosphate (IP_3) and diacylglycerol (DAG). IP_3 binds to the IP_3 -receptor to release calcium from intracellular stores, and DAG activates protein kinase C (PKC).

Other G_q -coupled GPCRs that signal through G_{α_q} , such as endothelin receptors (ETRs) and angiotensin receptors (ATRs), are believed to play an important role in the pathogenesis of heart failure. Hallmarks of cardiomyopathy with heart failure include contractile dysfunction (both systolic and diastolic), myocyte hypertrophy, fibrosis, and increased cardiac cell death (Anand and Florea, 2003), which can all be worsened by G_q -coupled receptors (Salazar et al., 2007).

However, it also needs to be recalled that the view that G_q -coupled receptor signaling is toxic is based in large part on a transgenic mouse model with G_q overexpression that markedly exceeds the 2-fold increase found in human heart failure (Adams et al., 1998; Ponicke et al., 1998; Sakata et al., 1998) and thus cannot be considered to simulate human pathophysiology. In human heart failure, the maximal increase in

G_q abundance is 2-fold (Ponicke et al., 1998), and transgenic mice with 2-fold cardiomyocyte-specific G_q overexpression have no discernible cardiac phenotype (Adams et al., 1998; Sakata et al., 1998).

Furthermore, α_1 -ARs differ from other G_q -coupled receptors in several important ways, including expression limited to myocytes within the heart (section II) and localization and signaling at the nucleus, as discussed in section III.

Thus, unlike what can be seen with some G_q -coupled receptors, α_1 -ARs protect the heart by activating an adaptive or physiologic hypertrophy, preventing cardiac myocyte death, augmenting contractile function in heart failure and inducing preconditioning (section IV). Finally, clinical trials indicate that blockade of α_1 -ARs exacerbates heart failure (section V), which could be explained by the cardioprotective functions of α_1 -ARs identified in cell and animal models. This review summarizes these data, which span decades, and emphasizes recent findings from our laboratories.

II. α_1 -Adrenergic Receptor Expression in the Heart

A. α_1 -Adrenergic Receptor Expression in the Heart in Animal Models

In mice and rats, all three α_1 -AR subtype mRNAs, α_1A , α_1B , and α_1D , are detected in the heart (Rokosh et al., 1994; Stewart et al., 1994; Cavalli et al., 1997; O'Connell et al., 2003). Interestingly, among most species, including mouse, guinea pig, rabbit, pig, and cow, heart α_1 -AR levels determined by ligand binding are relatively constant (mouse: mean of six studies, ~ 12 fmol/mg protein) (Steinfath et al., 1992a; Cavalli et al., 1997; Yang et al., 1998; Lin et al., 2001; O'Connell et al., 2003; Rokosh and Simpson, 2002), with the exception of rat heart, in which α_1 -AR levels are approximately 10-fold higher (rat: mean of four studies, ~ 114 fmol/mg) (Steinfath et al., 1992a; Michel et al., 1994; Noguchi et al., 1995; Stewart et al., 1994).

Determination of cell-type specific expression of α_1 -ARs in the heart, or any tissue, is hampered by the lack of validated, subtype-specific α_1 -AR antibodies (Jensen et al., 2009c), which is a general problem with antibodies for GPCRs, as reviewed (Michel et al., 2009). However, studies in α_1 -AR knockout mice demonstrate that cardiac myocytes express only the α_1A - and α_1B -subtypes, based on lack of [3H]prazosin binding or functional responses in hearts from α_1AB -double knockout mice (α_1ABKO) (McCloskey et al., 2003; O'Connell et al., 2003; Turnbull et al., 2003) as well as lack of binding to a fluorescent α_1 -AR antagonist or signaling in cardiac myocytes isolated from α_1ABKO hearts (O'Connell et al., 2003; Wright et al., 2008).

Ligand binding studies further indicate that the α_1B is predominant, with the α_1A - and α_1B -subtypes

expressed in a 1:2–4 ratio in cardiac myocytes (Rokosh and Simpson, 2002; O'Connell et al., 2003). Despite the presence of α_1 D-subtype mRNA, rodent cardiac myocytes do not appear to express the α_1 D-subtype protein by binding (O'Connell et al., 2003). However, the α_1 D might be expressed in the coronary vasculature, based on studies demonstrating α_1 -AR mediated reductions in coronary flow in isolated α_1 -AR knockout hearts (Chalothorn et al., 2003; Turnbull et al., 2003). This idea is supported by human studies (below). Conversely, rodent cardiac fibroblasts do not express α_1 -ARs (Stewart et al., 1994; O'Connell et al., 2001), and α_1 -agonist infusion induces hypertrophy without fibrosis (Marino et al., 1991), suggesting that α_1 -AR activation does not exacerbate fibrosis associated with heart failure. In contrast with α_1 -ARs, most ATRs and ET_BRs are in fibroblasts, not cardiac myocytes (Kim et al., 1995; Gray et al., 1998; Modesti et al., 1999).

Long-term activation of α_1 -ARs and other hypertrophic agonists increases the α_1 A-subtype, without desensitizing α_1 -mediated inositol phosphate (IP) turnover or growth, while decreasing α_1 B-subtype mRNA and protein levels in cultured neonatal rat cardiac myocytes (NRVM) and in rats subjected to aortic banding (Rokosh et al., 1996). Moreover, total α_1 -AR levels are not altered in vivo by hypertrophy or heart failure in rats (Rokosh et al., 1996; Sjaastad et al., 2003), and α_1 -AR inotropic effects are maintained or increased (Wang et al., 2010), in contrast to β -ARs that are desensitized and downregulated in heart failure (Bristow et al., 1982; Bristow et al., 1988).

Partial explanation for the differences in desensitization of α_1 -AR and β -ARs might reside in expression and regulation of G-protein receptor kinases (GRKs). GRK3 is found exclusively in myocytes, regulates α_1 -ARs, and is not upregulated in heart failure (Vinge et al., 2001, 2007; Agüero et al., 2012). In contrast, GRK2 and GRK5 that desensitize β -ARs but not α_1 -ARs are expressed in many myocardial cell types and are upregulated in heart failure (Rockman et al., 1996; Eckhart et al., 2000; Vinge et al., 2001, 2007; Agüero et al., 2012).

B. Unique Aspects of α_1 -Adrenergic Receptor Expression Profiles in Cardiac Myocytes

Recent studies provide unique information on the expression and distribution of α_1 -ARs in cardiac myocytes. First, both the α_1 A- and α_1 B-subtypes localize to and signal at the nuclear membrane, but not the plasma membrane, in adult mouse cardiac myocytes (Huang et al., 2007; Wright et al., 2008; Wright et al., 2012), as reviewed in section III. Second, α_1 A-subtype expression and function are graded in adult cardiac myocytes, from high levels to none, whereas the α_1 B-subtype is expressed in all cardiac myocytes (unpublished data).

C. α_1 -Adrenergic Receptor Expression in Human Heart

In human heart, all three α_1 -AR subtype mRNAs are detected (Jensen et al., 2009a). Furthermore, α_1 -AR expression levels in human heart determined by ligand binding are similar to mouse and most other species (human: mean of 6 studies, \sim 12 fmol/mg protein) (Bohm et al., 1988; Bristow et al., 1988; Vago et al., 1989; Steinfath et al., 1992b; Hwang et al., 1996; Jensen et al., 2009a). Human myocardium has the α_1 A- and α_1 B-subtypes, with the α_1 B predominant, similar to other species (Jensen et al., 2009a,b), and the α_1 A is functional in signaling (R. C. Thomas and P. C. Simpson, unpublished data). These data suggest that the mouse is a more appropriate model to approximate cardiac α_1 -AR function than the rat, which as mentioned above, has roughly 10-fold more α_1 -ARs.

Competition binding experiments do not detect the α_1 D-subtype in explanted human heart (Jensen et al., 2009a,b). However, the α_1 D-subtype is expressed and functional in coronary artery smooth muscle cells and might cause vasoconstriction (Jensen et al., 2009b). The α_1 B-subtype is expressed in coronary artery endothelial cells and might induce vasodilation and angiogenesis (Jensen et al., 2010).

D. α_1 -Adrenergic Receptor Levels Increase Proportionately in Human Heart Failure

In heart failure, β_1 -ARs are desensitized and downregulated. In contrast, radioligand-binding studies indicate that myocardial α_1 -AR levels are slightly increased in human heart failure (mean of six studies, increased from \sim 12 to \sim 19 fmol/mg protein) (Bohm et al., 1988; Bristow et al., 1988; Vago et al., 1989; Steinfath et al., 1992b; Hwang et al., 1996; Jensen et al., 2009a). This means that α_1 -AR levels in the heart, which normally represent approximately 11% of the total AR population at baseline (range 2–23%, mean of six studies), are proportionately increased to approximately 25% of the total AR population in heart failure (range 9–41%) (Bohm et al., 1988; Bristow et al., 1988; Vago et al., 1989; Steinfath et al., 1992b; Hwang et al., 1996; Jensen et al., 2009a). Given that sympathetic drive and catecholamine levels are increased in heart failure (Cohn et al., 1984), this could imply that α_1 -ARs sustain adrenergic function when β_1 -ARs are downregulated. In fact, α_1 -AR-induced positive inotropy, which at baseline is minimal, can be equal to β -AR-mediated inotropy in ventricular muscle strips isolated from human heart failure patients (Skomedal et al., 1997), as reviewed in more detail below.

E. Conclusions on α_1 -Adrenergic Receptor Heart Expression

In summary, α_1 -ARs constitute a minority of the total cardiac AR population in humans at baseline, and

this seems to hold across species, with the exception of rats where α_1 -ARs levels are ~ 10 -fold higher than any other species. This should be considered when interpreting results from studies of α_1 -ARs in rats, particularly in cultured NRVMs, the most common cardiac myocyte culture model.

Cardiac myocytes of all species have all three α_1 -AR subtype mRNAs, but only the α_1A - and α_1B -subtype receptor proteins are detected. In humans, the α_1D -subtype is present in coronary smooth muscle and might regulate coronary vasoconstriction, whereas the α_1B -subtype is in coronary endothelial cells and might regulate vasodilation and angiogenesis. In contrast, α_1 -ARs are not expressed by cardiac fibroblasts.

In heart failure, α_1 -ARs are not downregulated as are β_1 -ARs and thus become a greater share (25%) of ARs in the heart. This increase in α_1 -ARs could suggest that α_1 -ARs have a compensatory or adaptive role in heart failure, as suggested by studies showing that α_1 -mediated inotropy can be similar to β -AR-mediated inotropy in heart failure (Skomedal et al., 1997). The idea that α_1 -ARs might have an adaptive and protective role in the heart is a central theme of this review and is discussed in following sections.

III. α_1 -AR Signaling in Cardiac Myocytes

The following sections review the conventional models of α_1 -AR localization and signaling at the plasma membrane, or “outside-in” signaling, and evidence for novel models, suggesting that α_1 -ARs and other GPCRs signal from the cardiac myocyte nucleus, or “inside-out” signaling.

A. Conventional Models of α_1 -Adrenergic Receptor Signaling

Conventional models of GPCR signaling describe receptor activation at the plasma membrane leading to initiation of downstream signaling within the cell, commonly referred to as “outside-in” signaling. Furthermore, classic models of GPCR function suggest that GPCRs are expressed on the cell membrane and are only internalized after receptor phosphorylation and subsequent desensitization (Drake et al., 2006). α_1 -ARs signal through the $G_q/11$ class of G-proteins, leading to activation of $PLC\beta_1$ and increases in IP_3 /calcium signaling and activation of PKC (Graham et al., 1996; Piascik and Perez, 2001). The α_1B -subtype might also signal through G_i (Hu and Nattel, 1995; Steinberg et al., 1985; Akhter et al., 1997; Melien et al., 2000; Snabaitis et al., 2005). In NRVM, historically the primary cell model used to study cardiac α_1 -AR signaling, α_1 -AR-induced increases in IP_3 are readily observed, but in adult cardiac myocytes this is controversial. The general consensus is that α_1 -ARs signal through the $G_q/PLC\beta_1$ - IP_3 /PKC pathway, but downstream signaling pathways are diverse, as reviewed

elsewhere (Hein and Michel, 2007; Cotecchia, 2010; Jensen et al., 2011). To date, over 70 downstream signaling molecules have been implicated in cardiac α_1 -AR signaling, using the NRVM model of α_1 -AR-stimulated cardiac myocyte hypertrophy (Jensen et al., 2011). Some data suggest interactions with β -arrestin (Pediani et al., 2005; Stanasila et al., 2008; Hennenberg et al., 2011) and $G\beta\gamma$ (Vettel et al., 2012).

B. New Model for General G-Protein-Coupled Receptor Signaling: G-Protein-Coupled Receptors at the Nucleus

It is now clear that several GPCRs localize to and signal at the nucleus, or “inside-out” signaling. Nuclear signaling is seen in several cell types, including neurons, hepatocytes, and cardiac myocytes, as reviewed previously (Gobeil et al., 2006; Boivin et al., 2008; Bkaily et al., 2009; Tadevosyan et al., 2012). The GPCRs include receptors for prostaglandin E_2 in the brain (Gobeil et al., 2002), angiotensin II (AT1R) in the brain and in HEK and Chinese hamster ovary cells (Lu et al., 1998; Chen et al., 2000; Lee et al., 2004), platelet activating factor in the liver and brain (Marrache et al., 2002), apelin in the brain (Lee et al., 2004), bradykinin in HEK cells (Lee et al., 2004), and glutamate in neurons (O'Malley et al., 2003).

Several recent studies show that GPCRs localize to nuclei in binucleate adult cardiac myocytes, as reviewed previously (Tadevosyan et al., 2012). Specifically, ETRs are detected on nuclei isolated from adult cardiac myocytes, and endothelin stimulates nuclear calcium transients (Boivin et al., 2003). ATRs and β -ARs are also detected on nuclei isolated from adult cardiac myocytes and mediate increased RNA synthesis (Boivin et al., 2006; Tadevosyan et al., 2010; Vaniotis et al., 2011). These findings indicate that GPCR localization to the nucleus could regulate important physiologic functions in adult cardiac myocytes. However, the majority of these other receptors can localize also to the myocyte plasma membrane; for example, 95% of ETRs are on the sarcolemma (Boivin et al., 2003; Wright et al., 2012), so that the relative functional significance of nuclear versus surface localization is uncertain.

Despite the data reviewed above, the prevalent view is that GPCRs, including α_1 -ARs, are localized primarily to the plasma membrane in heart and myocytes. This impression is based predominantly on radioligand binding to membrane fractions, binding assays in whole cells (Filipeanu et al., 2006) and studies with α_1 -AR antibodies. Difficulties with these approaches are discussed in the next section.

C. α_1 -Adrenergic Receptors in the Nuclei in Cardiac Myocytes

Cellular localization of signaling molecules determines function, emphasizing the importance of α_1 -AR subcellular localization in cardiac myocytes. The following

sections review the limitations and advantages of different approaches to detect α_1 -AR subcellular localization and the evidence that α_1 -ARs are in the cardiac myocyte nucleus, derived from studies of localization, agonist uptake, and signaling. Physiologic implications of nuclear α_1 -ARs are also suggested. This novel nuclear α_1 -AR signaling paradigm in cardiac myocytes is illustrated in Fig. 1.

1. Nuclear Localization of α_1 -Adrenergic Receptors in Cardiac Myocytes. Limitations with the techniques used to detect α_1 -ARs, radioligand binding and α_1 -AR antibodies, might explain the conventional view that α_1 -ARs localize mainly to the plasma membrane. Ligand binding assays typically involve homogenization of heart tissue or cultured cells followed by a high-speed ultracentrifugation to isolate total membrane fractions. This high-speed ultracentrifugation pulls down all membranes, and if subcellular markers are not used, this technique does not distinguish between plasma, sarcoplasmic, and nuclear membranes (Lin et al., 2001; Rokosh and Simpson, 2002; O'Connell et al., 2003). Furthermore, most purified membrane preparations exclude over 65 to 85% of total heart α_1 -ARs that are found in "debris" and low-speed pellets

discarded normally (Simpson, 2006). Whole-cell binding assays are limited by the lack of radioligands that do not enter the cell (Filipeanu et al., 2006).

Immunochemical detection, either by immunoblot or cell/tissue staining, is another commonly used technique to detect α_1 -ARs. However, none of 10 commercial α_1 -AR antibodies are specific for α_1 -ARs in general or for any subtype, as documented by the fact that no antibody detects a band in wild-type (WT) tissue that is absent in tissue from α_1 -AR knockout (KO) mice (Jensen et al., 2009c). This nonspecificity of anti-GPCR antibodies is a general problem, reviewed recently, emphasizing that α_1 -AR antibodies need to be validated using KO tissue (Michel et al., 2009). Nonspecificity of α_1 -AR antibodies calls into question previous reports with these reagents, for example, work suggesting α_1 -AR localization to the plasma membrane and t-tubules in adult rat cardiac myocytes (O-Uchi et al., 2008) or a study using immunoprecipitation of α_1 -ARs with potential signaling partners (Fujita et al., 2001).

An antibody to the 1D4 epitope tag at the C terminus of the α_1A detects surface membrane expression in heart sections of a transgenic mouse (Lin et al., 2001).




Fig. 1. Model for α_1 -AR signaling at the nuclear membrane. In adult cardiac myocytes, catecholamine α_1 -AR agonists (NE/PE) are actively transported into the myocyte via organic cation transporter 3 (OCT), which can be inhibited by corticosterone. The membrane-permeable α_1 -AR antagonist prazosin (and similar derivatives) can cross the plasma membrane to inhibit signaling, whereas the membrane impermeable α_1 -AR antagonist CGP12177A fails to inhibit signaling. The model suggests that active α_1 -ARs localize to the inner nuclear membrane with the ligand-binding domain facing the space between the outer and inner nuclear membranes (ONM and INM, respectively). On the basis of this orientation, binding of agonist to α_1 -ARs induces signaling inside the nucleus, possibly through $G\alpha_q$, although downstream intranuclear signaling pathways remain to be defined. We propose that activation of nuclear α_1 -ARs can induce intranuclear hypertrophic signaling as well as extranuclear signaling, including activation of ERK in caveolae and survival signaling or phosphorylation of cardiac troponin I at the sarcomere and contractile function. HDAC, histone deacetylase; Ca Ch, calcium channel; RYR, ryanodine receptor; PTP, mitochondrial permeability transition pore; ER/SR, endoplasmic/sarcoplasmic reticulum; NR, nucleoplasmic reticulum; NPC, nuclear pore complex.

However, it is problematic whether receptor localization with 170-fold overexpression simulates that of endogenous α_1 -ARs (Lin et al., 2001).

A few studies use membrane fractionation combined with the caveolar marker caveolin-3 (*cav-3*) to detect α_1 -AR binding in caveolae in NRVMs (Fujita et al., 2001; Lanzafame et al., 2006). In NRVMs, a caveolar fraction defined in this way contains most or all α_1 -mediated IP turnover (Morris et al., 2006) and 27% of total α_1 -AR binding, both α_1A and α_1B (Lanzafame et al., 2006). The value of 27% α_1 -AR binding in caveolae in NRVMs agrees well with a more recent study finding 20% of total α_1 -ARs in adult myocyte membranes defined by high levels of *cav-3* (Wright et al., 2008).

The contrary notion that α_1 -ARs localize primarily to the nucleus arises from three main lines of evidence. First, 80% of total α_1 -AR binding in adult mouse cardiac myocytes is found in nuclear membranes defined by the marker LAP2 (Wright et al., 2008, 2012). In NRVMs, nuclear α_1 -AR binding is also observed (Buu et al., 1993), and 73% of total α_1 -ARs are in noncaveolar membranes that might be nuclear (Lanzafame et al., 2006), in good agreement with the results in adult myocytes.

Second, BODIPY-prazosin is a fluorescent analog of the α_1 -AR antagonist prazosin that binds all three α_1 -AR subtypes with equal affinity and fluoresces only when bound to receptor (Daly et al., 1998; Mackenzie et al., 2000; Pediani et al., 2005). BODIPY-prazosin staining of living adult cardiac myocytes identifies endogenous α_1 -ARs on the nuclear membrane but does not detect receptors at the plasma membrane (Wright et al., 2008). Nuclei isolated from adult cardiac myocytes confirm positive BODIPY-prazosin staining of endogenous nuclear α_1 -ARs (Wright et al., 2012).

Third, a reconstitution system, in which α_1 -AR-GFP fluorescent fusion proteins are expressed in cultured adult α_1 ABKO cardiac myocytes, recapitulates the nuclear localization of the endogenous α_1 -ARs (Huang et al., 2007; Wright et al., 2008, 2012).

Studies in other cells provide some support for the results in cardiac myocytes. In recombinant cells expressing α_1 -ARs, for example, HEK293 cells, all α_1 -AR subtypes show some intracellular localization (Daly et al., 1998; Mackenzie et al., 2000; Chalothorn et al., 2002). In primary cultures of smooth muscle cells, endogenous α_1 -ARs are also found on both the plasma membrane and intracellular, using a fluorescent ligand, BODIPY-FL prazosin (Mackenzie et al., 2000).

2. Mechanism of α_1 -Adrenergic Receptor Nuclear Localization. The mechanism for nuclear targeting involves nuclear localization sequences embedded in the protein. These nuclear localization sequences typically consist of mono- or bi-partite basic residues, usually lysines and arginines or glycine-arginine repeats (Dono et al., 1998; Hock et al., 1998; Lu et al., 1998). Nuclear

localization sequences are recognized by a class of proteins known as importins that bind these sequences and facilitate transport of the target protein to the nucleus. This importin-mediated nuclear localization not only occurs for proteins that target to the nucleoplasm but for proteins that target the inner nuclear membrane as well (King et al., 2006; Cook et al., 2007; Lusk et al., 2007). Importin-mediated nuclear localization was previously described for the type 1 parathyroid hormone receptor (Pickard et al., 2006, 2007) and more recently for the gonadotropin-releasing hormone type 1 receptor (Re et al., 2010). Recent experiments identify nuclear localization sequences in the α_1A - and α_1B -subtypes, and mutation of these sequences results in loss of nuclear localization for each subtype in adult mouse cardiac myocytes (Wright et al., 2012).

3. Receptor Orientation in the Inner Nuclear Membrane. As described above, nuclear membrane proteins are targeted to the inner nuclear membrane through nuclear localization sequences, similar to proteins in the nucleoplasm. Also important is the orientation of inner nuclear membrane proteins, which could affect how they signal. For GPCRs, such as α_1 -ARs, if the ligand-binding domain faces the inside of the nucleus, the ligand would have to enter the nucleus and signaling would be initiated on the cytoplasmic side in the space between the inner and outer nuclear membranes. Conversely, if the ligand-binding domain faces the space between the outer and inner nuclear membranes, then signaling would be activated inside the nucleus.

Recent studies with nuclear GPCRs detect signaling in isolated nuclei, implying that nuclear receptors are likely oriented with the ligand-binding domain facing outward and the C terminus facing the nucleoplasm. ETRs induce calcium transients in isolated nuclei (Boivin et al., 2003), and β -ARs and ATRs induce transcriptional responses in isolated nuclei (Tadevosyan et al., 2010; Vaniotis et al., 2011). These studies suggest that nuclear GPCR signaling is activated inside the nucleus, thereby indicating an orientation in the inner nuclear membrane similar to GPCRs at the plasma membrane where the C terminus faces the cytoplasm.

4. Catecholamine Uptake in Cardiac Myocytes. A prerequisite for nuclear α_1 -AR signaling is that NE and other α_1 -AR ligands must traverse the plasma membrane, transit to the nucleus, and bind to and activate receptors in a time course consistent with signaling. In nonneuronal cells, this process is known as NE "uptake-2" (Obst et al., 1996) and is facilitated by extraneuronal monoamine transporter/organic cation transporter 3 (EMT/OCT3) (Zwart et al., 2001; Schomig et al., 2006). EMT/OCT3 is expressed most abundantly in heart (Zwart et al., 2001), where it is present on both the plasma and nuclear membranes in

adult cardiac myocytes (Wright et al., 2008). In neonatal myocytes, uptake of [3 H]NE is observed, but the time scale of nearly an hour before NE is detected in the nucleus is not sufficiently rapid to account for α_1 -AR signaling (Buu et al., 1993).

However, a more sensitive fluorescent-based catecholamine uptake assay shows that catecholamines are taken up very rapidly in cultured adult mouse cardiac myocytes. In this system, catecholamine uptake begins within seconds, is clearly increased by 5 minutes, peaks at 30 minutes, and is antagonized by addition of unlabeled NE 15 minutes prior to catecholamine uptake measurement, indicating specificity (Wright et al., 2008).

Further consistent with rapid uptake, the intrinsic uptake kinetics of OCT3, which is the rate at which one transporter moves a cation, show that OCT3-mediated cation transport is in the time frame of seconds. Thus, the uptake kinetics of catecholamines by recombinant OCT3 expressed in HEK293 cells is a $V_{\max} \sim 30,000$ pmol/mg protein/min and a $K_m \sim 900$ μ M for NE, and a $V_{\max} \sim 13,000$ pmol/mg protein/min and a $K_m \sim 500$ μ M for epinephrine (Duan and Wang, 2010). Catecholamine uptake is observed in seconds, with half-maximum response seen in ~ 2 minute (Duan and Wang, 2010). It is likely that the kinetic properties of the transporter are relatively consistent from cell to cell and that expression level will dictate the absolute amount of uptake and OCT3 expression is highest in heart (Zwart et al., 2001).

In mice, OCT3-mediated heart uptake of the neurotoxin cation methyl-4-phenylpyridinium acetate is observed within minutes of infusion (~ 4000 ng/g tissue 5 minutes after infusion), and this uptake is inhibited by 75% in OCT3KO mice (Zwart et al., 2001), indicating a rapid and robust uptake system. The phenotype of OCT3KO mice is further interesting. Thus, OCT3KO mice have a trend toward reduced heart size in males [WT heart weight (HW) 160 mg, OCT3KO HW 145 mg, $n = 7$, $P = 0.138$, a 10% reduction] (Zwart et al., 2001), reminiscent of the small heart phenotype seen in male α_1 ABKO mice (WT HW 147 mg, α_1 ABKO 122 mg, $n = 33-27$, $P < 0.05$, a 17% reduction) (O'Connell et al., 2003), but the number of OCT3KO mice analyzed was small ($n = 7$) (Zwart et al., 2001).

Thus, the kinetics of catecholamine uptake in myocytes (Wright et al., 2008) and the biochemistry and biology of OCT3 (Zwart et al., 2001; Duan and Wang, 2010) are consistent with α_1 -AR responses initiated by agonist activation of nuclear receptors.

In agreement with the idea that α_1 -AR agonist must be transported into the myocyte for signaling, there is a long latency of α_1 -AR responses after agonist addition, in contrast to the rapid onset for β -AR agonism. Specifically, the latency for contractile or calcium responses to α_1 -agonism in isolated myocytes is 2 to 5 minutes in nine studies (Tohse et al., 1990;

Terzic et al., 1992; Gambassi et al., 1998; Zhang et al., 1998; Woo and Lee, 1999; Ross et al., 2003; O-Uchi et al., 2005; Luo et al., 2007; Ichishima et al., 2010), rather than seconds as would be expected for a receptor at the sarcolemma.

Finally, there is functional evidence that agonist uptake is required for α_1 -AR signaling. Inhibition of EMT/OCT3-mediated catecholamine uptake with corticosterone, an EMT/OCT3 antagonist, prevents α_1 -AR activation of ERK in cultured adult mouse cardiac myocytes (Wright et al., 2008).

In summary, the kinetics of agonist uptake in myocytes, EMT/OCT3 biochemistry and biology, including kinetics, heart expression, and inhibition by corticosterone, and the latency of α_1 -AR physiologic responses are all consistent with agonist uptake and activation of nuclear α_1 -ARs.

5. Functional Evidence for Nuclear α_1 -Adrenergic Receptor Signaling. α_1 -ARs have numerous signaling effects in the cytosol, as reviewed later. Thus, if α_1 -ARs signal in the nucleus, then that signal must be transduced out of the nucleus to reach these cytosolic targets, defining an inside-out (nuclear-to-cytoplasmic) signaling mechanism. Three sets of data support the idea that α_1 -signaling is initiated in the nucleus. First, CGP-12177A [4-[3-[(1,1-dimethylethyl)amino]2-hydroxypropoxy]-1,3-dihydro-2H-benzimidazol-2-one hydrochloride], an α_1 -antagonist that does not cross membranes (Staehelin et al., 1983; Levin et al., 2002; Brahmadevara et al., 2003, 2004), does not block α_1 -AR-ERK signaling in cultured adult mouse cardiac myocytes, whereas the prototypical α_1 -AR antagonist prazosin, which freely crosses the plasma membrane, does block α_1 -AR-ERK signaling (Wright et al., 2008). Second, mislocalization mutants of both the α_1A - and α_1B -subtype, in which the nuclear localization sequences are mutated, do not activate ERK in cultured adult mouse cardiac myocytes (Wright et al., 2012). These mislocalization mutants are not redirected to the plasma membrane, which would provide a more crucial test of the requirement for nuclear localization, but the mutants do show that nuclear localization is required for α_1 -AR signaling in adult cardiac myocytes. Finally, activation of nuclear α_1 -ARs leads to the activation of ERK in caveolae at the plasma membrane (Wright et al., 2008), and the nuclear export inhibitor leptomycin B blocks α_1 -AR-mediated activation of ERK, suggesting that α_1 -AR signaling to ERK at caveolae must originate in the nucleus (Wright et al., 2012). How signals are transported from the nucleus to cytosolic targets is uncertain. However, α_1 -ARs activate PKC, a molecule known to translocate upon activation, suggesting a possible mechanism to transmit a signal out of the nucleus. Taken together, these studies provide functional evidence for α_1 -AR signaling initiated in the nucleus.

6. *Localization of Signaling Partners with α_1 -Adrenergic Receptors in Cardiac Myocyte Nuclei.* To effect nuclear α_1 -AR signaling in cardiac myocytes, α_1 -ARs must colocalize with downstream signaling partners in the nuclear membrane. However, their identities so far remain unclear.

A fraction of G_{α_q} colocalizes with α_1 -ARs at the nucleus, based on immunocytochemistry in α_1 ABKO cardiac myocytes expressing α_1 -AR-GFP fluorescent fusion proteins, and on subcellular fractionation of WT adult cardiac myocytes (Wright et al., 2008). In NRVMs, approximately 56% of G_{α_q} is in caveolar membranes defined by cav-3, and the remainder is in noncaveolar membranes, some of which might be nuclear (Morris et al., 2006).

A role in nuclear signaling is uncertain for $PLC\beta_1$, the classic α_1 -coupled PLC. $PLC\beta_1$ is detected in the nuclei of adult cardiac myocytes using the G12 $PLC\beta_1$ antibody from Santa Cruz Biotechnology (Santa Cruz, CA) (Wright et al., 2008), but this and other commercial $PLC\beta_1$ antibodies are not proven specific for $PLC\beta_1$ using KO tissues. In NRVMs, 91% of $PLC\beta_1$ is detected in a caveolar fraction using a Santa Cruz antibody (Lanzafame et al., 2006), and forced expression of $PLC\beta_1b$ with an N-terminal enhanced GFP tag detects localization on the sarcolemma but not the nucleus (Grubb et al., 2008). Also in NRVMs, expression of a C-terminal peptide from $PLC\beta_1b$ blocks α_1 - and G_q -mediated IP turnover and aspects of hypertrophy (Grubb et al., 2008; Filtz et al., 2009). Finally, the substrate for $PLC\beta_1$, phosphatidylinositol 4,5-bisphosphate, is not detected in nuclear membranes (Zhang et al., 2013). Taken together, these data suggest that $PLC\beta_1$ might not be involved in α_1 -AR nuclear signaling.

An interesting alternate mediator of nuclear α_1 -AR signaling is $PLC\epsilon$, which might be regulated by small GTPases (Rho, Ras, Rap) and $G\beta\gamma$ subunits, but not by G_{α_q} (Lopez et al., 2001). $PLC\epsilon$ in NRVMs and heart is scaffolded to muscle-specific A kinase-anchoring protein at the nuclear envelope with PKD, one key hypertrophic signaling molecule (Zhang et al., 2011, 2013). Knockdown of $PLC\epsilon$ inhibits α_1 - and G_{α_q} -stimulated hypertrophy in NRVMs, but has no effect on IP turnover, and expression of $PLC\epsilon$ causes hypertrophy, an effect that requires $PLC\epsilon$ catalytic activity (Zhang et al., 2011, 2013). Myocyte-specific $PLC\epsilon$ KO also inhibits hypertrophy with pressure overload in vivo (Zhang et al., 2013). Extensive evidence suggests that $PLC\epsilon$ mediates hypertrophy by hydrolysis of phosphatidylinositol 4-phosphate (PI4P) at the nuclear envelope, with generation of DAG and activation of PKD, but upstream mechanisms are uncertain (Zhang et al., 2013). Clearly, definition of nuclear α_1 -AR signaling partners is a promising area for the future.

7. *Nuclear α_1 -Adrenergic Receptor Localization in Vivo.* It is possible that localization observed in isolated or cultured cardiac myocytes does not reflect the true

localization of α_1 -ARs in vivo. As mentioned, overexpressed α_1A -ARs in transgenic mice localize to the plasma membrane based on immunohistochemical staining for an epitope tag (Lin et al., 2001). However, very high receptor levels, about 170-fold over basal, might cause artifactual localization, and the lack of validated α_1 -AR antibodies (Jensen et al., 2009c) make conventional immunohistochemical approaches problematic.

Conversely, in a different α_1A -subtype transgenic model, in which an α_1A -subtype GFP fusion protein is expressed at a much lower level, approximately 5-fold over basal, α_1 -ARs are detected at the nuclei in ventricular tissue sections with a GFP antibody (Wright et al., 2008). This result with the α_1A -GFP transgenic mice suggests that α_1 -AR nuclear localization observed in cultured cardiac myocytes can represent α_1 -AR localization in vivo.

8. *Pathophysiologic Implications of Nuclear α_1 -Adrenergic Receptor Signaling.* ETRs, ATRs, and β -ARs signal in isolated nuclei from adult cardiac myocytes (Boivin et al., 2003; Tadevosyan et al., 2010; Vaniotis et al., 2011). However, it is difficult to assign a functional significance to nuclear signaling by these GPCRs in cardiac myocytes, because the majority of ETRs, ATRs, and β -ARs localize to the plasma membrane (although quantitative ligand binding in subcellular fractions for ATRs is not possible due to low level of expression). Conversely, approximately 80% of α_1 -ARs localize to the nuclei in adult mouse cardiac myocytes. Interestingly, in pathologic settings, α_1 -AR signaling is clearly protective (sections IV and V), whereas ETR and ATR signaling can exacerbate pathologic remodeling (Harada et al., 1999; Yang et al., 2004). This raises the possibility that differences in receptor localization might lead to differences between physiologic and pathologic signaling. In other words, nuclear receptors, like α_1 -ARs, might be protective, whereas ETRs and ATRs at the plasma membrane might induce pathologic signaling (Wright et al., 2012). Although these ideas remain to be tested, differential localization of G_q -coupled receptors could have significant implications for their physiologic functions and for therapeutic targeting of G_q -coupled receptors in heart disease.

9. *Summary of α_1 -Adrenergic Receptor Nuclear Localization.* Overall, the majority of current data supports the idea that α_1 -ARs localize to and signal from the nuclei in adult cardiac myocytes in vitro and in vivo. Identification of functional nuclear localization sequences in each α_1 -subtype provides a mechanistic basis to support nuclear α_1 -AR localization, oriented with the C-terminal tail in the nucleoplasm. Ligand uptake into the cell via EMT/OCT3 provides a mechanism for receptor activation. The signaling mechanisms of nuclear α_1 -ARs remain unclear, as do the physiologic implications of nuclear versus

sarcolemmal signaling by α_1 -ARs and other G_q -coupled receptors.

IV. α_1 -Adrenergic Receptor Physiologic Function in the Heart

In the heart, AR physiology is largely focused on acute β -AR mediated regulation of contractile function, whereas chronic β -AR signaling is maladaptive and β -AR antagonists are now standard therapy in heart failure. Short-term α_1 -AR signaling can increase contractility, as reviewed below, but this has not been studied in detail in vivo. On the other hand, many studies now indicate that chronic α_1 -AR signaling is adaptive, protecting the heart from pathologic stress through activation of physiologic hypertrophy, survival signaling, augmentation of contractility, and ischemic preconditioning. These data are described below.

A. α_1 -Adrenergic Receptors Activate Physiologic or Adaptive Hypertrophy

Cardiac myocyte hypertrophy is the most common cellular response in the heart to pathologic stress, but hypertrophy is not always maladaptive (Frey and Olson, 2003). Cardiac hypertrophy occurs during normal physiologic development and in response to exercise and also as an adaptive response to pathologic stress. Physiologic or adaptive hypertrophy is characterized by an increase in heart and cardiac myocyte size without fibrosis and an overall improvement in function.

In contrast, pathologic or maladaptive hypertrophy is characterized by an increase in heart and cardiac myocyte size accompanied by combinations of cardiac cell death, fibrosis, vessel loss, reduced innervation, and, most importantly, declining function. Clinically, cardiac hypertrophy in Framingham adults is correlated with a significantly increased risk of heart failure and sudden death (Levy et al., 1990). Thirty years of research from cell culture to genetically modified mice, summarized below, demonstrates clearly that α_1 -ARs can mediate a physiologic or adaptive form of cardiac hypertrophy that offsets pathologic remodeling in heart failure.

1. *α_1 -Adrenergic Receptor-Mediated Hypertrophy in Cell Culture Models.* Primary cultures of NRVMs are the most common cell culture model used to examine signaling in cardiac myocytes. The original and now classic experiments (Glembotski, 2013) demonstrated that catecholamines acting through α_1 -ARs produce a direct trophic response in NRVMs (Simpson et al., 1982; Simpson, 1983, 1985). At the time, there was debate as to whether catecholamines induced hypertrophy through increasing blood pressure or through a direct action on cardiac myocytes, that is, whether myocyte hypertrophy was regulated in some way only by "load" or whether growth factors and their receptors

were involved, as in other types of cells. These papers were the first demonstration that catecholamines induce cardiac myocyte hypertrophy directly. This finding was later confirmed in cultured adult rat and cat cardiac myocytes (Simpson, 1988; Fuller et al., 1990; Ikeda et al., 1991; Volz et al., 1991; Clark et al., 1993).

Cardiac hypertrophy is clearly linked to induction of gene transcription, and in NRVMs, α_1 -ARs induce a pattern of hypertrophic gene transcription characterized by re-expression of genes normally expressed only in the fetal heart (Simpson et al., 1989). Early studies identified a group of these "fetal genes" induced by α_1 -ARs, including c-myc (Starksen et al., 1986), atrial natriuretic factor (Knowlton et al., 1991, 1993), α -skeletal actin (α SkAct) (Bishopric et al., 1987; Long et al., 1989; Karns et al., 1995), and β -myosin heavy chain (β MyHC) (Waspe et al., 1990; Kariya et al., 1993, 1994).

A crucial study of endogenous transcription in intact NRVMs proved that α_1 -ARs stimulate transcription not only of a fetal gene (α SkAct), but also all RNA species, including the mRNA for an adult gene (cardiac actin), and total RNA (ribosomal and transfer) (Long et al., 1989). Several subsequent studies focusing on α_1 -AR-mediated transcriptional regulation delineated a host of transcriptional factors and modifiers activated by α_1 -ARs in cardiac myocytes, including TEF-1 (Kariya et al., 1993, 1994; Karns et al., 1995; McLean et al., 2003), GATA-4 (Morimoto et al., 2000; Liang et al., 2001a,b), Egr-1 (Jin et al., 2000), Elk 1 (McWhinney et al., 2000), Vgl-4 (Chen et al., 2004), Rlf (Post et al., 2002), CREB (Markou et al., 2004), Zfp260 (Debrus et al., 2005), and class 2 histone deacetylase (Vega et al., 2004; Liu et al., 2009).

Mechanistically, the α_1A -subtype is implicated in α_1 -AR mediated hypertrophy in NRVMs through the use of α_1 -AR subtype-specific pharmacologic agents (Autelitano and Woodcock, 1998). Several mechanisms for α_1 -AR-mediated hypertrophic signaling are proposed, and a multitude of signal transducers are implicated (Jensen et al., 2011). Certain molecules, based on frequency in the literature, might be considered essential or "core" molecules required for α_1 -AR-mediated hypertrophic signaling, including PLC (Filtz et al., 2009; Zhang et al., 2013), PKC (α , δ , and ϵ , three main isotypes activated by α_1 -ARs) (Henrich and Simpson, 1988; Kariya et al., 1991, 1993, 1994; Karns et al., 1995; Haworth et al., 2000; Rohde et al., 2000; Braz et al., 2002, 2004; Vega et al., 2004; Carnegie et al., 2008), PKD (Haworth et al., 2000; Vega et al., 2004; Harrison et al., 2006; Avkiran et al., 2008; Bossuyt et al., 2008, 2011; Carnegie et al., 2008; Liu et al., 2009), ERK (Bueno et al., 2000; Xiao et al., 2001; Barron et al., 2003; O'Connell et al., 2003), and class 2 histone deacetylase (Vega et al., 2004; Backs et al., 2006, 2008; Harrison et al., 2006; Liu et al., 2009).

Evidence also exists that transactivation of the EGFR is involved in α_1 -mediated hypertrophy (Morris et al., 2004; Guo et al., 2009; Li et al., 2011; Papay et al., 2013).

In total, α_1 -mediated hypertrophy in NRVMs is characterized by α_1 A-subtype-mediated activation of the "fetal gene program" along with general increases in transcription of all RNA species and protein synthesis (Simpson, 1985; Long et al., 1989). Because the fetal gene program is often associated with pathologic hypertrophy, it was believed originally that α_1 -ARs induce a pathologic hypertrophy, similar to high levels of G_q overexpression (Dorn and Brown, 1999). This idea proved to be incorrect.

2. α_1 -Adrenergic Receptor-Mediated Hypertrophy in Animal Models. Early studies in mice, cats, and dogs showed that long-term catecholamine infusion in various species and at doses that do not increase blood pressure produces cardiac hypertrophy in vivo, which is "physiological," in that cardiac function is normal or improved and there is no fibrosis (Laks et al., 1973; King et al., 1987; Marino et al., 1991; Patel et al., 1991; Stewart et al., 1992; Vecchione et al., 2002). Whereas these studies suggest clearly that activation of ARs induces cardiac hypertrophy directly, which was debated at the time, the lack of subtype-specific AR pharmacologic agents limits mechanistic insight. However, in preliminary experiments, infusion of a subpressor dose of an α_1 A agonist in mice can increase fetal gene expression (unpublished data).

The advent of transgenic mouse technology provided a platform to address AR subtype-specific function in vivo, and transgenic gain-of function and gene deletion loss-of-function models have mostly confirmed cell culture studies indicating that α_1 -ARs regulate hypertrophy, with some prominent exceptions.

Cardiac myocyte-specific transgenic overexpression of the WT α_1 A-subtype, even at very high levels (148- to 170-fold), does not alter heart size, although α_1 A transgenic mice eventually develop dilated cardiomyopathy and die prematurely (Lin et al., 2001; Chaulet et al., 2006). On the other hand, transgenic overexpression of constitutively active mutant (CAM) α_1 A with the endogenous α_1 A-promoter induces cardiac hypertrophy without an effect on systemic blood pressure (Papay et al., 2013).

Similarly, the α_1 B-subtype shows a variable ability to induce hypertrophy when overexpressed, depending on the model. Overexpression of a CAM of the α_1 B-subtype with the α -myosin heavy chain (α MyHC) promoter at low levels (2- to 3-fold) induces hypertrophy (Milano et al., 1994) and exacerbates pathologic remodeling after aortic constriction (Wang et al., 2000). Likewise, systemic overexpression of a CAM α_1 B with the endogenous α_1 B-promoter also induces cardiac hypertrophy, along with *hypotension*, and a decreased pressor response, clearly dissociating hypertrophy from

blood pressure (Zuscik et al., 2001). In the same study, overexpression of a wild-type (WT) α_1 B with the endogenous α_1 B-promoter shows a lesser degree of hypertrophy (Zuscik et al., 2001). More recently, a different result was found in that overexpression of the CAM α_1 B caused hypertrophy, but only in older mice, and hypertrophy was associated with fibrosis (Papay et al., 2013). Furthermore, hypertrophy seen in both the CAM α_1 A and α_1 B mice was not observed when the mice were interbred to derive a systemic CAM α_1 AB transgenic mouse (Papay et al., 2013). In contrast with these results, relatively high-level overexpression of the WT α_1 B with the α MyHC promoter (>40-fold) does not induce hypertrophy but results in dilated cardiomyopathy and death (Akhter et al., 1997; Grupp et al., 1998; Iaccarino et al., 2001; Lemire et al., 2001).

α_1 -AR gene-deletion models are reviewed in detail (Simpson, 2006). In brief, heart size is not different in the α_1 A-knockout on a mixed FVB/129SvJ background (α_1 AKO) (Rokosh and Simpson, 2002) or in the α_1 B-knockout on a mixed C57Bl/6/129SvJ background (α_1 BKO) (Cavalli et al., 1997). Knockout of the α_1 D-subtype, which is not expressed in cardiac myocytes, also has no effect on heart size (Tanoue et al., 2002; Chalothorn et al., 2003; Hosoda et al., 2005).

However, double knockout of both the α_1 A- and α_1 B-subtypes, which eliminates α_1 -AR binding in the heart, on a congenic C57Bl/6J background (α_1 ABKO) causes a 15% reduction in heart and cardiac myocyte size during normal postnatal development (O'Connell et al., 2003). Mechanistically, ERK activity is reduced 30% in α_1 ABKO hearts, and α_1 -mediated activation of ERK is absent in cultured α_1 ABKO cardiac myocytes, suggesting that α_1 -AR-ERK signaling might regulate hypertrophic growth during postnatal development (O'Connell et al., 2003). Importantly, α_1 ABKO mice have normal basal blood pressure, normal body and organ weights, normal home cage locomotor activity, and normal overall health (O'Connell et al., 2003, 2006). These data show that any effects of the α_1 A- and α_1 B-subtypes on maintenance of basal blood pressure can be compensated by other receptors, but that heart growth requires the α_1 A and/or α_1 B. Maximal contractile responses to phenylephrine in α_1 ABKO isolated arteries are reduced by 35% in carotid and by 77% in mesenteric, with little or no changes in pEC_{50} , indicating compensation by α_1 D (Methven et al., 2009). We have not tested α_1 -mediated pressor response in the intact α_1 ABKO, but they are reduced in both the α_1 AKO and α_1 BKO (Cavalli et al., 1997; Rokosh and Simpson, 2002).

Interestingly, a preliminary re-examination of the α_1 -AR single knockouts on a congenic C57Bl/6J background reveals that α_1 BKO mice have small hearts, similar to the α_1 ABKO, suggesting that the α_1 B-subtype alone is required for *physiologic* postnatal growth of the heart (unpublished data). In support of

this, subpressor catecholamine infusion does not cause hypertrophy in an α_1 BKO on a mixed genetic background (Vecchione et al., 2002).

Aortic constriction in the α_1 ABKO mice results in a worse dilated cardiomyopathy, with fibrosis, apoptosis, decreased contractility, and increased mortality contrasted to WT mice (O'Connell et al., 2006). Interestingly, the final degree of hypertrophy in α_1 ABKO hearts after aortic constriction is similar to that in WT hearts, but induction of the fetal gene program is lost (O'Connell et al., 2006). This shows that the absence of α_1 -ARs exacerbates *pathologic* hypertrophic responses and that induction of the fetal-gene program can be uncoupled from pathologic hypertrophy.

3. Summary of α_1 -Adrenergic Receptor in Hypertrophy. Early studies in NRVMs demonstrated that the α_1 A-subtype induces hypertrophy with activation of the fetal-gene program and overall RNA and protein synthesis. In vivo, infusions of subpressor doses of catecholamines cause a physiologic hypertrophy but generally do not pinpoint which α_1 -subtype might be responsible. Although studies from α_1 -AR transgenic mice provide inconsistent results, studies from α_1 -AR knockout mice suggest that the α_1 B-subtype is required for hypertrophic growth during postnatal cardiac development, a period of physiologic heart growth, and that α_1 -ARs are not required for *pathologic* hypertrophy after aortic constriction. In fact, pathologic hypertrophy is worse in the absence of α_1 -ARs.

Therefore, in vivo studies, either with catecholamine infusion or α_1 -AR knockout mice, collectively suggest that α_1 -ARs stimulate an adaptive or *physiologic* hypertrophy, with no decrease in contractile function. Interestingly, findings from NRVMs suggesting that α_1 -ARs induce pathologic hypertrophy based on activation of the fetal gene program are not supported by findings in α_1 ABKO mice. In α_1 ABKO mice, activation of the fetal gene program is absent despite significant hypertrophy and worse cardiomyopathy after aortic constriction.

A new consideration with regard to the fetal gene program is that a classic fetal gene, β -MyHC, is re-expressed only in a minor subpopulation of cardiac myocytes in the mouse heart after aortic constriction, and the myocytes with β -MyHC are smaller than the cells with α -MyHC, not larger (Lopez et al., 2011). These data question whether fetal genes are even markers of hypertrophy or pathology (Lopez et al., 2011).

B. α_1 -Adrenergic Receptors Prevent Cardiac Myocyte Death

Cell death, either apoptotic, necrotic, or autophagic, plays a significant role in the development of heart failure (Guerra et al., 1999; Kostin et al., 2003; Wencker et al., 2003; Baines et al., 2005; Foo et al., 2005; Nakayama et al., 2007; Nishida and Otsu, 2008;

Baines, 2010; Whelan et al., 2010; Nemchenko et al., 2011). Whereas substantial evidence indicates that β_1 -ARs induce cell death, a growing body of research summarized below indicates that α_1 -ARs prevent cardiac myocyte cell death in direct opposition to β_1 -ARs.

1. α_1 -Adrenergic Receptor-Mediated Myocyte Survival Signaling in Cell Culture Models. In cultured cardiac myocytes, NE stimulates apoptotic cell death through activation of β_1 -ARs, whereas β_2 -ARs are believed to be cytoprotective (Mann et al., 1992; Xiao et al., 2004). Interestingly, several studies indicate that α_1 -ARs are also cytoprotective and act antithetically to β_1 -ARs. In NRVM, the α_1 -AR agonist phenylephrine inhibits apoptosis induced by the β -agonist isoproterenol (Iwai-Kanai et al., 1999; Zhu et al., 2000), nonhydrolyzable cAMP analogs (Iwai-Kanai et al., 1999; Zhu et al., 2000), hypoxia (Zhu et al., 2000), serum starvation (Zhu et al., 2000), 2-deoxyglucose (Valks et al., 2002), and doxorubicin (Aries et al., 2004). Similar results are observed in cultured adult rat cardiac myocytes, where NE-induced apoptosis is abolished by the β -AR antagonist propranolol, but not the α_1 -AR antagonist prazosin (Communal et al., 1998; O'Connell et al., 2006).

Mechanistically, a variety of pathways are implicated in α_1 -AR-mediated survival signaling in cardiac myocytes. α_1 -AR survival signaling requires activation of ERK and subsequent regulation of Bcl-2 family members to stabilize the mitochondrial membrane (Iwai-Kanai et al., 1999; Zhu et al., 2000; Valks et al., 2002; Communal et al., 2003; Huang et al., 2007; Wright et al., 2008, 2012). In NRVM, phenylephrine inhibits apoptosis induced by serum starvation and hypoxia by preventing downregulation of Bcl-2 and Bcl-X mRNA and protein levels (Zhu et al., 2000). In addition, phenylephrine induces the phosphorylation of Bcl-2 family member Bad at Ser112 and Ser155, preventing 2-deoxyglucose-induced apoptosis (Valks et al., 2002). Interestingly, cAMP-dependent protein kinase (PKA), known to be downstream of β_1 -ARs, also stimulates phosphorylation of Bad at Ser136 (Valks et al., 2002). This finding might suggest that both β_1 -AR and α_1 -AR signaling converge on a single molecule to modulate cell survival, implying that the balance between β_1 - and α_1 -AR signaling could control cell fate. However, how the interplay between α_1 -AR and β -AR phosphorylation of Bad impacts cardiac myocyte survival is unclear.

Other studies propose a role for ERK as a regulator of α_1 -AR survival signaling. ERK mediates cytoprotective signaling in cardiac myocytes (Lips et al., 2004), and α_1 -AR mediated activation of ERK is a well-characterized signaling pathway involved in hypertrophy (Bueno et al., 2000; Xiao et al., 2001; Barron et al., 2003; O'Connell et al., 2003). In NRVM, the MEK-1 inhibitor PD098059 [2-(2-amino-3-methoxyphenyl)-4H-

1-benzopyran-4-one] negates phenylephrine-mediated survival signaling (Iwai-Kanai et al., 1999). Furthermore, in cultured adult rat cardiac myocytes, NE-mediated activation of ERK upregulates β_1 -integrin and protects cells against β_1 -AR-mediated apoptosis (Communal et al., 2003).

The most convincing evidence for ERK in α_1 -mediated survival signaling comes from α_1 ABKO myocytes. Cultured α_1 ABKO mouse myocytes have markedly increased necrosis and apoptosis with toxic stimuli, including hydrogen peroxide, doxorubicin, and β -AR stimulation (O'Connell et al., 2006; Huang et al., 2007). This sensitivity to death stimuli in α_1 ABKO myocytes is rescued by expression of the α_1A -subtype but not by the α_1B , indicating that the α_1A is necessary and sufficient for myocyte survival (Huang et al., 2007). Furthermore, rescue is mimicked by expression of constitutively activated MEK, which increases ERK activity, and rescue by the α_1A is prevented by dominant negative MEK, which inhibits ERK (Huang et al., 2007). Together, these experiments define an α_1A -ERK pathway for myocyte survival (Huang et al., 2007).

Potential mediators downstream of ERK include p90Rsk, a potential kinase for α_1 -AR-mediated phosphorylation of Ser112 in Bad (Valks et al., 2002), and the transcription factors GATA4 and nuclear factor of activated T cells (NFAT) (Pu et al., 2003; Aries et al., 2004).

Overall, the data from cultured cardiac myocytes indicate that α_1 -ARs mediate survival signaling, potentially through α_1A activation of ERK, leading to regulation of Bcl-2 family members and stabilization of the mitochondrial membrane, and/or by induction of GATA4 and NFAT.

2. α_1 -Adrenergic Receptor-Mediated Myocyte Survival Signaling in Animal Models. In mice, cats, and dogs, long-term infusion of subpressor doses of the mixed α_1/β -AR agonist NE produces an adaptive hypertrophy without increased cell death or fibrosis (Laks et al., 1973; King et al., 1987; Marino et al., 1991; Patel et al., 1991; Stewart et al., 1992; Vecchione et al., 2002). Furthermore, α_1 -AR stimulation in isolated, perfused hearts prevents ischemia-reperfusion-induced cell apoptosis and necrosis in mice (Tejero-Taldo et al., 2002), rats (Banerjee et al., 1993; Mitchell et al., 1995; Tosaki et al., 1995; Meng et al., 1996a,b, 1999; Meldrum et al., 1997; Imani et al., 2008), rabbits (Bankwala et al., 1994; Tsuchida et al., 1994; Cope et al., 1997; Baghelai et al., 1999a,b), and dogs (Kitakaze et al., 1987, 1991, 1994; Node et al., 1997).

However, gain-of-function α_1 -AR transgenic models are inconsistent in demonstrating that α_1 -ARs protect against cardiac cell death. Cardiac myocyte-specific transgenic overexpression of the α_1A -subtype at high levels (66-fold) protects against pathologic stress from pressure overload induced by aortic constriction and

ischemic injury induced by coronary artery ligation, although the mechanism is linked to a basal hypercontractile phenotype rather than prevention of cardiac myocyte death (Du et al., 2004, 2006). Moreover, by 1 year, ventricles from these α_1A -transgenic mice show increased fibrosis and apoptotic labeling, and the mice die prematurely, indicating that long-term, very high-level α_1A -subtype overexpression can be associated with cell death rather than survival signaling (Chaulet et al., 2006).

Although transgenic overexpression of the α_1B -subtype shows a variable ability to induce hypertrophy, prolonged overexpression of the α_1B -subtype in some models, although not directly linked to increased cell death, induces a pathologic remodeling (Grupp et al., 1998; Iaccarino et al., 2001; Lemire et al., 2001; Wang et al., 2000). The failure of these gain-of-function models to recapitulate the findings in NRVM or in other animal models might be linked to the high levels of overexpression in most of these models or failure of overexpressed receptors to recapitulate signaling by ligand-activated endogenous receptors.

Conversely, loss-of-function models clearly indicate that α_1 -ARs prevent cardiac myocyte cell death. In α_1 ABKO mice, which lack the two α_1 -AR subtypes expressed in cardiac myocytes, aortic constriction induces a worse dilated cardiomyopathy, accompanied by a significant increase in cardiac cell apoptosis and fibrosis, leading to decreased function and increased mortality compared with WT. Cultured α_1 ABKO cardiac myocytes have increased susceptibility to several pro-death agonists, as reviewed in the preceding section (O'Connell et al., 2006; Huang et al., 2007, 2008), and this is rescued by adenoviral mediated reconstitution of the α_1A -subtype, but not the α_1B -subtype, in a pathway that requires ERK (Huang et al., 2007). The absence of this α_1A -subtype ERK survival signaling might explain, at least partially, the negative outcome in α_1 ABKO mice subjected to aortic constriction (O'Connell et al., 2006).

In support of the finding that the α_1A -subtype is both sufficient and necessary to prevent cardiac myocyte death, long-term infusion of a subpressor concentration of the α_1A -subtype-specific agonist A61603 [*N*-[5-(4,5-dihydro-1*H*-imidazol-2-yl)-2-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl]methanesulphonamide hydrobromide] prevents cell death and pathologic remodeling associated with doxorubicin-induced cardiotoxicity (Chan et al., 2008; Dash et al., 2011).

3. Summary of α_1 -Adrenergic Receptor-Mediated Myocyte Survival Signaling. Studies in cultured neonatal and adult cardiac myocytes show that α_1 -ARs mediate survival signaling, most likely through activation of ERK and subsequent regulation of Bcl-2 family members to preserve mitochondrial membrane stability, as well as induction of protective transcription factors, such as GATA4 and NFAT, with their own

downstream effectors. These findings generally support early studies in which catecholamine infusion in several animal models stimulated an adaptive hypertrophy without cell death or fibrosis. However, gain-of-function studies in α_1 A- and α_1 B-transgenic mouse models are inconsistent with regard to survival signaling, possibly due to massive levels of overexpression and/or aberrant signaling by constitutively activated receptors. More importantly, loss-of-function models, particularly α_1 ABKO mice, support the notion that α_1 -ARs mediate survival signaling. Furthermore, studies in cultured α_1 ABKO cardiac myocytes with reconstitution of the α_1 A-subtype define an α_1 A-subtype ERK survival signaling pathway, the absence of which could at least partially explain the maladaptive responses to pathologic stress in α_1 ABKO mice.

C. α_1 -Adrenergic Receptors Augment Contractile Function

Contractile dysfunction is a major component of and a causative factor in heart failure progression. Furthermore, as β -AR-mediated inotropy declines in heart failure due to receptor desensitization and down-regulation, α_1 -AR mediated inotropy, which contributes little to basal contractile function, might function in a compensatory role to preserve contractile function in the failing heart.

1. *α_1 -Adrenergic Receptor Activation of Contraction in In Vitro Models.* In many species, α_1 -ARs induce a positive inotropic response in left ventricular myocytes, trabeculae, and the perfused heart (Endoh and Blinks, 1988; Terzic et al., 1992; Turnbull et al., 2003). However, in mice, the α_1 -AR inotropic response can be negative in a few left ventricular preparations, including isolated papillary muscles and a minority of isolated cardiac myocytes (Hirano et al., 2006; Chu et al., 2013). Interestingly, populations of cardiac myocytes from the right and left ventricle have a fraction of myocytes that have a positive inotropic response to phenylephrine and an increased Ca^{2+} transient, prominent in the left ventricle, and a fraction of myocytes that have a negative inotropic response to phenylephrine and a decreased Ca^{2+} transient, mainly in the right ventricle (Chu et al., 2013).

Another unexpected finding in mouse heart is that α_1 -ARs mediate negative inotropy in myocardium from the normal right ventricle but positive inotropy in left ventricular myocardium (Wang et al., 2010). Surprisingly, in heart failure caused by myocardial infarction, α_1 -AR inotropy in right ventricular myocardium switches from negative to positive, and α_1 -AR positive inotropy in left ventricular myocardium is preserved (Litwin et al., 1995; Wang et al., 2010). These two aspects of α_1 -AR-mediated positive inotropy in heart failure after myocardial infarction can be interpreted as adaptive, that is, undiminished inotropy in the left ventricle (Litwin et al., 1995; Wang et al., 2010) and

the appearance of positive inotropy in the right ventricle (Wang et al., 2010). Notably, right ventricular failure predicts worse outcomes in patients with left ventricular failure (Bleasdale and Frenneaux, 2002).

Interestingly, some find that the α_1 A- and α_1 B-subtypes differentially regulate contraction, with the α_1 A-subtype mediating a positive response and the α_1 B-subtype a negative response (Gambassi et al., 1998; Lin et al., 2001; Ross et al., 2003; O-Uchi et al., 2008).

A multitude of mechanisms are proposed to explain α_1 -AR-mediated positive or negative inotropy. Proposed mechanisms include inhibition of outward K^+ currents and increased action potential duration (Apkon and Nerbonne, 1988; Fedida et al., 1989, 1990, 1991; Ravens et al., 1989; Tohse et al., 1990, 1992; Wang et al., 1991, 2001; Braun et al., 1992; Sato and Koumi, 1995; Gaughan et al., 1998); inhibition of L-type Ca^{2+} channel current (Chen et al., 1996; Gaughan et al., 1998; Belevych et al., 2001) or activation (Zhang et al., 1998; Mohl et al., 2011; Chu et al., 2013); activation of the Na^+/H^+ exchanger and intracellular acidification (Gambassi et al., 1992; Terzic et al., 1992); and regulation of myofilament Ca^{2+} sensitivity through phosphorylation of myosin light chain and/or cardiac troponin I (Hartmann et al., 1995; Andersen et al., 2002; McCloskey et al., 2003; MacGowan et al., 2005; Wang et al., 2006, 2010).

2. *α_1 -Adrenergic Receptor-Mediated Contraction in Transgenic and Gene-Deletion Mouse Models.* Cardiac myocyte-specific transgenic overexpression of the α_1 A-subtype at high levels (148- to 170-fold) increases basal contractile function (Lin et al., 2001) and limits pathologic remodeling from pressure overload and ischemic injury (Du et al., 2004, 2006). These results suggest that the α_1 A-subtype mediates positive inotropic responses, in agreement with recent in vitro studies (Mohl et al., 2011; Chu et al., 2013). Conversely, transgenic overexpression of the α_1 B-subtype can be associated with depressed contractile function and pathologic remodeling in the heart (Grupp et al., 1998; Wang et al., 2000; Iaccarino et al., 2001; Lemire et al., 2001).

Loss-of-function models suggest that α_1 -AR-mediated inotropic responses are not required for basal contractile function, but might prevent contractile decline in response to pathologic stress. Basal contractile function by echocardiography is normal in both α_1 AKO and α_1 BKO mice (Rokosh and Simpson, 2002; Vecchione et al., 2002). In a similar fashion, basal contractile function assessed by echocardiography in α_1 ABKO mice is similar to WT, although cardiac output is decreased, due to a slight bradycardia and reduced left-ventricular volume, and exercise performance is impaired, presumably for the same reasons (O'Connell et al., 2003). Interestingly, calcium sensitivity is increased and maximal force is decreased in isolated

muscle strips from α_1 ABKO mice, suggesting subtle abnormalities in basal contractile function with long-term absence of α_1 -ARs (McCloskey et al., 2003). More strikingly, contractile function is impaired significantly by aortic constriction in α_1 ABKO mice (O'Connell et al., 2006). This contractile dysfunction might be caused by both the loss of α_1 -AR-mediated positive inotropy and the absence of α_1 -AR-mediated survival signaling and adaptive hypertrophic effects, such as increased myosin synthesis. The exaggerated contractile dysfunction in the α_1 ABKO confirms that α_1 -AR-mediated inotropy could play an important protective role in response to pathologic stress in the heart.

3. α_1 -Adrenergic Receptor Activation of Contraction in Humans. In healthy young women, systemic infusion of the α_1 -AR agonist methoxamine increases contractility determined noninvasively (Curiel et al., 1989). In both healthy patients and patients with New York Heart Association (NYHA) II–IV heart failure, infusion of the α_1 -AR agonist phenylephrine into the left main coronary artery increases contractility measured as dp/dt, demonstrating α_1 -AR inotropy in both healthy and heart failure patients (Landzberg et al., 1991). Interestingly, in the same patients, the α_1 -AR antagonist phentolamine shows no effect on baseline contractile function, suggesting that there is little contribution of α_1 -AR inotropy to basal contractile function in humans (Landzberg et al., 1991).

Surprisingly, α_1 -AR-mediated inotropy can equal β -AR-mediated inotropy in trabeculae isolated from failing human hearts (Skomedal et al., 1997). This suggests that in human heart failure, as β -ARs are desensitized and downregulated, α_1 -AR mediated contractility might act in a compensatory role to maintain contractile function. Similarly, inhalation of the α_1 -AR agonist methoxamine improves exercise performance in patients with significant left ventricular contractile dysfunction (Cabanès et al., 1992). Furthermore, in a small cohort of patients with hypotension due to end stage heart failure, the use of the α_1 -AR agonist midodrine is associated with increased contractile function (Zakir et al., 2009). The authors attributed the benefit to a modest increase in blood pressure that resulted from activation of vascular α_1 -ARs, permitting up-titration of recommended heart failure medications (Zakir et al., 2009). However, it is possible that direct activation of cardiac α_1 -ARs contributed to this beneficial effect.

4. Summary of α_1 -Adrenergic Receptor Activation of Contraction. Studies with isolated myocytes, muscle strips, and perfused hearts show that α_1 -ARs can induce either positive or negative inotropic responses by altering K^+ and Ca^{2+} currents, intracellular pH, and myofilament Ca^{2+} sensitivity. Mouse models confirm that the α_1A -subtype can induce positive inotropic responses in vitro and in vivo and that α_1A -subtype-mediated inotropy might protect the heart from pathologic

stress. Importantly, α_1 -AR-mediated positive inotropy is documented in intact humans and can be equal to β -AR inotropy in isolated trabeculae from human heart failure patients, identifying a clinically relevant adaptive function for α_1 -ARs.

D. α_1 -Adrenergic Receptors Induce Ischemic Preconditioning

Ischemic preconditioning is an intrinsic protective mechanism in the heart whereby transient periods of ischemia protect the myocardium from damage due to longer bouts of ischemia, and protection can be observed both early (minutes to hours after ischemia) and late (hours to days). Pharmacologic agents can also induce preconditioning, and α_1 -ARs are among the most effective (Jensen et al., 2011).

1. α_1 -Adrenergic Receptor-Mediated Preconditioning in Animal Models. In several animal models, including dog (Kitakaze et al., 1987, 1991, 1994; Node et al., 1997), rabbit (Bankwala et al., 1994; Tsuchida et al., 1994; Cope et al., 1997; Baghelai et al., 1999a,b), rats (Banerjee et al., 1993; Mitchell et al., 1995; Tosaki et al., 1995; Meng et al., 1996a,b, 1999; Meldrum et al., 1997; Imani et al., 2008) and mice (Tejero-Taldo et al., 2002), α_1 -ARs induce both early and late preconditioning, through a variety of mechanisms including adenosine release (Kitakaze et al., 1991), activation of 5'-nucleotidase activity (Kitakaze et al., 1994; Node et al., 1997), activation of PKC (Tsuchida et al., 1994; Mitchell et al., 1995; Node et al., 1997; Meng et al., 1999), regulation of Bcl2 family members (Baghelai et al., 1999a), induction of heat-shock proteins and protein synthesis (Meng et al., 1996a,b), activation of mitochondrial K-ATP channels (Imani et al., 2008), and induction of iNOS (Tejero-Taldo et al., 2002; Zhao et al., 2012). Once again though, poor specificity of antagonists for the α_1 -subtypes has made it difficult to define the α_1 -AR subtype(s) responsible for ischemic preconditioning.

However, transgenic overexpression of constitutively active mutants of the α_1A - and α_1B -subtypes reveals that the α_1A -subtype, but not the α_1B , mediates ischemic preconditioning that might not involve PKC (Rorabaugh et al., 2005). Preconditioning by the α_1A in a rat transgenic model could be mediated by MEK/ERK phosphorylation and iNOS (Zhao et al., 2012). Similarly, cardiac myocyte-specific transgenic overexpression of the α_1A -subtype suppresses ischemia-reperfusion-induced IP₃ generation in isolated, perfused hearts (Amirahmadi et al., 2008). Conversely, cardiac myocyte-specific transgenic overexpression of the α_1B -subtype does not prevent ischemic preconditioning or prevent ischemic-reperfusion injury (Gao et al., 2000).

2. α_1 -Adrenergic Receptor-Mediated Preconditioning in Humans. In human atrial and ventricular muscle strips, α_1 -ARs mediate ischemic preconditioning through activation of PKC, p38 MAPK, and opening of

mitochondrial K-ATP channels (Cleveland et al., 1996, 1997; Loubani and Galinanes, 2001, 2002).

3. *Summary of α_1 -Adrenergic Receptor-Mediated Preconditioning.* In short, the data indicate that α_1 -ARs, most likely the α_1 A-subtype based on studies in mouse and rat, induce ischemic preconditioning and prevent cardiac myocyte death from ischemic injury.

E. Conclusions: α_1 -Adrenergic Receptors Are Cardioprotective

Three fundamental conclusions emerge based on 30 years of studies examining the physiologic function of α_1 -ARs in the heart that tend to contradict the conventional wisdom regarding cardiac α_1 -ARs.

1. *α_1 -Adrenergic Receptors are Cardioprotective and Prevent Pathologic Remodeling in Heart Failure Unlike Other G_q -Coupled Receptors.* Specifically, the α_1 A-subtype induces cardioprotection and positive inotropy, whereas the α_1 B-subtype might be required for an adaptive, physiologic hypertrophy. Cardiac remodeling mediated by G_q -coupled receptors, such as α_1 -ARs, ETRs, and ATRs, is arguably the most significant physiologic function of these receptors in the heart. Currently, it is widely believed that all G_q -coupled receptors mediate a pathologic hypertrophic response. The idea that G_q -signaling is pathologic is based primarily on three lines of evidence, as reviewed previously (Dorn and Brown, 1999; Adams and Brown, 2001). First, G_q -agonists, such as phenylephrine (an α_1 -AR agonist), endothelin, and angiotensin, induce hypertrophy with expression of the “fetal genes” in cultured NRVM, and re-expression of the “fetal genes” is classically associated with pathologic ventricular remodeling, as reviewed previously (Dorn and Brown, 1999). Second, mouse models targeting overexpression of ETRs or ATRs suggest that these receptors generally induce pathologic remodeling (Ainscough et al., 2009; Paradis et al., 2000; Yang et al., 2004). Clinically, ATR blockers are used to treat heart failure (Chrysant, 2008), although current ETR antagonists have no proven efficacy for heart failure (Mylona and Cleland, 1999; McMurray et al., 2007). Third, cardiac-specific transgenic overexpression of $G\alpha_q$ in mice induces pathologic remodeling with increased cardiac myocyte apoptosis, as reviewed (Dorn and Brown, 1999; Dorn, 2005). However, high-level $G\alpha_q$ -overexpression is required for pathology, as reviewed previously (Jensen et al., 2011). Furthermore, the data summarized in this section clearly challenge the dogma that all G_q -coupled receptor signaling is pathologic. Instead, the data indicate α_1 -AR signaling can be cardioprotective.

2. *α_1 -Adrenergic Receptor-Mediated Cardioprotective Signaling Can Explain the Worsening of Heart Failure with α_1 -Blockers Observed in Clinical Trials.* As detailed in the next section (section V), α_1 -blockers exacerbate heart failure and are associated with worse outcomes in patients with hypertension (ALLHAT),

heart failure (V-HeFT), and benign prostatic hyperplasia (Cohn, 1993; Cohn et al., 1986; ALLHAT, 2000, 2003; Dhaliwal et al., 2009). Hallmarks of ventricular remodeling in heart failure include contractile dysfunction (both systolic and diastolic), pathologic hypertrophy, and increased cardiac cell death and fibrosis (Anand and Florea, 2003). Importantly, α_1 ABKO mice by virtue of their lack of cardiac myocyte α_1 -ARs approximate the use of α_1 -blockers (O’Connell et al., 2003). In the α_1 ABKO mouse model, pathologic stress from aortic constriction causes hypertrophy with failed gene transcription, increased cardiac cell death, increased fibrosis, and worsened contractile function, leading to dilated cardiomyopathy, heart failure, and ultimately 50% mortality (O’Connell et al., 2003, 2006). Follow-up studies in cultured α_1 ABKO cardiac myocytes define a cardioprotective α_1 A-subtype signaling pathway, identifying a direct requirement for protective α_1 -AR signaling in cardiac myocytes, the absence of which could at least partially explain the negative outcomes in α_1 ABKO mice (Huang et al., 2007, 2008). Support for the assertion that α_1 -ARs are cardioprotective can also be drawn from studies in gain-of-function models, where α_1 A-subtype overexpression protects against pathologic stress (Du et al., 2004, 2006; Lin et al., 2001). In combination, these studies indicate that α_1 -ARs are both sufficient to induce and required for cardioprotective signaling. In summary, these data provide a mechanistic basis to explain the negative outcomes in clinical trials with α_1 -blockers.

3. *α_1 -Agonist Therapies Might Improve Heart Failure Outcomes.* On the basis of the accumulated evidence, α_1 -AR activation of adaptive hypertrophy, prevention of cardiac myocyte death, augmentation of contractility, and induction of ischemic preconditioning could prevent worsened outcomes from systolic heart failure. This provides the foundation of the argument for α_1 -AR agonist therapy in heart failure. However, several counterarguments exist (Jensen et al., 2011). First, in transgenic models, the α_1 B-subtype can worsen function and induce dilated cardiomyopathy (Grupp et al., 1998; Wang et al., 2000; Iaccarino et al., 2001; Lemire et al., 2001). However, pharmacology and knockouts argue against the results seen in certain cardiac transgenics. Specifically, studies with α_1 -agonists in mouse (Chan et al., 2008; Dash et al., 2011) and human (Cleveland et al., 1996, 1997; Loubani and Galinanes, 2001, 2002), as well as in loss-of-function mouse models (O’Connell et al., 2003, 2006; Huang et al., 2007) as reviewed above, support α_1 -mediated cardioprotective effects.

Second, α_1 -ARs induce vasoconstriction, which is contraindicated in heart failure. However, in mice, suppressor doses of an α_1 A-subtype-specific ligand prevent doxorubicin cardiotoxicity (Chan et al., 2008; Dash et al., 2011); in humans, some small trials

indicate α_1 -ARs agonists might improve function in heart failure (Cabanès et al., 1992; Zakir et al., 2009); and numerous studies identify adaptive hypertrophy with suppressor α_1 -agonist infusion, as reviewed in the section on hypertrophy. These data provide a preliminary proof-of-principle demonstration of the efficacy of α_1 -agonists in heart failure, and justify further study.

A third argument against α_1 -agonist therapy is that the mixed α_1/β -blocker carvedilol has proven efficacy in heart failure. However, as discussed in the next section (section V), current evidence suggests that the α_1 -blocking properties of carvedilol are not sustained in long-term dosing (Kubo et al., 2001; Hryniewicz et al., 2003), and conversely carvedilol might potentiate α_1 -AR signaling (Van Tassel et al., 2008).

Fourth, α_1 -ARs are G_q -coupled receptors, and the conventional wisdom is that G_q -signaling exacerbates pathologic remodeling (Jensen et al., 2011). However, as mentioned already, α_1 -ARs clearly do not fit this paradigm. In summary, despite the caveats listed, α_1 -AR agonist therapy might present a novel effective treatment of heart failure.

V. α_1 -Adrenergic Receptors in Human Heart Disease

The classic physiologic function of α_1 -ARs is to increase vascular smooth muscle contractility and hence blood pressure. α_1 -ARs are found in vascular beds throughout the body, including smooth muscle cells in arteries of the heart, brain, kidneys, and gut, as well as in smooth muscle in the prostate and bladder (Michelotti et al., 2000). By virtue of their ability to block α_1 -AR-mediated smooth muscle contractions, α_1 -AR antagonists (α_1 -blockers) are used to treat hypertension (Lund-Johansen and Omvik, 1991; Frishman and Kotob, 1999; Sica, 2005) and benign prostatic hyperplasia (Caine et al., 1976, 1978; Schwinn and Roehrborn, 2008; Michel, 2010).

More recently, it has become clear that α_1 -ARs might play a significant role in preventing the clinical progression of heart failure, as reviewed above. Heart failure is a clinical syndrome of varied etiology in which the heart cannot pump enough blood to meet the body's needs. Heart failure is characterized by neuro-hormonal augmentation, leading to increased catecholamine levels, which are thought to play a causative role in pathologic ventricular remodeling through increased activation of ARs. Indeed, increased blood NE levels are a primary finding in heart failure patients and predict disease severity and mortality (Cohn et al., 1984). Whereas this does not establish a causal relationship between increased NE levels and induction of heart failure, it was part of the rationale to block NE activation of ARs for therapy in heart failure.

In fact, clinical trials with β -AR antagonists or β -blockers, such as Metoprolol CR/XL Randomized

Intervention Trial in Congestive Heart Failure (metoprolol) (MERIT, 1999), Cardiac Insufficiency Bisoprolol Study II (bucindolol) (CIBIS-II, 1999) and Carvedilol Prospective Randomized Cumulative Survival Trial (carvedilol) (Packer et al., 2001), show significant reductions in mortality in patients with heart failure (Foody et al., 2002; Teerlink and Massie, 1999; Chatterjee et al., 2013). On the basis of the success of β -blockers in improving outcomes in heart failure, β -blockers are standard of care in heart failure therapy (Hunt et al., 2005). This success has led to the notion that blocking all AR signaling in heart failure would be beneficial, and tests of this idea are reviewed below. Currently, 5.7 million Americans have heart failure, 1 million are admitted to the hospital each year, and the 5-year survival rate is only 50% (Roger et al., 2011). Thus, new drugs to treat heart failure are needed (Simpson, 2011). Here, we review the recent clinical trials suggesting that some AR signaling, particularly α_1 -AR signaling, might be beneficial in human heart failure. Table 1 summarizes the trials.

A. Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial: An α_1 -Adrenergic Receptor Antagonist in Hypertension Increases the Risk of Heart Failure. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT) was a large, randomized, double-blind, active controlled trial initiated in 1994 and funded by the National Heart, Lung, Blood Institute (ALLHAT, 2000, 2003). ALLHAT was designed to compare new treatments for hypertension versus older, standard treatments. Primary outcomes were fatal coronary heart disease and nonfatal myocardial infarction, and secondary outcomes included all-cause mortality, stroke, and combined cardiovascular disease. In one arm, 24,335 patients with hypertension and at least one other risk factor for coronary heart disease were randomized to the diuretic chlorthalidone (15,268) or the nonselective α_1 -AR antagonist (α_1 -blocker) doxazosin (9067) and were to be followed for 4–8 years.

In 2000, the ALLHAT data safety and monitoring board stopped this arm of the trial early, citing that patients on doxazosin had 25% more cardiovascular events and a significant doubling in the risk of heart failure versus patients on chlorthalidone (SoRelle, 2000). Although systolic blood pressure was approximately 3 mm Hg higher in the doxazosin group, the ALLHAT investigators concluded that this difference was unlikely to account for the doubling in the risk of heart failure (Davis et al., 2002; ALLHAT, 2003), and heart failure events were validated (Piller et al., 2002). Furthermore, approximately 60% of patients in both groups were on additional treatments to reduce blood pressure, but a follow up analysis revealed that the risk of heart failure was not reduced by this additional antihypertensive treatment, thereby confirming the initial findings (Davis et al., 2002). Subsequently, the

TABLE 1
Summary of clinical trials involving α_1 -blockers in cardiovascular disease

Trial	Treatment Groups and Patient Numbers		Inclusion Criteria	Result	Interpretation
	Control Groups	α_1 -Blocker Groups			
ALLHAT (2000) (ALLHAT, 2000, 2003; SoRelle, 2000; Davis et al., 2002; Piller et al., 2002)	Chlorthalidone (diuretic) [12.5–25 mg/day (<i>N</i> = 15,268)]	Doxazosin (α_1 -blocker) [2–8 mg/day (<i>N</i> = 9067)]	Men and women age 55 or older with hypertension (systolic 140 mm Hg and/or diastolic 90 mm Hg, or on medication for hypertension) and at least 1 other risk factor for coronary heart disease	Trial stopped early. Doxazosin increased cardiovascular events by 25% and doubled the risk of heart failure.	α_1 -Blockade worsens outcomes in hypertension with cardiac risk factors.
V-HeFT I and II (Cohn et al., 1986; Cohn, 1993)	V-HeFT I Placebo (<i>N</i> = 273); Hydralazine/ isosorbide dinitrate (direct vasodilators) [300/160 mg/day (<i>N</i> = 186)] V-HeFT II Hydralazine/ isosorbide dinitrate Enalapril (ACE inhibitor) (<i>N</i> = 403)	Prazosin (α_1 -blocker) [20 mg/day (<i>N</i> = 183)]	Male veterans mean age 58 with symptomatic heart failure due to dilated cardiomyopathy (ischemic or nonischemic), on digoxin and diuretics	Prazosin did not change left ventricular ejection fraction or mortality versus placebo; both were improved with hydralazine/isosorbide. Trend toward increased mortality at 5 years in prazosin group versus placebo, survival improved by other vasodilators.	α_1 -Blockade shows no benefit in heart failure and might worsen mortality.
α_1 -Blockers in BPH effect on HF (Dhaliwal et al., 2009)		Tamsulosin (58%), terazosin (40%), or doxazosin (2%) (<i>N</i> = 98)	Male veterans with dilated cardiomyopathy admitted with heart failure (<i>N</i> = 388 overall)	Among the 25% of patients taking α_1 -blockers, most for BPH, subsequent hospitalizations for heart failure were increased in patients receiving α_1 -blockers without β -blockers.	α_1 -Blockade worsens outcomes in BPH in the absence of β -blockade.
COMET (Carvedilol or Metoprolol European Trial) (Poole-Wilson et al., 2002, 2003)	Metoprolol tartrate (β_1 -selective antagonist) (50 mg bid; <i>N</i> = 1518)	Carvedilol ($\beta_{1/2}$ - α_1 -AR antagonist) [25 mg bid (<i>N</i> = 1511)]	Men (80%) and women mean age 62 with NYHA class II–IV heart failure due to dilated cardiomyopathy (ischemic or nonischemic) on stable therapy	Mortality reduced in carvedilol group versus metoprolol. Subsequent analyses indicates that benefit might not be due to α_1 -blockade (Kubo et al., 2001; Hryniewicz et al., 2003; Van Tassel et al., 2008)	Carvedilol reduces mortality, possibly related in part to potentiation of α_1 -AR signaling.
MOXSE (Swedberg et al., 2002)	Placebo (<i>N</i> = 38)	Moxonidine (sympatholytic) [0.3–1.5 mg bid (<i>N</i> = 227 total)]	Patients NYHA class II–IV heart failure due to dilated cardiomyopathy (ischemic or nonischemic)	Moxonidine reduced plasma NE and heart rate but increased adverse events	Some degree of AR signaling is cardioprotective.
MOXCON (Coats, 1999; Cohn et al., 2003; Pocock et al., 2004)	Placebo (<i>N</i> = 944)	Moxonidine (1.5 mg bid [<i>N</i> = 990])	As in MOXSE	Trial stopped early Moxonidine reduced NE and increased morbidity and mortality	Some degree of AR signaling is cardioprotective.
BEST (Bristow et al., 2004)	Placebo [3 month (<i>N</i> = 845); 12 month (<i>N</i> = 654)]; Total for BEST (<i>N</i> = 2708)	Bucindolol ($\beta_{1/2}$ -AR antagonist, sympatholytic) [3 month (<i>N</i> = 841); 12 month (<i>N</i> = 642)]	Patients NYHA class III–IV heart failure due to dilated cardiomyopathy (ischemic or nonischemic) on stable therapy, subgroup with NE measured at 3 and 12 months	Bucindolol reduced NE and increased mortality.	Some degree of AR signaling is cardioprotective.

ALLHAT results led to the recommendation against α_1 -blockers as a primary treatment of high blood pressure (Messerli, 2001); compliance with this recommendation has been modest (Stafford et al., 2004).

When it was initiated, ALLHAT was the largest trial yet to compare the efficacy of different methods for treating hypertension on cardiovascular outcomes. At the time, it was assumed that lowering blood pressure, regardless of mechanism, would by itself reduce morbidity and mortality (Messerli, 2000). However, ALLHAT shows that clinical trials involving new antihypertensive treatments must examine multiple cardiovascular outcomes.

More importantly, ALLHAT demonstrates that blocking α_1 -ARs has a negative impact on the heart, further challenging the established dogma that elevated AR signaling in heart disease is always pathologic. Although ALLHAT does not prove that the negative effect of α_1 -blockade is a direct effect on the heart, a direct effect is supported by studies on the α_1 ABKO mice and other animal and human data discussed above.

B. Vasodilator-Heart Failure Trial: An α_1 -Adrenergic Receptor Antagonist Does Not Improve Survival in Heart Failure. The Vasodilator-Heart Failure Trial (V-HeFT), phase I, was a small trial initiated in 1980 involving 642 men diagnosed with chronic congestive heart failure. V-HeFT was designed to evaluate the effects of vasodilators on mortality as the primary outcome (Cohn et al., 1986). Prior to V-HeFT, studies indicated that reducing systemic vascular resistance improved hemodynamics in patients with heart failure (Cohn and Franciosa, 1977a,b), and V-HeFT was designed to test whether this principal would translate into clinical benefit. Patients with heart failure and taking digoxin and a diuretic, a common therapeutic regimen for heart failure at the time, were randomized to receive placebo, the α_1 -blocker prazosin, or the combination of hydralazine and isosorbide dinitrate. At a mean follow-up of 2.3 years, the combination of hydralazine/isosorbide increased cardiac function (ejection fraction) and reduced mortality, but function and mortality were the same as placebo for prazosin (Cohn et al., 1986).

In phase II, the angiotensin converting enzyme inhibitor enalapril was included, and mortality was tracked in all groups from phase I and II for 5 years. As in phase I, prazosin showed no benefit in phase II, and at 5 years, a trend toward increased mortality was observed (Cohn, 1993). Interestingly, prazosin was the only vasodilator that failed to show any positive outcome. Although not as clear-cut as the results from ALLHAT, the negative results with prazosin in V-HeFT again suggest that α_1 -AR blockade is harmful to the heart and that α_1 -AR activation is indeed beneficial.

A caveat to consider with both prazosin (V-HeFT) and doxazosin (ALLHAT) is that both antagonists can cause myocyte apoptosis, independent of their α_1 -blocking activity (Gonzalez-Juanatey et al., 2003). However, the

dose required to cause apoptosis in culture (10 μ M) (Gonzalez-Juanatey et al., 2003) is far higher than the peak plasma levels attained in clinical use, e.g., \sim 200 nM for doxazosin (Fawzy et al., 1999). Furthermore, the concern that the α_1 -blockers prazosin and doxazosin might be maladaptive for the human heart via an off-target effect is largely obviated by the observation of maladaptive cardiac effects in the α_1 ABKO, as reviewed above.

C. α_1 -Adrenergic Receptor Antagonist Therapy in Benign Prostatic Hyperplasia Might Exacerbate Heart Failure. In the prostate, α_1 -ARs mediate smooth muscle contraction, which is the basis for α_1 -blocker use in benign prostatic hyperplasia (BPH). Currently, estimates indicate that 9.5 million men over 65 are diagnosed with benign prostatic hyperplasia (Vaughan, 2003), with common comorbidities for hypertension (87%) and a previous admission for heart failure (40%) (Dhaliwal et al., 2009). The U.S. Food and Drug Administration has approved five α_1 -blockers for BPH: silodosin (Rapaflo), an α_1 A-subtype-selective antagonist, and the nonselective α_1 -antagonists terazosin (Hytrin), doxazosin (Cardura), tamsulosin (Flomax), and alfuzosin (Uroxatral).

A recent meta-analysis of 388 men with heart failure revealed that cotreatment with an α_1 -blocker increased the risk of heart failure hospitalizations, unless patients were concurrently treated with a β -blocker for heart failure (Dhaliwal et al., 2009). Given that only two-thirds of patients in this trial were being treated with a β -blocker (Dhaliwal et al., 2009) and had the common comorbidities of BPH, hypertension, and heart failure, there is reason for concern regarding the safety of α_1 -blockers in BPH.

D. Carvedilol: A Nonselective $\beta_{1/2}$ -Adrenergic Receptor/ α_1 -Adrenergic Receptor Antagonist for Heart Failure. Carvedilol is a nonselective $\beta_{1/2}$ -AR/ α_1 -AR antagonist indicated for the treatment of heart failure based on its efficacy as established in several trials, for example, Carvedilol Prospective Randomized Cumulative Survival Trial (Packer et al., 2001, 2002) and US Carvedilol (Packer et al., 1996), as reviewed previously (Teerlink and Massie, 1999; Foody et al., 2002; Wollert and Drexler, 2002; Chatterjee et al., 2013). Interestingly, the Carvedilol or Metoprolol European Trial suggested that carvedilol extended survival relative to metoprolol, a β_1 -AR selective antagonist, in patients with NYHA Class II–IV heart failure (Poole-Wilson et al., 2002, 2003). Among other explanations for the benefit provided by carvedilol (Bristow et al., 2003), one hypothesis is that by blocking both β_2 - and α_1 -ARs, as well as β_1 -ARs, carvedilol confers additional benefit over selective β_1 -AR antagonism alone (Poole-Wilson et al., 2002).

As a mixed acting $\beta_1/\beta_2/\alpha_1$ -blocker, carvedilol is proposed to block α_1 -mediated vasoconstriction to account for the added benefit. Indeed, in HEK cells expressing the human α_1 -subtypes, carvedilol has higher

binding affinity for the α_1B and α_1D than for β -ARs or the α_1A and selectively inhibits α_1B - and α_1D -subtype-specific calcium transients (Koshimizu et al., 2004). However, the idea that benefit with carvedilol relies on its α_1 -blocking properties might seem counterintuitive, given the failure of the α_1 -blocker prazosin to improve mortality in V-HeFT (Cohn et al., 1986; Cohn, 1993). Indeed, α_1 -mediated vasopressor responses are not reduced during chronic treatment with carvedilol (Kubo et al., 2001; Hryniewicz et al., 2003). These studies show that chronic carvedilol treatment does not inhibit α_1 -AR-mediated vascular contraction and further suggest that the long-term benefits of carvedilol are not likely due to α_1 -AR antagonism.

In another smaller study of patients with heart failure, the blood pressure responses to α_1 -AR agonist infusion (phenylephrine) were actually increased in patients receiving carvedilol (Van Tassel et al., 2008). These data were interpreted to mean that chronic carvedilol treatment actually *potentiates* α_1 -mediated vasoconstriction (Van Tassel et al., 2008). Although this was a small study, an effect on the heart is also possible, and the implications are that carvedilol might provide an added benefit in heart failure through augmented α_1 -AR signaling.

E. Sympatholytics: Reducing Norepinephrine Levels Does Not Improve Heart Failure. If the central tenet of heart failure therapy for the last several decades is correct, namely that increased catecholamine signaling exacerbates heart failure, then by extension, reducing catecholamine levels should improve heart failure outcomes. Following that logic, both the Moxonidine Safety and Efficacy Trial (MOXSE) (Swedberg et al., 2002) and the Moxonidine Congestive Heart Failure Trial (MOXCON) (Cohn et al., 2003) examined the effects of the sympatholytic imidazoline receptor agonist moxonidine on mortality in heart failure. MOXSE examined 268 patients with New York Heart Association class II–IV systolic heart failure and found that moxonidine reduced heart rate and modestly improved ejection fraction but increased adverse events.

MOXCON, the larger of the two trials, also targeted patients with New York Heart Association class II–IV systolic heart failure. After enrolling roughly 1900 patients, the trial was stopped early due to increased mortality in the moxonidine group (Coats, 1999; Cohn et al., 2003; Pocock et al., 2004).

Another clinical trial showed that the β_1/β_2 -blocker/sympatholytic agent bucindolol increased mortality associated with pronounced NE reduction in the β -Blocker Evaluation of Survival Trial (BEST) (Bristow et al., 2004). In total, the failure of these trials suggests that some catecholamine signaling is beneficial in heart failure. Although the mechanism whereby sympatholysis leads to increased mortality is uncertain, it is possible that decreasing catecholamine levels abrogates the cardioprotective effects of α_1 -AR signaling.

F. Conclusions and Implications: Are Myocardial α_1 -Adrenergic Receptors Cardioprotective in Humans? In summary, one large clinical trial and several smaller studies show that α_1 -blockers or a reduction in NE levels worsens outcomes in patients with hypertension or heart failure, as summarized in Table 1. One implication of these results is to force a reconsideration of the notion that all AR signaling in heart failure is pathologic. Although β -AR blockade is clearly beneficial in heart failure, the negative results of trials involving sympatholytics indicate that reducing NE levels excessively can be harmful. Perhaps more importantly, another implication of these results would be to suggest that myocardial α_1 -ARs are protective, based on the negative results in trials with α_1 -blockers. None of the trials involving α_1 -blockers were designed to address mechanisms whereby α_1 -AR inhibition worsens heart failure. However, as discussed in section IV, α_1 -ARs are clearly cardioprotective in cell and animal models.

VI. Final Summary

The functional significance of cardiac α_1 -ARs has dramatically advanced over the last 30 years since α_1 -ARs were first demonstrated to have a direct trophic effect on cardiac myocytes (Simpson, 1983). Clinical data now show that α_1 -blockers cause heart failure in hypertensive patients and possibly in patients taking α_1 -blockers for BPH. These clinical data imply a protective function for cardiac α_1 -ARs, and data indicate that unlike β_1 -ARs, α_1 -ARs are not downregulated in human heart failure but are proportionally increased, available to mediate protective signaling in heart failure. In fact, studies in cultured myocytes and animal models show that α_1 -ARs are cardioprotective and prevent pathologic remodeling in heart failure. The mechanisms are multifactorial, including activation of adaptive or physiologic hypertrophy, prevention of cardiac myocyte death, augmentation of positive inotropic responses, and induction of ischemic preconditioning. Importantly, these studies provide a mechanistic basis to explain the failure of α_1 -blockers in humans. Perhaps most surprising is the finding that α_1 -ARs localize to and signal at the nuclear membrane in adult cardiac myocytes and engage “inside-out” signaling to regulate α_1 -AR cardioprotective signaling. Overall, these findings raise several new questions regarding the functional role of cardiac α_1 -ARs. Perhaps the most important are whether activation of cardiac α_1 -ARs, especially the α_1A subtype, might be a viable therapeutic strategy in heart failure and how nuclear α_1 -AR signaling might differ functionally from α_1 -AR signaling at the plasma membrane.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: O'Connell, Jensen, Baker, Simpson.

References

- Adams JW and Brown JH (2001) G-proteins in growth and apoptosis: lessons from the heart. *Oncogene* **20**:1626–1634.
- Adams JW, Sakata Y, Davis MG, Sah VP, Wang Y, Liggett SB, Chien KR, Brown JH, and Dorn GW 2nd (1998) Enhanced G_{α_q} signaling: a common pathway mediates cardiac hypertrophy and apoptotic heart failure. *Proc Natl Acad Sci USA* **95**:10140–10145.
- Agüero J, Almenar L, Montó F, Oliver E, Sánchez-Lázaro I, Vicente D, Martínez-Dolz L, D'Ocon P, Rueda J, and Salvador A (2012) Myocardial G protein receptor-coupled kinase expression correlates with functional parameters and clinical severity in advanced heart failure. *J Card Fail* **18**:53–61.
- Ainscough JF, Drinkhill MJ, Sedo A, Turner NA, Brooke DA, Balmforth AJ, and Ball SG (2009) Angiotensin II type-1 receptor activation in the adult heart causes blood pressure-independent hypertrophy and cardiac dysfunction. *Cardiovasc Res* **81**:592–600.
- Akhter SA, Milano CA, Shotwell KF, Cho MC, Rockman HA, Lefkowitz RJ, and Koch WJ (1997) Transgenic mice with cardiac overexpression of alpha1B-adrenergic receptors. In vivo alpha1-adrenergic receptor-mediated regulation of beta-adrenergic signaling. *J Biol Chem* **272**:21253–21259.
- ALLHAT Collaborative Research Group (2000) Major cardiovascular events in hypertensive patients randomized to doxazosin vs chlorthalidone: the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT) [published correction appears in *JAMA* (2002) **288**:2976]. *JAMA* **283**:1967–1975.
- Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial Collaborative Research Group (2003) Diuretic versus alpha-blocker as first-step antihypertensive therapy: final results from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Hypertension* **42**:239–246.
- Amirahmadi F, Turnbull L, Du XJ, Graham RM, and Woodcock EA (2008) Heightened alpha1A-adrenergic receptor activity suppresses ischaemia/reperfusion-induced Ins (1,4,5)P₃ generation in the mouse heart: a comparison with ischaemic preconditioning. *Clin Sci (Lond)* **114**:157–164.
- Anand IS and Florea VG (2003) Alterations in ventricular structure: Role of left ventricular remodeling, in *Heart Failure: A Companion to Braunwald's Heart Disease* (Mann DL, ed) pp 229–245, WB Saunders Co.
- Andersen GO, Qvigstad E, Schiander I, Aass H, Osnes JB, and Skomedal T (2002) Alpha(1)-AR-induced positive inotropic response in heart is dependent on myosin light chain phosphorylation. *Am J Physiol Heart Circ Physiol* **283**:H1471–H1480.
- Apkon M and Nerbonne JM (1988) Alpha 1-adrenergic agonists selectively suppress voltage-dependent K⁺ current in rat ventricular myocytes. *Proc Natl Acad Sci USA* **85**:8756–8760.
- Aries A, Paradis P, Lefebvre C, Schwartz RJ, and Nemer M (2004) Essential role of GATA-4 in cell survival and drug-induced cardiotoxicity. *Proc Natl Acad Sci USA* **101**:6975–6980.
- Autelitano DJ and Woodcock EA (1998) Selective activation of alpha1A-adrenergic receptors in neonatal cardiac myocytes is sufficient to cause hypertrophy and differential regulation of alpha1-adrenergic receptor subtype mRNAs. *J Mol Cell Cardiol* **30**:1515–1523.
- Avkiran M, Rowland AJ, Cuervo F, and Haworth RS (2008) Protein kinase d in the cardiovascular system: emerging roles in health and disease. *Circ Res* **102**:157–163.
- Backs J, Backs T, Bezprozvannaya S, McKinsey TA, and Olson EN (2008) Histone deacetylase 5 acquires calcium/calmodulin-dependent kinase II responsiveness by oligomerization with histone deacetylase 4. *Mol Cell Biol* **28**:3437–3445.
- Backs J, Song K, Bezprozvannaya S, Chang S, and Olson EN (2006) CaM kinase II selectively signals to histone deacetylase 4 during cardiomyocyte hypertrophy. *J Clin Invest* **116**:1853–1864.
- Baghelai K, Graham LJ, Wechsler AS, and Jakoi ER (1999a) Delayed myocardial preconditioning by alpha1-adrenoceptors involves inhibition of apoptosis. *J Thorac Cardiovasc Surg* **117**:980–986.
- Baghelai K, Graham LJ, Wechsler AS, and Jakoi ER (1999b) Phenylephrine induces delayed cardioprotection against necrosis without amelioration of stunning. *Ann Thorac Surg* **68**:1219–1224.
- Baines CP (2010) Role of the mitochondrion in programmed necrosis. *Front Physiol* **1**:156.
- Baines CP, Kaiser RA, Purcell NH, Blair NS, Osinska H, Hambleton MA, Brunskill EW, Sayen MR, Gottlieb RA, and Dorn GW, et al. (2005) Loss of cyclophilin D reveals a critical role for mitochondrial permeability transition in cell death. *Nature* **434**:658–662.
- Banerjee A, Locke-Winter C, Rogers KB, Mitchell MB, Brew EC, Cairns CB, Bensard DD, and Harken AH (1993) Preconditioning against myocardial dysfunction after ischemia and reperfusion by an alpha 1-adrenergic mechanism. *Circ Res* **73**:656–670.
- Bankwala Z, Hale SL, and Kloner RA (1994) Alpha-adrenoceptor stimulation with exogenous norepinephrine or release of endogenous catecholamines mimics ischemic preconditioning. *Circulation* **90**:1023–1028.
- Barron AJ, Finn SG, and Fuller SJ (2003) Chronic activation of extracellular-signal-regulated protein kinases by phenylephrine is required to elicit a hypertrophic response in cardiac myocytes. *Biochem J* **371**:71–79.
- Belevych AE, Nulton-Persson A, Sims C, and Harvey RD (2001) Role of tyrosine kinase activity in alpha-adrenergic inhibition of the beta-adrenergically regulated L-type Ca(2+) current in guinea-pig ventricular myocytes. *J Physiol* **537**:779–792.
- Bishopric NH, Simpson PC, and Ordahl CP (1987) Induction of the skeletal alpha-actin gene in alpha 1-adrenoceptor-mediated hypertrophy of rat cardiac myocytes. *J Clin Invest* **80**:1194–1199.
- Bkaily G, Avedanian L, and Jacques D (2009) Nuclear membrane receptors and channels as targets for drug development in cardiovascular diseases. *Can J Physiol Pharmacol* **87**:108–119.
- Bleasdale RA and Frenneaux MP (2002) Prognostic importance of right ventricular dysfunction. *Heart* **88**:323–324.
- Böhm M, Diet F, Feiler G, Kemkes B, and Erdmann E (1988) Alpha-adrenoceptors and alpha-adrenoceptor-mediated positive inotropic effects in failing human myocardium. *J Cardiovasc Pharmacol* **12**:357–364.
- Boivin B, Chevalier D, Villeneuve LR, Rousseau E, and Allen BG (2003) Functional endothelin receptors are present on nuclei in cardiac ventricular myocytes. *J Biol Chem* **278**:29153–29163.
- Boivin B, Lavoie C, Vanotis G, Baragli A, Villeneuve LR, Ethier N, Trieu P, Allen BG, and Hébert TE (2006) Functional beta-adrenergic receptor signalling on nuclear membranes in adult rat and mouse ventricular cardiomyocytes. *Cardiovasc Res* **71**:69–78.
- Boivin B, Vanotis G, Allen BG, and Hébert TE (2008) G protein-coupled receptors in and on the cell nucleus: a new signaling paradigm? *J Recept Signal Transduct Res* **28**:15–28.
- Bossuyt J, Chang CW, Helmstadter K, Kunkel MT, Newton AC, Campbell KS, Martin JL, Bossuyt S, Robia SL, and Bers DM (2011) Spatiotemporally distinct protein kinase D activation in adult cardiomyocytes in response to phenylephrine and endothelin. *J Biol Chem* **286**:33390–33400.
- Bossuyt J, Helmstadter K, Wu X, Clements-Jewery H, Haworth RS, Avkiran M, Martin JL, Pogwizd SM, and Bers DM (2008) Ca²⁺/calmodulin-dependent protein kinase II delta and protein kinase D overexpression reinforce the histone deacetylase 5 redistribution in heart failure. *Circ Res* **102**:695–702.
- Brahmadevara N, Shaw AM, and MacDonald A (2003) Evidence against beta 3-adrenoceptors or low affinity state of beta 1-adrenoceptors mediating relaxation in rat isolated aorta. *Br J Pharmacol* **138**:99–106.
- Brahmadevara N, Shaw AM, and MacDonald A (2004) Alpha1-adrenoceptor antagonist properties of CGP 12177A and other beta-adrenoceptor ligands: evidence against beta(3)- or atypical beta-adrenoceptors in rat aorta. *Br J Pharmacol* **142**:781–787.
- Braun AP, Fedida D, and Giles WR (1992) Activation of alpha 1-adrenoceptors modulates the inwardly rectifying potassium currents of mammalian atrial myocytes. *Pflügers Arch* **421**:431–439.
- Braz JC, Bueno OF, De Windt LJ, and Molkenin JD (2002) PKC alpha regulates the hypertrophic growth of cardiomyocytes through extracellular signal-regulated kinase1/2 (ERK1/2). *J Cell Biol* **156**:905–919.
- Braz JC, Gregory K, Pathak A, Zhao W, Sahin B, Klevisky R, Kimball TF, Lorenz JN, Nairn AC, and Liggett SB, et al. (2004) PKC-alpha regulates cardiac contractility and propensity toward heart failure. *Nat Med* **10**:248–254.
- Bristow MR (2000) beta-adrenergic receptor blockade in chronic heart failure. *Circulation* **101**:558–569.
- Bristow MR, Feldman AM, Adams KF Jr, and Goldstein S (2003) Selective versus nonselective beta-blockade for heart failure therapy: are there lessons to be learned from the COMET trial? *J Card Fail* **9**:444–453.
- Bristow MR, Ginsburg R, Minobe W, Cubicciotti RS, Sageman WS, Lurie K, Billingham ME, Harrison DC, and Stinson EB (1982) Decreased catecholamine sensitivity and beta-adrenergic-receptor density in failing human hearts. *N Engl J Med* **307**:205–211.
- Bristow MR, Krause-Steinrauf H, Nuzzo R, Liang CS, Lindenfeld J, Lowes BD, Hattler B, Abraham WT, Olson L, and Krueger S, et al. (2004) Effect of baseline or changes in adrenergic activity on clinical outcomes in the beta-blocker evaluation of survival trial. *Circulation* **110**:1437–1442.
- Bristow MR, Minobe W, Rasmussen R, Hershberger RE, and Hoffman BB (1988) Alpha-1 adrenergic receptors in the nonfailing and failing human heart. *J Pharmacol Exp Ther* **247**:1039–1045.
- Bueno OF, De Windt LJ, Tymitz KM, Witt SA, Kimball TR, Klevisky R, Hewett TE, Jones SP, Lefer DJ, and Peng CF, et al. (2000) The MEK1-ERK1/2 signaling pathway promotes compensated cardiac hypertrophy in transgenic mice. *EMBO J* **19**:6341–6350.
- Buu NT, Hui R, and Falardeau P (1993) Norepinephrine in neonatal rat ventricular myocytes: association with the cell nucleus and binding to nuclear alpha 1- and beta-adrenergic receptors. *J Mol Cell Cardiol* **25**:1037–1046.
- Cabanes L, Costes F, Weber S, Regnard J, Benvenuti C, Castaigne A, Guerin F, and Lockhart A (1992) Improvement in exercise performance by inhalation of methoxamine in patients with impaired left ventricular function. *N Engl J Med* **326**:1661–1665.
- Caine M, Perlberg S, and Meretyk S (1978) A placebo-controlled double-blind study of the effect of phenoxybenzamine in benign prostatic obstruction. *Br J Urol* **50**:551–554.
- Caine M, Pfau A, and Perlberg S (1976) The use of alpha-adrenergic blockers in benign prostatic obstruction. *Br J Urol* **48**:255–263.
- Carnegie GK, Soughayer J, Smith FD, Pedraja BS, Zhang F, Diviani D, Bristow MR, Kunkel MT, Newton AC, and Langeberg LK, et al. (2008) AKAP-Lbc mobilizes a cardiac hypertrophy signaling pathway. *Mol Cell* **32**:169–179.
- Cavalli A, Lattion AL, Hummler E, Nenniger M, Pedrazzini T, Aubert JF, Michel MC, Yang M, Lembo G, and Vecchione C, et al. (1997) Decreased blood pressure response in mice deficient of the alpha1b-adrenergic receptor. *Proc Natl Acad Sci USA* **94**:11589–11594.
- Chalothorn D, McCune DF, Edelmann SE, García-Cazarín ML, Tsujimoto G, and Piascik MT (2002) Differences in the cellular localization and agonist-mediated internalization properties of the alpha(1)-adrenoceptor subtypes. *Mol Pharmacol* **61**:1008–1016.
- Chalothorn D, McCune DF, Edelmann SE, Tobita K, Keller BB, Lasley RD, Perez DM, Tanoue A, Tsujimoto G, and Post GR, et al. (2003) Differential cardiovascular regulatory activities of the alpha 1B- and alpha 1D-adrenoceptor subtypes. *J Pharmacol Exp Ther* **305**:1045–1053.
- Chan T, Dash R, and Simpson PC (2008) An alpha-1A-adrenergic receptor subtype agonist prevents cardiomyopathy without increasing blood pressure. abstract *Circulation* **118**:S533.

- Chatterjee S, Biondi-Zoccai G, Abbate A, D'Ascenzo F, Castagno D, Van Tassel B, Mukherjee D, and Lichstein E (2013) Benefits of β blockers in patients with heart failure and reduced ejection fraction: network meta-analysis. *BMJ* **346**:f55.
- Chaulet H, Lin F, Guo J, Owens WA, Michalick J, Kesteven SH, Guan Z, Prall OW, Mearns BM, and Feneley MP, et al. (2006) Sustained augmentation of cardiac alpha1A-adrenergic drive results in pathological remodeling with contractile dysfunction, progressive fibrosis and reactivation of matricellular protein genes. *J Mol Cell Cardiol* **40**:540–552.
- Chen HH, Mullett SJ, and Stewart AF (2004) Vgl-4, a novel member of the vestigial-like family of transcription cofactors, regulates alpha1-adrenergic activation of gene expression in cardiac myocytes. *J Biol Chem* **279**:30800–30806.
- Chen L, el-Sherif N, and Boutjdir M (1996) Alpha 1-adrenergic activation inhibits beta-adrenergic-stimulated unitary Ca^{2+} currents in cardiac ventricular myocytes. *Circ Res* **79**:184–193.
- Chen R, Mukhin YV, Garnovskaya MN, Thielen TE, Iijima Y, Huang C, Raymond JR, Ullian ME, and Paul RV (2000) A functional angiotensin II receptor-GFP fusion protein: evidence for agonist-dependent nuclear translocation. *Am J Physiol Renal Physiol* **279**:F440–F448.
- Chrysant SG (2008) Angiotensin II receptor blockers in the treatment of the cardiovascular disease continuum. *Clin Ther* **30**:2181–2190.
- Chu C, Thai K, Park KW, Wang P, Makwana O, Lovett DH, Simpson PC, and Baker AJ (2013) Intraventricular and interventricular cellular heterogeneity of inotropic responses to $\alpha(1)$ -adrenergic stimulation. *Am J Physiol Heart Circ Physiol* **304**:H946–H953.
- CIBIS-II (1999) The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomized trial. *Lancet* **353**:9–13.
- Clark WA, Rudnick SJ, LaPres JJ, Andersen LC, and LaPointe MC (1993) Regulation of hypertrophy and atrophy in cultured adult heart cells. *Circ Res* **73**:1163–1176.
- Cleveland JC Jr, Meldrum DR, Rowland RT, Cain BS, Meng X, Gamboni-Robertson F, Banerjee A, and Harken AH (1997) Ischemic preconditioning of human myocardium: protein kinase C mediates a permissive role for alpha 1-adrenoceptors. *Am J Physiol* **273**:H902–H908.
- Cleveland JC Jr, Wollmering MM, Meldrum DR, Rowland RT, Rehiring TF, Sheridan BC, Harken AH, and Banerjee A (1996) Ischemic preconditioning in human and rat ventricle. *Am J Physiol* **271**:H1786–H1794.
- Coats AJ (1999) Heart Failure 99 — the MOXCON story. *Int J Cardiol* **71**:109–111.
- Cohn JN (1993) The Vasodilator-Heart Failure Trials (V-HeFT). Mechanistic data from the VA Cooperative Studies. Introduction. *Circulation* **87**(6, Suppl)VI1–V14.
- Cohn JN, Archibald DG, Ziesche S, Francis JA, Harston WE, Tristani FE, Dunkman WB, Jacobs W, Francis GS, and Flohr KH, et al. (1986) Effect of vasodilator therapy on mortality in chronic congestive heart failure. Results of a Veterans Administration Cooperative Study. *N Engl J Med* **314**:1547–1552.
- Cohn JN and Francis JA (1977a) Vasodilator therapy of cardiac failure (second of two parts). *N Engl J Med* **297**:254–258.
- Cohn JN and Francis JA (1977b) Vasodilator therapy of cardiac failure: (first of two parts). *N Engl J Med* **297**:27–31.
- Cohn JN, Levine TB, Olivari MT, Garberg V, Lura D, Francis GS, Simon AB, and Rector T (1984) Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N Engl J Med* **311**:819–823.
- Cohn JN, Pfeffer MA, Rouleau J, Sharpe N, Swedberg K, Straub M, Wiltse C, and Wright TJ; MOXCON Investigators (2003) Adverse mortality effect of central sympathetic inhibition with sustained-release moxonidine in patients with heart failure (MOXCON). *Eur J Heart Fail* **5**:659–667.
- Communal C, Singh K, Pimentel DR, and Colucci WS (1998) Norepinephrine stimulates apoptosis in adult rat ventricular myocytes by activation of the beta-adrenergic pathway. *Circulation* **98**:1329–1334.
- Communal C, Singh M, Menon B, Xie Z, Colucci WS, and Singh K (2003) beta1 integrins expression in adult rat ventricular myocytes and its role in the regulation of beta-adrenergic receptor-stimulated apoptosis. *J Cell Biochem* **89**:381–388.
- Cook A, Bono F, Jinek M, and Conti E (2007) Structural biology of nucleocytoplasmic transport. *Annu Rev Biochem* **76**:647–671.
- Cope JT, Mauney MC, Banks D, Binns OA, Moore CL, Rentz JJ, Shockey KS, King RC, Kron IL, and Tribble CG (1997) Intravenous phenylephrine preconditioning of cardiac grafts from non-heart-beating donors. *Ann Thorac Surg* **63**:1664–1668.
- Cotecchia S (2010) The α_1 -adrenergic receptors: diversity of signaling networks and regulation. *J Recept Signal Transduct Res* **30**:410–419.
- Curiel R, Pérez-González J, Brito N, Zerpa R, Téllez D, Cabrera J, Curiel C, and Cubeddu L (1989) Positive inotropic effects mediated by alpha 1 adrenoceptors in intact human subjects. *J Cardiovasc Pharmacol* **14**:603–615.
- Daly CJ, Milligan CM, Milligan G, Mackenzie JF, and McGrath JC (1998) Cellular localization and pharmacological characterization of functioning alpha-1 adrenoceptors by fluorescent ligand binding and image analysis reveals identical binding properties of clustered and diffuse populations of receptors. *J Pharmacol Exp Ther* **286**:984–990.
- Dash R, Chung J, Chan T, Yamada M, Barral J, Nishimura D, Yang PC, and Simpson PC (2011) A molecular MRI probe to detect treatment of cardiac apoptosis in vivo. *Magn Reson Med* **66**:1152–1162.
- Davis BR, Cutler JA, Furberg CD, Wright JT, Farber MA, Felicetta JV, and Stokes JD; ALLHAT Collaborative Research Group (2002) Relationship of antihypertensive treatment regimens and change in blood pressure to risk for heart failure in hypertensive patients randomly assigned to doxazosin or chlorthalidone: further analyses from the Antihypertensive and Lipid-Lowering treatment to prevent Heart Attack Trial. *Ann Intern Med* **137**:313–320.
- Debrus S, Rahbani L, Marttila M, Delorme B, Paradis P, and Nemer M (2005) The zinc finger-only protein Zfp260 is a novel cardiac regulator and a nuclear effector of alpha1-adrenergic signaling. *Mol Cell Biol* **25**:8669–8682.
- Dhalwal AS, Habib G, Deswal A, Verdugo M, Soucek J, Ramasubbu K, Aguilar D, Ma TS, Jneid HM, and Bolos M, et al. (2009) Impact of alpha 1-adrenergic antagonist use for benign prostatic hypertrophy on outcomes in patients with heart failure. *Am J Cardiol* **104**:270–275.
- Dono R, James D, and Zeller R (1998) A GR-motif functions in nuclear accumulation of the large FGF-2 isoforms and interferes with mitogenic signalling. *Oncogene* **16**:2151–2158.
- Dorn GW 2nd (2005) Physiologic growth and pathologic genes in cardiac development and cardiomyopathy. *Trends Cardiovasc Med* **15**:185–189.
- Dorn GW 2nd and Brown JH (1999) Gq signaling in cardiac adaptation and maladaptation. *Trends Cardiovasc Med* **9**:26–34.
- Drake MT, Shenoy SK, and Lefkowitz RJ (2006) Trafficking of G protein-coupled receptors. *Circ Res* **99**:570–582.
- Du XJ, Fang L, Gao XM, Kiriazis H, Feng X, Hotchkiss E, Finch AM, Chaulet H, and Graham RM (2004) Genetic enhancement of ventricular contractility protects against pressure-overload-induced cardiac dysfunction. *J Mol Cell Cardiol* **37**:979–987.
- Du XJ, Gao XM, Kiriazis H, Moore XL, Ming Z, Su Y, Finch AM, Hannan RA, Dart AM, and Graham RM (2006) Transgenic alpha1A-adrenergic activation limits post-infarct ventricular remodeling and dysfunction and improves survival. *Cardiovasc Res* **71**:735–743.
- Duan H and Wang J (2010) Selective transport of monoamine neurotransmitters by human plasma membrane monoamine transporter and organic cation transporter 3. *J Pharmacol Exp Ther* **335**:743–753.
- Eckhart AD, Duncan SJ, Penn RB, Benovic JL, Lefkowitz RJ, and Koch WJ (2000) Hybrid transgenic mice reveal in vivo specificity of G protein-coupled receptor kinases in the heart. *Circ Res* **86**:43–50.
- Endoh M and Blinks JR (1988) Actions of sympathomimetic amines on the Ca^{2+} transients and contractions of rabbit myocardium: reciprocal changes in myofibrillar responsiveness to Ca^{2+} mediated through alpha- and beta-adrenoceptors. *Circ Res* **62**:247–265.
- Fawzy A, Vashi V, Chung M, Dias N, and Gaffney M; Multicenter Study Group (1999) Clinical correlation of maximal urinary flow rate and plasma doxazosin concentrations in the treatment of benign prostatic hyperplasia. *Urology* **53**:329–335.
- Fedida D, Braun AP, and Giles WR (1991) Alpha 1-adrenoceptors reduce background K^+ current in rabbit ventricular myocytes. *J Physiol* **441**:673–684.
- Fedida D, Shimoni Y, and Giles WR (1989) A novel effect of norepinephrine on cardiac cells is mediated by alpha 1-adrenoceptors. *Am J Physiol* **256**:H1500–H1504.
- Fedida D, Shimoni Y, and Giles WR (1990) Alpha-adrenergic modulation of the transient outward current in rabbit atrial myocytes. *J Physiol* **423**:257–277.
- Filipeanu CM, Zhou F, Fugetta EK, and Wu G (2006) Differential regulation of the cell-surface targeting and function of beta- and alpha1-adrenergic receptors by Rab1 GTPase in cardiac myocytes. *Mol Pharmacol* **69**:1571–1578.
- Filtz TM, Grubb DR, McLeod-Dryden TJ, Luo J, and Woodcock EA (2009) Gq-initiated cardiomyocyte hypertrophy is mediated by phospholipase Cbeta1b. *FASEB J* **23**:3564–3570.
- Foo RS, Mani K, and Kitsis RN (2005) Death begets failure in the heart. *J Clin Invest* **115**:565–571.
- Foody JM, Farrell MH, and Krumholz HM (2002) beta-Blocker therapy in heart failure: scientific review. *JAMA* **287**:883–889.
- Frey N and Olson EN (2003) Cardiac hypertrophy: the good, the bad, and the ugly. *Annu Rev Physiol* **65**:45–79.
- Frishman WH and Kotob F (1999) Alpha-adrenergic blocking drugs in clinical medicine. *J Clin Pharmacol* **39**:7–16.
- Fujita T, Toya Y, Iwatsubo K, Onda T, Kimura K, Umemura S, and Ishikawa Y (2001) Accumulation of molecules involved in alpha1-adrenergic signal within caveolae: caveolin expression and the development of cardiac hypertrophy. *Cardiovasc Res* **51**:709–716.
- Fuller SJ, Gaitanaki CJ, and Sugden PH (1990) Effects of catecholamines on protein synthesis in cardiac myocytes and perfused hearts isolated from adult rats. Stimulation of translation is mediated through the alpha 1-adrenoceptor. *Biochem J* **266**:727–736.
- Gambassi G, Spurgeon HA, Lakatta EG, Blank PS, and Capogrossi MC (1992) Different effects of alpha- and beta-adrenergic stimulation on cytosolic pH and myofibrillar responsiveness to Ca^{2+} in cardiac myocytes. *Circ Res* **71**:870–882.
- Gambassi G, Spurgeon HA, Ziman BD, Lakatta EG, and Capogrossi MC (1998) Opposing effects of alpha 1-adrenergic receptor subtypes on Ca^{2+} and pH homeostasis in rat cardiac myocytes. *Am J Physiol* **274**:H1152–H1162.
- Gao XM, Wang BH, Woodcock E, and Du XJ (2000) Expression of active alpha1B-adrenergic receptors in the heart does not alleviate ischemic reperfusion injury. *J Mol Cell Cardiol* **32**:1679–1686.
- Gaughan JP, Hefner CA, and Houser SR (1998) Electrophysiological properties of neonatal rat ventricular myocytes with alpha1-adrenergic-induced hypertrophy. *Am J Physiol* **275**:H577–H590.
- Glembotski CC (2013) Classic studies of cultured cardiac myocyte hypertrophy: interview with a transformer. *Circ Res* **113**:1112–1116.
- Gobeil F Jr, Dumont I, Marrache AM, Vazquez-Tello A, Bernier SG, Abran D, Hou X, Beauchamp MH, Quiniou C, and Bouayad A, et al. (2002) Regulation of eNOS expression in brain endothelial cells by perinuclear EP(3) receptors. *Circ Res* **90**:682–689.
- Gobeil F, Fortier A, Zhu T, Bossolasco M, Leduc M, Grandbois M, Heveker N, Bkaily G, Chemtob S, and Barbaz D (2006) G-protein-coupled receptors signalling at the cell nucleus: an emerging paradigm. *Can J Physiol Pharmacol* **84**:287–297.
- González-Juanatey JR, Iglesias MJ, Alcaide C, Piñeiro R, and Lago F (2003) Doxazosin induces apoptosis in cardiomyocytes cultured in vitro by a mechanism that is independent of alpha1-adrenergic blockade. *Circulation* **107**:127–131.
- Graham RM, Perez DM, Hwa J, and Piascik MT (1996) Alpha 1-adrenergic receptor subtypes. Molecular structure, function, and signaling. *Circ Res* **78**:737–749.
- Gray MO, Long CS, Kalinyak JE, Li HT, and Karlner JS (1998) Angiotensin II stimulates cardiac myocyte hypertrophy via paracrine release of TGF-beta 1 and endothelin-1 from fibroblasts. *Cardiovasc Res* **40**:352–363.
- Grubb DR, Vasilevski O, Huynh H, and Woodcock EA (2008) The extreme C-terminal region of phospholipase Cbeta1 determines subcellular localization and function;

- the "b" splice variant mediates alpha1-adrenergic receptor responses in cardiomyocytes. *FASEB J* **22**:2768–2774.
- Grupp IL, Lorenz JN, Walsh RA, Boivin GP, and Rindt H (1998) Overexpression of alpha1B-adrenergic receptor induces left ventricular dysfunction in the absence of hypertrophy. *Am J Physiol* **275**:H1338–H1350.
- Guerra S, Leri A, Wang X, Finato N, Di Loreto C, Beltrami CA, Kajstura J, and Anversa P (1999) Myocyte death in the failing human heart is gender dependent. *Circ Res* **85**:856–866.
- Guo J, Gertsberg Z, Ozgen N, and Steinberg SF (2009) p66Shc links alpha1-adrenergic receptors to a reactive oxygen species-dependent AKT-FOXO3A phosphorylation pathway in cardiomyocytes. *Circ Res* **104**:660–669.
- Harada K, Sugaya T, Murakami K, Yazaki Y, and Komuro I (1999) Angiotensin II type 1A receptor knockout mice display less left ventricular remodeling and improved survival after myocardial infarction. see comments *Circulation* **100**:2093–2099.
- Harrison BC, Kim MS, van Rooij E, Plato CF, Papst PJ, Vega RB, McAnally JA, Richardson JA, Bassel-Duby R, and Olson EN, et al. (2006) Regulation of cardiac stress signaling by protein kinase d1. *Mol Cell Biol* **26**:3875–3888.
- Hartmann M, Stumpe T, and Schrader J (1995) alpha 1-Adrenoceptor stimulation inhibits the isoproterenol-induced effects on myocardial contractility and protein phosphorylation. *Eur J Pharmacol* **287**:57–64.
- Haworth RS, Goss MW, Rozengurt E, and Avkiran M (2000) Expression and activity of protein kinase D/protein kinase C mu in myocardium: evidence for alpha1-adrenergic receptor- and protein kinase C-mediated regulation. *J Mol Cell Cardiol* **32**:1013–1023.
- Hein P and Michel MC (2007) Signal transduction and regulation: are all alpha1-adrenergic receptor subtypes created equal? *Biochem Pharmacol* **73**:1097–1106.
- Hennenberg M, Schlenker B, Roosen A, Strittmatter F, Walther S, Stief C, and Gratzke C (2011) beta-arrestin-2 is expressed in human prostate smooth muscle and a binding partner of alpha1-adrenoceptors. *World J Urol* **29**:157–163.
- Henrich CJ and Simpson PC (1988) Differential acute and chronic response of protein kinase C in cultured neonatal rat heart myocytes to alpha 1-adrenergic and phorbol ester stimulation. *J Mol Cell Cardiol* **20**:1081–1085.
- Hirano S, Kusakari Y, O-Uchi J, Morimoto S, Kawai M, Hongo K, and Kurihara S (2006) Intracellular mechanism of the negative inotropic effect induced by alpha1-adrenoceptor stimulation in mouse myocardium. *J Physiol Sci* **56**:297–304.
- Hock R, Scheer U, and Bustin M (1998) Chromosomal proteins HMG-14 and HMG-17 are released from mitotic chromosomes and imported into the nucleus by active transport. *J Cell Biol* **143**:1427–1436.
- Hosoda C, Koshimizu TA, Tanoue A, Nasa Y, Oikawa R, Tomabechi T, Fukuda S, Shinoura H, Oshikawa S, and Takeo S, et al. (2005) Two alpha1-adrenergic receptor subtypes regulating the vasopressor response have differential roles in blood pressure regulation. *Mol Pharmacol* **67**:912–922.
- Hryniewicz K, Androne AS, Hudaihed A, and Katz SD (2003) Comparative effects of carvedilol and metoprolol on regional vascular responses to adrenergic stimuli in normal subjects and patients with chronic heart failure. *Circulation* **108**:971–976.
- Hu K and Nattel S (1995) Mechanisms of ischemic preconditioning in rat hearts. Involvement of alpha 1B-adrenoceptors, pertussis toxin-sensitive G proteins, and protein kinase C. *Circulation* **92**:2259–2265.
- Huang Y, Wright CD, Kobayashi S, Healy CL, Elgethun M, Cypher A, Liang Q, and O'Connell TD (2008) GATA4 is a survival factor in adult cardiac myocytes but is not required for (alpha)1A-adrenergic receptor survival signaling. *Am J Physiol* **295**:H699–H707.
- Huang Y, Wright CD, Merkwang CL, Baye NL, Liang Q, Simpson PC, and O'Connell TD (2007) An alpha1A-adrenergic-extracellular signal-regulated kinase survival signaling pathway in cardiac myocytes. *Circulation* **115**:763–772.
- Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini DM, and Michl K, et al.; American College of Cardiology; American Heart Association Task Force on Practice Guidelines; American College of Chest Physicians; International Society for Heart and Lung Transplantation; Heart Rhythm Society (2005) ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation* **112**:e154–e235.
- Hwang KC, Gray CD, Sweet WE, Moravec CS, and Im MJ (1996) Alpha 1-adrenergic receptor coupling with Gh in the failing human heart. *Circulation* **94**:718–726.
- Iaccarino G, Keys JR, Rapacciuolo A, Shotwell KF, Lefkowitz RJ, Rockman HA, and Koch WJ (2001) Regulation of myocardial betaARK1 expression in catecholamine-induced cardiac hypertrophy in transgenic mice overexpressing alpha1B-adrenergic receptors. *J Am Coll Cardiol* **38**:534–540.
- Ichishima K, Yamamoto S, Iwamoto T, and Ehara T (2010) alpha-Adrenoceptor-mediated depletion of phosphatidylinositol 4, 5-bisphosphate inhibits activation of volume-regulated anion channels in mouse ventricular myocytes. *Br J Pharmacol* **161**:193–206.
- Ikeda U, Tsuruya Y, and Yaginuma T (1991) Alpha 1-adrenergic stimulation is coupled to cardiac myocyte hypertrophy. *Am J Physiol* **260**:H953–H956.
- Imani A, Faghihi M, Sadr SS, Keshavaz M, and Niaraki SS (2008) Noradrenaline reduces ischemia-induced arrhythmia in anesthetized rats: involvement of alpha1-adrenoceptors and mitochondrial K ATP channels. *J Cardiovasc Electrophysiol* **19**:309–315.
- Iwai-Kanai E, Hasegawa K, Araki M, Kakita T, Morimoto T, and Sasayama S (1999) alpha- and beta-adrenergic pathways differentially regulate cell type-specific apoptosis in rat cardiac myocytes. *Circulation* **100**:305–311.
- Jensen BC, O'Connell TD, and Simpson PC (2011) Alpha-1-adrenergic receptors: targets for agonist drugs to treat heart failure. *J Mol Cell Cardiol* **51**:518–528.
- Jensen BC, Swigart PM, De Marco T, Hoopes C, and Simpson PC (2009a) alpha1-Adrenergic receptor subtypes in nonfailing and failing human myocardium. *Circ Heart Fail* **2**:654–663.
- Jensen BC, Swigart PM, Laden ME, DeMarco T, Hoopes C, and Simpson PC (2009b) The alpha-1D is the predominant alpha-1-adrenergic receptor subtype in human epicardial coronary arteries. *J Am Coll Cardiol* **54**:1137–1145.
- Jensen BC, Swigart PM, Montgomery MD, and Simpson PC (2010) Functional alpha-1B adrenergic receptors on human epicardial coronary artery endothelial cells. *Naunyn Schmiedebergs Arch Pharmacol* **382**:475–482.
- Jensen BC, Swigart PM, and Simpson PC (2009c) Ten commercial antibodies for alpha-1-adrenergic receptor subtypes are nonspecific. *Naunyn Schmiedebergs Arch Pharmacol* **379**:409–412.
- Jin Y, Sheikh F, Detillieux KA, and Cattini PA (2000) Role for early growth response-1 protein in alpha(1)-adrenergic stimulation of fibroblast growth factor-2 promoter activity in cardiac myocytes. *Mol Pharmacol* **57**:984–990.
- Kariya K, Farrance IK, and Simpson PC (1993) Transcriptional enhancer factor-1 in cardiac myocytes interacts with an alpha 1-adrenergic- and beta-protein kinase C-inducible element in the rat beta-myosin heavy chain promoter. *J Biol Chem* **268**:26658–26662.
- Kariya K, Karns LR, and Simpson PC (1991) Expression of a constitutively activated mutant of the beta-isozyme of protein kinase C in cardiac myocytes stimulates the promoter of the beta-myosin heavy chain isogene. *J Biol Chem* **266**:10023–10026.
- Kariya K, Karns LR, and Simpson PC (1994) An enhancer core element mediates stimulation of the rat beta-myosin heavy chain promoter by an alpha 1-adrenergic agonist and activated beta-protein kinase C in hypertrophy of cardiac myocytes. *J Biol Chem* **269**:3775–3782.
- Karns LR, Kariya K, and Simpson PC (1995) M-CAT, CarG, and Sp1 elements are required for alpha 1-adrenergic induction of the skeletal alpha-actin promoter during cardiac myocyte hypertrophy. Transcriptional enhancer factor-1 and protein kinase C as conserved transducers of the fetal program in cardiac growth. *J Biol Chem* **270**:410–417.
- Kim NN, Villarreal FJ, Printz MP, Lee AA, and Dillmann WH (1995) Trophic effects of angiotensin II on neonatal rat cardiac myocytes are mediated by cardiac fibroblasts. *Am J Physiol* **269**:E426–E437.
- King BD, Sack D, Kichuk MR, and Hintze TH (1987) Absence of hypertension despite chronic marked elevations in plasma norepinephrine in conscious dogs. *Hypertension* **9**:582–590.
- King MC, Lusk CP, and Blobel G (2006) Karyopherin-mediated import of integral inner nuclear membrane proteins. *Nature* **442**:1003–1007.
- Kitakaze M, Hori M, Morioka T, Minamino T, Takashima S, Sato H, Shinozaki Y, Chujo M, Mori H, and Inoue M, et al. (1994) Alpha 1-adrenoceptor activation mediates the infarct size-limiting effect of ischemic preconditioning through augmentation of 5'-nucleotidase activity. *J Clin Invest* **93**:2197–2205.
- Kitakaze M, Hori M, Sato H, Iwakura K, Gotoh K, Inoue M, Kitabatake A, and Kamada T (1991) Beneficial effects of alpha 1-adrenoceptor activity on myocardial stunning in dogs. *Circ Res* **68**:1322–1339.
- Kitakaze M, Hori M, Tamai J, Iwakura K, Koretsune Y, Kagiya T, Iwai K, Kitabatake A, Inoue M, and Kamada T (1987) Alpha 1-adrenoceptor activity regulates release of adenosine from the ischemic myocardium in dogs. *Circ Res* **60**:631–639.
- Knowlton KU, Baracchini E, Ross RS, Harris AN, Henderson SA, Evans SM, Glembocki CC, and Chien KR (1991) Co-regulation of the atrial natriuretic factor and cardiac myosin light chain-2 genes during alpha-adrenergic stimulation of neonatal rat ventricular cells. Identification of cis sequences within an embryonic and a constitutive contractile protein gene which mediate inducible expression. *J Biol Chem* **266**:7759–7768.
- Knowlton KU, Michel MC, Itani M, Shubeita HE, Ishihara K, Brown JH, and Chien KR (1993) The alpha 1A-adrenergic receptor subtype mediates biochemical, molecular, and morphologic features of cultured myocardial cell hypertrophy. *J Biol Chem* **268**:15374–15380.
- Koshimizu TA, Tsujimoto G, Hirasawa A, Kitagawa Y, and Tanoue A (2004) Carvedilol selectively inhibits oscillatory intracellular calcium changes evoked by human alpha1D- and alpha1B-adrenergic receptors. *Cardiovasc Res* **63**:662–672.
- Kostin S, Pool L, Elsäßer A, Hein S, Drexler HC, Arnon E, Hayakawa Y, Zimmermann R, Bauer E, and Klövekorn WP, et al. (2003) Myocytes die by multiple mechanisms in failing human hearts. *Circ Res* **92**:715–724.
- Kubo T, Azevedo ER, Newton GE, Parker JD, and Floras JS (2001) Lack of evidence for peripheral alpha(1)-adrenoceptor blockade during long-term treatment of heart failure with carvedilol. *J Am Coll Cardiol* **38**:1463–1469.
- Laks MM, Morady F, and Swan HJ (1973) Myocardial hypertrophy produced by chronic infusion of subhypertensive doses of norepinephrine in the dog. *Chest* **64**:75–78.
- Landzberg JS, Parker JD, Gauthier DF, and Colucci WS (1991) Effects of myocardial alpha 1-adrenergic receptor stimulation and blockade on contractility in humans. *Circulation* **84**:1608–1614.
- Lanzafame AA, Turnbull L, Amiramahdi F, Arthur JF, Huynh H, and Woodcock EA (2006) Inositol phospholipids localized to caveolae in rat heart are regulated by alpha1-adrenergic receptors and by ischemia-reperfusion. *Am J Physiol Heart Circ Physiol* **290**:H2059–H2065.
- Lee DK, Lança AJ, Cheng R, Nguyen T, Ji XD, Gobeil F Jr, Chemtob S, George SR, and O'Dowd BF (2004) Agonist-independent nuclear localization of the Apelin, angiotensin AT1, and bradykinin B2 receptors. *J Biol Chem* **279**:7901–7908.
- Lemire I, Ducharme A, Tardif JC, Poulin F, Jones LR, Allen BG, Hébert TE, and Rindt H (2001) Cardiac-directed overexpression of wild-type alpha1B-adrenergic receptor induces dilated cardiomyopathy. *Am J Physiol Heart Circ Physiol* **281**:H931–H938.
- Levin MC, Marullo S, Muntaner O, Andersson B, and Magnusson Y (2002) The myocardium-protective Gly-49 variant of the beta 1-adrenergic receptor exhibits constitutive activity and increased desensitization and down-regulation. *J Biol Chem* **277**:30429–30435.

- Levy D, Garrison RJ, Savage DD, Kannel WB, and Castelli WP (1990) Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N Engl J Med* **322**:1561–1566.
- Li Y, Zhang H, Liao W, Song Y, Ma X, Chen C, Lu Z, Li Z, and Zhang Y (2011) Transactivated EGFR mediates α_1 -AR-induced STAT3 activation and cardiac hypertrophy. *Am J Physiol Heart Circ Physiol* **301**:H1941–H1951.
- Liang Q, De Windt LJ, Witt SA, Kimball TR, Markham BE, and Molkenin JD (2001a) The transcription factors GATA4 and GATA6 regulate cardiomyocyte hypertrophy in vitro and in vivo. *J Biol Chem* **276**:30245–30253.
- Liang Q, Wiese RJ, Bueno OF, Dai YS, Markham BE, and Molkenin JD (2001b) The transcription factor GATA4 is activated by extracellular signal-regulated kinase 1- and 2-mediated phosphorylation of serine 105 in cardiomyocytes. *Mol Cell Biol* **21**:7460–7469.
- Lin F, Owens WA, Chen S, Stevens ME, Kesteven S, Arthur JF, Woodcock EA, Feneley MP, and Graham RM (2001) Targeted α_1 (A)-adrenergic receptor overexpression induces enhanced cardiac contractility but not hypertrophy. *Circ Res* **89**:343–350.
- Lips DJ, Bueno OF, Wilkins BJ, Purcell NH, Kaiser RA, Lorenz JN, Voisin L, Sabal-El-Leil MK, Meloche S, and Pouyssegur J, et al. (2004) MEK1-ERK2 signaling pathway protects myocardium from ischemic injury in vivo. *Circulation* **109**:1938–1941.
- Litwin SE, Vatner DE, and Morgan JP (1995) Inotropic effects of α_1 -adrenergic agonists in myocardium from rats with postinfarction heart failure. *Am J Physiol* **269**:H1553–H1563.
- Liu Y, Contreras M, Shen T, Randall WR, and Schneider MF (2009) α_1 -Adrenergic signaling activates protein kinase D and causes nuclear efflux of the transcriptional repressor HDAC5 in cultured adult mouse soleus skeletal muscle fibres. *J Physiol* **587**:1101–1115.
- Lohse MJ, Engelhardt S, and Eschenhagen T (2003) What is the role of β_1 -adrenergic signaling in heart failure? *Circ Res* **93**:896–906.
- Long CS, Ordahl CP, and Simpson PC (1989) α_1 -Adrenergic receptor stimulation of sarcomeric actin isogene transcription in hypertrophy of cultured rat heart muscle cells. *J Clin Invest* **83**:1078–1082.
- Lopez I, Mak EC, Ding J, Hamm HE, and Lomasney JW (2001) A novel bifunctional phospholipase c that is regulated by Galph α 12 and stimulates the Ras/mitogen-activated protein kinase pathway. *J Biol Chem* **276**:2758–2765.
- López JE, Myagmar BE, Swigart PM, Montgomery MD, Haynam S, Bigos M, Rodrigo MC, and Simpson PC (2011) β -myosin heavy chain is induced by pressure overload in a minor subpopulation of smaller mouse cardiac myocytes. *Circ Res* **109**:629–638.
- Loubani M and Galinanes M (2001) α_1 -Adrenoceptors during simulated ischemia and reoxygenation of the human myocardium: effect of the dose and time of administration. *J Thorac Cardiovasc Surg* **122**:103–112.
- Loubani M and Galinanes M (2002) Pharmacological and ischemic preconditioning of the human myocardium: mitoK(ATP) channels are upstream and p38MAPK is downstream of PKC. *BMC Physiol* **2**:10.
- Lu D, Yang H, Shaw G, and Raizada MK (1998) Angiotensin II-induced nuclear targeting of the angiotensin type 1 (AT1) receptor in brain neurons. *Endocrinology* **139**:365–375.
- Lund-Johansen P and Omvik P (1991) Acute and chronic hemodynamic effects of drugs with different actions on adrenergic receptors: a comparison between α_1 blockers and different types of β blockers with and without vasodilating effect. *Cardiovasc Drugs Ther* **5**:605–615.
- Luo DL, Gao J, Fan LL, Tang Y, Zhang YY, and Han QD (2007) Receptor subtype involved in α_1 -adrenergic receptor-mediated Ca^{2+} signaling in cardiomyocytes. *Acta Pharmacol Sin* **28**:968–974.
- Lusk CP, Blobel G, and King MC (2007) Highway to the inner nuclear membrane: rules for the road. *Nat Rev Mol Cell Biol* **8**:414–420.
- MacGowan GA, Rager J, Shroff SG, and Mathier MA (2005) In vivo α_1 -adrenergic responses and troponin I phosphorylation: anesthesia interactions. *J Appl Physiol* **98**:1163–1170.
- Mackenzie JF, Daly CJ, Pediani JD, and McGrath JC (2000) Quantitative imaging in live human cells reveals intracellular α_1 (1)-adrenoceptor ligand-binding sites. *J Pharmacol Exp Ther* **294**:434–443.
- Mann DL, Kent RL, Parsons B, and Cooper G 4th (1992) Adrenergic effects on the biology of the adult mammalian cardiocyte. *Circulation* **85**:790–804.
- Marino TA, Cassidy M, Marino DR, Carson NL, and Houser S (1991) Norepinephrine-induced cardiac hypertrophy of the cat heart. *Anat Rec* **229**:505–510.
- Markou T, Hadzopoulou-Cladaras M, and Lazou A (2004) Phenylephrine induces activation of CREB in adult rat cardiac myocytes through MSK1 and PKA signaling pathways. *J Mol Cell Cardiol* **37**:1001–1011.
- Marrache AM, Gobeil F Jr, Bernier SG, Stankova J, Rola-Pleszczynski M, Choufani S, Bkaily G, Bourdeau A, Sirois MS, and Vazquez-Tello A, et al. (2002) Proinflammatory gene induction by platelet-activating factor mediated via its cognate nuclear receptor. *J Immunol* **169**:6474–6481.
- McCloskey DT, Turnbull L, Swigart P, O'Connell TD, Simpson PC, and Baker AJ (2003) Abnormal myocardial contraction in α_1 (A)- and α_1 (B)-adrenoceptor double-knockout mice. *J Mol Cell Cardiol* **35**:1207–1216.
- McLean BG, Lee KS, Simpson PC, and Farrance IK (2003) Basal and α_1 -adrenergic-induced activity of minimal rat β MHC promoters in cardiac myocytes requires multiple TEF-1 but not NFAT binding sites. *J Mol Cell Cardiol* **35**:461–471.
- McMurray JJ, Teerlink JR, Cotter G, Bourge RC, Cleland JG, Jondeau G, Krum H, Metra M, O'Connor CM, and Parker JD, et al.; VERITAS Investigators (2007) Effects of tezosentan on symptoms and clinical outcomes in patients with acute heart failure: the VERITAS randomized controlled trials. *JAMA* **298**:2009–2019.
- McWhinney CD, Hansen C, and Robishaw JD (2000) α_1 -Adrenergic signaling in a cardiac murine atrial myocyte (HL-1) cell line. *Mol Cell Biochem* **214**:111–119.
- Meldrum DR, Cleveland JC Jr, Rowland RT, Banerjee A, Harken AH, and Meng X (1997) Early and delayed preconditioning: differential mechanisms and additive protection. *Am J Physiol* **273**:H725–H733.
- Melien O, Sandnes D, Johansen EJ, and Christoffersen T (2000) Effects of pertussis toxin on extracellular signal-regulated kinase activation in hepatocytes by hormones and receptor-independent agents: evidence suggesting a stimulatory role of G(i) proteins at a level distal to receptor coupling. *J Cell Physiol* **184**:27–36.
- Meng X, Brown JM, Ao L, Banerjee A, and Harken AH (1996a) Norepinephrine induces cardiac heat shock protein 70 and delayed cardioprotection in the rat through α_1 1 adrenoceptors. *Cardiovasc Res* **32**:374–383.
- Meng X, Cleveland JC Jr, Rowland RT, Mitchell MB, Brown JM, Banerjee A, and Harken AH (1996b) Norepinephrine-induced sustained myocardial adaptation to ischemia is dependent on α_1 1-adrenoceptors and protein synthesis. *J Mol Cell Cardiol* **28**:2017–2025.
- Meng X, Shames BD, Pulido EJ, Meldrum DR, Ao L, Joo KS, Harken AH, and Banerjee A (1999) Adrenergic induction of bimodal myocardial protection: signal transduction and cardiac gene reprogramming. *Am J Physiol* **276**:R1525–R1533.
- MERIT (1999) Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet* **353**:2001–2007.
- Messeri FH (2000) Implications of discontinuation of doxazosin arm of ALLHAT. Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. *Lancet* **355**:863–864.
- Messeri FH (2001) Doxazosin and congestive heart failure. *J Am Coll Cardiol* **38**:1295–1296.
- Methven L, Simpson PC, and McGrath JC (2009) α_1 A/B-knockout mice explain the native α_1 D-adrenoceptor's role in vasoconstriction and show that its location is independent of the other α_1 -subtypes. *Br J Pharmacol* **158**:1663–1675.
- Michel MC (2010) The forefront for novel therapeutic agents based on the pathophysiology of lower urinary tract dysfunction: α_1 -blockers in the treatment of male voiding dysfunction - how do they work and why do they differ in tolerability? *J Pharmacol Sci* **112**:151–157.
- Michel MC, Hanft G, and Gross G (1994) Radioligand binding studies of α_1 -adrenoceptor subtypes in rat heart. *Br J Pharmacol* **111**:533–538.
- Michel MC, Wieland T, and Tsujimoto G (2009) How reliable are G-protein-coupled receptor antibodies? *Naunyn-Schmiedeberg's Arch Pharmacol* **379**:385–388.
- Michelotti GA, Price DT, and Schwinn DA (2000) α_1 -Adrenergic receptor regulation: basic science and clinical implications. *Pharmacol Ther* **88**:281–309.
- Milano CA, Dolber PC, Rockman HA, Bond RA, Venable ME, Allen LF, and Lefkowitz RJ (1994) Myocardial expression of a constitutively active α_1 B-adrenergic receptor in transgenic mice induces cardiac hypertrophy. *Proc Natl Acad Sci USA* **91**:10109–10113.
- Mitchell MB, Meng X, Ao L, Brown JM, Harken AH, and Banerjee A (1995) Preconditioning of isolated rat heart is mediated by protein kinase C. *Circ Res* **76**:73–81.
- Modesti PA, Vanni S, Paniccia R, Bandinelli B, Bertolozzi I, Polidori G, Sani G, and Neri Serneri GG (1999) Characterization of endothelin-1 receptor subtypes in isolated human cardiomyocytes. *J Cardiovasc Pharmacol* **34**:333–339.
- Mohl MC, Iismaa SE, Xiao XH, Friedrich O, Wagner S, Nikolova-Krstevski V, Wu J, Yu ZY, Feneley M, and Fatkin D, et al. (2011) Regulation of murine cardiac contractility by activation of α_1 (A)-adrenergic receptor-operated Ca^{2+} entry. *Cardiovasc Res* **91**:310–319.
- Morimoto T, Hasegawa K, Kaburagi S, Kakita T, Wada H, Yanazume T, and Sasayama S (2000) Phosphorylation of GATA-4 is involved in α_1 -adrenergic agonist-responsive transcription of the endothelin-1 gene in cardiac myocytes. *J Biol Chem* **275**:13721–13726.
- Morris JB, Huynh H, Vasilevski O, and Woodcock EA (2006) α_1 -Adrenergic receptor signaling is localized to caveolae in neonatal rat cardiomyocytes. *J Mol Cell Cardiol* **41**:17–25.
- Morris JB, Pham TM, Kenney B, Sheppard KE, and Woodcock EA (2004) UTP transactivates epidermal growth factor receptors and promotes cardiomyocyte hypertrophy despite inhibiting transcription of the hypertrophic marker gene, atrial natriuretic peptide. *J Biol Chem* **279**:8740–8746.
- Mylona P and Cleland JG (1999) Update of REACH-1 and MERIT-HF clinical trials in heart failure. Cardio.net Editorial Team. *Eur J Heart Fail* **1**:197–200.
- Naga Prasad SV, Nienaber J, and Rockman HA (2001) β -adrenergic axis and heart disease. *Trends Genet* **17**:S44–S49.
- Nakayama H, Chen X, Baines CP, Kleivitsky R, Zhang X, Zhang H, Jaleel N, Chua BH, Hewett TE, and Robbins J, et al. (2007) Ca^{2+} - and mitochondrial-dependent cardiomyocyte necrosis as a primary mediator of heart failure. *J Clin Invest* **117**:2431–2444.
- Nemchenko A, Chiong M, Turer A, Lavandero S, and Hill JA (2011) Autophagy as a therapeutic target in cardiovascular disease. *J Mol Cell Cardiol* **51**:584–593.
- Nishida K and Otsu K (2008) Cell death in heart failure. *Circ J* **72** Suppl A:A17–21.
- Node K, Kitakaze M, Sato H, Minamino T, Komamura K, Shinozaki Y, Mori H, and Hori M (1997) Role of intracellular Ca^{2+} in activation of protein kinase C during ischemic preconditioning. *Circulation* **96**:1257–1265.
- Noguchi H, Muraoka R, Kigoshi S, and Muramatsu I (1995) Pharmacological characterization of α_1 1-adrenoceptor subtypes in rat heart: a binding study. *Br J Pharmacol* **114**:1026–1030.
- O'Connell TD, Ishizaka S, Nakamura A, Swigart PM, Rodrigo MC, Simpson GL, Cotecchia S, Rokosh DG, Grossman W, and Foster E, et al. (2003) The α_1 (A/C)- and α_1 (B)-adrenergic receptors are required for physiological cardiac hypertrophy in the double-knockout mouse. *J Clin Invest* **111**:1783–1791.
- O'Connell TD, Rokosh DG, and Simpson PC (2001) Cloning and characterization of the mouse α_1 C/A-adrenergic receptor gene and analysis of an α_1 C promoter in cardiac myocytes: role of an MCAT element that binds transcriptional enhancer factor-1 (TEF-1). *Mol Pharmacol* **59**:1225–1234.
- O'Connell TD, Swigart PM, Rodrigo MC, Ishizaka S, Joho S, Turnbull L, Tecott LH, Baker AJ, Foster E, and Grossman W, et al. (2006) α_1 -Adrenergic receptors prevent a maladaptive cardiac response to pressure overload. *J Clin Invest* **116**:1005–1015.

- O'Malley KL, Jong YJ, Gonchar Y, Burkhalter A, and Romano C (2003) Activation of metabotropic glutamate receptor mGlu5 on nuclear membranes mediates intracellular Ca^{2+} changes in heterologous cell types and neurons. *J Biol Chem* **278**: 28210–28219.
- Obst OO, Rose H, and Kammermeier H (1996) Characterization of catecholamine uptake2 in isolated cardiac myocytes. *Mol Cell Biochem* **163-164**:181–183.
- O-Uchi J, Komukai K, Kusakari Y, Obata T, Hongo K, Sasaki H, and Kurihara S (2005) Alpha1-Adrenoceptor stimulation potentiates L-type Ca^{2+} current through Ca^{2+} /calmodulin-dependent PK II (CaMKII) activation in rat ventricular myocytes. *Proc Natl Acad Sci U S A* **102**:9400–9405.
- O-Uchi J, Sasaki H, Morimoto S, Kusakari Y, Shinji H, Obata T, Hongo K, Komukai K, and Kurihara S (2008) Interaction of alpha1-adrenoceptor subtypes with different G proteins induces opposite effects on cardiac L-type Ca^{2+} channel. *Circ Res* **102**:1378–1388.
- Packer M, Bristow MR, Cohn JN, Colucci WS, Fowler MB, Gilbert EM, and Shusterman NH; U.S. Carvedilol Heart Failure Study Group (1996) The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. *N Engl J Med* **334**:1349–1355.
- Packer M, Coats AJ, Fowler MB, Katus HA, Krum H, Mohacsi P, Rouleau JL, Tendera M, Castaigne A, and Roecker EB, et al.; Carvedilol Prospective Randomized Cumulative Survival Study Group (2001) Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med* **344**:1651–1658.
- Packer M, Fowler MB, Roecker EB, Coats AJ, Katus HA, Krum H, Mohacsi P, Rouleau JL, Tendera M, and Staiger C, et al.; Carvedilol Prospective Randomized Cumulative Survival (COPERNICUS) Study Group (2002) Effect of carvedilol on the morbidity of patients with severe chronic heart failure: results of the carvedilol prospective randomized cumulative survival (COPERNICUS) study. *Circulation* **106**:2194–2199.
- Papay RS, Shi T, Piascik MT, Naga Prasad SV, and Perez DM (2013) α_1A -adrenergic receptors regulate cardiac hypertrophy in vivo through interleukin-6 secretion. *Mol Pharmacol* **83**:939–948.
- Paradis P, Dali-Youcef N, Paradis FW, Thibault G, and Nemer M (2000) Overexpression of angiotensin II type I receptor in cardiomyocytes induces cardiac hypertrophy and remodeling. *Proc Natl Acad Sci USA* **97**:931–936.
- Patel MB, Stewart JM, Loud AV, Anversa P, Wang J, Fiegel L, and Hintze TH (1991) Altered function and structure of the heart in dogs with chronic elevation in plasma norepinephrine. *Circulation* **84**:2091–2100.
- Pediani JD, Colston JF, Caldwell D, Milligan G, Daly CJ, and McGrath JC (2005) Beta-arrestin-dependent spontaneous alpha1A-adrenoceptor endocytosis causes intracellular transportation of alpha-blockers via recycling compartments. *Mol Pharmacol* **67**:992–1004.
- Piascik MT and Perez DM (2001) Alpha1-adrenergic receptors: new insights and directions. *J Pharmacol Exp Ther* **298**:403–410.
- Pickard BW, Hodsman AB, Fraher LJ, and Watson PH (2006) Type 1 parathyroid hormone receptor (PTH1R) nuclear trafficking: association of PTH1R with importin alpha1 and beta. *Endocrinology* **147**:3326–3332.
- Pickard BW, Hodsman AB, Fraher LJ, and Watson PH (2007) Type 1 parathyroid hormone receptor (PTH1R) nuclear trafficking: regulation of PTH1R nuclear-cytoplasmic shuttling by importin-alpha/beta and chromosomal region maintenance 1. *Endocrinology* **148**:2282–2289.
- Piller LB, Davis BR, Cutler JA, Cushman WC, Wright JT Jr, Williamson JD, Leenen FH, Einhorn PT, Randall OS, and Golden JS, et al. The ALLHAT Collaborative Research Group (2002) Validation of Heart Failure Events in the Antihypertensive and Lipid Lowering Treatment to Prevent Heart Attack Trial (ALLHAT) Participants Assigned to Doxazosin and Chlorthalidone. *Curr Control Trials Cardiovasc Med* **3**:10.
- Pocock S, Wilhelmsen L, Dickstein K, Francis G, and Wittes J (2004) The data monitoring experience in the MOXCON trial. *Eur Heart J* **25**:1974–1978.
- Pönicke K, Vogelsang M, Heinroth M, Becker K, Zolk O, Böhm M, Zerkowski HR, and Brodde OE (1998) Endothelin receptors in the failing and nonfailing human heart. *Circulation* **97**:744–751.
- Poole-Wilson PA, Cleland JG, Di Lenarda A, Hanrath P, Komajda M, Metra M, J Remme W, Swedberg K, and Torp-Pedersen C (2002) Rationale and design of the carvedilol or metoprolol European trial in patients with chronic heart failure: COMET. *Eur J Heart Fail* **4**:321–329.
- Poole-Wilson PA, Swedberg K, Cleland JG, Di Lenarda A, Hanrath P, Komajda M, Lubsen J, Lutiger B, Metra M, and Remme WJ, et al.; Carvedilol Or Metoprolol European Trial Investigators (2003) Comparison of carvedilol and metoprolol on clinical outcomes in patients with chronic heart failure in the Carvedilol Or Metoprolol European Trial (COMET): randomised controlled trial. *Lancet* **362**:7–13.
- Post GR, Swiderski C, Waldrop BA, Salty L, Glembotski CC, Wolthuis RM, and Mochizuki N (2002) Guanine nucleotide exchange factor-like factor (Rlf) induces gene expression and potentiates α 1-adrenergic receptor-induced transcriptional responses in neonatal rat ventricular myocytes. *J Biol Chem* **277**: 15286–15292.
- Pu WT, Ma Q, and Izumo S (2003) NFAT transcription factors are critical survival factors that inhibit cardiomyocyte apoptosis during phenylephrine stimulation in vitro. *Circ Res* **92**:725–731.
- Ravens U, Wang XL, and Wettwer E (1989) Alpha adrenoceptor stimulation reduces outward currents in rat ventricular myocytes. *J Pharmacol Exp Ther* **250**:364–370.
- Re M, Pampillo M, Savard M, Dubuc C, McArdle CA, Millar RP, Conn PM, Gobeil F Jr, Bhattacharya M, and Babwah AV (2010) The human gonadotropin releasing hormone type I receptor is a functional intracellular GPCR expressed on the nuclear membrane. *PLoS ONE* **5**:e11489.
- Rockman HA, Choi DJ, Rahman NU, Akhter SA, Lefkowitz RJ, and Koch WJ (1996) Receptor-specific in vivo desensitization by the G protein-coupled receptor kinase-5 in transgenic mice. *Proc Natl Acad Sci USA* **93**:9954–9959.
- Roger VL, Go AS, Lloyd-Jones DM, Adams RJ, Berry JD, Brown TM, Carnethon MR, Dai S, de Simone G, and Ford ES, et al.; American Heart Association Statistics Committee and Stroke Statistics Subcommittee (2011) Heart disease and stroke statistics—2011 update: a report from the American Heart Association. *Circulation* **123**:e18–e209.
- Rohde S, Sabri A, Kamasamudran R, and Steinberg SF (2000) The alpha(1)-adrenoceptor subtype- and protein kinase C isoform-dependence of Norepinephrine's actions in cardiomyocytes. *J Mol Cell Cardiol* **32**:1193–1209.
- Rokosh DG, Bailey BA, Stewart AF, Karns LR, Long CS, and Simpson PC (1994) Distribution of alpha 1C-adrenergic receptor mRNA in adult rat tissues by RNase protection assay and comparison with alpha 1B and alpha 1D. *Biochem Biophys Res Commun* **200**:1177–1184.
- Rokosh DG and Simpson PC (2002) Knockout of the alpha 1A/C-adrenergic receptor subtype: the alpha 1A/C is expressed in resistance arteries and is required to maintain arterial blood pressure. *Proc Natl Acad Sci USA* **99**:9474–9479.
- Rokosh DG, Stewart AF, Chang KC, Bailey BA, Karliner JS, Camacho SA, Long CS, and Simpson PC (1996) Alpha1-adrenergic receptor subtype mRNAs are differentially regulated by alpha1-adrenergic and other hypertrophic stimuli in cardiac myocytes in culture and in vivo. Repression of alpha1B and alpha1D but induction of alpha1C. *J Biol Chem* **271**:5839–5843.
- Rorabaugh BR, Ross SA, Gaivin RJ, Papay RS, McCune DF, Simpson PC, and Perez DM (2005) alpha1A- but not alpha1B-adrenergic receptors precondition the ischemic heart by a staurosporine-sensitive, chelerythrine-insensitive mechanism. *Cardiovasc Res* **65**:436–445.
- Ross SA, Rorabaugh BR, Chalothorn D, Yun J, Gonzalez-Cabrera PJ, McCune DF, Piascik MT, and Perez DM (2003) The alpha(1B)-adrenergic receptor decreases the inotropic response in the mouse Langendorff heart model. *Cardiovasc Res* **60**: 598–607.
- Sakata Y, Hoit BD, Liggett SB, Walsh RA, and Dorn GW 2nd (1998) Decompensation of pressure-overload hypertrophy in G alpha q-overexpressing mice. *Circulation* **97**: 1488–1495.
- Salazar NC, Chen J, and Rockman HA (2007) Cardiac GPCRs: GPCR signaling in healthy and failing hearts. *Biochim Biophys Acta* **1768**:1006–1018.
- Sato R and Koumi S (1995) Modulation of the inwardly rectifying K^{+} channel in isolated human atrial myocytes by alpha 1-adrenergic stimulation. *J Membr Biol* **148**:185–191.
- Schomig E, Lazar A, and Grundemann D (2006) Extraneuronal monoamine transporter and organic cation transporters 1 and 2: A review of transport efficiency. *Handb Exp Pharmacol* **175**:151–180.
- Schwinn DA and Roehrborn CG (2008) Alpha1-adrenoceptor subtypes and lower urinary tract symptoms. *Int J Urol* **15**:193–199.
- Sica DA (2005) Alpha1-adrenergic blockers: current usage considerations. *J Clin Hypertens (Greenwich)* **7**:757–762.
- Simpson P (1983) Norepinephrine-stimulated hypertrophy of cultured rat myocardial cells is an alpha 1 adrenergic response. *J Clin Invest* **72**:732–738.
- Simpson P (1985) Stimulation of hypertrophy of cultured neonatal rat heart cells through an alpha 1-adrenergic receptor and induction of beating through an alpha 1- and beta 1-adrenergic receptor interaction. Evidence for independent regulation of growth and beating. *Circ Res* **56**:884–894.
- Simpson P, McGrath A, and Savion S (1982) Myocyte hypertrophy in neonatal rat heart cultures and its regulation by serum and by catecholamines. *Circ Res* **51**: 787–801.
- Simpson PC (1988) Comments on "Load regulation of the properties of adult feline cardiocytes: the role of substrate adhesion." *Circ Res* **62**: 864–869.
- Simpson PC (2006) Lessons from Knockouts: the alpha-1-ARs, in *The Adrenergic Receptors in the 21st Century* (Perez DM, ed) pp 207–240, Humana Press, Totowa, NJ.
- Simpson PC (2011) Where are the new drugs to treat heart failure? Introduction to the special issue on "key signaling molecules in hypertrophy and heart failure". *J Mol Cell Cardiol* **51**:435–437.
- Simpson PC, Long CS, Waspe LE, Henrich CJ, and Ordahl CP (1989) Transcription of early developmental isogenes in cardiac myocyte hypertrophy. *J Mol Cell Cardiol* **21** (Suppl 5):79–89.
- Sjaastad I, Schiander I, Sjetnan A, Qvigstad E, Bøkenes J, Sandnes D, Osnes JB, Sejersted OM, and Skomedal T (2003) Increased contribution of alpha 1- vs. beta-adrenoceptor-mediated inotropic response in rats with congestive heart failure. *Acta Physiol Scand* **177**:449–458.
- Skomedal T, Borthne K, Aass H, Geiran O, and Osnes JB (1997) Comparison between alpha-1 adrenoceptor-mediated and beta adrenoceptor-mediated inotropic components elicited by norepinephrine in failing human ventricular muscle. *J Pharmacol Exp Ther* **280**:721–729.
- Snabaitis AK, Muntendorf A, Wieland T, and Avkiran M (2005) Regulation of the extracellular signal-regulated kinase pathway in adult myocardium: differential roles of G(q/11), Gi and G(12/13) proteins in signalling by alpha1-adrenergic, endothelin-1 and thrombin-sensitive protease-activated receptors. *Cell Signal* **17**: 655–664.
- SoRelle R (2000) National Heart, Lung, and Blood Institute halts part of antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *Circulation* **101**:E9025.
- Staehelein M, Simons P, Jaeggi K, and Wigger N (1983) CGP-12177. A hydrophilic beta-adrenergic receptor radioligand reveals high affinity binding of agonists to intact cells. *J Biol Chem* **258**:3496–3502.
- Stafford RS, Furberg CD, Finkelstein SN, Cockburn IM, Alehegn T, and Ma J (2004) Impact of clinical trial results on national trends in alpha-blocker prescribing, 1996-2002. *JAMA* **291**:54–62.
- Stanasila L, Abuin L, Dey J, and Cotecchia S (2008) Different internalization properties of the alpha1a- and alpha1b-adrenergic receptor subtypes: the potential role of receptor interaction with beta-arrestins and AP50. *Mol Pharmacol* **74**: 562–573.
- Starkens NF, Simpson PC, Bishopric N, Coughlin SR, Lee WM, Escobedo JA, and Williams LT (1986) Cardiac myocyte hypertrophy is associated with c-myc protooncogene expression. *Proc Natl Acad Sci USA* **83**:8348–8350.

- Steinberg SF, Drugge ED, Bilezikian JP, and Robinson RB (1985) Acquisition by innervated cardiac myocytes of a pertussis toxin-specific regulatory protein linked to the alpha 1-receptor. *Science* **230**:186–188.
- Steinfath M, Chen YY, Lavický J, Magnussen O, Nose M, Rosswag S, Schmitz W, and Scholz H (1992a) Cardiac alpha 1-adrenoceptor densities in different mammalian species. *Br J Pharmacol* **107**:185–188.
- Steinfath M, Danielsen W, von der Leyen H, Mende U, Meyer W, Neumann J, Nose M, Reich T, Schmitz W, and Scholz H, et al. (1992b) Reduced alpha 1- and beta 2-adrenoceptor-mediated positive inotropic effects in human end-stage heart failure. *Br J Pharmacol* **105**:463–469.
- Stewart AF, Rokosh DG, Bailey BA, Karns LR, Chang KC, Long CS, Kariya K, and Simpson PC (1994) Cloning of the rat alpha 1C-adrenergic receptor from cardiac myocytes. alpha 1C, alpha 1B, and alpha 1D mRNAs are present in cardiac myocytes but not in cardiac fibroblasts. *Circ Res* **75**:796–802.
- Stewart JM, Patel MB, Wang J, Ochoa M, Gewitz M, Loud AV, Anversa P, and Hintze TH (1992) Chronic elevation of norepinephrine in conscious dogs produces hypertrophy with no loss of LV reserve. *Am J Physiol* **262**:H331–H339.
- Swedberg K, Bristow MR, Cohn JN, Dargie H, Straub M, Wiltse C, and Wright TJ; Moxonidine Safety and Efficacy (MOXSE) Investigators (2002) Effects of sustained-release moxonidine, an imidazoline agonist, on plasma norepinephrine in patients with chronic heart failure. *Circulation* **105**:1797–1803.
- Tadevosyan A, Maguy A, Villeneuve LR, Babin J, Bonnefoy A, Allen BG, and Nattel S (2010) Nuclear-delimited angiotensin receptor-mediated signaling regulates cardiomyocyte gene expression. *J Biol Chem* **285**:22338–22349.
- Tadevosyan A, Vaniotis G, Allen BG, Hébert TE, and Nattel S (2012) G protein-coupled receptor signalling in the cardiac nuclear membrane: evidence and possible roles in physiological and pathophysiological function. *J Physiol* **590**:1313–1330.
- Tanoue A, Nasa Y, Koshimizu T, Shinoura H, Oshikawa S, Kawai T, Sunada S, Takeo S, and Tsujimoto G (2002) The alpha(1D)-adrenergic receptor directly regulates arterial blood pressure via vasoconstriction. *J Clin Invest* **109**:765–775.
- Teerlink JR and Massie BM (1999) Beta-adrenergic blocker mortality trials in congestive heart failure. *Am J Cardiol* **84** (9A):94R–102R.
- Tejero-Taldo MI, Gursoy E, Zhao TC, and Kukreja RC (2002) Alpha-adrenergic receptor stimulation produces late preconditioning through inducible nitric oxide synthase in mouse heart. *J Mol Cell Cardiol* **34**:185–195.
- Terzic A, Pucéat M, Clément O, Scamps F, and Vassort G (1992) Alpha 1-adrenergic effects on intracellular pH and calcium and on myofilaments in single rat cardiac cells. *J Physiol* **447**:275–292.
- Tohse N, Nakaya H, Hattori Y, Endou M, and Kanno M (1990) Inhibitory effect mediated by alpha 1-adrenoceptors on transient outward current in isolated rat ventricular cells. *Pflügers Arch* **415**:575–581.
- Tohse N, Nakaya H, and Kanno M (1992) Alpha 1-adrenoceptor stimulation enhances the delayed rectifier K⁺ current of guinea pig ventricular cells through the activation of protein kinase C. *Circ Res* **71**:1441–1446.
- Tosaki A, Behjet NS, Engelman DT, Engelman RM, and Das DK (1995) Alpha-1 adrenergic receptor agonist-induced preconditioning in isolated working rat hearts. *J Pharmacol Exp Ther* **273**:689–694.
- Tsuchida A, Liu Y, Liu GS, Cohen MV, and Downey JM (1994) alpha 1-adrenergic agonists precondition rabbit ischemic myocardium independent of adenosine by direct activation of protein kinase C. *Circ Res* **75**:576–585.
- Turnbull L, McCloskey DT, O'Connell TD, Simpson PC, and Baker AJ (2003) Alpha 1-adrenergic receptor responses in alpha 1A-AR knockout mouse hearts suggest the presence of alpha 1D-AR. *Am J Physiol Heart Circ Physiol* **284**:H1104–H1109.
- Vago T, Bevilacqua M, Norbatio G, Baldi G, Chebat E, Bertora P, Baroldi G, and Accinni R (1989) Identification of alpha 1-adrenergic receptors on sarcolemma from normal subjects and patients with idiopathic dilated cardiomyopathy: characteristics and linkage to GTP-binding protein. *Circ Res* **64**:474–481.
- Valks DM, Cook SA, Pham FH, Morrison PR, Clerk A, and Sugden PH (2002) Phenylephrine promotes phosphorylation of Bad in cardiac myocytes through the extracellular signal-regulated kinases 1/2 and protein kinase A. *J Mol Cell Cardiol* **34**:749–763.
- Van Tassel BW, Rondina MT, Huggins F, Gilbert EM, and Munger MA (2008) Carvedilol increases blood pressure response to phenylephrine infusion in heart failure subjects with systolic dysfunction: evidence of improved vascular alpha1-adrenoreceptor signal transduction. *Am Heart J* **156**:315–321.
- Vaniotis G, Del Duca D, Trieu P, Rohlicek CV, Hébert TE, and Allen BG (2011) Nuclear β -adrenergic receptors modulate gene expression in adult rat heart. *Cell Signal* **23**:89–98.
- Vaughan ED Jr (2003) Medical management of benign prostatic hyperplasia—are two drugs better than one? *N Engl J Med* **349**:2449–2451.
- Vecchione C, Fratta L, Rizzoni D, Notte A, Poulet R, Porteri E, Frati G, Guelfi D, Trimarco V, and Mulvany MJ, et al. (2002) Cardiovascular influences of alpha1b-adrenergic receptor defect in mice. *Circulation* **105**:1700–1707.
- Vega RB, Harrison BC, Meadows E, Roberts CR, Papst PJ, Olson EN, and McKinsey TA (2004) Protein kinases C and D mediate agonist-dependent cardiac hypertrophy through nuclear export of histone deacetylase 5. *Mol Cell Biol* **24**:8374–8385.
- Vettel C, Wittig K, Vogt A, Wuertz CM, El-Armouche A, Lutz S, and Wieland T (2012) A novel player in cellular hypertrophy: Gi β /PI3K-dependent activation of the RacGEF TIAM-1 is required for α_1 -adrenoceptor induced hypertrophy in neonatal rat cardiomyocytes. *J Mol Cell Cardiol* **53**:165–175.
- Vinge LE, Andressen KW, Attramadal T, Andersen GO, Ahmed MS, Peppel K, Koch WJ, Freedman NJ, Levy FO, and Skomedal T, et al. (2007) Substrate specificities of G protein-coupled receptor kinase-2 and -3 at cardiac myocyte receptors provide basis for distinct roles in regulation of myocardial function. *Mol Pharmacol* **72**:582–591.
- Vinge LE, Øie E, Andersson Y, Grøgaard HK, Andersen G, and Attramadal H (2001) Myocardial distribution and regulation of GRK and beta-arrestin isoforms in congestive heart failure in rats. *Am J Physiol Heart Circ Physiol* **281**:H2490–H2499.
- Volz A, Piper HM, Siegmund B, and Schwartz P (1991) Longevity of adult ventricular rat heart muscle cells in serum-free primary culture. *J Mol Cell Cardiol* **23**:161–173.
- Wang BH, Du XJ, Autelitano DJ, Milano CA, and Woodcock EA (2000) Adverse effects of constitutively active alpha(1B)-adrenergic receptors after pressure overload in mouse hearts. *Am J Physiol Heart Circ Physiol* **279**:H1079–H1086.
- Wang GY, McCloskey DT, Turcato S, Swigart PM, Simpson PC, and Baker AJ (2006) Contrasting inotropic responses to alpha1-adrenergic receptor stimulation in left versus right ventricular myocardium. *Am J Physiol Heart Circ Physiol* **291**:H2013–H2017.
- Wang GY, Yeh CC, Jensen BC, Mann MJ, Simpson PC, and Baker AJ (2010) Heart failure switches the RV alpha1-adrenergic inotropic response from negative to positive. *Am J Physiol Heart Circ Physiol* **298**:H913–H920.
- Wang H, Yang B, Zhang Y, Han H, Wang J, Shi H, and Wang Z (2001) Different subtypes of alpha1-adrenoceptor modulate different K⁺ currents via different signaling pathways in canine ventricular myocytes. *J Biol Chem* **276**:40811–40816.
- Wang XL, Wettwer E, Gross G, and Ravens U (1991) Reduction of cardiac outward currents by alpha-1 adrenoceptor stimulation: a subtype-specific effect? *J Pharmacol Exp Ther* **259**:783–788.
- Waspe LE, Ordahl CP, and Simpson PC (1990) The cardiac beta-myosin heavy chain isogene is induced selectively in alpha 1-adrenergic receptor-stimulated hypertrophy of cultured rat heart myocytes. *J Clin Invest* **85**:1206–1214.
- Wencker D, Chandra M, Nguyen K, Miao W, Garantziotis S, Factor SM, Shirani J, Armstrong RC, and Kitsis RN (2003) A mechanistic role for cardiac myocyte apoptosis in heart failure. *J Clin Invest* **111**:1497–1504.
- Whelan RS, Kaplinskiy V, and Kitsis RN (2010) Cell death in the pathogenesis of heart disease: mechanisms and significance. *Annu Rev Physiol* **72**:19–44.
- Wollert KC and Drexler H (2002) Carvedilol prospective randomized cumulative survival (COPERNICUS) trial: carvedilol as the sun and center of the beta-blocker world? *Circulation* **106**:2164–2166.
- Woo SH and Lee CO (1999) Role of PKC in the effects of alpha1-adrenergic stimulation on Ca²⁺ transients, contraction and Ca²⁺ current in guinea-pig ventricular myocytes. *Pflügers Arch* **437**:335–344.
- Wright CD, Chen Q, Baye NL, Huang Y, Healy CL, Kasinathan S, and O'Connell TD (2008) Nuclear alpha1-adrenergic receptors signal activated ERK localization to caveolae in adult cardiac myocytes. *Circ Res* **103**:992–1000.
- Wright CD, Wu SC, Dahl EF, Szazama AJ, and O'Connell TD (2012) Nuclear localization drives α_1 -adrenergic receptor oligomerization and signaling in cardiac myocytes. *Cell Signal* **24**:794–802.
- Xiao L, Pimental DR, Amin JK, Singh K, Sawyer DB, and Colucci WS (2001) MEK1/2-ERK1/2 mediates alpha1-adrenergic receptor-stimulated hypertrophy in adult rat ventricular myocytes. *J Mol Cell Cardiol* **33**:779–787.
- Xiao RP, Zhu W, Zheng M, Chakir K, Bond R, Lakatta EG, and Cheng H (2004) Subtype-specific beta-adrenoceptor signaling pathways in the heart and their potential clinical implications. *Trends Pharmacol Sci* **25**:358–365.
- Yang LL, Gros R, Kabir MG, Sadi A, Gotlib AI, Husain M, and Stewart DJ (2004) Conditional cardiac overexpression of endothelin-1 induces inflammation and dilated cardiomyopathy in mice. *Circulation* **109**:255–261.
- Yang M, Reese J, Cotecchia S, and Michel MC (1998) Murine alpha1-adrenoceptor subtypes. I. Radioligand binding studies. *J Pharmacol Exp Ther* **286**:841–847.
- Zakir RM, Folefack A, Saric M, and Berkowicz RL (2009) The use of midodrine in patients with advanced heart failure. *Congest Heart Fail* **15**:108–111.
- Zhang L, Malik S, Kelley GG, Kapiloff MS, and Smrcka AV (2011) Phospholipase C epsilon scaffolds to muscle-specific A kinase anchoring protein (mAKAPbeta) and integrates multiple hypertrophic stimuli in cardiac myocytes. *J Biol Chem* **286**:23012–23021.
- Zhang L, Malik S, Pang J, Wang H, Park KM, Yule DI, Blaxall BC, and Smrcka AV (2013) Phospholipase C ϵ hydrolyzes perinuclear phosphatidylinositol 4-phosphate to regulate cardiac hypertrophy. *Cell* **153**:216–227.
- Zhang S, Hiraoka M, and Hirano Y (1998) Effects of alpha1-adrenergic stimulation on L-type Ca²⁺ current in rat ventricular myocytes. *J Mol Cell Cardiol* **30**:1955–1965.
- Zhao X, Park J, Ho D, Gao S, Yan L, Ge H, Iismaa S, Lin L, Tian B, and Vatner DE, et al. (2012) Cardiomyocyte overexpression of the α_1 A-adrenergic receptor in the rat phenocopies second but not first window preconditioning. *Am J Physiol Heart Circ Physiol* **302**:H1614–H1624.
- Zhu H, McElwee-Witmer S, Perrone M, Clark KL, and Zilberstein A (2000) Phenylephrine protects neonatal rat cardiomyocytes from hypoxia and serum deprivation-induced apoptosis. *Cell Death Differ* **7**:773–784.
- Zuscik MJ, Chalothorn D, Hellard D, Deighan C, McGee A, Daly CJ, Waugh DJ, Ross SA, Gaivin RJ, and Morehead AJ, et al. (2001) Hypotension, autonomic failure, and cardiac hypertrophy in transgenic mice overexpressing the alpha 1B-adrenergic receptor. *J Biol Chem* **276**:13738–13743.
- Zwart R, Verhaagh S, Buitelaar M, Popp-Snijders C, and Barlow DP (2001) Impaired activity of the extraneuronal monoamine transporter system known as uptake-2 in Orct3/Slc22a3-deficient mice. *Mol Cell Biol* **21**:4188–4196.