Lawrence Berkeley National Laboratory
Recent Work

Title
PRODUCTION OF pi+ -MESONS BY X-RAYS AS A FUNCTION OF ATOMIC NUMBER

Permalink
https://escholarship.org/uc/item/6zx404hg

Author
Mozley, R.F.

Publication Date
1950-06-22

eScholarship.org Powered by the California Digital Library
University of California

UCRL- 706 REV
UNCLASSIFIED

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

UNIVERSITY OF CALIFORNIA — BERKELEY

RADIATION LABORATORY

UCRL-706 Rev.
C.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-706 Revised
Unclassified Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

Production of π^+-Mesons by X-Rays as a Function of Atomic Number

R. F. Mozley

June 22, 1950

Berkeley, California

INSTALLATION	No. of Copies
Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brush Beryllium Company	1
Brookhaven National Laboratory	8
Bureau of Medicine and Surgery	1
Bureau of Ships	1
Carbide and Carbon Chemicals Div., Union Carbide and Carbon Corp. (K-25 Plant)	4
Carbide and Carbon Chemicals Div., Union Carbide and Carbon Corp. (Y-12 Plant)	4
Chicago Operations Office	1
Cleveland Area Office, AEC	1
Columbia University (J. R. Dunning)	2
Columbia University (G. Failla)	1
Dow Chemical Company	1
H. K. Ferguson Company	1
General Electric Company, Richland	3
Harshaw Chemical Corporation	1
Idaho Operations Office	1
Iowa State College	2
Kansas City Operations Branch	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos Scientific Laboratory	3
Mallinckrodt Chemical Works	1
Massachusetts Institute of Technology (A. Gaudin)	1
Massachusetts Institute of Technology (A. R. Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	2
National Bureau of Standards	2
Naval Radiological Defense Laboratory	2
New Brunswick Laboratory	1
New York Operations Office	5
North American Aviation, Inc.	1
Oak Ridge National Laboratory	8
Patent Branch, Washington	1
Rand Corporation	1
Sandia Laboratory	1
Sante Fe Operations Office	1
Sylvania Electric Products, Inc.	1
Technical Information Division, Oak Ridge	15
USAF, Air Surgeon (R. H. Blount)	1
USAF, Director of Armament (C. I. Browne)	1
USAF, Director of Plans and Operations (R. L. Applegate)	1
USAF, Director of Research and Development (F. W. Bruner and R. J. Mason)	2
USAF, Eglin Air Force Base (A. C. Field)	1

INSTALLATION	No. of Copies
USAF, Kirtland Air Force Base (M. F. Cooper)	1
USAF, Maxwell Air Force Base (F. N. Moyers)	1
USAF, NEPA Office	2
USAF, Office of Atomic Energy (A. A. Fickel and H. C. Donnelly)	2
USAF, Offutt Air Force Base (H. R. Sullivan, Jr.)	1
USAF, Wright-Patterson Air Force Base (Rodney Nudenberg)	1
U. S. Army, Atomic Energy Branch (A. W. Betts)	1
U. S. Army, Army Field Forces (James Kerr)	1
U. S. Army, Commanding General, Chemical Corps Technical Command (J. A. MacLaughlin thru Mrs. G. Benjamin)	1
U. S. Army, Chief of Ordnance (A. R. Del Campo)	1
U. S. Army, Commanding Officer Watertown Arsenal (C. H. Deitrick)	1
U. S. Army, Director of Operations Research (Ellis Johnson)	1
U. S. Army, Office of Engineers (Allen O'Leary)	1
U. S. Army, Office of the Chief Signal Officer (Curtis T. Clayton thru G. C. Hunt)	1
U. S. Army, Office of the Surgeon General (W. S. Stone)	1
U. S. Geological Survey (T. B. Nolan)	1
U. S. Public Health Service	1
University of California at Los Angeles	1
University of California Radiation Laboratory	5
University of Rochester	2
University of Washington	1
Western Reserve University	2
Westinghouse Electric Company	4
University of Rochester (R. E. Marshak)	1
California Institute of Technology (R. F. Bacher)	1
Total	144

Information Division
Radiation Laboratory
University of California
Berkeley, California

Production of π^+-Mesons by X-Rays as a Function of Atomic Number

R. F. Mozley

Radiation Laboratory, Department of Physics
University of California, Berkeley, California

June 22, 1950

The electronic method of meson detection developed by Steinberger and Bishop[1] has been used to study the relative cross sections of various elements for production of π^+-mesons by x-rays.

This study has been made at an angle of 90 $\pm$ 8 degrees to the x-ray beam, at approximately 317 Mev synchrotron energy, and at meson energies of 42 $\pm$ 7 Mev and 76 $\pm$ 6 Mev. The targets used were of identical shape and approximately 1.5 grams/cm^2 for both x-ray beam and mesons. With two exceptions the targets were of approximately equivalent meson range. The tin and lead targets were of about 1/10 the range and corrections of about 10 percent were required for this. The other corrections required for meson range, attenuation of the x-ray beam, impurities,[2] and decay of mesons in flight were of the order of 2 percent or less in most cases. No correction has been made for nuclear scattering or nuclear absorption of the mesons in the absorbers. The hydrogen cross section was obtained by a polyethylene, carbon subtraction.

The values of the relative cross sections per proton in the nucleus are tabulated below in arbitrary units.


---

(1) J. L. Steinberger and A. S. Bishop, Phys. Rev. 78, 493 (1950)

(2) Analysis of purity done by Conway and Moore.

Element	42 Mev	Standard Deviation (percent)	76 Mev	Standard Deviation (percent)
H	6.6	17	8.07	11
Li	3.32	10	2.80	11
Be	2.82	11	2.13	10
B	3.02	11	2.28	15
C	2.60	6	1.93	5
Al	2.50	11	1.68	9
Cu	1.92	19	1.17	15
Sn	1.66	25	0.51	55
Pb	0.51	91	0.80	65

Additional non-statistical errors may be as large as 10 percent for hydrogen, tin, and lead and 5 percent for the others.



Steinberger, Panofsky, and Steller have made similar measurements as yet unpublished[3] on the yields of 75 Mev neutral mesons, using the method described in their recent paper.[4] Relative cross sections were obtained for hydrogen, lithium, beryllium, carbon, aluminum, copper, and lead. When these are tabulated in terms of the cross section per nucleon, $\frac{\sigma_{\pi^0}}{A}$, and arbitrarily normalized to the $\frac{\sigma_{\pi^+}}{Z}$ data at beryllium, all except hydrogen agree within the statistical error. Their hydrogen value is approximately the same as the lithium cross section per nucleon.

It is apparent from the above that at least two factors contribute to the decrease of $\frac{\sigma}{Z}$ with atomic number, one dependent on and the other independent of the meson charge. Drs. Chew and Lewis[5] have suggested that for π^+-mesons $\frac{\sigma}{Z}$ may be higher for hydrogen than for other elements since the exclusion principle limits the phase space available for the residual neutron. For other elements lower neutron states are occupied. Such an effect would be less with neutral mesons since the residual nucleon does not have a changed sign.

The further decrease of $\frac{\sigma_{\pi^+}}{Z}$ and $\frac{\sigma_{\pi^0}}{A}$ with atomic number is not incompatible with the possibility that only the nucleons on the surface of the nucleus take part in the production of mesons. The decrease cannot readily be explained by screening of the internal nucleons from the x-rays and hence possibly may be caused by an interaction of the outgoing π-mesons with the nucleons.

---

(3) Steinberger, Panofsky, and Steller have been very kind in allowing me to quote their as yet unpublished work.

(4) J. Steinberger, W. K. H. Panofsky, and J. Steller, Phys. Rev. 78, 802, (1950)

(5) G. Chew, Private communication.

If it is assumed that such an interaction is the entire cause of the effect, a calculation can be made of the mean free path of mesons in nuclear matter. Although the calculated decrease is relatively insensitive to the mean free path assumed, the data are incompatible with a mean free path greater than approximately five times the proton radius.

It is a pleasure to acknowledge the assistance of Dr. J. Steinberger in all parts of this experiment. I wish also to express my gratitude to Professor E. McMillan and W. K. H. Panofsky for their interest and advice. My sincere thanks also go to Mr. Walter Gibbins and the entire synchrotron crew. This work was sponsored by the Atomic Energy Commission.

Information Division
7/3/50 md