

UC Irvine

UC Irvine Previously Published Works

Title

ICTV Virus Taxonomy Profile: Papillomaviridae.

Permalink

<https://escholarship.org/uc/item/702311px>

Journal

Journal of General Virology, 99(8)

Authors

Van Doorslaer, Koenraad

Chen, Zigui

Chan, Paul

et al.

Publication Date

2018-08-01

DOI

10.1099/jgv.0.001105

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



ICTV Virus Taxonomy Profile: *Papillomaviridae*

Koenraad Van Doorslaer,^{1,*} Zigu Chen,² Hans-Ulrich Bernard,³ Paul K. S. Chan,² Rob DeSalle,⁴ Joakim Dillner,⁵ Ola Forslund,⁶ Takeshi Haga,⁷ Alison A. McBride,⁸ Luisa L. Villa,⁹ Robert D. Burk¹⁰ and ICTV Report Consortium

Abstract

The *Papillomaviridae* is a family of small, non-enveloped viruses with double-stranded DNA genomes of 5 748 to 8 607 bp. Their classification is based on pairwise nucleotide sequence identity across the L1 open reading frame. Members of the *Papillomaviridae* primarily infect mucosal and keratinised epithelia, and have been isolated from fish, reptiles, birds and mammals. Despite a long co-evolutionary history with their hosts, some papillomaviruses are pathogens of their natural host species. This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the taxonomy of the *Papillomaviridae*, which is available at <http://www.ictv.global/report/papillomaviridae>.

Table 1. Characteristics of the family *Papillomaviridae*

Typical member:	human papillomavirus 16 (K02718), species <i>Alphapapillomavirus 9</i>, genus <i>Alphapapillomavirus</i>, subfamily <i>Firstpapillomavirinae</i>
Virion	Non-enveloped, 55 nm, icosahedral
Genome	Circular dsDNA genome of 5 748 to 8 607 bp
Replication	Bidirectional (theta) replication
Translation	Early and late transcripts, alternative splicing, alternative open reading frames
Host Range	Mammals, birds, reptiles and fish
Taxonomy	Two subfamilies include >50 genera and >130 species

VIRION

The non-enveloped viral capsid is ~600 Å in diameter and consists of 72 pentamers of the major capsid protein, L1, and ~12 molecules of the L2 minor capsid protein (Table 1, Fig. 1) [1].

GENOME

The viral genome varies from 5 748 to 8 607 bp. The genome comprises three functional regions. The early region encodes proteins involved in transcription, replication, and manipulation of the cellular milieu. The late region encodes

the capsid proteins L1 and L2. The upstream regulatory region, located between the L1 and E6 open reading frames, contains the origin of replication as well as binding sites for viral and cellular transcription factors (Fig. 2) [2].

REPLICATION

Papillomaviruses primarily infect epithelial cells. Following a micro-abrasion, the incoming virion complexes with extracellular heparin sulfate proteoglycans on the basement membrane. This interaction results in conformational changes in the L1 and L2 capsid proteins, in turn allowing

Received 23 May 2018; Accepted 31 May 2018

Author affiliations: ¹School of Animal and Comparative Biomedical Sciences, Cancer Biology Graduate Interdisciplinary Program, Genetics Graduate Interdisciplinary Program, BIO5 Institute, and the University of Arizona Cancer Center, University of Arizona, Tucson, AZ, USA; ²Department of Microbiology, The Chinese University of Hong Kong, Hong Kong SAR, PR China; ³Department of Molecular Biology and Biochemistry, School of Biological Sciences, University of California Irvine, Irvine, CA, USA; ⁴Sackler Institute for Comparative Genomics, American Museum of Natural History, Central Park West and 79th St., New York, NY, USA; ⁵International HPV Reference Center, Department of Laboratory Medicine, Karolinska Institutet, 14186 Stockholm, Sweden; ⁶Department of Medical Microbiology, Laboratory Medicine, Lund University, Sölvegatan 23, Sjukhusområdet, 221 85 Lund, Sweden; ⁷Division of Infection Control and Disease Prevention, Department of Veterinary Medical Science, The University of Tokyo, Tokyo, Japan; ⁸Laboratory of Viral Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA; ⁹Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, Instituto do Câncer do Estado de São Paulo, Centro de Investigação Translacional em Oncologia, Universidade de São Paulo, São Paulo, SP, Brazil; ¹⁰Departments of Epidemiology and Population Health, Pediatrics, Microbiology and Immunology, and Obstetrics and Gynecology and Women's Health, Albert Einstein College of Medicine, Bronx, NY, USA.

*Correspondence: Koenraad Van Doorslaer, vandoorslaer@email.arizona.edu

Keywords: *Papillomaviridae*; ICTV Report; taxonomy.




Fig. 1. Atomic rendering of a papillomavirus capsid. Derived from an image reconstruction from cryo-electron microscopy of human papillomavirus type 16 at 4.5 Å resolution and colored according to the radial coloring scheme shown (PDB: 5KEP; [7]).

for transfer of the virion to an unknown entry receptor. Following furin cleavage of L2, the virion becomes internalised using a process that shares similarities with macropinocytosis. The L2-DNA complex traffics to the trans-Golgi network, until mitosis. By metaphase, the viral DNA can be seen to be associated with host chromosomes [3]. The viral replication cycle consists of three distinct phases of replication. Initial limited viral DNA amplification is supported by the viral E1 and E2 replication proteins. This initial burst of replication is followed by maintenance replication, during which the viral genome is maintained at a relatively low, but constant copy number in the proliferating cells of a lesion. Finally, as an infected cell completes cellular differentiation there is a switch towards differentiation-dependent genome amplification, and eventual generation of progeny virions [4]. During maintenance replication, the viral E6 and E7 proteins are able to usurp the cellular environment, allowing




Fig. 2. Diagram of the human papillomavirus 16 genome. The viral dsDNA is indicated. The outer boxes indicate the protein-coding open reading frames. Dotted lines represent intron sequences. The black circle represents the viral origin of replication (ori).

for viral replication in differentiated cells [5]. In the top layers of the differentiated epithelia, viral DNA is amplified to a high copy number and the capsid proteins self-assemble into particles encapsidating the viral DNA. As the cells slough off into the environment, infectious virions are released, completing the viral replication cycle.

TAXONOMY

Classification of the *Papillomaviridae* is based on sequence identity across the L1 open reading frame [6]. The family includes two subfamilies, *Firstpapillomavirinae*, which includes >50 genera and >130 species, and *Secondpapillomavirinae*, with a single genus and species. Genera are named according to the Greek alphabet (e.g. *Alphapapillomavirus*), with the prefixes ‘Dyo-’ and ‘Treis-’ indicating additional cycles through the alphabet (i.e. Greek for ‘a second or third time’).

RESOURCES

- Full ICTV Online (10th) Report: <http://www.ictv.global/report/papillomaviridae>.
- The Papillomavirus Episteme (PaVE): <http://pave.niaid.nih.gov>.
- The human papillomavirus reference center: <http://www.hpvcenter.se>.
- The animal papillomavirus reference center: <http://www.animalpv.org>.

Funding information

Production of this summary, the online chapter, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA). The work in the McBride lab is in part supported by the Intramural Research Program of the NIAID, NIH.

Acknowledgements

Current and previous members of the ICTV papillomavirus study group. Members of the ICTV Report Consortium are Elliot J. Lefkowitz, Andrew J. Davison, Stuart G. Siddell, Peter Simmonds, Sead Sabanadzovic, Donald B. Smith, Richard J. Orton and Balázs Harrach.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Finnen RL, Erickson KD, Chen XS, Garcea RL. Interactions between papillomavirus L1 and L2 capsid proteins. *J Virol* 2003;77:4818–4826.
2. van Doorslaer K. Evolution of the *Papillomaviridae*. *Virology* 2013; 445:11–20.
3. Campos SK. Subcellular trafficking of the papillomavirus genome during initial infection: the remarkable abilities of minor capsid protein L2. *Viruses* 2017;9:370.
4. McBride AA. Replication and partitioning of papillomavirus genomes. *Adv Virus Res* 2008;72:155–205.
5. Harden ME, Munger K. Human papillomavirus molecular biology. *Mutat Res Rev Mutat Res* 2017;772:3–12.
6. de Villiers EM, Fauquet C, Broker TR, Bernard HU, Zur Hausen H. Classification of papillomaviruses. *Virology* 2004;324:17–27.
7. Guan J, Bywaters SM, Brendle SA, Ashley RE, Makhov AM et al. Cryoelectron microscopy maps of human papillomavirus 16 reveal L2 densities and heparin binding site. *Structure* 2017;25: 253–263.