

UC-70
LBL-23637
c.1



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

EARTH SCIENCES DIVISION

RECEIVED
LAWRENCE
BERKELEY LABORATORY

AUG 21 1987

LIBRARY AND
DOCUMENTS SECTION

To be presented at the American Nuclear Society
Winter Meeting, Los Angeles, CA, November 15-19, 1987

PRESSURE-INDUCED BRINE MIGRATION IN CONSOLIDATED SALT IN A REPOSITORY

Y. Hwang, P.L. Chambré, W.W.-L. Lee, and T.H. Pigford

June 1987

For Reference

Not to be taken from this room



LBL-23637
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Pressure-Induced Brine Migration in Consolidated Salt in a Repository*

Y. Hwang, P. L. Chambre, W. W.-L. Lee and T. H. Pigford
Department of Nuclear Engineering & Lawrence Berkeley Laboratory
University of California, Berkeley, CA 94720

Within a few years after the emplacement of waste packages in a geologic repository of nuclear waste in salt, salt creep is expected to close the air gap between a waste container and the borehole wall.¹ In this paper we present an analysis of brine migration completely inside consolidated salt, considered here to be a thermoelastic, porous rock.

Salt creep causes consolidation of the crushed salt in the annulus between the waste package and the borehole, resulting in monolithic consolidation of salt around the waste package. Thereafter, neglecting the consumption of brine by container corrosion, brine in grain boundaries near the waste package can only migrate outward into the surrounding salt, under the influence of pressure gradients caused by transient heating of the salt. Figure 1 shows qualitatively the profiles of temperature and brine pressure expected in the salt after consolidation². Hot salt near the waste package expands against the waste package and surrounding salt, resulting in high compressive stresses near the waste package. Grain-boundary brine expands more than does the salt and further increases the local pressure and pressure gradients that cause brine to flow outward into the cooler salt. Such outward flow of brine relieves the pressure gradient on the fluid, which finally relaxes to near-lithostatic pressure. If the waste containers are corroded, then this outward brine movement may become a mechanism for radionuclide transport. To determine the extent to which advection by brine in grain boundaries is an important transport mechanism for released radionuclides, we have estimated the time-dependent migration of brine after salt consolidation.

We consider a spherical-equivalent spent-fuel waste package of 0.72-meter radius in an infinite salt medium. The governing equations for brine migration treating salt as a thermoelastic, porous rock follow those of McTigue.³ Chambré obtained analytic solutions to the governing equations⁴, and numerical illustrations of the solutions are given here. The parameter values used in the calculations are from McTigue, and shown in Table 1. The heat flux used was 928 watts per square meter.

*Work supported in part by U. S. Department of Energy via Contract DE-AC03-76SF00098

Figure 2 shows the superficial brine velocity near a waste package as a function of time. Pressure-induced brine migration within consolidated salt is a transient phenomenon. After about ten years the velocity is nearly zero. Even the highest velocity is of the order of *millimeters per year*, and occurs only within a meter or so of a waste package. At these velocities molecular diffusion may be the dominant transport process for radionuclides in salt. In a companion paper⁵ we show the application of diffusional mass transfer to estimating release rates in a salt repository.

References

1. T. Brandshaug, 1987, *Estimate of Consolidation of Crushed Salt Around a Spent Fuel Waste Package*, RE/SPEC Report RSI-315
2. T. H. Pigford and P. L. Chambré, "Mass Transfer in a Salt Repository," LBL-19918, 1985
3. D. T. McTigue, 1986, "Thermoelastic Response of Fluid-Saturated, Porous Rock," *J. Geophys. Res.*, 91, B9, 9533
4. P. L. Chambré, to be published
5. P. L. Chambré, Y. Hwang, W. W.-L. Lee and T. H. Pigford, 1987 "Release Rates from Waste Packages in a Salt Repository," UCB-NE-4102, Paper submitted for the 1987 Winter Meeting, American Nuclear Society

Table 1 Parameter Values Used in Calculations
(After McTigue,³ for the Salado Formation, Delaware Basin, New Mexico)

Property	Value	Units
Conductivity (\mathcal{K})	6.60	$W \cdot m^{-1} \cdot K^{-1}$
Heat Capacity (ρc_v)	1.89×10^6	$J \cdot m^{-3} \cdot K^{-1}$
Drained Bulk Modulus (K)	20.7	<i>GPa</i>
Fluid Bulk Modulus (K_f)	2.0	<i>GPa</i>
Solid Bulk Moduli (K'_s, K''_s)	23.5	<i>GPa</i>
Shear Modulus (G)	12.4	<i>GPa</i>
Porosity (ϕ_o)	0.001	
Fluid Expansivity (α_f)	3.0×10^{-4}	K^{-1}
Solid Expansivities (α'_s, α''_s)	1.2×10^{-4}	K^{-1}
Permeability (k)	10^{-21}	m^2
Fluid Viscosity (μ)	1.0×10^{-3}	<i>Pa · s</i>
B	0.93	
Poisson's Ratio (ν)	0.25	
Undrained Poisson's Ratio (ν_u)	0.27	
b'	29.0*	<i>kPa · K⁻¹</i>
Fluid Diffusivity (c)	0.16×10^{-6} *	$m^2 \cdot s^{-1}$
Thermal Diffusivity (κ)	3.5×10^{-6}	$m^2 \cdot s^{-1}$
R ($\sqrt{c/\kappa}$)	0.21	

* Calculated for zero mean stress.




Figure 1. Pressure Profile on Intergranular Brine after Consolidation of Salt in a Salt Repository (from Pigford and Chambré, 1985)




Figure 2. Brine Migration Velocity Away from a Waste Package Due to Pressure Gradient

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*