

UC Irvine

UC Irvine Previously Published Works

Title

Feasibility of an emotional health curriculum for elementary school students in an underserved Hispanic community.

Permalink

<https://escholarship.org/uc/item/7147d0ck>

Journal

Journal of child and adolescent psychiatric nursing : official publication of the Association of Child and Adolescent Psychiatric Nurses, Inc, 30(3)

ISSN

1073-6077

Authors

Guo, Yuqing
Rousseau, Julie
Renno, Patricia
et al.

Publication Date


2017-08-01

DOI

10.1111/jcap.12185

Peer reviewed

Feasibility of an emotional health curriculum for elementary school students in an underserved Hispanic community

Yuqing Guo PhD, RN¹  | Julie Rousseau PhD, CNM, RN¹ | Patricia Renno PhD² | Priscilla Kehoe PhD¹ | Monique Daviss BA³ | Sara Flores MA³ | Kathleen Saunders MSN, RN, CNS¹ | Susanne Phillips DNP, APRN, FNP-BC, FAANP¹ | Mindy Chin BS, RN¹ | Lorraine S. Evangelista PhD, RN, FAAN¹

¹Sue & Bill Gross School of Nursing, University of California, Irvine, CA, USA

²Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles Semel Institute for Neuroscience & Human Behavior, Los Angeles, CA, USA

³El Sol Science and Arts Academy of Santa Ana Santa Ana, CA, USA

Correspondence

Yuqing Guo, PhD, RN, Sue & Bill Gross School of Nursing, University of California, Irvine, CA 92697-3959, USA.

Email: gyuqing@uci.edu

Abstract

Problem: Hispanic children have greater mental health challenges but fewer received mental health services than other ethnic groups. A classroom-based Emotional Health Curriculum (EHC) was developed to address mental health disparities in an underserved Hispanic community.

Methods: A quasi-experimental design with one group pre- and post-intervention was used to test the feasibility of an 8-week EHC for one hundred 3rd and 4th grade children in a dual-immersion Spanish-English elementary school. Limited efficacy was measured by changes in depression and anxiety scores reported by children and teachers. Acceptance was evaluated by a child-reported satisfaction survey and a focus group in which the four teachers shared their experiences. Implementation was measured by participation, retention, and fidelity rates.

Findings: The child-reported depression and anxiety and teacher-reported depression were significantly decreased in at-risk children with the effect size ranging from 0.60 to 1.16 ($ps < 0.05$). The majority of children (89.7%) enjoyed the EHC and teachers observed that children had acquired skills to manage their emotional distress. The participation, retention, and fidelity rates were 98%, 94%, and 99.13%, respectively.

Conclusions: The results provide promising evidence that the EHC has the potential to improve depression and anxiety symptoms in at-risk children.

1 | INTRODUCTION

1.1 | Mental health disparities

Studies document that anxiety and depression (i.e., internalizing behaviors) are the most common psychiatric conditions in children (Ahlen, Lenhard, & Ghaderi, 2015; Bennett et al., 2015). In the United States, among children aged 8–15 years, 3.7% had major depression and 0.7% generalized anxiety disorder (Merikangas et al., 2010). Toppelberg, Hollinshead, Collins, and Nieto-Castañón (2013) found the prevalence rate of 4.9% of internalizing problems reported by teachers in a community sample of 5- to 7-year-old Hispanic children. Some studies showed that Hispanic children experienced higher depression rates than Whites, Blacks, and Asian Americans (Anderson & Mayes,

2010; Joiner, Perez, Wagner, Berenson, & Marquina, 2001; Siegel, Yancey, Aneshensel, & Schuler, 1999). In another study in the United States, Saluja et al. (2004) reported that 22% of Hispanic youth had depressive symptoms compared with 18% of White, 17% of Asian American, and 15% of Black youth.

Although Hispanic children are at greater risk for mental disorders, Hispanic youth receive fewer and lower quality mental health services than non-Hispanic, White children (Alegria, Vallas, & Pumariega, 2010; Merikangas et al., 2011; Toppelberg et al., 2013). This type of mental health service gap is especially pronounced for internalizing behaviors because it has been observed that Hispanic children often do not perceive “feeling depressed or afraid” as rationale for treatment (Cumings, Case, Ji, Chae, & Druss, 2014). Research has identified multilevel barriers to mental health services in Hispanic families including financial constraints, transportation issues, lack of mental health

services in the neighborhood, limited English language fluency, low health literacy, and persistent stigmas against mental health (Bear, Finer, Guo, & Lau, 2014; Stewart, Simmons, & Habibpour, 2012; Uebelacker et al., 2012). School-based programs have the potential to reduce these barriers.

1.2 | School-based mental health program

Preventing anxiety and depression are a public health priority (World Health Organization [WHO], 2008), as these disorders in childhood have deleterious consequences for academic success, emotional well-being, and peer relationships throughout childhood and adolescence (Bennett et al., 2015; Cerdá et al., 2013; Hughes, Lourea-Waddell, & Kendall, 2008). Once these disorders are established they are more likely to persist in adulthood (Patton et al., 2014). Schools become promising settings to provide mental health wellness programs not only because children spend most of their day there, but also because they are rife with anxiety-provoking academic and social situations (Herzig-Anderson, Colognori, Fox, Stewart, & Masia Warner, 2012). School-based programs also reduce barriers to mental health care through increasing accessibility and decreasing stigma (Bear et al., 2014; Keeton, Soleimanpour, & Brindis, 2012; Mason-Jones et al., 2012). Juszczak, Melinkovich, and Kaplan (2003) found that adolescents were 21 times more likely to seek school-based mental health services than community-based mental health care. Comparing mental health service utilization in adolescents across clinic and school settings, Cummings, Ponce, and Mays (2010) showed that Hispanic youth had lower rates of receiving counseling in clinical settings relative to White youth, whereas there were no ethnic differences in counseling service use in school settings.

A growing body of evidence supports the effectiveness of school-based cognitive behavioral therapy (CBT) or mindfulness in preventing depression and anxiety in children and adolescents (Barrett & Turner, 2001; Callear & Christensen, 2010; Gillham et al., 2007; Mendelson et al., 2010; Neil & Christensen, 2009). However, there are very few culturally sensitive CBT or mindfulness interventions focusing on preventing depression and anxiety in underserved Hispanic children. To address the gap of knowledge in prevention and early intervention at a school setting with few mental health resources, our research team created a classroom-based Emotional Health Curriculum (EHC) by integrating previous classroom-based CBT principles and structure, mindfulness skills, and also utilizing a coordinated nursing care implementation approach. Additionally, Hispanic cultural terms and concepts were threaded throughout the EHC to increase the cultural relevance of the intervention implemented in a dual-immersion Spanish-English elementary school.

1.3 | Theoretical framework

Cognitive behavioral theory guides the development of the EHC as CBT is the first-line recommended approach to mental health disorders in pediatrics (Forman-Hoffman et al., 2016). This theory emphasizes identifying affective, cognitive, and behavioral responses to distressing situations and modifying anxious or depressed experiences by

changing maladaptive beliefs and generating positive thoughts (Beck, 2005; Fujii et al., 2013; Kendall & Hedtke, 2006). Acquisition of self-control skills is the key CBT principle, including skills such as identifying negative feelings and restructuring depressed or anxious feelings by generating positive thoughts (Rey, Marin, & Silverman, 2011). Mindfulness refers to the ability to focus moment-to-moment attention on thoughts, feelings, or perceptions in nonjudgmental and non-reactive ways (Schonert-Reichl et al., 2015). Mindfulness practice was found to promote cognitive and emotional self-regulation of rumination or intrusive thoughts that are common symptoms of internalizing problems (Mendelson et al., 2010; Weijer-Bergsma, van de, Langenberg, Brandsma, Oort, & Bögels, 2014). In order to enhance self-control and self-awareness skills, mindfulness was integrated into the EHC as an additional coping strategy.

A recent meta-analysis showed that professional healthcare providers using a school-based curriculum delivery model had a stronger positive impact overall, compared to those given by classroom teachers (Werner-Seidler, Perry, Callear, Newby, & Christensen, 2017). Given the scarcity of mental healthcare providers in the community of Santa Ana, the EHC was implemented through a coordinated nursing team model (i.e., trained licensed registered nurses who were in a Master of Science Family Nurse Practitioner program together with trained senior Bachelor of Science in Nursing students under the supervision of University Nursing Professors).

The primary aim of this study was to examine the feasibility of using the EHC as a preventive intervention with 3rd and 4th grade children in a dual-immersion Spanish-English elementary school in terms of limited efficacy, acceptability, and implementation. Specifically, the teachers' perspectives and children's self-report were used to assess of the limited efficacy of the EHC on depression and anxiety as well as the acceptability of the intervention.

2 | METHODS

2.1 | Design and setting

A quasi-experimental design with one group pre- and post-intervention was used to examine the limited efficacy of the EHC. A limited efficacy refers to the impact of an intervention in a convenience sample with intermediate outcomes and limited statistical power (Bowen et al., 2009). Acceptability describes experiences in targeted individuals or those involved in the programs (Bowen et al., 2009). Children's satisfaction and teachers' experiences were used to describe their acceptance of the EHC. Intervention Fidelity Checklist (i.e., how the intervention was consistently delivered), participation, and retention rates (i.e., how the children were engaged in the intervention) are indicators of implementation (Bowen et al., 2009). The timeline of this study was designed to align with the school's schedule: children attended the EHC once a week with each session lasting approximately 45 minutes for 8 weeks. The research team took 15 minutes to collect the pre- and post-intervention data with the 3rd and 4th grade children during the first and the last sessions. Teachers' data were collected one week before the intervention and one month after the

TABLE 1 Content of the emotional health curriculum

Session	Focus
Emotional awareness	Use words to describe different emotions in both English and Spanish
Body signals	Recognize behavioral and physiological reactions to positive and negative emotions
Negative self-talk	Identify negative thoughts and their relation to emotions
Brave thoughts	Generate positive thoughts to replace negative self-talk
Deep breathing	Breathe in with their belly rather than their chest and breathing out slowly through their mouth
Mindfulness exercises	Integrate emotional awareness, body signals, and deep breathing techniques to non-judgmentally and non-reactively respond to emotional distress
Action steps	Calmly facing challenging situations by practicing mindfulness and breaking them into manageable steps
Steps to friendship	Communicate feelings to a friend and empathetically respond to a peer's feelings

intervention, respectively. The study was implemented in the classrooms of a dual-immersion Spanish–English elementary school in the community of Santa Ana, California between January and June, 2016.

2.2 | Intervention

The classroom-based EHC was composed of eight sessions in which children acquired skills to become aware of emotions, recognize body signals of emotional distress, practice mindfulness to manage distress, understand the impact of both negative and positive self-talk on feelings, and communicate their feelings with friends, teachers, and parents as well as show empathy toward friends' distress (See Table 1). Each session had the same format starting with reviewing homework assigned the previous week, then reading a story to introduce a new concept that relates to emotional health, followed by a demonstration of the skill necessary to deal with emotional distress and communication. Children were then divided into small groups of four to five children to one nursing student facilitator where they were able to practice the concept or skill.

The EHC was adapted to the Hispanic community by adding cultural metaphors, linguistic elements, and social contexts (Castro, Barrera, & Holleran Steiker, 2010; Macklem, 2014). Given that this is a dual-immersion Spanish–English school, 3rd and 4th grade children are fluent in both Spanish and English and also have a shared understanding of Hispanic culture through their class curriculums. A bilingual MD/MPH student facilitated cultural adaptation of the EHC through integrating the feedback from reviews of two school teachers. First, Hispanic cultural symbols were built into the teaching materials and rewards. For example, a horse, a symbol of strength and wealth in Mexican culture, was designed as the mascot of the curriculum. Second, Spanish words were included to enhance the

children's understanding of the curriculum. Children were taught to identify their emotions with words in both English and Spanish. Third, scenarios specific to the children's school life were captured in role-play and small group activities, such as managing test anxiety and trying out for the school play or band. One scenario recommended by school teachers was the first day of school distress, "On the first day of school, I don't have all of my school supplies." The students were then encouraged to come up with positive thoughts and calming strategies to address their distressed and anxious feelings.

The teacher was present when the EHC was delivered in the classroom and encouraged children to complete the weekly homework related to that session. Each class had the same coordinated nursing team so that the weekly sessions were delivered with consistency in terms of explaining the concepts, facilitating small groups, checking homework, and reinforcing positive classroom behaviors. Additionally, the research team had developed an Intervention Fidelity Checklist to further ensure the consistent delivery of the EHC. Nursing Professors directly supervised every aspect of the implementation procedures in each classroom and completed the Fidelity Checklist to record the completion of the consistent curriculum and activities.

2.3 | Procedure

All stages and tools utilized in this study had University Institutional Review Board (IRB) approval. The elementary school administrator, designated as the study coordinator, supported recruitment. Parents of 3rd and 4th grade students were informed about the study through a bilingual Spanish/English Parent Newsletter and Parent Study Information Sheet. The information was provided to all parents, so they could decide whether their child would participate in or withdraw from the study (i.e., allow the individual data to be collected and reported). Parents were notified that their children were expected to attend the EHC as part of their general health class. Also, they were told that it was entirely voluntarily for their child to fill out surveys and questionnaires. No parents withdrew or objected to their children's participation in data collection.

Using a written script, the first author trained the coordinated nursing team to consistently explain the study procedures to the children. Verbal assent was obtained from all children. At baseline demographics and pre-intervention questionnaires were collected and after the intervention, the questionnaires and satisfaction surveys were collected. To support children's comprehension of the questionnaires, the trained research assistants read the questions aloud and provided additional assistance for those children who had difficulty understanding or filling them out. Children received school supplies as rewards for their participation.

The research team and the elementary school collaboratively developed a comprehensive plan to minimize risks and potential emotional distress, and to clinically screen any children self-identified or observed to be at risk; this plan was part of our approved IRB protocol. Specifically, if a child self-reported distressed feelings, abuse, or risky family/social circumstances any time or exhibited signs of distress, a licensed nurse faculty who supervised the intervention in the classroom each session immediately notified and referred

this child to the school administrator who was part of our research team. This designated school administrator functioned as the school's social support coordinator in the study and was responsible for direct referral to the school counselor and coordination with community mental health services according to the individual child's needs.

The four teachers of the 3rd and 4th grade classrooms received a Teacher Study Information Sheet and were asked to fill out an individual assessment for each of their students. Each teacher received \$200 in appreciation for their significant time and effort to complete the demographic survey as well as pre- and post-intervention questionnaires (24–26 students in a classroom). At the end of the intervention, the study coordinator invited the four teachers to attend a focus group to share their experiences with the EHC. The bilingual MD/MPH student, trained by the first author, facilitated the focus group using semistructured interview questions. Beginning with the consent process, all the teachers agreed to be audio-recorded using a digital recorder. Each teacher received a \$30 gift card as compensation for their time participating in the focus group.

2.4 | Participants

Attendance of the 102 3rd and 4th grade children was obligatory as the EHC was included in the required health curriculum classes for 8 weeks. Two children dropped from the curriculum due to family crisis necessitating direct school counselor interventions. Of the 100 children, 82 were Hispanic, 13 American Indian or Alaska Native, and four non-Hispanic Whites. The girl to boy ratio was almost equal with 45.0% girls to 55.0% boys. The average age was 8.71 years old with 50% children being in the 3rd and 50% in the 4th grades. Most of the students (80.0%) lived with both their parents and 83.0% spoke both English and Spanish at home. All four school teachers (three females and one male), who participated in this study, were Hispanic, bilingual educators. Three of them had master's degrees and had taught at this dual-immersion Spanish–English elementary school for five years or more; the fourth teacher had a bachelor's degree and taught at the school for two years.

2.5 | Measures

The Revised Child Anxiety and Depression Scale-25 (RCADS-25; Ebesutani et al., 2012) is a child self-reported questionnaire including 10 items to measure major depression symptoms and 15 items to measure general anxiety symptoms on a four-point scale (0 = never, 1 = sometimes, 2 = often, 3 = always). The RCADS-25 is an instrument with acceptable content and concurrent validity for both school-based and clinic-referred samples for children in grades 3–12 in various ethnicity groups including Whites and Hispanics (Ebesutani et al., 2012). The Cronbach's alpha for depression, anxiety, and total problems in this sample ranged from 0.74 to 0.91. According to the RCADS guidelines, children with RCADS-25 total scores equal to or above 65 are considered to be in the borderline/clinical range or at risk; children with total scores lower than 65 are within the normal range.

The Teacher's Report Form (TRF; Achenbach, Dumenci, & Rescorla, 2003) was used by classroom teachers to evaluate their students'

internalizing and externalizing behaviors for the past month in children aged 6–18 years. Teachers reported their perceptions of students' depression and anxiety symptoms before and after the EHC. The TRF is a culturally validated assessment in Hispanic populations and consists of 33 items to measure internalizing broad symptoms including depression, anxiety, and somatic complaints rated on a three-point scale (0 = not true, 1 = somewhat or sometimes true, and 2 = very true or often true) (Bird, 1996; Toppelberg et al., 2013). Internalizing behaviors were the focus and thus reported in this study. According to the TRF broad symptom standard, children with TRF internalizing symptoms equal to or above 60 are considered to be in the borderline/clinical range or at risk; children with internalizing symptoms lower than 60 are within the normal range. In order to match child RCADS-25 scores (depression, anxiety), we reported TRF depression and anxiety separately. The Cronbach's alpha for depression and anxiety in this sample ranged from 0.77 to 0.85.

The children's satisfaction survey was developed by the first author who had training in and experience with instrument development. Prior to administration, the research team and school teachers provided feedback on the survey in terms of content and language use. The survey was composed of two sections including an overall assessment of the EHC and a specific evaluation of each of the eight skills taught during the EHC. For instance, an overall question "Did you enjoy the emotional health classes?" included the responses, "Yes," "A little," and "No." And a specific question "Did you find deep breathing useful?" included the responses, "Useful," "Somewhat useful," and "Not useful." The Cronbach's alpha for this instrument was 0.85. We also included an open-ended question at the end of the survey asking students to provide written feedback or draw a picture reflecting their experiences with the EHC.

2.6 | Analysis

We used IBM SPSS, version 22.0 (SPSS, Armonk, NY) to conduct analyses. As recommended by the instrument manuals, the three raw RCADS-25 scores (depression, anxiety, and total scores) and the three raw TRF scores (depression, anxiety, and internalizing symptoms) were converted to T scores for analyses. First, descriptive analyses were used to examine characteristics of children and teachers and also to evaluate children's satisfaction. Second, repeated measures analysis of variance (ANOVA) and post hoc paired t-tests were conducted to examine the changes in depression and anxiety before and after the EHC in at-risk children versus normal children. Children's and teachers' assessments were analyzed separately due to differences in scales and few at-risk children overlapped between them. The Cohen's *d* statistics were used to report effect sizes.

The audio recordings of the teachers' interviews were transcribed verbatim by a nursing student who was trained by the first author to accurately convert speech to a written document. Content analysis was conducted according to constant comparison methods identifying differences and seeking commonalities (Dye, Schatz, Rosenberg, & Coleman, 2000). Three steps were used in the analysis: (1) two researchers independently analyzed two transcripts by identifying line-by-line themes with manual coding; (2) the same two researchers

TABLE 2 Changes in child-reported and teacher-reported depression and anxiety

^a RCADS	At-Risk Group (n = 7)		Normal Group (n = 87)		P (Time)	P (Group)	P (T x G)
	Pre-intervention (M±SD)	Post-intervention (M±SD)	Pre-intervention (M±SD)	Post-intervention (M±SD)			
Depression	65.69 ± 6.11	53.28 ± 10.93	45.52 ± 7.95	44.56 ± 10.63	< 0.000	<0.000	0.001
Anxiety	66.23 ± 3.98	52.42 ± 10.24	46.70 ± 7.76	46.37 ± 9.85	< 0.000	<0.000	<0.000
^b TRF	At-Risk Group (n = 16)		Normal Group (n = 84)		P (Time)	P (Group)	P (T x G)
	Pre-intervention (M±SD)	Post-intervention (M±SD)	Pre-intervention (M±SD)	Post-intervention (M±SD)			
Depression	69.50 ± 9.75	65.00 ± 8.49	52.42 ± 3.72	52.49 ± 3.75	< 0.000	<0.000	<0.000
Anxiety	62.75 ± 5.78	61.56 ± 8.46	52.31 ± 3.41	53.39 ± 4.50	0.93	<0.000	0.07

Note. ^aRCADS, Revised Child Anxiety Depression Scales-25. ^bTRF, Teacher Report Form.

discussed coding to reach consensus about themes; and (3) the same two researchers categorized the themes and summarized the key overarching themes. In order to maintain teachers' voices, codings generated in the analyses were based on teachers' direct language, rather than the researchers' paraphrases (Saldaña, 2009).

3 | RESULTS

3.1 | Limited efficacy of the emotional health curriculum

The repeated measures ANOVA showed that for child-reported depression there was a significant interaction effect between group and time ($F(1, 92) = 12.04, p = .001$). The same significant interaction effect was found for child-reported anxiety ($F(1, 92) = 18.32, p < .000$). Post hoc paired t-tests found that there were significant decreases in child-reported depression ($t(1, 6) = 3.08, p = 0.02, d = 1.16$) and anxiety ($t(1, 6) = 3.02, p = 0.02, d = 1.14$) with a large effect size in children who rated themselves at risk, while depression ($t(1, 86) = 1.09, p = 0.28, d = 0.12$) and anxiety scores ($t(1, 86) = 0.40, p = 0.69, d = 0.04$) in children self-rated as normal did not change significantly (see Table 2).

The repeated measures ANOVA showed that for teacher-reported depression there was a significant interaction effect between group and time ($F(1, 98) = 16.53, p < .000$). Post hoc paired t-tests found that teacher-reported depression ($t(1, 15) = 2.38, p = 0.03, d = 0.60$) was significantly decreased with a moderate effect size in children who were rated at risk by teachers, while depression scores ($t(1, 83) = -0.21, p = 0.83, d = 0.02$) in children who were rated as normal by teachers did not change significantly. The interaction effect for teacher-reported anxiety was marginally significant ($F(1, 98) = 3.38, p = .07$) and thus post hoc paired t-tests were not conducted (see Table 2).

3.2 | Acceptability of the emotional health curriculum

3.2.1 | Children's satisfaction with the EHC

Overall, the majority of the school children (89.7%) enjoyed the EHC and 72.9% reported that the EHC taught them skills to manage stress and worries. Table 3 summarizes children's assessment of each skill

TABLE 3 Child assessment of each session of emotional health curriculum (N = 97)

Emotional Health Skills	Not Useful	Somewhat Useful	Useful
Emotional awareness	2 (2.1%)	16 (16.5%)	79 (81.4%)
Body signals	3 (3.1%)	19 (19.6%)	75 (77.3%)
Negative self-talk	3 (3.1%)	29 (29.9%)	65 (67.0%)
Brave thoughts	1 (1.0%)	10 (10.3%)	86 (88.7%)
Deep breathing	5 (5.1%)	9 (9.3%)	83 (85.6%)
Mindfulness exercises	2 (2.1%)	11 (11.3%)	84 (86.6%)
Action steps	7 (7.2%)	29 (29.9%)	61 (62.9%)
Steps to friendship	6 (6.2%)	21 (21.6%)	70 (72.2%)

taught during the EHC sessions. Approximately, two-thirds of the students perceived every skill to be "useful" (62.9%–88.7%). Brave thoughts (88.7%), mindfulness exercises (86.6%), and deep breathing (85.6%) were reported as the most useful skills. Approximately, two thirds of the children (62, 63.92%) responded to the open-ended question by sharing their experiences with the EHC. Of these children, the majority (56, 90.32%) expressed positive experiences with the EHC such as "I like that you taught us about our emotions" and "It was the best time I had in school". Six students (9.68%) reported neutral comments such as "the classes were so-so".

3.2.2 | Teachers' experience with the EHC

All four teachers noticed a change after the intervention in students' coping skills for managing their emotional distress and interpersonal conflicts with peers and linked the acquisition of these emotional regulation skills with the specific techniques taught in the curriculum. One teacher described her observation: "The students before had a hard time problem solving their emotions. Like if they were angry or if they were sad or feeling upset for whatever reason, they had a hard time handling it and putting it into perspective in order to move on in the day...but a lot of them are able to now have some strategy like the mindfulness exercises or just breathing...or different things to help them."

All four teachers reported that the greatest benefit was that this intervention opened a door and created a safe environment for students to communicate their worries, fears, and stresses. One male teacher described his observation of how girls benefited from com-

TABLE 4 Summary of feasibility outcomes

Feasibility	Indicators	Results
Limited efficacy	<ul style="list-style-type: none"> Change in child-reported depression/anxiety in children with elevated symptoms Change in teacher-reported depression/anxiety in children with elevated symptoms 	<ul style="list-style-type: none"> Significant decreases in child-reported depression and anxiety symptoms in at-risk children Significant decreases in teacher-reported depression in at-risk children
Acceptability	<ul style="list-style-type: none"> Child satisfaction Teacher's experiences with the EHC 	<ul style="list-style-type: none"> 89.7% children enjoyed the EHC Four teachers observed that children acquired coping skills to manage their emotional distress after attending the EHC, and perceived the coordinated nursing care delivery approach empowered these children
Implementation	<ul style="list-style-type: none"> Participation Retention Fidelity 	<ul style="list-style-type: none"> 98% children completed a pre-intervention questionnaire 94% children completed both pre- and post-questionnaires Coordinated nursing team maintained 99.13% fidelity rate

municating with the Nursing Professors: "...I know it's hard for them [girls] to communicate with me...and they do need somebody to talk to and I feel that sometimes, myself, coming from a Hispanic family can be a little more conservative and I think having someone to talk to about those issues, somebody [Nursing Professors] that's in the medical field, somebody that's not going to judge, somebody that's going to listen to them...they'll talk to somebody and they'll come back and I don't ask them, but I do notice that they'll come back with a different language...their body language...they feel more comfortable..."

The teachers endorsed the specific content of the intervention and noted that the delivery model was acceptable; overall, they were positive about how the curriculum was designed and implemented. They emphasized that the cultural adaptation of the intervention made it more relevant for the students. One teacher stated: "It is very important culturally; it is important to see that aspect, to make connections to it." All the teachers embraced the coordinated team approach to delivering the curriculum: "Everybody there brought in so much energy. They came in really prepared... I think the kids were feeding off that energy and feeding off that the presenters had that emotion they cared for what they were doing..." One female teacher pointed out: "He [one senior BS nursing student] related to the kids. He's young and had a lot of energy...he always talked a lot about his feelings...I think it is important especially with boys because socially, it's not accepted for boys to cry or express their emotions..." The male teacher echoed his colleague's observation by commenting on the Hispanic cultural perspective of feelings: "...coming from a Hispanic background it is difficult to talk about emotions and particularly, being male is even more difficult."

3.3 | Implementation of the emotional health curriculum

With two children dropping out, 100 remained in the curriculum and were included in the final analysis. A total of 98 children filled out the baseline questionnaire and 94 children completed both pre- and post-questionnaires. With the use of the Intervention Fidelity Checklist, we found a consistency rate of 99.13%.

4 | DISCUSSION

The primary aim of this study was to examine the limited efficacy, acceptability, and implementation of using the EHC with 3rd and 4th grade children in a dual-immersion Spanish-English school (see Table 4). The key results revealed that following the EHC child-reported depression and anxiety symptoms and teacher-reported depression symptoms were reduced in at-risk children with the effect size ranging from medium to large. These findings with 3rd and 4th grade children were similar to other studies with older at-risk children who significantly benefited from a school-based CBT preventive program (Cardemil, Reivich, Beevers, Seligman, & James, 2007; Iizuka, Barrett, Gillies, Cook, & Miller, 2014; Mazurek Melnyk, Kelly, & Lusk, 2014). Together, these findings suggest that school-based interventions using cognitive behavioral theory had a positive effect on children who demonstrated elevated internalizing symptoms.

There were no significant changes in normal children in terms of both child-reported and teacher-reported depression and anxiety. Prior evidence showed that a school-based mindfulness curriculum generated beneficial effects on self-regulation and prosociality (Schonert-Reichl et al., 2015; Weijer-Bergsma et al., 2014). Broader measures of emotional and social adjustment data should be included in future studies to further investigate the impact of the EHC on coping and psychosocial competencies in addition to depression and anxiety symptoms.

Our results indicate that the EHC was well accepted by both children and teachers. The majority of children were satisfied with the EHC. Teachers observed that children openly communicated their emotional distress to them, applied deep breathing and mindfulness exercises to manage their test anxiety, and practiced ways to solve playground social conflicts. Teacher observations were congruent with children self-reported experiences that the skills taught as part of the EHC were useful. Furthermore, teachers highlighted that it was valuable to have the coordinated nursing team deliver this curriculum because often children sought guidance from the school nurses to assist them with their emotional challenges.

The high participation and retention rates suggest that the research team was able to successfully implement the EHC with the 3rd and

4th grade children in this disadvantaged community. The strong fidelity rate implies that the coordinated nursing team consistently implemented the core elements of the curriculum.

4.1 | Limitations

There are four limitations in this study. First, a pre-experimental design without an attention control group was used. We thus cannot conclude that it is the EHC that reduced the depression and/or anxiety symptoms. Second, depression and anxiety outcomes without psychosocial competency data were collected. It is unclear how the EHC will impact developmental outcomes (e.g., cognitive, social, academic outcomes) in at-risk and normal children, respectively. Third, no follow-up data were collected. Without longitudinal data, it is unknown how long the decreased depression and anxiety in children with clinically elevated symptoms would be sustainable. Fourth, mental health problems were not evaluated by clinicians. Clinical diagnosis would increase the validity of the results. A future study is needed to replicate the results with a longitudinal randomized clinical trial (RCT). This RCT should include an attention control with cognitive, emotional, social, behavioral, and academic outcomes from multiple perspectives including clinicians, children, teachers, and parents.

5 | CONCLUSIONS

The EHC is an innovative classroom-based curriculum which integrated CBT, mindfulness, Hispanic cultural elements, and a coordinated nursing team delivery approach. Our study provides preliminary evidence supporting that the EHC has potential to improve depression and anxiety symptoms in primarily Hispanic children with elevated internalizing problems. Children and teachers were satisfied with the content of the curriculum and responded well to the coordinated nursing team delivery approach. Our study suggests that it is essential to implement culturally sensitive interventions in dual immersion Spanish–English elementary schools where at-risk children would benefit most. Future research with a more rigorous design is needed to demonstrate that the EHC is a cost-effective early prevention and intervention model to address mental health disparities in underserved Hispanic communities.

The clinical implications are primarily for school nurses who are champions for early mental health programs. The culturally sensitive EHC was developed through collaborating with the school personnel including administrators, teachers, and nurses. The implementation of the EHC was thus well received by the students. School nurses should actively engage school administrators and teachers when developing a mental health prevention or intervention program. Furthermore, the study suggests that the delivery of the EHC in a classroom format to all students and supported by the school nurse provides a feasible approach to reducing mental health stigma and promoting a safe school environment to openly communicate emotional distress. The EHC is thus aligned with the new school health model entitled “Whole School, Whole Community and Whole Child (WSCC)” (Lewallen, Hunt, Potts-Datema, Zaza, & Giles, 2015). The WSCC

model emphasizes that it is essential for school nurses to build a healthy social and emotional school environment to foster children's whole development and academic success (Lewallen et al., 2015).

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

ACKNOWLEDGEMENTS

This study was funded by University of California Irvine (UCI) Institute for Clinical and Translational Sciences Campus-Community Research Pilot Grant (NIH: IU54RR031271), UCI Advance Spirit Grant, and Health Resources and Services Administration Grant (HRSA: D11HP22200). The authors would like to express their gratitude to the administrators and staff of Sol Science and Arts Academy of Santa Ana for their support; the authors thank Nancy Neudorf, FNP, MS, RN and Raquel Campos, WHNP, MS, RN for supervising the nursing students. The authors thank their research assistants for facilitating intervention implementation and collecting data, especially Irene Chen, Mindy Chin, Mindaugas Galins, Fabiola Movius, Felice Ng, and Eduardo Rivera,

ORCID

Yuqing Guo PhD, RN  <http://orcid.org/0000-0002-7637-1276>

REFERENCES

- Achenbach, T. M., Dumenci, L., & Rescorla, L. A. (2003). DSM-oriented and empirically based approaches to constructing scales from the same item pools. *Journal of Clinical Child and Adolescent Psychology: The Official Journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division 53*, 32(3), 328–340. https://doi.org/10.1207/S15374424JCCP3203_02
- Ahlen, J., Lenhard, F., & Ghaderi, A. (2015). Universal prevention for anxiety and depressive symptoms in children: A meta-analysis of randomized and cluster-randomized trials. *The Journal of Primary Prevention*, 36(6), 387–403. <https://doi.org/10.1007/s10935-015-0405-4>
- Alegria, M., Vallas, M., & Pumariega, A. J. (2010). Racial and ethnic disparities in pediatric mental health. *Child and Adolescent Psychiatric Clinics of North America*, 19(4), 759–774. <https://doi.org/10.1016/j.jchc.2010.07.001>
- Anderson, E., & Mayes, L. (2010). Race/ethnicity and internalizing disorders in youth: A review. *Clinical Psychology Review*, 30(3), 338–348.
- Barrett, P., & Turner, C. (2001). Prevention of anxiety symptoms in primary school children: Preliminary results from a universal school-based trial. *The British Journal of Clinical Psychology*, 40(Pt 4), 399–410.
- Bear, L., Finer, R., Guo, S., & Lau, A. S. (2014). Building the gateway to success: An appraisal of progress in reaching underserved families and reducing racial disparities in school-based mental health. *Psychological Services*, 11(4), 388–397. <https://doi.org/10.1037/a0037969>
- Beck, A. T. (2005). The current state of cognitive therapy: A 40-year retrospective. *Archives of General Psychiatry*, 62(9), 953–959. <https://doi.org/10.1001/archpsyc.62.9.953>
- Bennett, K., Manassis, K., Duda, S., Bagnell, A., Bernstein, G. A., Garland, E. J., ... Wilansky, P. (2015). Preventing child and adolescent anxiety disorders: Overview of systematic reviews. *Depression and Anxiety*, 32(12), 909–918. <https://doi.org/10.1002/da.22400>

- Bird, H. R. (1996). Epidemiology of childhood disorders in a cross-cultural context. *Journal of Child Psychology and Psychiatry*, 37(1), 35–49. <https://doi.org/10.1111/j.1469-7610.1996.tb01379.x>
- Bowen, D. J., Kreuter, M., Spring, B., Cofta-Woerpel, L., Linnan, L., Weiner, D., ... Fernandez, M. (2009). How we design feasibility studies. *American Journal of Preventive Medicine*, 36(5), 452–457. <https://doi.org/10.1016/j.amepre.2009.02.002>
- Calear, A. L., & Christensen, H. (2010). Systematic review of school-based prevention and early intervention programs for depression. *Journal of Adolescence*, 33(3), 429–438. <https://doi.org/10.1016/j.adolescence.2009.07.004>
- Cardemil, E. V., Reivich, K. J., Beevers, C. G., Seligman, M. E. P., & James, J. (2007). The prevention of depressive symptoms in low-income, minority children: Two-year follow-up. *Behaviour Research and Therapy*, 45(2), 313–327. <https://doi.org/10.1016/j.brat.2006.03.010>
- Castro, F. G., Barrera, M., & Holleran Steiker, L. K. (2010). Issues and challenges in the design of culturally adapted evidence-based interventions. *Annual Review of Clinical Psychology*, 6, 213–239. <https://doi.org/10.1146/annurev-clinpsy-033109-132032>
- Cerdá, M., Bordelois, P. M., Keyes, K. M., Galea, S., Koenen, K. C., & Pardini, D. (2013). Cumulative and recent psychiatric symptoms as predictors of substance use onset: Does timing matter? *Addiction*, 108(12), 2119–2128. <https://doi.org/10.1111/add.12323>
- Cummings, J. R., Case, B. G., Ji, X., Chae, D. H., & Druss, B. G. (2014). Racial/ethnic differences in perceived reasons for mental health treatment in US adolescents with major depression. *Journal of the American Academy of Child and Adolescent Psychiatry*, 53(9), 980–990. <https://doi.org/10.1016/j.jaac.2014.05.016>
- Cummings, J. R., Ponce, N. A., & Mays, V. M. (2010). Comparing racial/ethnic differences in mental health service use among high-need subpopulations across clinical and school-based settings. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine*, 46(6), 603–606. <https://doi.org/10.1016/j.jadohealth.2009.11.221>
- Dye, J. F., Schatz, I. M., Rosenberg, B. A., & Coleman, S. T. (2000). Constant comparison method: A kaleidoscope of data. *The Qualitative Report*, 4(1), 1–10. Retrieved from <http://www.nova.edu/ssss/QR/QR4-1/dye.html>.
- Ebesutani, C., Reise, S. P., Chorpita, B. F., Ale, C., Regan, J., Young, J., ... Weisz, J. R. (2012). The Revised Child Anxiety and Depression Scale-Short Version: Scale reduction via exploratory bifactor modeling of the broad anxiety factor. *Psychological Assessment*, 24(4), 833–845. <https://doi.org/10.1037/a0027283>
- Forman-Hoffman, V. L., McClure, E., McKeeman, J., Wood, C. T., Middleton, J. C., Skinner, A. C., Perrin, E. M., & Viswanathan, M. (2016). Screening for major depressive disorder among children and adolescents: A systematic review for the U.S. Preventive Services Task Force. *Evidence Synthesis No. 116. AHRQ Publication No. 13-05192-EF-1*. Rockville, MD: Agency for Healthcare Research and Quality.
- Fujii, C., Renno, P., McLeod, B., Lin, C. E., Decker, K., Zielinski, K., & Wood, J. J. (2013). Intensive cognitive behavioral therapy for anxiety disorders in school-aged children with autism: A preliminary comparison with treatment-as-usual. *School Mental Health*, 5, 25–37.
- Gillham, J. E., Reivich, K. J., Freres, D. R., Chaplin, T. M., Shatté, A. J., Samuels, B., ... Seligman, M. E. P. (2007). School-based prevention of depressive symptoms: A randomized controlled study of the effectiveness and specificity of the Penn Resiliency Program. *Journal of Consulting and Clinical Psychology*, 75(1), 9–19. <https://doi.org/10.1037/0022-006X.75.1.9>
- Herzig-Anderson, K., Colognori, D., Fox, J. K., Stewart, C. E., & Masia Warner, C. (2012). School-based anxiety treatments for children and adolescents. *Child and Adolescent Psychiatric Clinics of North America*, 21(3), 655–668. <https://doi.org/10.1016/j.chc.2012.05.006>
- Hughes, A. A., Lourea-Waddell, B., & Kendall, P. C. (2008). Somatic complaints in children with anxiety disorders and their unique prediction of poorer academic performance. *Child Psychiatry and Human Development*, 39(2), 211–220. <https://doi.org/10.1007/s10578-007-0082-5>
- Iizuka, C. A., Barrett, P. M., Gillies, R., Cook, C. R., & Miller, D. (2014). The FRIENDS emotional health program for minority groups at risk. *The Journal of School Health*, 84(2), 124–132. <https://doi.org/10.1111/josh.12127>
- Joiner, T. E., Perez, M., Wagner, K. D., Berenson, A., & Marquina, G. S. (2001). On fatalism, pessimism, and depressive symptoms among Mexican-American and other adolescents attending an obstetrics-gynecology clinic. *Behaviour Research and Therapy*, 39(8), 887–896.
- Juszcak, L., Melinkovich, P., & Kaplan, D. (2003). Use of health and mental health services by adolescents across multiple delivery sites. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine*, 32(6 Suppl), 108–118.
- Keeton, V., Soleimanpour, S., & Brindis, C. D. (2012). School-based health centers in an era of health care reform: Building on history. *Current Problems in Pediatric and Adolescent Health Care*, 42(6), 132–156. <https://doi.org/10.1016/j.cppeds.2012.03.002>
- Kendall, P. C., & Hedtke, K. (2006). *Coping cat workbook* (2nd ed.). Ardmore, PA: Workbook Publishing.
- Lewallen, T. C., Hunt, H., Potts-Datema, W., Zaza, S., & Giles, W. (2015). The whole school, whole community, whole child model: A new approach for improving educational attainment and healthy development for students. *Journal of School Health*, 85(11), 729–739. <https://doi.org/10.1111/josh.12310>
- Macklem, G. L. (2014). *Preventive mental health at school: Evidence-based services for students*. New York, NY: Springer-Verlag New York Inc.
- Mason-Jones, A. J., Crisp, C., Momberg, M., Koech, J., De Koker, P., & Mathews, C. (2012). A systematic review of the role of school-based healthcare in adolescent sexual, reproductive, and mental health. *Systematic Reviews*, 1:49. <https://doi.org/10.1186/2046-4053-1-49>. <https://systematicreviewsjournal.biomedcentral.com/about>
- Mazurek Melnyk, B., Kelly, S., & Lusk, P. (2014). Outcomes and feasibility of a manualized cognitive-behavioral skills building intervention: Group COPE for depressed and anxious adolescents in school settings. *Journal of Child and Adolescent Psychiatric Nursing*, 27(1), 3–13. <https://doi.org/10.1111/jcap.12058>
- Mendelson, T., Greenberg, M. T., Dariotis, J. K., Gould, L. F., Rhoades, B. L., & Leaf, P. J. (2010). Feasibility and preliminary outcomes of a school-based mindfulness intervention for urban youth. *Journal of Abnormal Child Psychology*, 38(7), 985–994. <https://doi.org/10.1007/s10802-010-9418-x>
- Merikangas, K. R., He, J. P., Brody, D., Fisher, P. W., Bourdon, K., & Koretz, D. S. (2010). Prevalence and treatment of mental disorders among US children in the 2001–2004 NHANES. *Pediatrics*, 125(1), 75–81. <https://doi.org/10.1542/peds.2008-2598>
- Neil, A. L., & Christensen, H. (2009). Efficacy and effectiveness of school-based prevention and early intervention programs for anxiety. *Clinical Psychology Review*, 29(3), 208–215. <https://doi.org/10.1016/j.cpr.2009.01.002>
- Patton, G. C., Coffey, C., Romaniuk, H., Mackinnon, A., Carlin, J. B., Degenhardt, L., ... Moran, P. (2014). The prognosis of common mental disorders in adolescents: A 14-year prospective cohort study. *Lancet*, 383(9926), 1404–1411. [https://doi.org/10.1016/S0140-6736\(13\)62116-9](https://doi.org/10.1016/S0140-6736(13)62116-9)
- Rey, Y., Marin, C. E., & Silverman, W. K. (2011). Failures in cognitive-behavior therapy for children. *Journal of Clinical Psychology*, 67(11), 1140–1150. <https://doi.org/10.1002/jclp.20848>

- Saldaña J. (2009). *The Coding Manual for Qualitative Researchers*. Thousand Oaks, CA: Sage.
- Saluja, G., Iachan, R., Scheidt, P. C., Overpeck, M. D., Sun, W., & Giedd, J. N. (2004). Prevalence of and risk factors for depressive symptoms among young adolescents. *Archives of Pediatrics & Adolescent Medicine*, 158(8), 760–765. <https://doi.org/10.1001/archpedi.158.8.760>
- Schonert-Reichl, K. A., Oberle, E., Lawlor, M. S., Abbott, D., Thomson, K., Oberlander, T. F., & Diamond, A. (2015). Enhancing cognitive and social-emotional development through a simple-to-administer mindfulness-based school program for elementary school children: A randomized controlled trial. *Developmental Psychology*, 51(1), 52–66. <https://doi.org/10.1037/a0038454>
- Siegel, J. M., Yancey, A. K., Aneshensel, C. S., & Schuler, R. (1999). Body image, perceived pubertal timing, and adolescent mental health. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine*, 25(2), 155–165.
- Stewart, S. M., Simmons, A., & Habibpour, E. (2012). Treatment of culturally diverse children and adolescents with depression. *Journal of Child and Adolescent Psychopharmacology*, 22(1), 72–79. <https://doi.org/10.1089/cap.2011.0051>
- Toppelberg, C. O., Hollinshead, M. O., Collins, B. A., & Nieto-Castañon, A. (2013). Cross-sectional study of unmet mental health need in 5- to 7-Year Old Latino children in the United States: Do teachers and parents make a difference in service utilization? *School Mental Health*, 5(2), 59–69. <https://doi.org/10.1007/s12310-012-9089-6>
- Uebelacker, L. A., Marootian, B. A., Pirraglia, P. A., Primack, J., Tigue, P. M., Haggarty, R., ... Miller, I. W. (2012). Barriers and facilitators of treatment for depression in a Latino community: A focus group study. *Community Mental Health Journal*, 48(1), 114–126. <https://doi.org/10.1007/s10597-011-9388-7>
- Weijer-Bergsma, E., van de, Langenberg, G., Brandsma, R., Oort, F. J., & Bögels, S. M. (2014). The effectiveness of a school-based mindfulness training as a program to prevent stress in elementary school children. *Mindfulness*, 5(3), 238–248. <https://doi.org/10.1007/s12671-012-0171-9>
- Werner-Seidler, A., Perry, Y., Calear, A. L., Newby, J. M., & Christensen, H. (2017). School-based depression and anxiety prevention programs for young people: A systematic review and meta-analysis. *Clinical Psychology Review*, 51, 30–47. <https://doi.org/10.1016/j.cpr.2016.10.005>
- World Health Organization. (2008). Mental health gap action programme: Scaling up care for mental, neurological and substance use disorders. Geneva: World Health Organization Press. Retrieved from http://www.who.int/mental_health/mhgap_final_english.pdf

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Guo Y, Rousseau J, Renno P, et al. Feasibility of an emotional health curriculum for elementary school students in an underserved Hispanic community. *J Child Adolesc Psychiatr Nurs*. 2018;1–9. <https://doi.org/10.1111/jcap.12185>