

UC San Diego

UC San Diego Previously Published Works

Title

Assessing the motivational effects of ethanol in mice using a discrete-trial current-intensity intracranial self-stimulation procedure

Permalink

<https://escholarship.org/uc/item/7156q7bh>

Authors

Barkley-Levenson, Amanda M
Der-Avakian, Andre
Palmer, Abraham A

Publication Date

2020-02-01

DOI

10.1016/j.drugalcdep.2019.107806

Peer reviewed

Assessing the motivational effects of ethanol in mice using a discrete-trial current-intensity intracranial
self-stimulation procedure

Amanda M. Barkley-Levenson¹, Andre Der-Avakian¹, Abraham A. Palmer^{1,2}

1. Department of Psychiatry, University of California San Diego, La Jolla, CA 92093, USA
2. Institute for Genomic Medicine, University of California San Diego, La Jolla, CA 92093, USA

Corresponding author information: **Corresponding Author:** Abraham A. Palmer; Department of
Psychiatry, University of California San Diego, La Jolla, CA 92093, USA. Voice 858-534-2093. Fax 858-
534-7653. aap@ucsd.edu

Funding support: This work was supported by NIH-NIAAA grant R01 AA026281. AMB-L was
supported by NIH-NIAAA grant F32 AA025515.

Abstract

Background: Alcohol (ethanol) produces both rewarding and aversive effects, and sensitivity to these effects is associated with risk for an alcohol use disorder (**AUD**). Measurement of these motivational effects in animal models is an important but challenging aspect of preclinical research into the neurobiology of AUD. Here, we evaluated whether a discrete-trial current-intensity intracranial self-stimulation (**ICSS**) procedure can be used to assess both reward-enhancing and aversive responses to ethanol in mice.

Methods: Male and female C57BL/6J mice were surgically implanted with bipolar stimulating electrodes targeting the medial forebrain bundle and trained on a discrete-trial current-intensity ICSS procedure. Mice were tested for changes in response thresholds after various doses of ethanol (0.5 g/kg-1.75 g/kg; n=5-7 per dose), using a Latin square design.

Results: A 1 g/kg dose of ethanol produced a significant reward-enhancement (i.e., lowered response thresholds), whereas a 1.75 g/kg dose produced an aversive effect (elevated response thresholds). Ethanol doses from 1-1.75 g/kg increased response latencies as compared to saline treatment.

Conclusions: The discrete-trial current-intensity ICSS procedure is an effective assay for measuring both reward-enhancing responses to ethanol as well as aversive responses in the same animal. This should prove to be a useful tool for assessing the effects of experimental manipulations on the motivational effects of ethanol in mice.

Key words: Intracranial self-stimulation, brain stimulation reward, ethanol, mice, reward, aversion

1. Introduction

Alcohol (ethanol) is a compound stimulus with both rewarding and aversive effects (Verendeev and Riley 2013), and motivation to consume ethanol is influenced by sensitivity to these effects. In humans, the rewarding effects of ethanol include euphoria, stimulation, and drug-liking; in contrast, aversive effects include sedation, dysphoria, and drug-disliking (Fridberg et al. 2017; Rueger and King 2013). Both greater sensitivity to ethanol's rewarding effects and reduced sensitivity to ethanol's aversive effects have been associated with a greater risk of an alcohol use disorder (AUD) (e.g. King et al. 2014; Schuckit et al. 2007). Understanding the neurobiological basis of ethanol's motivational effects may therefore provide critical insight into excessive ethanol consumption and AUD. Furthermore, drugs that are intended to treat AUD could potentially act via blocking the rewarding effects of ethanol or by enhancing the aversive effects of ethanol.

The intracranial self-stimulation (ICSS; sometimes called brain stimulation reward) procedure provides a direct measure of brain reward function, where responding is reinforced by electrical stimulation of the mesolimbic reward circuitry. Response thresholds can be established by determining the level of stimulation that maintains operant responding. Drugs of abuse can enhance the rewarding properties of this stimulation, partly through dopaminergic activation (Carlezon and Chartoff 2007). This reward-enhancement effect is characterized by a lowering of ICSS response thresholds, wherein less stimulation is required to maintain responding. In contrast, increases in ICSS response thresholds are interpreted as an aversive or anhedonic response because a greater stimulation intensity is required to maintain the same level of responding. ICSS has been widely used to study the reward-enhancing properties of drugs of abuse, with nearly all classes of abused drugs showing enhancement of reward in ICSS procedures (for review, see Kenny et al. 2018; Negus and Miller 2014). However, ICSS has not been routinely used with ethanol in rodents, and especially has been underutilized in mice. To the best of our knowledge, the only previous efforts in mice have also exclusively used rate-frequency procedures (Fish et al. 2010, 2012) that measure drug-induced increases in response rates for low stimulation

frequencies. These procedures can be more sensitive to locomotor changes (both stimulation and sedation), and therefore may present challenges for interpreting results across ethanol doses. Here, we demonstrate that a discrete-trial current-intensity procedure to measure ICSS response threshold current can be used to assess both reward-enhancing and aversive responses to ethanol in the same animal across a range of doses.

2. Methods

2.1 Animals and husbandry

Mice were tested in two cohorts approximately six months apart. Adult male and female C57BL/6J mice were bred in house (cohort 1) or purchased from Jackson Laboratories (cohort 2). Mice were housed 2-5 per cage and had *ad libitum* access to food (Envigo 8604, Indianapolis, IN) and water. Mice were housed on a 12h:12h reverse light-dark cycle with lights off at 07:00h, and all testing occurred during the dark phase. All procedures were conducted at the University of California San Diego (UCSD) and were approved by the UCSD Institutional Animal Care and Use Committees and were conducted in accordance with the NIH Guidelines for the Care and Use of Laboratory Animals.

2.2 Surgical procedures

ICSS surgical procedures have been described previously (Stoker et al. 2008). Briefly, anesthetized mice were surgically implanted with a bipolar stimulating electrode (Plastics One Inc., Roanoke, VA) aimed at the medial forebrain bundle (A/P: -1.58, M/L: ± 1.0 , D/V: -5.3; Paxinos and Franklin 2001). Left/right position of the electrode was counterbalanced across animals. Electrodes were secured to the skull with dental acrylic and two stainless steel anchor screws. All animals were allowed to recover for one week prior to the start of behavioral training. Electrode placement was confirmed behaviorally by acquisition of the ICSS procedure described below, as this has been shown previously to correlate with electrode positioning in the medial forebrain bundle (Markou and Frank 1987; Matthews et al. 1996).

2.3 ICSS apparatus

ICSS training and testing were conducted in eight Plexiglas operant chambers ($30.5 \times 24 \times 27$ cm; Med Associates, St. Albans, VT). Each operant chamber was enclosed within a light and sound-attenuated chamber ($40 \times 60 \times 63.5$ cm; San Diego Instruments, San Diego, CA). Intracranial stimulation was delivered by constant current stimulators (Stimtek model 1200c, San Diego Instruments, San Diego, CA). The mice were connected to the stimulation circuit through flexible bipolar leads (Plastics One, Roanoke, VA) attached to gold-contact swivel commutators (model SL2C, Plastics One, Roanoke, VA) mounted above the operant chamber. The operant response required by the subjects was a quarter turn of a wheel manipulandum (5.5 cm diameter, 4 cm width) that extended 1.5 cm from one wall of the operant chamber.

2.4 ICSS procedure

The ICSS procedure was adapted from a discrete-trial current-intensity threshold procedure used in rats (Kornetsky and Esposito 1979; Markou and Koob 1992) and has been described previously for mice in conjunction with other drugs (Gill et al. 2004; Stoker et al. 2008; Stoker and Markou 2011). Briefly, mice were first trained to turn a wheel manipulandum on a fixed-ratio 1 (**FR1**) schedule of reinforcement to receive electrical stimulation. The frequency of the electrical stimulation was fixed (100 hz), and the current was adjusted for each animal to maintain responding (120-180 μ A). After acquisition of the FR1 schedule (i.e., two consecutive test sessions with 200 reinforcers earned in less than 10 min), animals were trained on the discrete-trial current threshold procedure. Each trial was initiated by the delivery of a noncontingent electrical stimulation followed by a 7.6 s response window within which the subject could make a response to receive a second contingent stimulation. The electrical current was varied across descending and ascending series of trials to determine the minimal current level that maintained responding (i.e., response threshold). Mean response latency (time between trial start and response for the contingent stimulation) was also recorded. Of the 26 mice that underwent electrode

implantation, three failed to acquire the ICSS procedure and eleven were lost due to technical issues with the head cap or electrode prior to starting the Latin square (supplemental Table S1). Twelve mice completed all training and were tested on the Latin square as described below.

2.5 Experimental design

Following training, mice received one test session per day. Mice were given at least 10 sessions to establish stable baseline response thresholds before the start of ethanol testing. An animal was considered to have a stable initial baseline when there was less than 10% variation in thresholds over five consecutive sessions. Mice were tested on a series of ethanol doses using a Latin square design. Because of the counterbalanced design, mice were not habituated to injections. On ethanol test days, mice were weighed and given an i.p. injection of the assigned ethanol dose. They were returned to the home cage for 5 min and then tested in the discrete-trial current-intensity ICSS procedure, which lasted approximately 20-35 min. Stable baseline thresholds were reestablished between each ethanol test session (less than 10% variation in threshold over at least three consecutive sessions; supplemental figure S1).

2.6 Drugs

Ethanol solution (Deacon Laboratories Inc., King of Prussia, PA) was made fresh each test day (20% v/v in saline) and administered i.p. at doses of 0 g/kg [saline], 0.5 g/kg, 0.75 g/kg, 1 g/kg, 1.5 g/kg, and 1.75 g/kg. This dose range was based on the effective dose range seen in previous studies (Fish et al. 2010) and pilot experiments in our laboratory that indicated higher doses produced significant locomotor sedation that interfered with task performance.

2.7 Analyses

All analyses were conducted using SPSS (version 25; IBM, Armonk, NY). Response threshold on each ethanol test day was converted to a percent of baseline threshold for each individual animal. Because we were specifically interested in which ethanol doses would produce a significant change compared to

control (saline), we used planned comparisons (two-sample t-tests) of each ethanol dose vs. saline for both thresholds and response latencies. Unpaired t-tests were used because not all animals completed each stage of the Latin square due to technical issues (e.g. loss of head cap, electrode chewed by cagemates; see Supplemental Table S1) and dose groups therefore did not necessarily consist of the same animals. Analysis of Variance (ANOVA) was used first to determine whether there were significant interactions with sex or cohort and ethanol dose that would make collapsing on these variables statistically inappropriate. We analyzed the data by 3-way ANOVA with ethanol dose, sex, and cohort as between-subjects factors for both response threshold and mean response latency. The threshold for one mouse at the 1 g/kg dose was found to be a statistical outlier (>2 standard deviations higher than the mean) and this animal's 1g/kg data were excluded from analyses. Final group sizes were $n=5$ for 1.5 g/kg, $n=7$ for 1 g/kg, and $n=6$ for all other doses. The level of significance was 0.05.

3. Results

Figure 1 shows the ethanol dose-response curve for ICSS response thresholds. Analysis of response thresholds showed a main effect of ethanol dose ($F_{5,18}=4.74$, $p=0.006$). There were no main effects of sex ($F_{1,18}=1.71$, $p=0.207$) or cohort ($F_{1,18}=0.31$, $p=0.585$), and no significant interactions (two-way interactions: $F_{1-3,18}\leq 1.65$, $p\geq 0.213$; three-way interaction: $F_{1,18}=0.26$, $p=0.615$). Planned comparisons of response threshold for each ethanol dose to saline showed that the 1 g/kg dose significantly reduced thresholds compared to saline (two-tailed: $t_{11}=2.27$, $p=0.045$). This is consistent with a reward-enhancing effect. In contrast, the 1.75 g/kg dose significantly increased thresholds compared to baseline (two-tailed: $t_{10}=-2.48$, $p=0.033$), which is consistent with an aversive effect. No other dose groups were found to differ significantly from saline ($p\geq 0.189$ for all).

Figure 2 shows response latencies at each dose. There was a main effect of dose on response latencies ($F_{5,18}=9.53$, $p<0.001$). Planned comparisons showed that the 1 g/kg ($t_{11}=-2.74$, $p=0.019$), 1.5 g/kg ($t_9=-2.94$, $p=0.016$), and 1.75 g/kg ($t_{10}=-3.81$, $p=0.003$) doses of ethanol significantly increased latencies compared to saline. No other doses significantly differed from saline ($p\geq 0.209$ for all), and there

were no other main effects ($F_{1,18} \leq 1.69$, $p \geq 0.21$) or significant interactions (two-way interactions: $F_{1,3,18} \leq 1.85$, $p \geq 0.174$; three-way interaction: $F_{1,18} = 3.60$, $p = 0.074$).

4. Discussion

This experiment provides the first data showing that a discrete-trial current-intensity ICSS procedure can assess both reward-enhancing and aversive effects of ethanol in mice. We demonstrated that a 1 g/kg dose of ethanol produces a modest but significant decrease in response thresholds (approximately 15%), indicating a reward-enhancing effect. This effect is similar in magnitude to what has been reported before for ethanol (Fish et al. 2010) and diazepam (Straub et al. 2010) when tested in a rate-frequency ICSS procedure in mice. Treatment with a 1.75 g/kg dose produced a significant increase in response thresholds (approximately 40%), which is consistent with an aversive response. We observed longer response latencies at the 1 g/kg, 1.5 g/kg, and 1.75 g/kg doses, which may indicate locomotor sedation or the presence of competing behaviors. However, we saw opposite effects on response thresholds at two ethanol doses that both increased response latency (1 g/kg and 1.75 g/kg), indicating that ethanol's effects on performance-dependent variables such as response latency are dissociable from its motivational effects.

We are only aware of two prior mouse studies that involved ethanol and ICSS. One previous study using a rate-frequency procedure in C57BL/6J and DBA/2J mice found that a 0.6 g/kg dose had a reward-enhancing effect at 0-15 min post-treatment in C57BL/6J mice but had no effect at later time points (Fish et al. 2010). Doses from 0.6 g/kg-1.7 g/kg were found to be reward-enhancing in DBA/2J mice for up to 30 min post-treatment., but also produced significant increases in maximum response rate. In a different study using mice that had been selectively bred for high (FAST) or low (SLOW) locomotor response to ethanol, doses up to 2.4 g/kg were found to be reward-enhancing for the FAST mice, whereas there were no doses that produced significant threshold changes in the SLOW mice (Fish et al. 2012). The effective dose range for assessing reward-enhancing vs. aversive effects is likely genotype-specific and

therefore the doses identified in our study should be re-established when working with non-C57BL/6J mice.

There are also several procedural variations between our study and the few previous published reports on ethanol ICSS in mice that should be noted. In our experiment, we used an i.p. injection as the route of administration, whereas others have used intragastric administration via oral gavage (Fish et al. 2010, 2012). This could potentially explain some of the differences we noted in the dose-response and time course, although one previous study in rats found no difference between i.p. and intragastric administration, with both routes of administration producing no effect on thresholds (Schaefer and Michael 1987). Our study also used the discrete-trial current-intensity procedure instead of a rate-frequency procedure. Both procedures are intended to assess the same underlying process, but the discrete-trial current-intensity procedure may be more resistant to drug-induced changes in rate of responding and nonspecific performance deficits (Markou and Koob 1992). In the present experiment we also included both male and female mice; however, final group sizes ($n=2-3/\text{sex}/\text{dose}$) were not sufficient to determine if there are sex differences.

Previous work with rats has shown mixed effects of ethanol on ICSS, with some studies showing decreases in thresholds (Lewis and June 1994; Moolten and Kornetsky 1990) and some studies showing no change or increases in thresholds (Carlson and Lydic 1976, Schaefer and Michael 1987). Previous mouse studies have shown decreases in thresholds or no effect depending on the genotype tested, although some doses do appear to produce non-statistically significant increases in thresholds (Fish et al. 2010, 2012). Our findings appear to be unique in identifying different doses that can either decrease (1.0 g/kg) or increase (1.75 g/kg) response thresholds. This suggests that our procedure may be useful for evaluating both reward-enhancing and aversive sensitivity to ethanol in the same animal. It should also be noted that the time course of ethanol effects on ICSS response threshold may prove to be an important variable. In the present study, response thresholds were averaged over the entire session which covered roughly 25-40 min. However, it is possible that the strength and/or direction of ethanol's effects on

response thresholds may vary within this time period. Future experiments will be needed to determine whether this is the ideal time window for observing these effects.

In conclusion, we have presented the first evidence that ethanol modulates response thresholds in a discrete-trial current-intensity ICSS procedure in a dose-dependent manner in mice. This procedure has the potential for broad application in ethanol research in order to better understand the neurobiology of ethanol's motivational effects.

References

- Carlezon Jr, W. A., Chartoff, E. H., 2007. Intracranial self-stimulation (ICSS) in rodents to study the neurobiology of motivation. *Nature Protocols* 2(11), 2987–2995. <https://doi.org/10.1038/nprot.2007.441>
- Carlson, R. H., Lydic, R., 1976. The effects of ethanol upon threshold and response rate for self-stimulation. *Psychopharmacology* 50(1), 61–64.
- Fish, E. W., Riday, T. T., McGuigan, M. M., Faccidomo, S., Hodge, C. W., Malanga, C. J., 2010. Alcohol, cocaine, and brain stimulation-reward in C57Bl6/J and DBA2/J mice. *Alcoholism, Clinical and Experimental Research* 34(1), 81–89. <https://doi.org/10.1111/j.1530-0277.2009.01069.x>
- Fish, E. W., Robinson, J. E., Krouse, M. C., Hodge, C. W., Reed, C., Phillips, T. J., Malanga, C. J., 2012. Intracranial self-stimulation in FAST and SLOW mice: Effects of alcohol and cocaine. *Psychopharmacology* 220(4), 719–730. <https://doi.org/10.1007/s00213-011-2523-x>
- Fridberg, D. J., Rueger, S. Y., Smith, P., King, A. C., 2017. Association of Anticipated and Laboratory-Derived Alcohol Stimulation, Sedation, and Reward. *Alcoholism, Clinical and Experimental Research* 41(7), 1361–1369. <https://doi.org/10.1111/acer.13415>
- Gill, B. M., Knapp, C. M., Kornetsky, C., 2004. The effects of cocaine on the rate independent brain stimulation reward threshold in the mouse. *Pharmacology Biochemistry and Behavior* 79(1), 165–170. <https://doi.org/10.1016/j.pbb.2004.07.001>
- Kenny, P. J., Hoyer, D., Koob, G. F., 2018. Animal Models of Addiction and Neuropsychiatric Disorders and Their Role in Drug Discovery: Honoring the Legacy of Athina Markou. *Biological Psychiatry* 83(11), 940–946. <https://doi.org/10.1016/j.biopsych.2018.02.009>
- King, A. C., McNamara, P. J., Hasin, D. S., Cao, D., 2014. Alcohol challenge responses predict future alcohol use disorder symptoms: A 6-year prospective study. *Biological Psychiatry* 75(10), 798–806. <https://doi.org/10.1016/j.biopsych.2013.08.001>
- Kornetsky, C., Esposito, R. U., 1979. Euphorogenic drugs: Effects on the reward pathways of the brain. *Federation Proceedings* 38(11), 2473–2476.
- Lewis, M. J., June, H. L., 1994. Synergistic effects of ethanol and cocaine on brain stimulation reward. *Journal of the Experimental Analysis of Behavior* 61(2), 223–229. <https://doi.org/10.1901/jeab.1994.61-223>
- Markou, A., Frank, R.A., 1987. The effect of operant and electrode placement on self-stimulation train duration response functions. *Physiology & Behavior*, 41, 303-308.
- Markou, A., Koob, G. F., 1992. Construct validity of a self-stimulation threshold paradigm: Effects of reward and performance manipulations. *Physiology & Behavior* 51(1), 111–119. [https://doi.org/10.1016/0031-9384\(92\)90211-J](https://doi.org/10.1016/0031-9384(92)90211-J)
- Matthews, K., Baldo, B.A., Markou, A., Lown, O., Overstreet, D.H., Koob, G.F., 1996. Rewarding electrical brain stimulation: similar thresholds for flinders sensitive line hypercholinergic and flinders resistant line hypochoinergic rats. *Physiology & Behavior*, 59(6), 1155-1162.
- Moolten, M., Kornetsky, C., 1990. Oral self-administration of ethanol and not experimenter-administered ethanol facilitates rewarding electrical brain stimulation. *Alcohol* 7(3), 221–225.

- Negus, S. S., Miller, L. L., 2014. Intracranial self-stimulation to evaluate abuse potential of drugs. *Pharmacological Reviews* 66(3), 869–917. <https://doi.org/10.1124/pr.112.007419>
- Paxinos, G., Franklin, K.B.J., 2001. *The Mouse Brain in Stereotaxic Coordinates*. 2nd Edition, Academic Press, San Diego.
- Rueger, S. Y., King, A. C., 2013. Validation of the brief Biphasic Alcohol Effects Scale (B-BAES). *Alcoholism, Clinical and Experimental Research* 37(3), 470–476. <https://doi.org/10.1111/j.1530-0277.2012.01941.x>
- Schaefer, G. J., Michael, R. P., 1987. Ethanol and current thresholds for brain self-stimulation in the lateral hypothalamus of the rat. *Alcohol* 4(3), 209–213.
- Schuckit, M. A., Smith, T. L., Danko, G. P., Pierson, J., Hesselbrock, V., Bucholz, K. K., Kramer, J., Kuperman, S., Dietiker, C., Brandon, R., Chan, G., 2007. The Ability of the Self-Rating of the Effects of Alcohol (SRE) Scale to Predict Alcohol-Related Outcomes Five Years Later. *Journal of Studies on Alcohol and Drugs* 68(3), 371–378. <https://doi.org/10.15288/jsad.2007.68.371>
- Stoker, A. K., Markou, A., 2011. Withdrawal from chronic cocaine administration induces deficits in brain reward function in C57BL/6J mice. *Behavioural Brain Research* 223(1), 176–181. <https://doi.org/10.1016/j.bbr.2011.04.042>
- Stoker, A. K., Semenova, S., Markou, A., 2008. Affective and somatic aspects of spontaneous and precipitated nicotine withdrawal in C57BL/6J and BALB/cByJ mice. *Neuropharmacology* 54(8), 1223–1232. <https://doi.org/10.1016/j.neuropharm.2008.03.013>
- Straub, C.J., Carlezon Jr., W.A., Rudolph, U., 2010. Diazepam and cocaine potentiate brain stimulation reward in C57BL/6J mice. *Behavioural Brain Research*, 206(1), 17-20. <https://doi.org/10.1016/j.bbr.2009.08.025>
- Verendeev, A., Riley, A. L., 2013. The role of the aversive effects of drugs in self-administration: Assessing the balance of reward and aversion in drug-taking behavior. *Behavioural Pharmacology* 24(5–6), 363–374. <https://doi.org/10.1097/FBP.0b013e32836413d5>




Figure 1. ICSS response thresholds across ethanol dose. Data are presented as a percent of average baseline thresholds from non-injected control trials (dashed line). The inset shows the average baseline threshold currents for each dose, with a scatterplot of values for individual mice. * indicates statistically significant difference from saline group ($p < 0.05$). $N = 5-7$ per dose.




Figure 2. Mean response latencies across ethanol doses. Mean response latencies across all trials in a session are shown for each dose of ethanol. * indicates statistically significant difference from saline group ($p < 0.05$). $N = 5-7$ per dose.

Table S1. Attrition information for all mice that underwent electrode implantation

<i>Attrition prior to starting Latin square</i>						
	Did not acquire ICSS	Electrode chewed by cagemates			Head cap lost	
Number of mice	3	5			6	
<i>Attrition during Latin square (number of ethanol doses completed)</i>						
	1 dose	2 doses	3 doses	4 doses	5 doses	6 doses
Number of mice	2	3	3	1	1	2




Figure S1. Average baseline response thresholds for each test session across time. The mean three-day response thresholds used as baselines for each test session are shown, with scatterplot showing values for individual mice.