

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

The S/L Branching Ratio of Y \*1

### Permalink

<https://escholarship.org/uc/item/71c38013>

### Authors

Alston, Margaret H

Alvarez, Luis W

Eberhard, Philippe

et al.

### Publication Date

1961-04-01

UNIVERSITY OF  
CALIFORNIA

*Ernest O. Lawrence*

*Radiation  
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California  
Contract No. W-7405-eng-48

THE  $\Sigma/\Lambda$  BRANCHING RATIO OF  $\gamma_1^{*+}$

Margaret H. Alston, Luis W. Alvarez, Philippe Eberhard, Myron L. Good  
William Graziano, Harold K. Ticho, and Stanley G. Wojcicki

April 25, 1961

THE  $\Sigma/\Lambda$  BRANCHING RATIO OF  $Y_1^{*+}$

Margaret H. Alston, Luis W. Alvarez, Philippe Eberhard,<sup>§</sup> Myron L. Good,<sup>\*\*</sup>  
William Graziano, Harold K. Ticho,<sup>††</sup> and Stanley G. Wojcicki

Lawrence Radiation Laboratory and Department of Physics  
University of California, Berkeley, California

April 25, 1961

Recently a  $T = 1$  resonance in the  $\Lambda\pi$  system called  $Y_1^{*+}$  has been observed with a mass of 1385 Mev.<sup>1-6</sup> Two types of resonances have been predicted that might relate this observation to other elementary-particle interactions: (1)  $P_{3/2}$  resonances in the  $\Lambda\pi$  and  $\Sigma\pi$  systems predicted by global symmetry<sup>7,8</sup> corresponding to the  $(3/2, 3/2)$  resonance of the  $\pi N$  system, (2) a spin-1/2  $Y-\pi$  resonance resulting from a bound state in the  $\bar{K}N$  system.<sup>9,10</sup> The position and width of the observed  $Y_1^{*+}$  resonance agree with both theories, but since the spin and parity have not yet been determined, it is impossible at present to distinguish between the two theoretical interpretations.

Global symmetry<sup>11</sup> predicts a theoretical branching ratio  $(Y_1^{*+} \rightarrow \Sigma^0 + \pi^+) / (Y_1^{*+} \rightarrow \Lambda + \pi^+) = 1/4$  for the  $T = 1$  resonance. The phase-space factor  $(P_\Sigma / P_\Lambda)^3 = (126/207)^3 = 0.225$  reduces the expected branching ratio for this process to  $R = (1/4) \times 0.225 \sim 5\%$ . Furthermore, as a consequence of charge independence the rates  $Y_1^{*+} \rightarrow \Sigma^\pm + \pi^0$ ,  $Y_1^{*+} \rightarrow \Sigma^0 + \pi^\pm$ , and  $Y_1^{*0} \rightarrow \Sigma^\pm + \pi^\mp$  are equal. In addition to the  $T = 1$  resonance, a  $T = 2$   $\Sigma-\pi$  resonance with a total energy of 1540 Mev and a half width,  $\Gamma/2$ , of 60 Mev is predicted by global symmetry.<sup>8</sup>

\* Work done under the auspices of the U. S. Atomic Energy Commission.

§ Presently at Laboratoire de Physique Atomique, College de France, Paris, France.

\*\* Presently at University of Wisconsin, Madison, Wisconsin.

†† Presently at the University of California at Los Angeles, Los Angeles, California

The  $\bar{K}$ -N bound-state model suggests values of R considerably larger than 5%. However, when non-zero effective ranges are taken into account<sup>1,2</sup>, R can become quite small, especially if the  $(\Sigma\Lambda)$  parity should be odd.

To investigate these possibilities, we have continued our study of  $K^- - p$  interactions at 1.15 Bev/c in the Lawrence Radiation Laboratory 15-in. hydrogen bubble chamber by studying events in which a  $\Sigma$  is observed. The total cross sections for these interactions are shown in Table I; only statistical errors are indicated. The separation of  $\Sigma^\pm + \pi^\mp + \pi^0$  and  $\Sigma^\pm + \pi^\mp + 2\pi^0$  events was difficult because many of the latter events can also be fitted to the first hypothesis. The numbers given in Table I and in the Dalitz and mass plots below were corrected to account for this ambiguity. The correction factor was estimated by using our  $\Sigma^\pm + \pi^\mp + \pi^+ + \pi^-$  events.

Dalitz plots for the three-body reactions are shown in Fig. 1. The  $Y_1^*$  resonance of mass 1385 Mev should appear as a bunching of events about both horizontal and vertical lines corresponding to  $T_\pi = 282$  Mev. To obtain an upper limit for the branching ratio R, we combined the events into different charge states of the  $\Sigma\pi$  system. All charged  $\Sigma$  were observed; however, in the  $\Sigma^0$  cases only two-thirds of the events were observable because of the neutral decays of the  $\Lambda^0$ . Furthermore, we had estimated that about one-third of the  $\Sigma^0 \pi^+ \pi^-$  events also fitted a  $\Lambda\pi^+ \pi^-$  interpretation and had been included in already published data.<sup>1</sup> Consequently each  $\Sigma^0 \pi^+ \pi^-$  event was given a weight of 2.25. The resultant mass spectra are shown in Fig. 2. In the cases of  $(\Sigma\pi)^+$  and  $(\Sigma\pi)^-$  there appears to be no excess of events in the region of  $M = 1385$  Mev. Using the number of  $(\Lambda\pi^+)$  and  $(\Lambda\pi^-)$  events with  $1355 \text{ Mev} < M_{\Lambda\pi} < 1415 \text{ Mev}$  from reference 1, and assuming that all  $\Sigma\pi$  events in the same regions of Fig. 2 are  $Y_1^*$ , we obtain  $R_{\text{max}} \leq 8\%$ .

This treatment yields an unrealistic upper limit, since there is no evidence of any peaking above background. The results are consistent with  $R = 0$ . The  $\Sigma^\pm + \pi^+ + 2\pi^0$  events possibly misidentified as  $\Sigma^\pm + \pi^\mp + \pi^0$  (or vice versa) do not fall into the mass band used in this analysis, since they yield apparently high masses of the  $\Sigma^\pm \pi^0$  system.

We conclude that the  $\Sigma/\Lambda$  branching ratio  $R$  for the strong decay of the  $T = 1$   $Y_1^*$  is at most a few percent and is consistent with zero. This result agrees with the value of  $R$  obtained by Berge.<sup>3</sup> As indicated above this value of  $R$  does not rule out either the global symmetry or the  $\bar{K}N$  bound-state model of the  $Y_1^*$  resonance. No evidence for the resonance with  $T = 2$  predicted by global symmetry at  $M = 1540$  Mev is observed; however, this wide resonance would be hard to separate from background.

The authors wish to thank the many members of the Bevatron and 15-in. bubble chamber crews and the scanners who made this experiment possible. One of us, Philippe Eberhard, wishes to thank the Philippe Foundation, Inc. and the Commissariat à l'Énergie Atomique for a fellowship.

Table I. Cross sections for the  $\Sigma$  producing interactions at 1.15 Bev/c

<u>Reaction</u>	<u>No. of events (uncorrected)</u>	<u>Cross sections (mb)</u>
$K^- + p \rightarrow \Sigma^- + \pi^+$	87	$1.40 \pm 0.16$
$\rightarrow \Sigma^+ + \pi^-$	84	$1.34 \pm 0.18$
$\rightarrow \Sigma^+ + \pi^- + \pi^0$	57	$0.97 \pm 0.16$
$\rightarrow \Sigma^- + \pi^+ + \pi^0$	54	$0.83 \pm 0.20$
$\rightarrow \Sigma^0 + \pi^+ + \pi^-$	27	$0.97 \pm 0.20$
$\rightarrow \Sigma^+ + \pi^- + \pi^0 + \pi^0$	13	$0.18 \pm 0.06$
$\rightarrow \Sigma^- + \pi^+ + \pi^0 + \pi^0$	9	$0.12 \pm 0.05$
$\rightarrow \Sigma^+ + \pi^+ + \pi^- + \pi^-$	19	$0.19 \pm 0.06$
$\rightarrow \Sigma^- + \pi^- + \pi^+ + \pi^+$	13	$0.12 \pm 0.05$



FOOTNOTES

1. M. Alston, L. Alvarez, P. Eberhard, M. Good, W. Graziano, H. Ticho, and S. Wojcicki, *Phys. Rev. Letters* 5, 520 (1960).
2. O. Dahl, N. Horwitz, D. Miller, J. Murray, and P. White, *Phys. Rev. Letters* 6, 142 (1961).
3. J. P. Berge, P. Bastien, O. Dahl, M. Ferro-Luzzi, J. Kirz, D. H. Miller, J. J. Murray, A. H. Rosenfeld, R. D. Tripp, and M. Watson, (submitted to *Phys. Rev. Letters*).
4. H. Martin, L. Leipuner, W. Chinowsky, F. Shively, and R. Adair, *Phys. Rev. Letters* 6, 283 (1961).
5. M. M. Block, E. B. Brucker, R. Gessaroli, T. Kikuchi, A. Kovacs, C. M. Meltzer, R. Kraemer, M. Nussbaum, A. Pevsner, P. Schlein, R. Strand, H. O. Cohn, E. M. Harth, J. Leitner, L. Lendinara, L. Monari, and G. Puppi, (submitted to *Nuovo cimento*).
6. R. Ely, S. Fung, G. Gidal, Y. Pan, W. Powell, and H. White, *Bull. Am. Phys. Soc.* 6, 291 (1961).
7. M. Gell-Mann, *Phys. Rev.* 106, 1297 (1957).
8. P. Amati, A. Stanghellini, and B. Vitali, *Phys. Rev. Letters* 5, 524 (1960), *Nuovo cimento* 13, 1143 (1959).
9. R. Dalitz and S. F. Tuan, *Phys. Rev. Letters* 2, 425 (1959).
10. R. Dalitz, *Phys. Rev. Letters* 6, 239 (1961).
11. Ph. Meyer, J. Prentki, and Y. Yamaguchi, *Phys. Rev. Letters* 5, 442 (1960).
12. J. L. Shaw, University of California at La Jolla (private communication).

## FIGURE LEGENDS

Fig. 1. Dalitz plots for the reactions:

- (a)  $K^- + p \rightarrow \Sigma^+ + \pi^- + \pi^0$  (57 events)  
 (b)  $K^- + p \rightarrow \Sigma^- + \pi^+ + \pi^0$  (54 events)  
 (c)  $K^- + p \rightarrow \Sigma^0 + \pi^+ + \pi^-$  (27 events).

Fig. 2. Mass plots of the charged and neutral  $\Sigma$ - $\pi$  systems, including curves representing phase-space distributions.

- (a) Mass of  $(\Sigma\pi)^-$ , from the reactions:  $K^- + p \rightarrow \Sigma^0 + \pi^- + \pi^+$   
 $\rightarrow \Sigma^- + \pi^0 + \pi^+$
- (b) Mass of  $(\Sigma\pi)^+$ , from the reactions:  $K^- + p \rightarrow \Sigma^0 + \pi^+ + \pi^-$   
 $\rightarrow \Sigma^+ + \pi^0 + \pi^-$
- (c) Mass of  $(\Sigma\pi)^0$ , from the reactions:  $K^- + p \rightarrow \Sigma^+ + \pi^- + \pi^0$   
 $\rightarrow \Sigma^- + \pi^+ + \pi^0$






FIG. 1




FIG. 1




FIG. 1

