## Lawrence Berkeley National Laboratory

**LBL Publications** 

## Title

High Tc Superconducting Infrared Detectors

## Permalink


https://escholarship.org/uc/item/728026ff

### Author

Richards, P L

# Publication Date

1989-05-01



二條

Prepared for the U.S. Department of Energy under Contract DE-AC03-76SF00098

#### DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

#### HIGH T<sub>o</sub> SUPERCONDUCTING INFRARED DETECTORS

P.L. RICHARDS Department of Physics, University of California and Center for Advanced Materials, Materials and Chemical Sciences Division Lawrence Berkeley Laboratory, One Cyclotron Road, Berkeley, California 94720, U.S.A.

#### ABSTRACT

101

 $\langle \rangle$ 

A brief review will be given of published data from searches for a high  $\rm T_C$  photon detection mechanism. Critical comments will be made about the tests used to distinguish photon and thermal detection modes. An analysis of the prospects for a high  $\rm T_C$  bolometric detector will be presented.

#### RESULTS AND DISCUSSION

Much work has been stimulated by the idea that pair breaking by infrared photons in high  $T_c$  superconductors could cause modifications in film conductivity and thus lead to useful infrared photon detectors. Data have been published by many groups on the response of high  $T_c$  films with a variety of morphologies to visible and infrared photons. In general the values of specific detectivity  $D^{*}$  obtained are  $\leq 10^{\circ}$  cmHz<sup>1</sup>/<sup>2</sup>W<sup>-1</sup> which is too low for most practical applications. A common practice is to identify as thermal that fraction of the responsivity that scales as dR/dT, where R(T) is the film resistance. Since the responsivity also depends on temperature dependent thermal conductivities and specific heats, this practice is not reliable. Uncertainties remain as to what fraction of the observed response is thermal.

When a high T<sub>C</sub> infrared detector is designed specifically to enhance the thermal response, values of D<sup>\*</sup> approaching 10<sup>11</sup> cmHz<sup>1/2</sup>W<sup>-1</sup> can be obtained. Such performance will be very useful, especially for infrared wavelengths longer than -10  $\mu$ m. A theoretical analysis of this high T<sub>C</sub> bolometer will be presented.

#### CONCLUSIONS

- (1) The thermal component of the infrared responsivity of high  $T_C$  films cannot be reliably identified as the component that scales as dR/dT.
- (2) A properly optimized high T<sub>c</sub> infrared bolometer will have useful properties.

#### ACKNOWLEDGMENT

Work supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Materials Sciences Division of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

.

LAWRENCE BERKELEY LABORATORY CENTER FOR ADVANCED MATERIALS 1 CYCLOTRON ROAD BERKELEY, CALIFORNIA 94720