

# UC Irvine

## Faculty Publications

### Title

Catastrophic loss of stratospheric ozone in dense volcanic clouds

### Permalink

<https://escholarship.org/uc/item/741037pk>

### Journal

Journal of Geophysical Research: Atmospheres, 97(D9)

### ISSN

01480227

### Author

Prather, Michael

### Publication Date

1992-06-20

### DOI

10.1029/92JD00845

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

## Catastrophic Loss of Stratospheric Ozone in Dense Volcanic Clouds

MICHAEL PRATHER

*Department of Geosciences, University of California, Irvine*

Rapid, localized loss of ozone is predicted to occur in the mid-latitude and tropical stratosphere in the presence of very large concentrations of sulfate aerosols. Volcanic eruptions can increase the effective surface area of sulfuric acid so that heterogeneous reactions involving ClONO<sub>2</sub>, and secondarily N<sub>2</sub>O<sub>5</sub>, are able to suppress NO<sub>x</sub> abundances by more than a factor of 10 relative to gas phase chemistry. When NO<sub>x</sub> levels fall below a threshold, e.g., 0.6 ppb at 24 km in mid-latitudes, the chlorine-catalyzed loss of O<sub>3</sub> proceeds at rates comparable to those during the formation of the Antarctic ozone hole, more than 50 ppb per day. If such losses occurred following the eruption of Mount Pinatubo in the most volcanically perturbed regions over the tropics and mid-latitudes, this model predicts that they are driven primarily by the suppression of NO<sub>x</sub> below these critical levels. The increase in stratospheric chlorine since El Chichon has made Mount Pinatubo more than twice as effective in causing rapid O<sub>3</sub> loss. Overall global losses associated with a volcanic eruption are approximately linear in the amount of sulfate surface area, but depend critically on the rate of the ClONO<sub>2</sub>-sulfate reaction.

## 1. INTRODUCTION



Ever since laboratory measurements first showed that chemical reactions of stratospheric concern occur on sulfuric acid - water mixtures [Tolbert *et al.*, 1988; Mozurkewich and Calvert, 1988], photochemical models have been used to demonstrate that large increases in the stratospheric sulfate layer due to volcanic eruptions could lead to measurable ozone depletion by enhancing the chlorine-catalyzed destruction of ozone [Hofmann and Solomon, 1989]. Since then, further laboratory studies have confirmed that heterogeneous chemistry occurs on sulfate aerosols [Van Doren *et al.*, 1991; Hanson and Ravishankara, 1991; also J. A. Manion *et al.*, Heterogeneous reaction of N<sub>2</sub>O<sub>5</sub> and ClONO<sub>2</sub> on sulfuric acid surfaces representative of global stratospheric particulate, submitted to *Journal of Geophysical Research*, 1992 (hereinafter Manion *et al.*, submitted manuscript, 1992)]. Additional model studies continue to predict the importance of these reactions to stratospheric ozone [e.g., Mather and Brune, 1990; World Meteorological Organization (WMO), 1992] by affecting long-term trends due to the buildup of atmospheric chlorine [Rodriguez *et al.*, 1991], volcanic eruptions [Brasseur *et al.*, 1990], or projected aircraft emissions [Weissenstein *et al.*, 1991].

I show here the existence of a nonlinear, catastrophic (i.e., rapid) response in stratospheric photochemistry to the addition of sulfate aerosols. There exists a threshold, above which the odd-nitrogen (NO<sub>x</sub>) chemistry shuts down, as shown previously for the case of multiple solutions [Prather *et al.*, 1979] or in the case when chlorine levels (Cl<sub>y</sub>) globally exceed NO<sub>x</sub> abundances [Prather *et al.*, 1984]. If the sulfate aerosol loading reaches these critical levels, as might have occurred in some regions during the months following a major volcanic eruption such as Mount Pinatubo in June 1991 [McCormick, 1992], then the chemistry shifts to a predominantly Cl<sub>y</sub> plus HO<sub>x</sub> (= OH + HO<sub>2</sub>) system. Ozone loss rates become equivalent to those over springtime Antarctica when the ozone


hole is created, more than 50 ppb (1%) per day [e.g., Schoeberl *et al.*, 1989; Solomon, 1990]. In this case, however, such ozone losses are predicted to occur in the middle latitudes or even the tropics under bright sunlit conditions.

## 2. HETEROGENEOUS CHEMISTRY

The critical impact of sulfate layer chemistry, as noted by Rodriguez *et al.* [1991], is to move the odd-nitrogen species from NO<sub>x</sub> (= NO + NO<sub>2</sub>) into HNO<sub>3</sub> by two heterogeneous reactions:



The rate coefficient for each reaction (s<sup>-1</sup>) is calculated as the product of the reaction probability (γ, dimensionless), the aerosol area (α, cm<sup>2</sup> cm<sup>-3</sup> or cm<sup>-1</sup>), and the mean speed of the gaseous species (cm s<sup>-1</sup>) divided by 4. Reduction in the NO concentration leads indirectly to increases in ClO by changing the balance of HCl to ClO through the reduced efficiency of the reaction sequence,



The photochemical model [Prather and Jaffe, 1990; DeMore *et al.*, 1990; Prather, 1992] is used to derive a steady state cycle of all chemical species over 24 hours. The primary case study shown here is for the mid-latitude, middle stratosphere: 24 km altitude (29 mbar) at 45°N in March. Results from the two limiting cases of gas phase only (γ<sub>1</sub> = γ<sub>2</sub> = 0) and the aerosol-saturated case (γ<sub>1</sub> = 0.1, γ<sub>2</sub> = 0.01, α = 1000 × 10<sup>-8</sup> cm<sup>-1</sup>) are given in Table 1. The two limits demonstrate clearly the difference between the gas phase NO<sub>x</sub>-dominated chemistry and the aerosol-saturated Cl<sub>y</sub>-HO<sub>x</sub> chemistry. The transition from gas phase to aerosol-saturated depends critically on the reaction probability of (2) and is examined in Figures 1-3. Release of ClO from the HCl reservoir (by reaction with OH) will be rapid, less than 20 days, once the HO<sub>x</sub> levels rise in

Copyright 1992 by the American Geophysical Union.

Paper number 92JD00845.  
0148-0227/92/92JD-00845\$02.00




Fig. 1. Cumulative production rate of  $\text{HNO}_3$  (parts per billion per day) as a function of aerosol surface area ( $\alpha$ ,  $10^{-8} \text{ cm}^2 \text{ cm}^{-3}$ ) for the three reactions:  $\text{NO}_2 + \text{OH}$ ,  $\text{N}_2\text{O}_5 + \text{aerosol}$ ,  $\text{ClONO}_2 + \text{aerosol}$ . Calculations apply to  $45^\circ\text{N}$  in March at 24 km altitude (see Table 1). The panels show results for different values for the reaction probability of  $\text{ClONO}_2 + \text{H}_2\text{O}/\text{H}_2\text{SO}_4$ : (a)  $\gamma_2 = 0.0$ , (b)  $\gamma_2 = 0.0012$ , (c)  $\gamma_2 = 0.01$ .

response to the reduced  $\text{NO}_x$  abundances. Therefore, the assumed steady state balance among  $\text{NO}_y$  and  $\text{Cl}_y$  species is readily achieved.

The relative importance of reactions (1) and (2) in producing  $\text{HNO}_3$  is shown in Figure 1. For  $\alpha < 0.1 \times 10^{-8} \text{ cm}^{-1}$  the gas phase formation of  $\text{HNO}_3$  dominates:



As the area increases, the pathway through reaction (1) becomes comparable to (5). Without reaction (2), as in Figure 1a, the impact of heterogeneous chemistry saturates for  $\alpha > 10 \times 10^{-8} \text{ cm}^{-1}$ . When reaction (2) is included with probability equal to 0.0012, as in Figure 1b, the reaction path through  $\text{ClONO}_2$  quickly becomes the dominant source of  $\text{HNO}_3$  (and  $\text{HOCl}$  as well). If the probability of reaction (2) is as large as 0.01 (Figure 1c) then it becomes important even at background levels of aerosols.

The abundance of  $\text{NO}_x$  is shown in Figure 2b for these same three cases. Even at background levels,  $\text{NO}_x$  is suppressed by a factor of 2 relative to the gas phase limit [e.g., Mather and Brune, 1990; Rodriguez et al., 1991]. For large surface areas associated with volcanic eruptions [Hofmann and Solomon, 1989; McCormick and Veiga, 1992], locally  $\text{NO}_x$  mixing ratios can be reduced by more than a factor of 20. Measurements of  $\text{NO}_2$  after Pinatubo show substantial reductions, greater than 20% [Johnston et al., 1992], but these column integrals cannot resolve the perturbations within the volcanic layers. Under




Fig. 2. (a) Net ozone loss (parts per billion per day) and (b) noontime  $\text{NO}_x$  (parts per billion) as a function of aerosol surface area ( $\alpha$ ,  $10^{-8} \text{ cm}^2 \text{ cm}^{-3}$ ) for different  $\gamma_2$ . Calculations apply to  $45^\circ\text{N}$  in March at 24 km altitude (see Table 1). Also denoted are the loss rates typical during the formation of the Antarctic ozone hole, and the sulfate aerosol areas typical of background levels and after large volcanic eruptions, such as Mount Pinatubo. Net  $\text{O}_3$  loss for  $\gamma_2 = 0.0012$  with a 25% reduction in  $\text{Cl}_y$  (2.1 ppb) is shown in Figure 2a as a dotted line.

these conditions the reduction in  $\text{NO}_x$  depends critically on the reaction probability of (2). Reaction (1) alone cannot reduce the  $\text{NO}_x$  abundance below 1.2 ppb (at 24 km,  $45^\circ\text{N}$  in March) because the rate is essentially quadratic in  $\text{NO}_x$  densities (i.e., the formation of  $\text{N}_2\text{O}_5$  is proportional to the product of  $\text{NO}_2$  and  $\text{NO}_3$ ). When  $\text{NO}_x$  is less than 1.2 ppb the formation of  $\text{N}_2\text{O}_5$  overnight (and its subsequent reaction on sulfate aerosols) does not compete with the daylight photochemistry. With the addition of reaction (2), the enhancement of aerosol surface area can further suppress  $\text{NO}_x$  (i.e., formation of  $\text{ClONO}_2$  is linear in the  $\text{NO}_x$  abundance, being proportional to the product of  $\text{ClO}$  and  $\text{NO}_2$ ) to a new asymptotic value of 0.1 ppb.

### 3. RAPID OZONE LOSS

Net loss rates for  $\text{O}_3$  as functions of  $\alpha$  and  $\gamma_2$  are shown in Figure 2a. At 24 km,  $45^\circ\text{N}$  in March for gas phase chemistry, the ozone is almost in balance: production of 25.4 ppb/day is offset by losses of 27.4 ppb/day (see Table 1). Ozone losses increase linearly at first with aerosols for areas greater than  $1 \times 10^{-8} \text{ cm}^{-1}$  (background); this would be apparent if Figure 2a were plotted with a linear abscissa. The rate of increase with aerosol area depends sensitively on  $\gamma_2$ . For  $\gamma_2 = 0.0$ , the  $\text{O}_3$  loss becomes constant for  $\alpha$  between 10 and  $30 \times 10^{-8} \text{ cm}^{-1}$  because  $\text{NO}_x$  no longer decreases. For  $\gamma_2 = 0.01$ , the  $\text{O}_3$  loss asymptotes (for  $\alpha$  between 100 and  $300 \times 10^{-8} \text{ cm}^{-1}$ ) at 70 ppb per day, equivalent to loss rates during formation of the

Antarctic ozone hole [Schoeberl *et al.*, 1989]. In the cases shown here, the ozone loss rate can be derived simply as a function, albeit a highly nonlinear function, of the  $\text{NO}_x$  abundance. Rapid ozone loss only begins when the noontime  $\text{NO}_x$  mixing ratios fall below 0.6 ppb, and Antarcticlike losses of greater than 50 ppb per day occur when  $\text{NO}_x$  levels reach 0.2 ppb. Cumulative loss rates due to the  $\text{O}_x$ ,  $\text{NO}_x$ ,  $\text{HO}_x$ ,  $\text{ClO}_x$ , and  $\text{BrO}_x$  catalytic cycles (see definitions in Table 1) are shown in Figure 3 as a function of  $\alpha$  for the case  $\gamma_2 = 0.01$ . For smaller  $\gamma_2$  the asymptotic limit would shift to large  $\alpha$ . Total losses under these conditions (24 km at 45°N in March) exceed 90 ppb per day as the area approaches  $1000 \times 10^{-8} \text{ cm}^{-1}$ : the combined total of  $\text{NO}_x + \text{HO}_x$  loss remains approximately

constant as the total of  $\text{ClO}_x + \text{BrO}_x$  loss increases by more than a factor of 10.

This potential for rapid ozone loss extends over much of the stratosphere. Beyond the example shown here, similar behavior with  $\text{O}_3$  losses of 20 to 40 ppb per day extends down to 20 km altitude at 45°N. In the tropical stratosphere, the net  $\text{O}_3$  tendency is largely positive (primary production by  $\text{O}_2$  photodissociation greatly exceeds chemical losses). Nevertheless, the addition of rapid heterogeneous processing to the altitude range 24–28 km at 15°N in March can increase the  $\text{ClO}_x$  plus  $\text{HO}_x$ -catalyzed losses by as much as 40 ppb per day, with the net still remaining positive. At higher altitudes in the stratosphere, the effect of heterogeneous chemistry is more complex: the initial suppression of  $\text{NO}_x$ -catalyzed loss dominates and leads to reduced  $\text{O}_3$  losses with increasing aerosol area; only at very high aerosol areas would  $\text{NO}_x$  levels fall below the critical threshold as described above. The maintenance of such large surface areas of sulfuric acid - water mixtures at higher altitudes and warmer temperatures is difficult and would require significantly greater sulfur and water abundances [e.g., Hamill *et al.*, 1988].

A question remains as to why this effect, the nonlinear suppression of  $\text{NO}_x$  followed by the catastrophic loss rate of  $\text{O}_3$ , has not been reported in previous studies on the subject of volcano-induced ozone loss. Brasseur *et al.* [1990] assumed large aerosol loadings, reaching  $35 \times 10^{-8} \text{ cm}^{-1}$  at 24 km over the equator, and a value of  $\gamma_2 = 0.0026$ , which stops just short of the rapid losses shown in Figure 2a. However, their Figure 2 shows clearly that global ozone loss becomes sensitive to  $\gamma_2$  rather than  $\gamma_1$ , reaching -25% (globally!) for  $\gamma_2 = 1$ . Hofmann and Solomon [1989] used  $\gamma_2$  values of 0.0065 and  $\alpha$  peak aerosol areas of about  $40 \times 10^{-8} \text{ cm}^{-1}$ , less at 24 km. They predicted large increases in HOCl (as here) but did not report on the nonlinearity where the high ClO becomes a positive feedback in the  $\text{NO}_x$  cycles and the chlorine-catalyzed loss rapidly escalates. Rodriguez *et al.* [1991] considered only reaction (1) and would not have seen this phenomenon.

In the presence of such large aerosol content, what other modifications of the stratospheric environment might affect  $\text{O}_3$  chemistry? For one, the enhanced scattering from the aerosols [Michelangeli *et al.*, 1989] will increase photolysis rates at the top of the cloud and reduce them near the bottom. Further, net radiative heating by the aerosols may also perturb the temperature [Labitzke and McCormick, 1992], and possibly lead to a change in the mean diabatic circulation (S. Kinne *et al.*, Radiative impact of Pinatubo aerosols on tropical ozone, manuscript in preparation, 1992). These perturbations are too complex to analyze with this model. Their impacts on stratospheric  $\text{O}_3$  may be important, but on a localized parcel scale they are likely to be small compared with the dramatic shift in the chlorine-catalyzed ozone loss discussed here.

#### 4. LABORATORY AND ATMOSPHERIC CONSTRAINTS

The different laboratory measurements are in qualitative agreement regarding the sharp dependence of reaction (2) on the water content of the sulfuric acid - water mix, although absolute values for  $\gamma_2$  vary by a factor of 2 [Tolbert *et al.*, 1988; Hanson and Ravishankara, 1991; also Manion *et al.*, submitted manuscript, 1992]. A  $\gamma_2$  of 0.01 or greater corresponds to extremely wet mixtures with less than 52% by weight of  $\text{H}_2\text{SO}_4$ . Such wet sulfuric acid droplets would occur in the lower-middle stratosphere only if temperatures dropped

TABLE 1. Limiting Cases of Gas Phase and Sulfate Aerosol Chemistry at 45°N, 24 km (219 K, 29 mbar)

		Initial Concentration	
$\text{O}_3$ ,	ppm	4.5	
$\text{H}_2\text{O}$ ,	ppm	4.0	
$\text{H}_2$ ,	ppm	0.5	
$\text{CH}_4$ ,	ppm	1.1	
$\text{NO}_y$ ,	ppb	10.6	
$\text{Cl}_y$ ,	ppb	2.8	
$\text{Br}_y$ ,	ppt	15.	
Gas Phase Only $\gamma_1 = \gamma_2 = 0$ $\alpha = 0$		Aerosol-Saturated Case $\gamma_1 = 0.1$ , $\gamma_2 = 0.01^*$ $\alpha = 1000 \times 10^{-8} \text{ cm}^{-1}$	
<i>Noontime Mixing Ratios</i>			
OH,	ppt	2.22	5.24
$\text{HO}_2$ ,	ppt	10.9	25.0
NO,	ppb	1.22	0.02
$\text{NO}_2$ ,	ppb	1.70	0.08
$\text{N}_2\text{O}_5$ ,	ppb	0.61	0.00
$\text{HNO}_3$ ,	ppb	5.32	10.42
$\text{ClONO}_2$ ,	ppb	0.93	0.07
HCl,	ppb	1.76	0.44
HOCl,	ppb	0.03	1.01
ClO,	ppb	0.09	1.23
Cl,	ppt	0.069	0.050
<i>Production (24-hour Average Rates), ppb/day</i>			
$\text{O}_2 + \text{h}\nu$ ( $\times 2$ )		21.8	21.8
$\text{HO}_2 + \text{NO}$ ( $\times 1$ )		3.6	0.2
<i><math>\text{O}_x</math> Loss, ppb/day</i>			
$\text{O}_3 + \text{O}$ ( $\times 2$ )		2.4	2.4
<i><math>\text{NO}_x</math> Loss, ppb/day</i>			
$\text{NO}_2 + \text{O}$ ( $\times 2$ )		14.3	0.5
$\text{NO}_3 + \text{h}\nu$ ( $\times 2$ )		0.8	0.1
<i><math>\text{HO}_x</math> Loss, ppb/day</i>			
$\text{HO}_2 + \text{O}_3$ ( $\times 1$ )		5.4	16.7
$\text{OH} + \text{O}_3$ ( $\times 1$ )		1.5	4.1
<i><math>\text{Cl}_y + \text{Br}_y</math> Loss, ppb/day</i>			
$\text{ClO} + \text{O}$ ( $\times 2$ )		2.4	38.7
$\text{Cl}_2\text{O}_2 + \text{h}\nu$ ( $\times 2$ )		0.0	6.1
$\text{HOCl} + \text{h}\nu$ ( $\times 1$ )		0.3	12.6
$\text{BrO} + \text{ClO}$ ( $\times 2$ )		0.3	11.1
<i>Net Production minus Loss, ppb/day</i>			
$\text{O}_3$		-2.0	-70.3

\* The large values for  $\gamma_2$  and  $\alpha$  are chosen here to illustrate the upper limit in the heterogeneous  $\text{NO}_x$  processing; the suppression of  $\text{NO}_x$  depends on their product. Note that  $\gamma_2 = 0.01$  would apply to colder temperatures or to enhanced water vapor.




Fig. 3. Cumulative O<sub>3</sub> loss (parts per billion per day) as a function of aerosol surface area ( $\alpha$ ,  $10^{-8}$  cm<sup>2</sup>cm<sup>-3</sup>) for different catalytic cycles with  $\gamma_2 = 0.01$ . Calculations apply to 45°N in March at 24 km altitude (see Table 1).

to 197 K or less (approaching the threshold for condensation of nitric acid - water mixtures [Hanson and Mauersberger, 1988]), or if water vapor were greatly enhanced. Values of  $\gamma_2 = 0.0012$  correspond to 66%wt H<sub>2</sub>SO<sub>4</sub> and stratospheric temperatures of about 210 K; while  $\gamma_2 = 2 \times 10^{-4}$  (not shown in calculations above) is applicable to 75%wt solutions (about 224 K) and may be an effective rate for reaction (2) throughout much of the middle stratosphere.

The uncertainty in  $\gamma_2$  from the laboratory data ( $\pm 50\%$ ) is matched by the uncertainty in averaging  $\gamma$  over a range of stratospheric temperatures. The exponential dependence of  $\gamma_2$  on %wt H<sub>2</sub>SO<sub>4</sub> and thus on temperature (i.e., a factor of 2 for each change of 5 K), makes the temperature history of an air parcel an extremely important factor in averaging the effective rate of reaction (2). The range in  $\gamma_1$  from the laboratory data, 0.05 to 0.14, would not greatly affect these results because reaction (1) is ineffective at low levels of NO<sub>x</sub>.

The available surface area for reactions (1) and (2) throughout the stratosphere in the form of liquid sulfuric acid aerosols is highly variable [Hofmann, 1990] and not easy to define [Thomason, 1991]. Background levels in the middle-lower stratosphere at mid-latitudes (i.e., the cleanest periods observed over the last decade) are about  $1 \times 10^{-8}$  cm<sup>-1</sup>, with a range of at least a factor of 2. Several months after the eruption of El Chichon in 1983, surface areas in excess of  $40 \times 10^{-8}$  cm<sup>-1</sup> were observed [Hofmann and Solomon, 1989], but values above 24 km were a factor of 8 less. Current estimates indicate that the sulfur injected by Mount Pinatubo (June 1991) exceeded that of El Chichon by a factor of 2 to 3 [Bluth et al., 1992] and reached higher altitudes [McCormick and Veiga, 1992]. Early observations of aerosol optical extinction (proportional to the surface area of the larger particles) show large regions in the tropical stratosphere with  $\alpha > 100 \times 10^{-8}$  cm<sup>-1</sup> at altitudes near 24 km. It is likely that significant volumes of the tropical and mid-latitude stratosphere were exposed to such large effective surface areas of sulfuric acid droplets.

In summary, the uncertainty in laboratory measurements and stratospheric conditions suggest a range for  $\gamma_2$  of 0.0001 to 0.0012 for the middle stratosphere, with larger values in the

lower stratosphere or where temperatures are much colder. If the effective  $\gamma_2$  were of order 0.01 in the lower stratosphere, then this proposed mechanism should have been (but was apparently not) triggered by El Chichon [Hofmann and Solomon, 1989; Chandra and Stolarski, 1991]. If the averaged  $\gamma_2$  were to exceed 0.002 in some regions, then Mount Pinatubo may have injected enough sulfur into the stratosphere for an unambiguous test.

## 5. IMPLICATIONS AND TESTS

The chlorine-catalyzed ozone loss predicted here for mid-latitudes is similar in magnitude to polar processes causing the Antarctic ozone hole, but is driven predominantly by the O+ClO and HOCl+hv cycles, rather than the Cl<sub>2</sub>O<sub>2</sub> cycle. The loss is described here as catastrophic in that large, isolated parcels of perturbed air may experience measurable ozone loss, of order 1% per day, in the 2-3 weeks before they mix with the background stratosphere [Prather and Jaffe, 1990]. Such a property makes this type of O<sub>3</sub> loss more readily detectable than if the equivalent loss were spread globally over a season. Average global loss of ozone, on the other hand, will depend less on the details of rapid loss in these clouds and more critically on the rate of this heterogeneous reaction and its ability to suppress NO<sub>x</sub>. Calculations of global ozone response to large volcanic injections must also include dynamical transports as in the global chemical transport models of Brasseur et al. [1990] and Rodriguez et al. [1991].

Hofmann and Solomon [1989] point out that the volcano-modulated O<sub>3</sub> loss they ascribe to El Chichon (however, see Chandra and Stolarski [1991]) might be much greater for large volcanic injections in the future when chlorine levels are greater. Pittock's [1965] observation of ozone depletion within a volcanic aerosol layer associated with the eruption of Mount Agung (March 1963) cannot be easily ascribed to this mechanism since background Cl<sub>y</sub> levels at this time were less than 1 ppb. Michelangeli et al. [1991] argue that NaCl from volcanic ash injected into the stratosphere would react, enhancing ClO values, and lead to ozone depletion. Any additional source of Cl<sub>y</sub> could tip the balance between the chlorine and odd-nitrogen species at lower levels of sulfate aerosols.

Stratospheric chlorine loading (i.e., Cl<sub>y</sub> abundances in the lower-middle stratosphere where the bulk of the ozone loss is observed to date [see WMO, 1992]) was about 2.8 ppb in 1991, the year of the Mount Pinatubo eruption, and about 2.1 ppb in 1983, the time of El Chichon. Figure 2a includes, as a dotted line, a calculation for  $\gamma_2 = 0.0012$  with Cl<sub>y</sub> = 2.1 ppb. The chlorine levels are indeed critical; the 33% increase in Cl<sub>y</sub> from 1983 to 1991 results in equivalent ozone destruction for sulfate aerosol areas that are lower by a factor of 2.5! Two effects amplify: the asymptotic loss rates increase with Cl<sub>y</sub>, and, more importantly, the suppression of NO<sub>x</sub> through reaction (2) is enhanced at lower surface areas. Additional tests, in the laboratory and atmosphere, of the efficiency of the reaction of ClONO<sub>2</sub> on sulfate aerosols are needed.

Ozone depletion is approximately linear with volcanic aerosol area, but highly nonlinear with regard to chlorine abundances and temperatures. It should therefore be extremely patchy, occurring in regions with sufficient Cl<sub>y</sub> and large concentrations of sulfate aerosols. The initial properties of the plume (i.e., tropospheric air with entrained stratospheric air) would look similar to perturbed stratospheric air with low O<sub>3</sub>

and low NO<sub>x</sub>; however, the other properties of the freshly injected air would appear quite different (e.g., low ClO and Cl<sub>2</sub>, low NO<sub>x</sub>, high N<sub>2</sub>O, adiabatic lapse rates, extremely high aerosol concentrations with volcanic ash). In order for significant O<sub>3</sub> loss to occur, the initial plume must be mixed on a molecular scale with air containing stratospheric concentrations of Cl<sub>2</sub> and O<sub>3</sub>.

Compared with El Chichon, Mount Pinatubo (2-3 times the amount of sulfur [Bluth *et al.*, 1992], in an atmosphere with 33% more ambient chlorine) is many times more likely to have created some regions in the stratosphere that crossed the threshold of catastrophic chlorine-driven ozone loss during the first year following the eruption. Pinatubo offers a unique opportunity, and we await the measurements [Grant *et al.*, 1992; also M. R. Schoeberl *et al.*, Tropical ozone loss following the eruption of Mount Pinatubo, manuscript in preparation, 1992; J. Waters *et al.*, MLS observations of tropical ozone following Mount Pinatubo, manuscript in preparation, 1992].

**Acknowledgments.** NASA's Goddard Institute for Space Studies supported the initial work on this paper; revisions were completed at UC Irvine. This work derives from a collaboration with M. Tolbert investigating the impact of heterogeneous (sulfuric acid - water) reactions involving methane oxidation products (M. Tolbert *et al.*, Uptake of formaldehyde by sulfuric acid solutions: Impact on stratospheric ozone, manuscript in preparation, 1992). Inclusion of these reactions in the current work has a noticeable effect on the odd-hydrogen cycles, but no consequential impact on the results. I thank J. Waters and M. Schoeberl for preliminary discussions of their observations. This research is supported by grants to Goddard Institute for Space Studies, Columbia University, and UC Irvine from the Atmospheric Chemistry Programs of NASA and the National Science Foundation.

#### REFERENCES

- Bluth, G. J. S., S. D. Doiron, C. C. Schnetzler, A. J. Krueger, and L. S. Walter, Global tracking of the SO<sub>2</sub> clouds from the June 1991 Mount Pinatubo eruptions, *Geophys. Res. Lett.*, **19**, 151-154, 1992.
- Brasseur, G. P., C. Granier, and S. Walters, Future changes in stratospheric ozone and the role of heterogeneous chemistry, *Nature*, **348**, 626-628, 1990.
- Chandra, S. and R. S. Stolarski, Recent trends in stratospheric total ozone: Implications of dynamical and El Chichon perturbations, *Geophys. Res. Lett.*, **19**, 2278-2280, 1991.
- DeMore, W. B., *et al.*, Chemical kinetics and photochemical data for use in stratospheric modeling, Rep. *JPL 90-1*, NASA Jet Propulsion Lab., Pasadena, Calif., 1990.
- Grant, W. B., *et al.*, Observations of reduced ozone concentrations in the tropical stratosphere after the eruption of Mount Pinatubo, *Geophys. Res. Lett.*, **20**, in press, 1992.
- Hamill, P., R. P. Turco, and O. B. Toon, On the growth of nitric and sulfuric acid aerosol particles under stratospheric conditions, *J. Atmos. Chem.*, **7**, 287-315, 1988.
- Hanson, D., and K. Mauersberger, Laboratory studies of the nitric acid trihydrate: Implications for the south polar stratosphere, *Geophys. Res. Lett.*, **15**, 855-858, 1988.
- Hanson, D. R., and A. R. Ravishankara, The reaction probabilities of ClONO<sub>2</sub> and N<sub>2</sub>O<sub>5</sub> on 40 to 75% sulfuric acid solutions, *J. Geophys. Res.*, **96**, 17307-17314, 1991.
- Hofmann, D. J., Increase in the stratospheric background sulfuric acid aerosol mass in the past 10 years, *Science*, **248**, 996-1000, 1990.
- Hofmann, D. J., and S. Solomon, Ozone destruction through heterogeneous chemistry following the eruption of El Chichon, *J. Geophys. Res.*, **94**, 5029-5041, 1989.
- Johnston, P. V., R. L. McKenzie, J. G. Keys, and W. A. Matthews, Observations of depleted stratospheric NO<sub>2</sub> following the Pinatubo eruption, *Geophys. Res. Lett.*, **19**, 211-213, 1992.
- Labitzke, K., and M. P. McCormick, Stratospheric temperature increases due to Pinatubo aerosols, *Geophys. Res. Lett.*, **19**, 207-210, 1992.
- Mather, J. H., and W. H. Brune, Heterogeneous chemistry on liquid sulfate aerosols: A comparison of in situ measurements with zero-dimensional model calculations, *Geophys. Res. Lett.*, **17**, 1283-1286, 1990.
- McCormick, M. P., Initial assessment of the stratospheric and climatic impact of the 1991 Mount Pinatubo eruption: Prologue, *Geophys. Res. Lett.*, **19**, 149, 1992.
- McCormick, M. P., and R. E. Veiga, SAGE II measurements of early Pinatubo aerosols, *Geophys. Res. Lett.*, **19**, 155-158, 1992.
- Michelangeli, D. V., M. Allen, and Y.L. Yung, El Chichon volcanic aerosols: Impact of radiative, thermal, and chemical perturbations, *J. Geophys. Res.*, **94**, 18429-18443, 1989.
- Michelangeli, D. V., M. Allen, and Y.L. Yung, Heterogeneous reactions with NaCl in the El Chichon volcanic aerosols, *Geophys. Res. Lett.*, **18**, 673-676, 1991.
- Mozurkewich, M., and J. G. Calvert, Reaction probability of N<sub>2</sub>O<sub>5</sub> on aqueous aerosols, *J. Geophys. Res.*, **93**, 15889-15896, 1988.
- Pitcock, A.B., Possible destruction of ozone by volcanic material at 50 mbar, *Nature*, **207**, 182, 1965.
- Prather, M. J., More rapid polar ozone depletion through reaction of HOCl with HCl on polar stratospheric clouds, *Nature*, **355**, 534-537, 1992.
- Prather, M. J., and A. H. Jaffe, Global impact of the Antarctic ozone hole: Chemical propagation, *J. Geophys. Res.*, **95**, 3473-3492, 1990.
- Prather, M. J., M. B. McElroy, S. C. Wofsy, and J. A. Logan, Stratospheric chemistry: Multiple solutions, *Geophys. Res. Lett.*, **6**, 163-164, 1979.
- Prather, M. J., M. B. McElroy, and S. C. Wofsy, Reductions in ozone at high concentrations of stratospheric halogens, *Nature*, **312**, 227-231, 1984.
- Rodriguez, J. M., M. K. W. Ko, and N.-D. Sze, Role of heterogeneous conversion of N<sub>2</sub>O<sub>5</sub> on sulphate aerosols in global ozone losses, *Nature*, **352**, 134-137, 1991.
- Schoeberl, M. R., *et al.*, Reconstruction of the constituent distribution and trends in the Antarctic polar vortex from ER-2 flight observations, *J. Geophys. Res.*, **94**, 16815-16846, 1989.
- Solomon, S., Progress towards a quantitative understanding of Antarctic ozone depletion, *Nature*, **347**, 347-354, 1990.
- Thomason, L. W., A diagnostic stratospheric aerosol size distribution inferred from SAGE II measurements, *J. Geophys. Res.*, **96**, 22501-22508, 1991.
- Tolbert, M. A., M. J. Rossi, and D. M. Golden, Heterogeneous interactions of ClONO<sub>2</sub>, HCl and HNO<sub>3</sub> with sulfuric acid surfaces at stratospheric temperatures, *Geophys. Res. Lett.*, **15**, 847-850, 1988.
- Van Doren, J. M., *et al.*, Uptake of N<sub>2</sub>O<sub>5</sub> and HNO<sub>3</sub> by aqueous sulfuric acid droplets, *J. Phys. Chem.*, **95**, 1684-1689, 1991.
- Weisenstein, D. K., M. K. W. Ko, J. M. Rodriguez, and N.-D. Sze, Impact of heterogeneous chemistry on model-calculated ozone change due to high speed civil transport aircraft, *Geophys. Res. Lett.*, **18**, 1991-1994, 1991.
- World Meteorological Organization (WMO), Scientific assessment of stratospheric ozone: 1991, Rep. 25, Global Ozone Res. and Monit. Proj., Geneva, 1992.

M. J. Prather, Department of Geosciences, UC Irvine, CA 92717

(Received December 20, 1991;  
revised April 2, 1992;  
accepted April 2, 1992.)