

UC Berkeley

UC Berkeley Previously Published Works

Title

Smoking-Cessation Interventions in Appalachia: A Systematic Review and Meta-Analysis

Permalink

<https://escholarship.org/uc/item/7464d5x0>

Journal

American Journal of Preventive Medicine, 58(2)

ISSN

0749-3797

Authors

Gupta, Shalika
Scheuter, Claudia
Kundu, Arti
[et al.](#)

Publication Date

2020-02-01

DOI

10.1016/j.amepre.2019.09.013

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Smoking-Cessation Interventions in Appalachia: A Systematic Review and Meta-analysis

Shalika Gupta, MA,¹ Claudia Scheuter, MD,^{1,2} Arti Kundu, PhD,¹ Naina Bhat, MBBS,¹ Alasdair Cohen, PhD, MPH, MSc,¹ Shelley N. Facente, MPH,^{1,3}

From the ¹Division of Epidemiology, School of Public Health, University of California, Berkeley, Berkeley, California; ²Clinical Excellence Research Center, Stanford University, California; and ³Facente Consulting, Richmond, California

Address correspondence to: Shelley N. Facente, MPH, School of Public Health, Department of Epidemiology, University of California, Berkeley, 2121 Berkeley Way #5302, Berkeley CA 94720. E-mail: sfacente@berkeley.edu.

Context: Appalachia, a socioeconomically disadvantaged rural region in the eastern U.S., has one of the nation's highest prevalence rates of smoking and some of the poorest health outcomes. Effective interventions that lower smoking rates in Appalachia have great potential to reduce health disparities and preventable illness; however, a better understanding of effective interventions is needed.

Evidence acquisition: This review included trials that evaluated the impact of smoking-cessation programs among populations living in Appalachia. The search was carried out on October 9, 2018, and comprised the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, Embase, and Scopus for academic journal articles published in English, with no date restrictions. After preliminary screening, potentially relevant full-text articles were independently reviewed by the authors with a Cohen's κ of 0.72, leading to final inclusion of nine articles.

Evidence synthesis: Eligible studies were assessed qualitatively for heterogeneity and risk of bias. Six of the nine included studies had extractable data related to dichotomous smoking status and reported a measure of association suitable for inclusion in a meta-analysis. For those six studies, the pooled RR and pooled OR were estimated using random effects models, with an I^2 index demonstrating substantial heterogeneity. A funnel plot of the six trials appeared relatively symmetric.

Conclusions: Participation in smoking-cessation interventions increased the probability of smoking abstinence among Appalachian smokers an estimated 2.33 times (pooled RR=2.33, 95% CI=1.03, 5.25, $p=0.04$). Given the low number of studies, their substantial heterogeneity, and high risk of bias, the evidence of the effectiveness of smoking-cessation interventions in Appalachia must be interpreted with caution.

CONTEXT

Smoking is widely recognized as the leading, yet preventable risk factor for the development of disease as well as premature mortality in the U.S.¹ Owing to concerted public health campaigns and legislative efforts, smoking rates among adults have dropped from about 43% in the 1960s to 16% in 2016.² However, this tremendous success has not been distributed uniformly. Smoking rates have remained substantially higher in populations of lower SES, such as those with lower educational attainment and individuals living in poverty.² Appalachia is a heterogeneous and socioeconomically disadvantaged region in the eastern U.S. spanning several states. Although Appalachia includes large urban and suburban areas like Pittsburgh, PA, 42% of Appalachia's population are rural, compared with 20% nationally.³ There is considerable economic heterogeneity within Appalachia with poverty rates approaching 25% of the population in Central Appalachian counties, whereas counties in metropolitan areas of Appalachia are close to the national average.⁴ Appalachia has one of the highest prevalence rates of smoking in the U.S.,⁵ and some of the worst health outcomes in the U.S. as well, associated with a 2.4-year gap in life expectancy compared with the rest of the U.S.⁶

In national surveys, up to 69% of smokers are interested in quitting smoking and 42% attempted to quit smoking in the previous year.⁷ Yet, only 55% of these attempts lead to sustained abstinence.⁷ Effective interventions that lower smoking rates in Appalachia therefore have great potential to reduce health disparities and preventable illness in this socioeconomically disadvantaged population. Given the challenges of reducing smoking rates in populations of lower SES, a better understanding of effective interventions is needed. The aim of this systematic review is to collect and summarize the evidence on smoking-cessation programs specifically

targeting individuals living in Appalachia and associated changes in (smoking-related) health outcomes.

EVIDENCE ACQUISITION

The methods described for this review were based on the guidelines provided by the Cochrane Handbook for Systematic Reviews of Interventions.⁸

This review included studies published in English that evaluated the impact of smoking-cessation programs in populations living in Appalachia as defined by the Appalachia Regional Commission as belonging to this area.³ According to this definition, Appalachia includes all of West Virginia and select rural counties in Alabama, Georgia, Kentucky, Maryland, Mississippi, New York, North Carolina, Ohio, Pennsylvania, South Carolina, Tennessee, and Virginia (Figure 1). A complete list of the counties comprising Appalachia can be found on the Appalachia Regional Commission's website.⁹

Studies that examined individuals in other states as well as those belonging to the Appalachian region were included only if the data was reported separately by region. Studies included all types of smoking-cessation interventions (i.e., programs that were administered in person or online) administered to any population (i.e., no restrictions by age, gender, or ethnicity of participants) in any setting (i.e., including community-based or hospital-based interventions). Appendix Table 1 provides more details on the eligibility criteria according to the PICOS framework. Both studies with smoking-cessation programs as the primary intervention of interest, and as well as studies with smoking-cessation interventions as part of a package of

health-related interventions, were included if smoking rates or a smoking-related health outcome were reported in the publication as primary or secondary outcomes. This review was limited to individual and cluster randomized interventions, and excluded observational studies from this review.

The search included The Cochrane Central Register of Controlled Trials (CENTRAL) as well as MEDLINE (1966–2018), Embase (1980–2018), and Scopus (2004–2018), for academic journal articles published in English, from the start of each database to October 9, 2018 (i.e., no restrictions on publication dates), searching title, abstract, and MeSH Terms/author keywords. Prior to conducting the search, the systematic review protocol was preregistered on PROSPERO (registration number CRD42018112135). Appendix Table 2 shows search sets and terms, Appendix Table 3 details the search code used, and Appendix Tables 4–7 provide detailed results by database.

The following stepwise approach was used during the screening and reviewing process. First, the database search was conducted as outlined above, generating a total number of articles generated from each repository. All duplicates were then removed from searches across multiple repositories, using Zotero, version 5.0.569, followed by a manual review of titles and abstracts to determine remaining duplicate entries. In the next step, one member of the team (SG) randomly assigned 50% of the articles to NB, SF, CS, and AG, who independently screened titles and abstracts to identify papers potentially related to the research question and eligibility criteria. Thus, each article was screened separately by two team members. Once preliminary screening was completed, one team member (SG) received the results of each rater's screening and

calculated inter-rater reliability using a κ statistic for the screening to this point. It was determined in advance that a poor inter-rater reliability ($\kappa < 0.6$)¹⁰ would result in a re-examination of eligibility criteria and screening by the whole team, and a repeat of preliminary screening procedures until inter-rater reliability improved above that set threshold. Team member SG served as a tiebreaker for any articles for which the two preliminary screeners were not in agreement about inclusion. After screening was complete, the full text of the articles identified in the previous step was retrieved; any articles for which full text could not be obtained were excluded from the review. Each full-text article was reviewed independently by two team members to determine if the study met eligibility criteria. Whenever there was disagreement about whether or not a study should be included, SG read the article and made a final decision. Finally, after the full-text review of the initial articles, references of all eligible articles were hand searched for additional, potentially eligible articles. These articles were then reviewed and eligibility determined according to the process outlined above.

Risk of bias was assessed using the Cochrane “Risk of Bias 2.0” tool from the Cochrane Handbook for Systematic Reviews of Interventions.⁸ For each study selected for review, two team members independently rated the risk of bias by applying three categories of risk (“high risk,” “some concern,” or “low risk” of bias) across the different domains of bias found in the tool. If there was disagreement between the two team members, a third team member performed an assessment using the Cochrane “Risk of Bias” tool and served as a tiebreaker.

After assessing eligible articles for homogeneity of outcomes and outcome assessment methods, a meta-analysis focused on evaluating the effect of smoking-cessation interventions on smoking

abstinence was performed using the six eligible full-text articles with extractable tabular data. Where information on both self-reported and biochemically validated smoking abstinence outcomes were available, data for the biochemically validated outcomes were used in the meta-analysis. Because both cluster and individually randomized trials were included in this systematic review, the ratio estimator approach developed by Rao and Scott¹¹ was used to adjust for clustering before estimating the pooled parameter of interest in the meta-analysis. In the ratio estimator approach, observed cell frequencies are divided by the estimated design effect of the study to account for dependence among observations belonging to the same cluster.^{11, 12} The design effect was estimated as $1 + (\bar{m} - 1)\hat{\rho}$, where $\bar{m}$ represents the average cluster size across study arms and $\hat{\rho}$ is an estimate of the intraclass correlation coefficient (ICC).¹²

For the three individually randomized trials included in the meta-analysis (Harris et al.,¹³ Wewers and colleagues,¹⁴ and Stoops et al.¹⁵), average cluster size was equal to 1 and the ICC was also equal to 1. As a result, observed cell frequencies were unchanged after adjusting for clustering in these three studies. Because average cluster size and the sample ICCs were not reported for the three cluster randomized trials, the number of participants the authors aimed to enroll in each cluster and the ICC described in the sample size estimation sections of each paper were used to estimate design effects. No information regarding the ICC was provided in Schoenberg and colleagues,¹⁶ so the ICC estimate from Wewers et al.¹⁴ was used to estimate the ICC for this study (overview provided in Appendix Table 8). Sensitivity analyses were performed to explore if using a different estimate of the ICC in Schoenberg and colleagues¹⁶ affected the results of the meta-analysis (see Appendix Table 9). The range of external ICC estimates used in the sensitivity analyses were selected from a school-based smoking prevention

trial¹⁸ and a community trial evaluating cardiovascular health that included smoking-related health outcomes.¹⁹

Fixed and random effects models with the Mantel–Haenszel method were used to estimate both the pooled RR and pooled OR of smoking abstinence. Homogeneity was assessed qualitatively using data extracted about each study’s population, intervention, and outcome (Appendix Table 10). Homogeneity was also assessed quantitatively using Cochran’s Q statistic and the I^2 index. Because both the Q statistic and I^2 index have low power to detect heterogeneity among a small number of studies,²⁰ a L’Abbé plot was also used to attempt to graphically evaluate homogeneity.^{21, 22} If there was evidence of a violation of the homogeneity assumption, the pooled effect of smoking-cessation intervention programs on smoking abstinence was estimated from a random effects model. In addition, sensitivity analyses were performed to explore if any one study had an outsize impact on the results of the meta-analysis by iteratively leaving out each trial and estimating the pooled RR of smoking abstinence.²³ Funnel plots were used to attempt to identify potential publication bias. All meta-analysis procedures were performed in R, version 3.5.0 using the package meta, version 4.9-2.²⁴

EVIDENCE SYNTHESIS

The search was conducted on October 9, 2018 and resulted in 74 records identified via MEDLINE (PubMed), 14 records identified via Embase, 24 records identified via Scopus, and three records identified via the Cochrane Library (CENTRAL).

Records were exported from each database and imported into Zotero to organize and remove

duplicates. Twenty-one duplicate records were removed using Zotero's automatic tool for detecting duplicates and an additional two duplicate entries were manually removed after sorting by article title and author names.

The remaining 92 articles were randomly assigned to four members of the research team (AK, CS, NB, SF) who independently scanned titles and abstracts to identify papers related to the research question based on the eligibility criteria. The resulting overall agreement was 85.9% (Cohen's $\kappa=0.72$), indicating a moderate level of interrater reliability.¹⁰ Of the 44 included articles, a further four articles were excluded owing to non-availability of full text. The remaining 40 articles were then subjected to a full-text review.

The full-text articles were retrieved and randomly assigned to four team members (AK, CS, NB, SF), with each article independently screened by two team members. There was 82.5% agreement (Cohen's $\kappa=0.62$) at this stage, again demonstrating moderate interrater reliability.¹⁰ All disagreements were referred to a third team member (SG) who independently read the articles and made a final decision. Thirty-one full-text articles were excluded at this stage, as these articles did not meet the predefined eligibility criteria. The most common reason for exclusion was the study population living in geographic areas outside the rural Appalachian region, or the trial being conducted in states within and outside the Appalachian region, without data separately reported such that Appalachian subjects could be isolated for this review. Two studies used quasi- and non-randomized trial designs and hence were excluded from the review. In four studies, the measured outcomes were not directly related to smoking cessation or smoking-related health (e.g., the main outcomes of interest for which data were reported were

physical activity, with smoking status only noted as a covariate), thereby leading to their exclusion. One study was excluded during the data extraction phase when it was determined to be a substudy of another trial already included. A total of nine RCTs^{13-17,25-28} evaluating the effect of smoking-cessation interventions in populations residing in rural Appalachia were included in the review. The screening process and reasons for exclusion are summarized in Figure 2.

The characteristics of the nine studies included in the review are summarized in Table 1 (with details provided in Appendix Table 10). Most studies were conducted in rural Ohio, Kentucky, Alabama, and West Virginia. Intervention outcomes included smoking-cessation rates, biochemical validation, adherence to nicotine patches, self-reported cigarettes per day, abstinence rates, and nicotine dependency scores. The number of participants in the studies varied from 17 to 707. Most of the articles were published from 2009 to 2016.

The studies were relatively heterogeneous, each using different interventions to promote smoking abstinence and each focusing on different populations in Appalachia, including pregnant women, non-pregnant women, people living with HIV/AIDS, and adolescents. Five of the studies were individually randomized whereas four were cluster randomized, each with different units of randomization (church, school, county, and physician). However, six of the nine included studies had extractable data related to dichotomous smoking status (abstinent versus not abstinent) and each reported a measure of association or provided enough information in their study publication to allow for creation of a contingency table of smoking abstinence by intervention arm. These six studies were therefore included in a quantitative meta-analysis.

The six trials of smoking-cessation intervention programs resulted in increased smoking abstinence overall; however, the effect was significant in only one study (Schoenberg et al.¹⁶). The pooled RR (Figure 3) and pooled OR (Appendix Figure 1) were estimated using random effects models because the validity of summary estimates from fixed effect models requires an assumption of homogeneity, which did not appear to be satisfied for these studies. The trials included in this meta-analysis were relatively heterogeneous with respect to specific interventions, populations, and unit of randomization (see Appendix Table 10). In addition, the Q test provided evidence of substantial heterogeneity across the six studies ($\chi^2_{(5)}=18.86$, $p<0.01$), and the I^2 index indicated that between-study variability constituted approximately 74% of the total variability across the six estimated RRs ($I^2=73.5\%$, 95% CI=39.3%, 88.4%). There was also heterogeneity in the percentage of trial participants who achieved abstinence in intervention groups (2.86%–31.07%) and control groups (0%–28.57%) among the trials (Appendix Figure 2). However, given the small number of studies, the authors also used a fixed effects model to estimate the pooled parameters, which yielded similar results to the random effects model described below (Appendix Figures 3 and 4 provide Forest plots of pooled RR and OR estimates using fixed effects).

The results of the random effects model indicated that participation in smoking-cessation interventions increased the probability of smoking abstinence among Appalachian smokers an estimated 2.33 times (RR=2.33, 95% CI=1.03, 5.25, $p=0.04$). The pooled OR estimate was also significant, but the point estimate was slightly further from the null (Appendix Figure 1; OR=2.69, 95% CI=1.05, 6.87, $p=0.04$). In the sensitivity analyses used to understand the impact

of changing the imputed ICC value for one of the three cluster randomized trials, participation in smoking-cessation intervention programs significantly increased the probability of smoking abstinence as well; as expected, the magnitude of the effectiveness of smoking-cessation programs decreased as the imputed ICC increased (Appendix Table 9).

Potential publication bias was assessed with the use of a funnel plot of the six trials included in this meta-analysis (Appendix Figure 5). Although the plot appears to be relatively symmetric, a larger number of studies would improve the ability to discern if there is evidence of publication bias. The precision of Stoops and colleagues¹⁵ was much lower than that of the other five trials.

All but one of the six trials included in the meta-analysis had at least some concerns related to bias in study design and results reporting (Appendix Table 11). This was especially true with regard to apparent deviations from intended interventions, and missing outcome data, though there were also concerns in four of six studies related to randomization processes.

DISCUSSION

In this systematic review of nine randomized trials of smoking-cessation programs with a total of 2,233 participants carried out in Appalachia, a moderately beneficial effect of various types of interventions was observed on smoking abstinence, across multiple populations.

Six trials, with a total of 1,845 participants, reported data in sufficient detail for meta-analysis to be performed. The estimated pooled RR of 2.33 indicated that smoking-cessation programs in this rural and socioeconomically disadvantaged population are an effective tool to achieve

abstinence. This reflects a number needed to treat of nine. In other words, to achieve abstinence in one participant, nine individuals need to participate in a smoking-cessation program.

These findings are in agreement with previous systematic reviews on the efficacy of smoking-cessation programs that focused on the general adult population. For example, Internet-based interventions such as used in three trials included in this review (Harris et al.,¹³ Reynolds and colleagues,²⁵ and Stoops et al.¹⁵) have been shown to be moderately more effective than non-active controls.²⁹ Similarly, individual behavioral counseling, as employed in three trials included in this review (Cropsey and colleagues,²⁶ Wewers et al.,¹⁴ and Wewers and colleagues¹⁷) has been shown to be useful in helping smokers quit.³⁰ Finally, nicotine-replacement therapy, which was employed in three trials (Wewers et al.,¹⁴ Cropsey and colleagues,²⁶ and Schoenberg et al.¹⁶), has been found to increase quit rates by 50%–60%, regardless of setting.³¹

Appalachia has one of the highest smoking prevalence rates⁵ and simultaneously one of the worst health outcomes in the U.S., resulting in a mean gap in life expectancy of 2.4 years compared with the national average.⁶ Indeed, smoking-related illnesses account for more than half of the life expectancy gap between Appalachia and the rest of the U.S.⁶ Reducing smoking prevalence should remain as a critical component of efforts to reduce health disparities in this vulnerable population.

Limitations

There are several limitations to this systematic review that merit discussion. First, there was substantial heterogeneity on several levels among the included studies. Despite the narrow

inclusion by geography, the studied populations were diverse and varied widely (e.g., some studies were adults only and others were adolescents only; others included only pregnant women, or only people living with HIV). Another important source of heterogeneity arose from varying definitions and ascertainment of smoking status and abstinence (e.g., some studies biochemically validated smoking status and others were entirely self-report). More consistency in ascertaining and defining individuals' smoking status would improve the validity of the estimates and one's ability to make inferences about them. Furthermore, the nature of the smoking-cessation interventions varied substantially, ranging from web-based interventions with economic incentives over traditional counseling to screening, brief intervention, and referral for treatment; telephone counseling, and nicotine-replacement therapy. Heterogeneity also stemmed from differing control interventions, ranging from usual care to active control interventions such as telephone counseling services (i.e., "quitlines"). The fact that the majority of trials included an active control arm rather than usual care or a sham intervention likely underestimates the effect of the smoking-cessation programs on abstinence rates.

Second, the number of studies included in this systematic review was rather low. As a consequence of the narrow geographic inclusion criterion, a substantial number of studies had to be excluded, as they pooled data across several states and did not report outcomes in sufficient regional granularity. However, as Appalachia is uniquely disadvantaged in economic and social terms that greatly influence health behaviors, a regional focus is justified. For future reviews, a larger number of studies, ideally with a greater number of participants, would help clarify and increase the certainty of the potential size of the effect estimate of smoking-cessation interventions on smoking abstinence in this area. As a result of the small number of studies

included in the meta-analysis, each study had a substantial effect on the pooled parameter estimate as illustrated by the results of the leave-one-out sensitivity analysis. Omitting any one of four studies (Schoenberg and colleagues,¹⁶ Ferketich et al.,²⁷ Wewers and colleagues,¹⁴ or Stoops et al.¹⁵) from the meta-analysis resulted in a pooled RR of smoking cessation that was not statistically significant (Appendix Figure 6).

A third limitation originated from the difficulty in pooling the results of cluster randomized trials with individually randomized trials, particularly when the ICCs and average cluster sizes were reported inadequately. This uncertainty was addressed by performing sensitivity analyses with a range of potential ICCs derived from the literature for similar outcomes, which left the pooled estimate largely unchanged. If there had been a larger number of studies that satisfied the exclusion criteria for this systematic review, pooled effect estimates could have been estimated separately for cluster and individually randomized trials or subgrouped the data by the unit of randomization to better address concerns about heterogeneity among the studies included in the meta-analysis.³²

Fourth, the majority of the studies included in this review reported outcomes over a limited follow-up period ranging from 6 to 17 weeks. Only two trials (Wewers and colleagues¹⁴ and Wewers et al.¹⁷) followed participants for 12 months. Thus, short-term follow-up restricts any predictions about the long-term efficacy of anti-smoking interventions in Appalachian smokers. Yet, given substantial relapse rates, smoking cessation must be a long-term commitment to substantially impact cardiovascular disease and cancer prevalence rates.^{33, 34}

Fifth, the trials included in this review addressed combustible tobacco only. Given the elevated rates of smokeless tobacco use (particularly among men) in rural areas³⁵ and the high co-use of other forms of tobacco,³⁶ up to 12% of nicotine users might not be included in the studies identified in this review. However, the broad term of “tobacco” employed in the search should have captured any randomized trials studying smokeless tobacco and its various products (such as chewing tobacco, dry snuff, and moist snuff [snus]). Future cessation trials should therefore aim to include smokeless tobacco users.

Finally, although the funnel plot appears symmetric, thus indicating minimal to no publication bias, this finding should be interpreted with caution given the low number of included studies. It is possible that a larger number of studies might reveal evidence for publication bias, affecting the ability to interpret the external generalizability of this review.

These limitations highlight the need for more rigorous research to identify successful smoking cessation interventions for the Appalachian population. Notably, standardized definitions of smoking status, routine ascertainment of nicotine dependence and concurrent use of other tobacco products, and standardized control arms (“quitlines”) might reduce the heterogeneity of future studies and facilitate inferences about the best interventions available.

CONCLUSIONS

The pooled estimate of the meta-analysis showed significant effectiveness of smoking-cessation interventions in rural Appalachia, and this qualitative systematic review supported this finding. This estimate is in line with previous systematic reviews in general adult populations, which

emphasize the important potential of these interventions to improve population health.

Particularly in disadvantaged populations, such as in Appalachia, successful smoking-cessation programs hold the promise to decrease health disparities. However, taken together with the low number of studies included in this meta-analysis, as well as the substantial heterogeneity and high risk of bias among these studies, the evidence of the effectiveness of smoking-cessation interventions in Appalachia appears to be limited and must be interpreted with caution.

ACKNOWLEDGMENTS

Shalika Gupta and Claudia Scheuter are co-first authors.

The authors wish to thank John M. Colford, Jr., MD, for his input in trial design methods and systematic reviews, and his encouragement to submit this work for publication, as well as Maureen Lahiff, PhD, for her comments and suggestions.

No funding was provided for this research and none of the authors have any conflicts of interest to report. No financial disclosures were reported by the authors of this paper.

REFERENCES

1. GBD 2015 Tobacco Collaborators. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990–2015: a systematic analysis from the Global Burden of Disease Study 2015. *Lancet*. 2017;389(10082):1885–1906.
[https://doi.org/10.1016/S0140-6736\(17\)30819-X](https://doi.org/10.1016/S0140-6736(17)30819-X).
2. CDC. *Current Cigarette Smoking Among U.S. Adults Aged 18 Years and Older*. Atlanta, GA: CDC; 2018.
3. Appalachian Regional Commission. *Appalachia Then and Now: Examining Changes to the Appalachian Region Since 1965*. Washington, DC: Appalachian Regional Commission; 2015.
4. Appalachian Regional Development Initiative. Economic Assessment of Appalachia.
www.arc.gov/images/programs/ardi/EconomicAssessmentofAppalachiaJune2010.pdf.
Published June 2010. Accessed September 13, 2019.
5. Pickle LW, Su Y. Within-state geographic patterns of health insurance coverage and health risk factors in the United States. *Am J Prev Med*. 2002;22(2):75–83.
[https://doi.org/10.1016/s0749-3797\(01\)00402-0](https://doi.org/10.1016/s0749-3797(01)00402-0).
6. Singh GK, Kogan MD, Slifkin RT. Widening disparities in infant mortality and life expectancy between Appalachia and the rest of the United States, 1990–2013. *Health Aff (Millwood)*. 2017;36(8):1423–1432. <https://doi.org/10.1377/hlthaff.2016.1571>.
7. HHS. *The Health Consequences of Smoking—50 years of Progress: A Report of the Surgeon General*. Atlanta, GA: HHS, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014.

- www.ncbi.nlm.nih.gov/books/NBK179276/pdf/Bookshelf_NBK179276.pdf. Accessed September 13, 2019.
8. Higgins J, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration; 2011.
 9. Appalachian Regional Commission. Counties in Appalachia. www.arc.gov/appalachian_region/CountiesinAppalachia.asp. Published 2019. Accessed September 13, 2019.
 10. McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med (Zagreb)*. 2012;22(3):276–282. <https://doi.org/10.11613/bm.2012.031>.
 11. Rao JN, Scott AJ. A simple method for the analysis of clustered binary data. *Biometrics*. 1992;48(2):577–585. <https://doi.org/10.2307/2532311>.
 12. Donner A, Klar N. Issues in the meta-analysis of cluster randomized trials. *Stat Med*. 2002;21(19):2971–2980. <https://doi.org/10.1002/sim.1301>.
 13. Harris M, Reynolds B. A pilot study of home-based smoking cessation programs for rural, Appalachian, pregnant smokers. *J Obstet Gynecol Neonatal Nurs*. 2015;44(2):236–45. <https://doi.org/10.1111/1552-6909.12547>.
 14. Wewers ME, Ferketich AK, Harness J, Paskett ED. Effectiveness of a nurse-managed, lay-led tobacco cessation intervention among Ohio Appalachian women. *Cancer Epidemiol Biomarkers Prev*. 2009;18(12):3451–3458. <https://doi.org/10.1158/1055-9965.epi-09-0952>.
 15. Stoops WW, Dallery J, Fields NM, et al. An internet-based abstinence reinforcement smoking cessation intervention in rural smokers. *Drug Alcohol Depend*. 2009;105(1–2):56–62. <https://doi.org/10.1016/j.drugalcdep.2009.06.010>.

16. Schoenberg NE, Studts CR, Shelton BJ, et al. A randomized controlled trial of a faith-placed, lay health advisor delivered smoking cessation intervention for rural residents. *Prev Med Rep.* 2016;3:317–323. <https://doi.org/10.1016/j.pmedr.2016.03.006>.
17. Wewers ME, Shoben A, Conroy S, et al. Effectiveness of two community health worker models of tobacco dependence treatment among community residents of Ohio Appalachia. *Nicotine Tob Res.* 2017;19(12):1499–1507. <https://doi.org/10.1093/ntr/ntw265>.
18. Siddiqui O, Hedeker D, Flay BR, Hu FB. Intraclass correlation estimates in a school-based smoking prevention study: outcome and mediating variables, by sex and ethnicity. *Am J Epidemiol.* 1996;144(4):425–433. <https://doi.org/10.1093/oxfordjournals.aje.a008945>.
19. Hannan PJ, Murray DM, Jacobs DR Jr, McGovern PG. Parameters to aid in the design and analysis of community trials: intraclass correlations from the Minnesota Heart Health Program. *Epidemiology.* 1994;5(1):88–95. <https://doi.org/10.1097/00001648-199401000-00013>.
20. Huedo-Medina TB, Sanchez-Meca, J, Marin-Martinez F, Botella J. Assessing heterogeneity in meta-analysis: Q statistic or I2 index? *Psychol Methods.* 2006;11(2):193–206. <https://doi.org/10.1037/1082-989x.11.2.193>.
21. L'Abbe KA, Detsky AS, O'Rourke K. Meta-analysis in clinical research. *Ann Intern Med.* 1987;107(2):224–233. <https://doi.org/10.7326/0003-4819-107-2-224>.
22. Song F. Exploring heterogeneity in meta-analysis: is the L'Abbe plot useful? *J Clin Epidemiol.* 1999;52(8):725–730. [https://doi.org/10.1016/s0895-4356\(99\)00066-9](https://doi.org/10.1016/s0895-4356(99)00066-9).

23. Rothman KJ, Greenland S, Lash TL. *Modern Epidemiology*. Philadelphia, PA: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2008.
24. Schwarzer G. meta: An R package for meta-analysis. *R News*. 2007;7(3):40–45.
25. Reynolds B, Harris M, Slone SA, et al. A feasibility study of home-based contingency management with adolescent smokers of rural Appalachia. *Exp Clin Psychopharmacol*. 2015;23(6):486–493. <https://doi.org/10.1037/pha0000046>.
26. Cropsey KL, Hendricks PS, Jardin B, Clark CB, Katiyar N, Willig J, et al. A pilot study of screening, brief intervention, and referral for treatment (SBIRT) in non-treatment seeking smokers with HIV. *Addict Behav*. 2013;38(10):2541–2546. <https://doi.org/10.1016/j.addbeh.2013.05.003>.
27. Ferketich AK, Pennell M, Seiber EE, et al. Provider-delivered tobacco dependence treatment to Medicaid smokers. *Nicotine Tob Res*. 2014;16(6):786–793. <https://doi.org/10.1093/ntr/ntt221>.
28. Horn K, Branstetter S, Zhang J, et al. Understanding physical activity outcomes as a function of teen smoking cessation. *J Adolesc Health*. 2013;53(1):125–131. <https://doi.org/10.1016/j.jadohealth.2013.01.019>.
29. Taylor GMJ, Dalili MN, Semwal M, Civljak M, Sheikh A, Car J. Internet-based interventions for smoking cessation. *Cochrane Database Syst Rev*. 2017;9:CD007078. <https://doi.org/10.1002/14651858.cd007078.pub5>.
30. Lancaster T, Stead LF. Individual behavioural counselling for smoking cessation. *Cochrane Database Syst Rev*. 2017;3:CD001292. <https://doi.org/10.1002/14651858.CD001292.pub3>.

31. Hartmann-Boyce J, Chepkin SC, Ye W, Bullen C, Lancaster T. Nicotine replacement therapy versus control for smoking cessation. *Cochrane Database Syst Rev*. 2018;5:CD000146. <https://doi.org/10.1002/14651858.cd000146.pub5>.
32. Richardson M, Garner P, Donegan S. Cluster randomised trials in Cochrane Reviews: evaluation of methodological and reporting practice. *PLoS One*. 2016;11(3):e0151818. <https://doi.org/10.1371/journal.pone.0151818>.
33. Cornuz J. Smoking cessation interventions in clinical practice. *Eur J Vasc Endovasc Surg*. 2007;34(4):397–404. <https://doi.org/10.1016/j.ejvs.2007.06.009>.
34. The Clinical Practice Guideline Treating Tobacco Use and Dependence 2008 Update Panel, Liaisons, and Staff. A clinical practice guideline for treating tobacco use and dependence: 2008 update: a U.S. Public Health Service report. *Am J Prev Med*. 2008;35(2):158–176. <https://doi.org/10.1016/j.amepre.2008.04.009>.
35. Roberts ME, Doogan NJ, Stanton CA, et al. Rural versus urban use of traditional and emerging tobacco products in the United States, 2013–2014. *Am J Public Health*. 2017;107(10):1554–1559. <https://doi.org/10.2105/ajph.2017.303967>.
36. Cheng YC, Rostron BL, Day HR, et al. Patterns of use of smokeless tobacco in U.S. adults, 2013–2014. *Am J Public Health*. 2017;107(9):1508–1514. <https://doi.org/10.2105/ajph.2017.303921>.

LIST OF FIGURES

Figure 1. Map of the Appalachian region (as defined by the Appalachian Regional Commission³).

Figure 2. Flowchart of search process and results.


Figure 3. Results of random effects meta-analysis of smoking cessation intervention effects on smoking abstinence (RRs).

Table 1. Summary of Eligible Study Characteristics


Study characteristics	Number of studies	Total participants
Type of randomization		
Individual	5	489
Cluster	4	1,744
Bias assessment		
Low	1	590
Some concerns	3	1,077
High	2	231
Not assessed	3	335
Study outcomes ^a		
Self-reported smoking abstinence	8	2,165
Biochemical validation of smoking abstinence	6	1,370
Nicotine dependence (e.g., Fagerstrom)	4	709
Reported measure of association ^a		
OR	6	1,898
RR	0	0
Other (difference in means, eta-squared)	3	692
None	1	233
Total	9	2,233

^aStudies reported multiple outcomes/measures; total >9.

1




2



3

Source	RR (95% CI)
Schoenberg et al. 2016	9.57 [3.75; 24.41]
Harris et al. 2015	1.05 [0.23; 4.73]
Ferketich et al. 2014	3.19 [0.70; 14.62]
Wewers et al. 2009	1.73 [0.84; 3.53]
Wewers et al. 2017	1.24 [0.79; 1.96]
Stoops et al. 2009	2.83 [0.12; 67.10]
Total	2.33 [1.03; 5.25]
Heterogeneity: $\chi^2_5 = 18.86$ ($P < .01$), $I^2 = 73.5\%$	



Appendix Table 1. Eligibility Criteria Framed in Terms of the PICO's Framework

Item	Detailed eligibility criteria
P - patients	● Currently smoking individuals of any age and gender ● Appalachia: definition according to the 2008 Appalachian Regional Commission
I - intervention	● Any type of program designed to promote smoking cessation, including those administered in person and online ● Any settings, including community-based and hospital-based smoking cessation interventions
C - comparison	● Include studies that look at smoking cessation programs as the primary intervention of interest as well as studies that consider smoking cessation along with other health-related interventions
O - outcomes	● Any smoking-related behavioral change or health outcome as the primary or secondary outcomes of interest for the study ● Specific outcomes: smoking cessation rate; type, frequency, or duration of smoking
S - study designs	● Randomized trials, where the unit of randomization can be individuals or clusters ● Excluding observational studies ● English language only

Appendix Table 2. Search Sets and Terms Used for Title, Abstract, and MeSH Terms/Author Keyword Searches

Location	Type of study	Intervention focus
Appalachia or Appalachian -OR- Rural AND West Virginia or Alabama or Georgia or Kentucky or Maryland or Mississippi or New York or North Carolina or Ohio or Pennsylvania or South Carolina or Tennessee or Virginia	Trial or Randomized or Randomised	Smoking or Tobacco or Smoker or Smokers or Cigarette or Cigarettes or E-cigarette or E-cigarettes or Electronic cigarettes or Electronic cigarette Vaping AND Cessation -OR- Anti-smoking

Appendix Table 3. Search Code

Database	Code
MEDLINE (PubMed)	<pre> ((((((((((((((((((((smoking[Title/Abstract]) OR smoking[MeSH Terms]) OR tobacco[Title/Abstract]) OR tobacco[MeSH Terms]) OR smoker[Title/Abstract]) OR smoker[MeSH Terms]) OR smokers[Title/Abstract]) OR smokers[MeSH Terms]) OR cigarette[Title/Abstract]) OR cigarette[MeSH Terms]) OR cigarettes[Title/Abstract]) OR cigarettes[MeSH Terms]) OR e-cigarette[Title/Abstract]) OR e-cigarette[MeSH Terms]) OR e-cigarettes[Title/Abstract]) OR e-cigarettes[MeSH Terms]) OR electronic cigarettes[Title/Abstract]) OR electronic cigarettes[MeSH Terms]) OR electronic cigarette[Title/Abstract]) OR electronic cigarette[MeSH Terms]) OR vaping[Title/Abstract]) OR vaping[MeSH Terms]) AND ((cessation[Title/Abstract]) OR cessation[MeSH Terms])) OR ((anti- smoking[Title/Abstract]) OR anti-smoking[MeSH Terms])) AND (((((trial[Title/Abstract]) OR trial[MeSH Terms]) OR randomized[Title/Abstract]) OR randomized[MeSH Terms]) OR randomised[Title/Abstract]) OR randomised[MeSH Terms]) AND (((((Appalachia[Title/Abstract]) OR Appalachia[MeSH Terms]) OR Appalachian[Title/Abstract]) OR Appalachian[MeSH Terms]) OR ((((((((((((((((((((West Virginia[Title/Abstract]) OR West Virginia[MeSH Terms]) OR Alabama[Title/Abstract]) OR Alabama[MeSH Terms]) OR Georgia[Title/Abstract]) OR Georgia[MeSH Terms]) OR Kentucky[Title/Abstract]) OR Kentucky[MeSH Terms]) OR Maryland[Title/Abstract]) OR Maryland[MeSH Terms]) OR Mississippi[Title/Abstract]) OR Mississippi[MeSH Terms]) OR New York[Title/Abstract]) OR New York[MeSH Terms]) OR North Carolina[Title/Abstract]) OR North Carolina[MeSH Terms]) OR Ohio[Title/Abstract]) OR Ohio[MeSH Terms]) OR Pennsylvania[Title/Abstract]) OR Pennsylvania[MeSH Terms]) OR South Carolina[Title/Abstract]) OR South Carolina[MeSH Terms]) OR Tennessee[Title/Abstract]) OR Tennessee[MeSH Terms]) OR Virginia[Title/Abstract]) OR Virginia[MeSH Terms]) AND ((rural[Title/Abstract]) OR rural[MeSH Terms])]]]])) </pre>
Embase	<pre> (('smoking':ti,ab,kw OR 'tobacco':ti,ab,kw OR 'smoker':ti,ab,kw OR 'smokers':ti,ab,kw OR 'cigarette':ti,ab,kw OR 'cigarettes':ti,ab,kw OR 'e-cigarette':ti,ab,kw OR 'e-cigarettes':ti,ab,kw OR 'electronic cigarettes':ti,ab,kw OR 'electronic cigarette':ti,ab,kw OR 'vaping':ti,ab,kw) AND 'cessation':ti,ab,kw OR 'anti-smoking':ti,ab,kw) AND ('trial':ti,ab,kw OR 'randomized':ti,ab,kw OR 'randomised':ti,ab,kw) AND ('appalachia':ti,ab,kw OR 'appalachian':ti,ab,kw OR (('west virginia':ti,ab,kw OR 'alabama':ti,ab,kw OR 'georgia':ti,ab,kw OR 'kentucky':ti,ab,kw OR 'maryland':ti,ab,kw OR 'mississippi':ti,ab,kw OR 'new york':ti,ab,kw OR 'north carolina':ti,ab,kw OR 'ohio':ti,ab,kw OR </pre>

'pennsylvania':ti,ab,kw OR 'south carolina':ti,ab,kw OR 'tennessee':ti,ab,kw OR 'virginia':ti,ab,kw) AND 'rural':ti,ab,kw))

Scopus

((((TITLE-ABS-KEY (west AND virginia) OR TITLE-ABS-KEY (alabama) OR TITLE-ABS-KEY (georgia) OR TITLE-ABS-KEY (kentucky) OR TITLE-ABS-KEY (maryland) OR TITLE-ABS-KEY (mississippi) OR TITLE-ABS-KEY (new AND york) OR TITLE-ABS-KEY (north AND carolina) OR TITLE-ABS-KEY (ohio) OR TITLE-ABS-KEY (pennsylvania) OR TITLE-ABS-KEY (south AND carolina) OR TITLE-ABS-KEY (tennessee) OR TITLE-ABS-KEY (virginia))) AND ((TITLE-ABS-KEY (rural)))) OR ((TITLE-ABS-KEY (appalachia) OR TITLE-ABS-KEY (appalachian)))) AND ((TITLE-ABS-KEY (trial) OR TITLE-ABS-KEY (randomized) OR TITLE-ABS-KEY (randomised))) AND ((((TITLE-ABS-KEY (smoking) OR TITLE-ABS-KEY (tobacco) OR TITLE-ABS-KEY (smoker) OR TITLE-ABS-KEY (smokers) OR TITLE-ABS-KEY (cigarette) OR TITLE-ABS-KEY (cigarettes) OR TITLE-ABS-KEY (e-cigarette) OR TITLE-ABS-KEY (e-cigarettes) OR TITLE-ABS-KEY (electronic AND cigarettes) OR TITLE-ABS-KEY (electronic AND cigarette) OR TITLE-ABS-KEY (taping)))) AND (TITLE-ABS-KEY (cessation)))) OR (TITLE-ABS-KEY (anti-smoking))))

Cochrane Library
(CENTRAL)

(((((smoking in Title Abstract Keyword OR tobacco in Title Abstract Keyword OR smoker in Title Abstract Keyword OR smokers in Title Abstract Keyword OR cigarette in Title Abstract Keyword OR cigarettes in Title Abstract Keyword OR e-cigarette in Title Abstract Keyword OR e-cigarettes in Title Abstract Keyword OR electronic cigarettes in Title Abstract Keyword OR electronic cigarette in Title Abstract Keyword OR vaping in Title Abstract Keyword) AND (cessation in Title Abstract Keyword)) OR (anti-smoking in Title Abstract Keyword)) AND (trial in Title Abstract Keyword OR randomized in Title Abstract Keyword OR randomised in Title Abstract Keyword) AND ((Appalachia in Title Abstract Keyword OR Appalachian in Title Abstract Keyword) OR ((West Virginia in Title Abstract Keyword OR Alabama in Title Abstract Keyword OR Georgia in Title Abstract Keyword OR Kentucky in Title Abstract Keyword OR Maryland in Title Abstract Keyword OR Mississippi in Title Abstract Keyword OR New York in Title Abstract Keyword OR North Carolina in Title Abstract Keyword OR Ohio in Title Abstract Keyword OR Pennsylvania in Title Abstract Keyword OR South Carolina in Title Abstract Keyword OR Tennessee in Title Abstract Keyword OR Virginia in Title Abstract Keyword) AND (rural in Title Abstract Keyword))))))

Appendix Table 4. Records Identified via MEDLINE (PubMed)

Item number	Search terms	Number of records
1	West Virginia or Alabama or Georgia or Kentucky or Maryland or Mississippi or New York or North Carolina or Ohio or Pennsylvania or South Carolina or Tennessee or Virginia	223,561
2	Rural	124,312
3	Appalachia or Appalachian	127,944
Set 1 - location	(#1 AND #2) OR #3	129,387
Set 2 - type of study	Trial or Randomized or Randomised	1,067,203
5	Smoking or tobacco or Smoker or Smokers or Cigarette or Cigarettes or E-cigarette or or E-cigarettes or Electronic cigarettes or Electronic cigarette or Vaping	329,986
6	Cessation	67,431
7	Anti-smoking	1,191
Set 3 - intervention focus	(#5 AND #6) OR #7	29,116
Set 4 - TOTAL SEARCH	Set 1 AND Set 2 AND Set 3	74

Appendix Table 5. Records Identified via EMBASE

Item number	Search terms	Number of records
1	West Virginia or Alabama or Georgia or Kentucky or Maryland or Mississippi or New York or North Carolina or Ohio or Pennsylvania or South Carolina or Tennessee or Virginia	200,952
2	Rural	151,179
3	Appalachia or Appalachian	2,067
Set 1 - location	(#1 AND #2) OR #3	7,170
Set 2 - type of study	Trial or Randomized or Randomised	1,163,018
5	Smoking or tobacco or Smoker or Smokers or Cigarette or Cigarettes or E-cigarette or or E-cigarettes or Electronic cigarettes or Electronic cigarette or Vaping	409,150
6	Cessation	93,567
7	Anti-smoking	1,540
Set 3 - intervention focus	(#5 AND #6) OR #7	37,394
Set 4 - TOTAL SEARCH	Set 1 AND Set 2 AND Set 3	14

Appendix Table 6. Records Identified via Scopus

Item number	Search terms	Number of records
1	West Virginia or Alabama or Georgia or Kentucky or Maryland or Mississippi or New York or North Carolina or Ohio or Pennsylvania or South Carolina or Tennessee or Virginia	792,770
2	Rural	332,790
3	Appalachia or Appalachian	13,341
Set 1 - location	(#1 AND #2) OR #3	25,435
Set 2 - type of study	Trial or Randomized or Randomised	2,282,755
5	Smoking or tobacco or Smoker or Smokers or Cigarette or Cigarettes or E-cigarette or or E-cigarettes or Electronic cigarettes or Electronic cigarette or Vaping	480,795
6	Cessation	115,551
7	Anti-smoking	1,679
Set 3 - intervention focus	(#5 AND #6) OR #7	60,511
Set 4 - TOTAL SEARCH	Set 1 AND Set 2 AND Set 3	24

Appendix Table 7. Records Identified via Cochrane Library (CENTRAL)

Item number	Search terms	Number of records
1	West Virginia or Alabama or Georgia or Kentucky or Maryland or Mississippi or New York or North Carolina or Ohio or Pennsylvania or South Carolina or Tennessee or Virginia	140
2	Rural	47
3	Appalachia or Appalachian	1
Set 1 - location	(#1 AND #2) OR #3	19
Set 2 - type of study	Trial or Randomized or Randomised	852
5	Smoking or tobacco or Smoker or Smokers or Cigarette or Cigarettes or E-cigarette or E-cigarettes or Electronic cigarettes or Electronic cigarette or Vaping	121
6	Cessation	101
7	Anti-smoking	9
Set 3 - intervention focus	(#5 AND #6) OR #7	56
Set 4 - TOTAL SEARCH	Set 1 AND Set 2 AND Set 3	3

Appendix Table 8. Quantities Used to Estimate Design Effects in Cluster Randomized Trials

Source	Unit of randomization	Estimated average cluster size	Estimated ICC
Schoenberg et al. 2016	Church	30	Not provided (used 0.005)
Ferketich et al. 2014	Physician	30	0.03
Wewers et al. 2017	County	59	0.005

ICC, Intraclass Correlation Coefficient.

Appendix Table 9. Results of Meta-Analysis Using Different Values to Estimate the ICC in Schoenberg 2016

Imputed ICC	Pooled RR estimate^a	95% CI	<i>p</i>
0.005 ^b	2.33	1.03, 5.25	0.04
0.01	2.30	1.05, 5.04	0.04
0.05	2.03	1.07, 3.83	0.03
0.10	1.77	1.05, 2.98	0.03

^aUsing a random effects model.

^bICC value used in the primary analysis.

ICC, Intraclass Correlation Coefficient.

Appendix Table 10. Details About the Included/Eligible Studies²²⁻³⁰

First author	Pub. year	Years of data collection	Study setting	Participant number	Intervention(s)	Study duration (weeks)	Study outcome(s)	How measured	Reported measure
Schoenberg NE	2016	2013–2016	Rural Appalachian Kentucky	590 enrolled 537 completed the study and were thus included in analysis	12-week behavioral smoking cessation program	17	smoking status nicotine dependence	self-report, dichotomous direct measure (Fagerstrom), continuous	OR difference in means
Harris M	2015	not reported	Rural Appalachia Ohio and Kentucky	17	web-based contingency management program compared with phone-delivered cessation counseling program	not reported	smoking abstinence reduction in smoking by > 50% breath CO and urinary cotinine nicotine dependence	self-report, dichotomous self-report, dichotomous direct verification of self-report, dichotomous direct measure (Fagerstrom), continuous	OR OR OR none


Ferketich AK	2014	not reported	Appalachian Ohio	214	12 weeks of phone counseling from a trained treatment specialist and systems-level physician training	36	smoking abstinence cotinine-confirmed abstinence	self-report, dichotomous direct, dichotomous	OR (adj.) OR (crude)
Cropsey KL	2013	2011	Birmingham, AL	40	8-weeks of nicotine replacement therapy plus Screening, Brief Intervention, and Referral for Treatment	not reported	cigarettes smoked per day physical nicotine dependence	self-report, continuous self-report and direct (Fagerstrom), continuous	eta-squared eta-squared
Horn K	2013	2006–2009	West Virginia	233	Not-On-Tobacco (N-O-T) teen cessation program; second intervention arm received N-O-T plus a physical activity module	36	smoking abstinence total number of cigarettes smoked per day	self-report, dichotomous self-report, continuous	no point estimate or CI reported, only “no sig. difference”
Wewers	200	2005–	Rural	302	Nurse-	52	smoking	self-report,	OR (adj.

ME	9	2009	Appalachian Ohio		managed, lay-led protocol that incorporated nicotine replacement and behavioral counseling.		abstinence	dichotomous	and unadj.) for verified abstinence
							cotinine-confirmed abstinence	direct verification of self-report, dichotomous	
Wewers ME	2017	2010–2014	12 Ohio Appalachian counties	707	1) community health worker (CHW) face-to-face counseling; vs. 2) CHW phone quitline	52	smoking abstinence at 12 months	self-report, with biochemical validation, dichotomous	OR
Reynolds B	2015	not reported	Appalachian Kentucky, Ohio, and West Virginia	62	Web-based contingency management (CM) program	12	cigarettes per day	self-report, continuous	difference in means
							cotinine values	direct, continuous	difference in means
							breath CO level	direct, continuous	difference in means
							nicotine dependence	direct measure (Fagerstrom), continuous	
Stoops WW	2009	not reported	Appalachian Kentucky	68	Internet-based abstinence	12	breath CO level	direct, dichotomous	OR


reinforcement
intervention


Appendix Table 11. Risk of Bias Analysis


Author, year	Bias arising from the randomization process	Bias due to timing of identification and recruitment of individuals in relation to timing of randomization (cluster RCTs only)	Bias due to deviations from intended interventions	Bias due to missing outcome data	Bias in measurement of the outcome	Bias in selection of the reported result	Overall judgement for risk of bias
Ferketich, 2014	some concerns	low	high	low	low	low	high
Harris, 2015	low	n/a	high	high	some concerns	some concerns	high
Schoenberg, 2016	low	low	low	low	low	low	low
Stoops, 2009	some concerns	n/a	some concerns	high	low	some concerns	high
Wewers, 2009	some concerns	n/a	some concerns	high	low	some concerns	high
Wewers, 2017	some concerns	low	some concerns	low	low	low	some concerns

Appendix Figure 1. Forest plot of pooled OR estimates using a random effects model.


Appendix Figure 2. L'abbé plot where point size corresponds to the magnitude of the weights applied to each study in the random effects meta-analysis using the Mantel-Haenszel method.




Appendix Figure 3. Forest plot of pooled risk ratio estimates using a fixed effects model.

Appendix Figure 4. Forest plot of pooled OR estimates using a fixed effects model.

Appendix Figure 5. Funnel plot of trial results.



Appendix Figure 6. Forest plot of results from leave-one-out analysis.

Notes: In the leave-one-out analyses, estimated pooled RRs ranged from 1.42 to 2.85. In two of the six analyses, the probability of smoking abstinence was significantly higher in the intervention groups than the control groups.

APPENDIX REFERENCES

1. Schoenberg NE, Studts CR, Shelton BJ, et al. A randomized controlled trial of a faith-placed, lay health advisor delivered smoking cessation intervention for rural residents. *Prev Med Rep.* 2016;3:317–323. <https://doi.org/10.1016/j.pmedr.2016.03.006>.
2. Harris M, Reynolds B. A pilot study of home-based smoking cessation programs for rural, Appalachian, pregnant smokers. *J Obstet Gynecol Neonatal Nurs.* 2015;44(2):236–45. <https://doi.org/10.1111/1552-6909.12547>.
3. Ferketich AK, Pennell M, Seiber EE, et al. Provider-delivered tobacco dependence treatment to Medicaid smokers. *Nicotine Tob Res.* 2014;16(6):786–793. <https://doi.org/10.1093/ntr/ntt221>.
4. Cropsey KL, Hendricks PS, Jardin B, Clark CB, Katiyar N, Willig J, et al. A pilot study of screening, brief intervention, and referral for treatment (SBIRT) in non-treatment seeking smokers with HIV. *Addict Behav.* 2013;38(10):2541–2546. <https://doi.org/10.1016/j.addbeh.2013.05.003>.
5. Horn K, Branstetter S, Zhang J, et al. Understanding physical activity outcomes as a function of teen smoking cessation. *J Adolesc Health.* 2013;53(1):125–131. <https://doi.org/10.1016/j.jadohealth.2013.01.019>.
6. Wewers ME, Ferketich AK, Harness J, Paskett ED. Effectiveness of a nurse-managed, lay-led tobacco cessation intervention among Ohio Appalachian women. *Cancer Epidemiol Biomarkers Prev.* 2009;18(12):3451–3458. <https://doi.org/10.1158/1055-9965.epi-09-0952>.

7. Wewers ME, Shoben A, Conroy S, et al. Effectiveness of two community health worker models of tobacco dependence treatment among community residents of Ohio Appalachia. *Nicotine Tob Res.* 2017;19(12):1499–1507.
<https://doi.org/10.1093/ntr/ntw265>.
8. Reynolds B, Harris M, Slone SA, et al. A feasibility study of home-based contingency management with adolescent smokers of rural Appalachia. *Exp Clin Psychopharmacol.* 2015;23(6):486–493. <https://doi.org/10.1037/pha0000046>.
9. Stoops WW, Dallery J, Fields NM, et al. An internet-based abstinence reinforcement smoking cessation intervention in rural smokers. *Drug Alcohol Depend.* 2009;105(1–2):56–62. <https://doi.org/10.1016/j.drugalcdep.2009.06.010>.