

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Coupled Sensing of Hunger and Thirst Signals Balances Sugar and Water Consumption

### Permalink

<https://escholarship.org/uc/item/74p1n6f4>

### Journal

Cell, 166(4)

### ISSN

0092-8674

### Authors

Jourjine, Nicholas  
Mullaney, Brendan C  
Mann, Kevin  
et al.

### Publication Date

2016-08-01

### DOI

10.1016/j.cell.2016.06.046

Peer reviewed



# HHS Public Access

Author manuscript

Cell. Author manuscript; available in PMC 2017 August 11.

Published in final edited form as:

Cell. 2016 August 11; 166(4): 855–866. doi:10.1016/j.cell.2016.06.046.

## Coupled sensing of hunger and thirst signals balances sugar and water consumption

Nicholas Jourjine<sup>#1</sup>, Brendan C. Mullaney<sup>#1</sup>, Kevin Mann<sup>1,#</sup>, and Kristin Scott<sup>1,3</sup>

<sup>1</sup>Department of Molecular and Cell Biology and Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA 94720

<sup>#</sup> These authors contributed equally to this work.

### Abstract

Hunger and thirst are ancient homeostatic drives for food and water consumption. Although molecular and neural mechanisms underlying these drives are currently being uncovered, less is known about how hunger and thirst interact. Here, we use molecular genetic, behavioral, and anatomical studies in *Drosophila* to identify four neurons that modulate food and water consumption. Activation of these neurons promotes sugar consumption and restricts water consumption, whereas inactivation promotes water consumption and restricts sugar consumption. By calcium imaging studies, we show that these neurons are directly regulated by a hormone signal of nutrient levels and by osmolality. Finally, we identify a hormone receptor and an osmolality-sensitive ion channel that underlie this regulation. Thus, a small population of neurons senses internal signals of nutrient and water availability to balance sugar and water consumption. Our results suggest an elegant mechanism by which interoceptive neurons oppositely regulate homeostatic drives to eat and drink.

### Graphical abstract

---


<sup>3</sup> kscott@berkeley.edu.

<sup>#</sup>present address: Stanford University, Neurobiology Dept., Stanford, CA 94305

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

#### Author Contributions

N.J. performed the RNAi screen and characterized the role of Nanchung. B.M. performed the neural activation screen and characterized the role of AKHR. N.J. and B.M. performed calcium imaging, behavioral, and osmolality measurement studies. K.M. performed electrophysiology. N.J., B.M., and K.S. wrote the manuscript and K.S. supervised the study.



## Introduction

To achieve homeostasis, animals must regulate consumption of external nutrients based on internal metabolic needs. Remarkably, animals whose nervous systems differ dramatically in organization exhibit some of the same homeostatic consumption behaviors, such as orally consuming food or water in response to starvation or dehydration. While these similarities suggest conserved underlying mechanisms, how nervous systems regulate consumption in a manner that reflects internal state remains an open question.

A key requirement for homeostatic regulation is the ability to sense internal nutrient abundance and either promote consumption in nutrient-deprived states or inhibit consumption in nutrient-replete states. In mammals, a major site of internal nutrient sensing is the hypothalamus, a conserved forebrain region that senses nutrients and coordinates behavioral responses to changes in their abundance (Sternson, 2013). Neurons in the hypothalamus include direct sensors of circulating sugars like glucose, as well as sensors of metabolic cues, like insulin, ghrelin, and glucagon. The hypothalamus also contains neurons whose activity is regulated by extracellular osmolality and are therefore sensitive to internal water abundance (Bourque, 2008). However, how the nervous system uses information encoded by hypothalamic sensors to regulate consumption of food and water remains unresolved.

Like mammals, the fruit fly *Drosophila melanogaster* regulates consumption of food and water depending on internal metabolic state. Although flies lack a direct homolog of the hypothalamus, neural populations in the *Drosophila* brain function as internal nutrient sensors, including glucose, fructose, and amino acid sensors that regulate feeding decisions (Dus et al., 2015; Bjordal et al., 2014; Miyamoto et al., 2012). Flies also regulate water consumption based on internal water abundance (Dethier, 1976), although internal sensors underlying this behavior have not previously been characterized. Thus, mammals and insects both regulate food and water consumption based on internal metabolic state, and in many instances accomplish this regulation by similar mechanisms. However, the neurons and molecules that regulate homeostatic consumption remain incompletely understood. In

particular, mechanisms that coordinate the consumption of different essential nutrients, such as sugar or water, have been largely unexplored.

Here we report findings from two behavioral screens for neurons that regulate food or water consumption in *Drosophila*. Surprisingly, these screens independently identified the same four neurons as regulators of both food and water consumption. The neurons are located in the subesophageal zone (SEZ), a key relay for feeding regulation in the fly brain, and we name them Interoceptive SEZ Neurons (ISNs). Using genetic tools, behavioral assays, and calcium imaging, we show that ISNs are sensitive both to an internal signal of nutrient deprivation, the glucagon-like peptide adipokinetic hormone (AKH), and an internal signal of water abundance, extracellular osmolality. We identify the G-protein coupled receptor, adipokinetic hormone receptor (AKHR), and a conserved TRPV channel, Nanchung (Nan), as underlying responses to AKH and osmolality, respectively. Finally, we show that ISNs oppositely regulate sugar and water consumption, suggesting that they function to restore internal homeostasis. The convergence of internal signals of nutrient and water availability onto interoceptive neurons suggests an unexpected principle by which the nervous system might coordinate homeostatic behaviors.

## Results

### A behavioral screen for neurons that regulate feeding

To identify neurons that regulate feeding, we transiently activated candidate neurons with the heat-activated cation channel, dTRPA1, and determined the effect on feeding in adult flies. Consumption was monitored by scoring the amount of blue dye in the abdomens of flies with access to 200mM sucrose containing blue dye for 30 minutes. Approximately 600 Gal4 lines from the InSite collection (Gohl et al., 2011) with expression in the central nervous system were crossed to flies carrying the *UAS-dTRPA1* transgene (Hamada et al., 2008), allowing for Gal4-dependent neural activation. Fed flies were tested for sucrose consumption during heat-induced depolarization of Gal4-expressing neurons (Figure 1A; Table S1 reports transgenic flies used in this study). Four lines exhibited dramatically increased feeding, with the *954-Gal4* line showing the strongest consumption.

### Four neurons in the *954-Gal4* line promote ingestive behaviors to sucrose

To identify neurons causal for increased feeding in the *954-Gal4* line, we began by characterizing its expression pattern. In the central brain, *954-Gal4* drove *UAS-mCD8::GFP* expression in neurons of the *pars intercerebralis*, dorsal lateral protocerebrum, subesophageal zone (SEZ), and ventral nerve cord (VNC) (Figure 1B). To identify which neurons contribute to the feeding phenotype of the *954-Gal4* line, we employed an intersectional approach using the Gal4 inhibitor Gal80 to restrict Gal4 expression to smaller neural populations. The *Tshirt-Gal80* transgene (Clyne and Miesenbock, 2008) blocked Gal4 expression in the VNC but did not eliminate increased consumption upon dTRPA1 activation of *954-Gal4* neurons, demonstrating that VNC neurons are not required (Figure S1A). Next, we identified one line, *149-Gal80* (Gordon, M.D., unpublished), that eliminated SEZ GFP expression without affecting *pars intercerebralis* or dorsal lateral protocerebrum

expression (Figure S1B). This transgene eliminated the increased feeding phenotype in the *954-Gal4* line, arguing that *954-Gal4* SEZ neurons are necessary for increased consumption.

We screened existing Gal4 collections (Jenett et al., 2012; Dickson, B.J., unpublished) to identify lines that exhibit Gal4 expression in these SEZ neurons and identified two Gal4 lines, *R34G02* and *VT011155*, that labeled neurons that resembled the *954-Gal4* SEZ cluster. The *R34G02* line also drove expression in a pair of VNC neurons and abdominal ganglia projections (Figure 1C). Remarkably, the *VT011155* line exclusively labeled the SEZ neurons (Figure 1D). Both lines exhibited the feeding phenotype identified in *954-Gal4* (Figure 1E-G). In addition, activation of *954-Gal4*, *R34G02* or *VT011155* neurons with *UAS-dTRPA1* increased rates of proboscis extension to sugar stimuli, a non-ingestive behavior that flies exhibit to an appetitive taste stimulus (Figure S1C). We generated a *R34G02-LexA* line and performed double labeling experiments with the Gal4 lines to test for co-expression and found that *954-Gal4*, *VT011155*, and *R34G02-Gal4* are all co-expressed in the same two SEZ neurons per hemisphere, with *954-Gal4* also showing expression in two additional SEZ neurons per hemisphere (Figure S1D, E). These findings demonstrate that activation of four neurons in the SEZ causes increased feeding and promotes proboscis extension in well-fed animals. We name these interoceptive SEZ neurons (ISNs) and the *VT011155* line which specifically labels them *ISN-Gal4*.

### ISNs directly respond to AKH and are indirectly inhibited by insulin

Pre- and post-synaptic sites on ISNs overlap in the dorsal SEZ (Figure S1F). ISNs might therefore be components of feeding sensorimotor circuits, or they may modulate activity in these circuits in response to internal cues. To distinguish between these models, we tested whether ISNs respond to sensory detection of taste compounds. Tastants were applied to the proboscis while monitoring activity in ISNs by GCaMP5G or GCaMP6s calcium imaging in live flies (Akerboom et al., 2012; Harris et al., 2015). Although stimulation with sucrose, water, or the bitter compound denatonium triggered sensory neuron responses, no activation was seen in ISNs (Figure S2), indicating that the ISNs are unlikely to report detection of taste compounds.

An alternative hypothesis is that these neurons encode information about hunger state. As hormones often signal metabolic status, we tested whether existing hormone receptor Gal4 lines marked the ISNs. One line, *adipokinetic hormone receptor (AKHR)-Gal4* (Bharucha et al., 2008) exhibited strong labeling of ISNs, confirmed by double labeling with *R34G02-LexA*, suggesting these neurons may be regulated by AKH (Figure 2A).

AKH is a peptide hormone that is synthesized exclusively by neurosecretory cells in the corpus cardiacum and secreted into the circulating hemolymph, where it acts in a similar manner to mammalian glucagon. AKH secretion is stimulated under low nutrient conditions, which in turn leads to lipolysis, glycogenolysis, and release of sugar and lipid nutrients into the hemolymph from the fat body, the primary site of nutrient storage in *Drosophila* (Kim and Rulifson, 2004; Lee and Park, 2004). The endocrine role of AKH in regulation of insect metabolism is established, and this hormone is well positioned to signal nutrient status to the brain (Bharucha et al., 2008).

To directly test whether AKH modulates ISNs, calcium levels in these neurons were monitored by GCaMP5G fluorescence upon AKH perfusion in a dissected brain preparation (Figure 2B). Brief pulses of AKH produced rapid, robust, and dose-dependent GCaMP5G fluorescence increases with picomolar to nanomolar AKH concentrations, the physiological range of AKH measured in locust hemolymph (Candy, 2002). The ISNs of animals lacking AKHR did not respond to AKH, verifying that the AKH-induced responses were a result of activation of AKHR (Figure 2C).

In principle, the AKH-induced response might be cell-autonomous (AKH might directly bind to AKHR on ISNs to increase calcium) or non-autonomous (AKH might bind a receptor on other neurons that increase ISN activity via synaptic transmission). To test whether AKH directly modulates ISN activity, we applied the voltage-gated sodium channel blocker tetrodotoxin (TTX) to inhibit action potentials. ISNs responded to AKH even in the presence of TTX (Figure 2C), arguing that AKH directly activates ISNs.

AKH plays a role analogous to glucagon in signaling nutrient depletion and promoting release of stored nutrients. Insulin plays an opposing role, signaling nutrient abundance and promoting storage of circulating nutrients. Given the opposing endocrine roles of insulin and AKH, we tested whether insulin might regulate ISNs in a manner opposite to AKH. Insulin application alone did not induce a calcium response in ISNs (not shown). To test whether insulin affects the ability of ISNs to respond to AKH, dissected brains were perfused with two spaced pulses of AKH, separated by a 3-minute perfusion with insulin (Figure 2D). Insulin reduced the response of ISNs to AKH in a concentration-dependent manner. To test whether insulin acts cell-autonomously, TTX was applied to eliminate action potentials and non-autonomous effects. We found that insulin no longer reduced AKH responses in the presence of TTX (Figure 2D). This argues that ISNs do not directly sense insulin, but receive synaptic input from other neurons that provide an inhibitory drive onto ISNs. These studies demonstrate that ISNs are capable of receiving and integrating inputs reflecting both low and high nutrient levels, directly responding to AKH and receiving inhibition from an insulin-sensitive pathway.

To directly test whether ISNs report physiological need, we reasoned that the activity in ISNs would likely be different in starved versus fed animals. We monitored ISN activity by cell-attached electrophysiological recordings in the living fly under fed or starved conditions. ISN activity decreased in the fed state and increased in the starved state, and this state-dependent activity of ISNs was absent in the AKHR mutant (Figure 2E). These experiments show that the ISNs are modulated in the living animal based on nutritional state and that this requires AKHR.

### **A behavioral screen for molecules that regulate water consumption**

Our studies indicate that ISNs respond to signals of internal energy state and that their activity is sufficient to promote feeding, separating metabolic and neural functions of AKH. In a different behavioral screen to identify molecules that regulate water consumption, we ultimately identified additional molecules that are expressed in ISNs, regulate their activity, and provide unexpected insight into their function.

With the goal of identifying molecules that might signal thirst or water satiety in *Drosophila*, we screened a panel of 94 candidate ion channels and neuropeptide receptors for water consumption defects following RNAi knockdown with the pan-neuronal *nSynaptobrevin-Gal4* (*nSyb-Gal4*) line. Flies were placed in less than 20% relative humidity for 2 hours, which specifically increased water consumption (Lin et al., 2014) (Figure 3A). Upon pan-neuronal RNAi expression, the majority of RNAi lines showed behavior similar to GFP RNAi controls (109% control consumption). As expected, RNAi to PPK28, an ion channel expressed in sensory neurons and essential for water taste detection (Cameron et al., 2010; Chen et al., 2010), reduced water consumption (12% control consumption). In addition, RNAi against a number of genes reproducibly increased or decreased water consumption compared to sibling controls. These included the insulin-like receptor (5% control consumption) and the follicle stimulating hormone receptor (315% control consumption), which have been implicated in water homeostasis in insects (Liu et al., 2015; Paluzzi et al., 2014; Sellami et al., 2011).

Among the candidates with no known role in water consumption behavior was the TRPV family member, Nanchung (Nan) (184% and 197% control consumption, independent RNAi lines). Nan is a non-selective cation channel that participates in *Drosophila* proprioception, hearing, and hygrosensation (Gong et al., 2004; Kim et al., 2003; Liu et al., 2007; Zhang et al., 2013). Interestingly, TRPV channels function as osmosensors in *C. elegans* and mammals and likely play a role in water consumption regulation in mice (Colbert et al., 1997; Liedtke and Friedman, 2003; Liedtke et al., 2003). We therefore chose to further investigate the role of Nan in *Drosophila* water consumption.

### ***nanchung* is required to restrict water consumption**

To examine whether *nan*-expressing neurons regulate water consumption, we tested the behavioral response of *nan* mutants as well as animals expressing *nan* RNAi specifically in *nan-Gal4* neurons, using described lines (Kim et al., 2003). Water consumption time was measured in single flies following two hours of acute desiccation (Figure 3B). Both *nan*<sup>36a</sup> and *nan*<sup>dy5</sup> mutants as well as *nan-Gal4, UAS-nan RNAi* flies consumed significantly more water than controls, demonstrating that *nanchung* is necessary to restrict water consumption.

To examine how activity in *nanchung* neurons influences water consumption, we inducibly activated these neurons with dTRPA1. Flies expressing *UAS-dTRPA1* in *nan-Gal4* neurons reduced water consumption upon dTRPA1 activation. Transient activation of *nan* neurons was also sufficient to rescue elevated water consumption observed in *nan*<sup>dy5</sup> heterozygotes (Figure 3C), suggesting that *nanchung* neurons function to restrict water consumption.

### **Nanchung neurons in the SEZ are osmosensitive**

The expression pattern of *nanchung* has been described previously using *nan-Gal4* (Kim et al., 2003; Liu et al., 2007) (Figure 3D). These studies found expression in chordotonal organs of the legs and antennae, as well as in a small number of putative interneurons in the higher brain. Because *nan* has been proposed to detect water vapor in antennal neurons, we tested whether antennae were required for *dTRPA1*-mediated activation of *nan* neurons to reduce water consumption in *nan*<sup>dy5</sup> heterozygotes and found no role for antennae (not

shown). These data argue that Nanchung's function as an antennal sensor of water vapor is independent of its role in water consumption.

An alternative possibility is that Nanchung functions in central brain neurons labeled by *nan-Gal4* to regulate water consumption. Nan belongs to a class of mechanosensitive TRPV channels and is sufficient to confer osmosensitivity to cultured mammalian cells (Kim et al., 2003). Moreover, TRPV channels have been proposed to confer osmosensitivity to central neurons in the mammalian brain (Bourque, 2008; Liedtke and Friedman, 2003). We therefore tested whether central neurons labeled by the *nan-Gal4* line were directly osmosensitive.

*nan-Gal4, UAS-GCaMP6s* brains were perfused with artificial hemolymph (AHL) of different osmolalities while GCaMP6s fluorescence was monitored (Chen et al., 2013). Decreasing osmolality caused robust calcium responses from bilaterally symmetric *nan-Gal4* neurons in the ventrolateral SEZ (Figure 4A). These responses were dose-dependent and observed following osmolality decreases but not increases (Figure 4B, C). To test whether these responses are specific to *nan* SEZ neurons, brains expressing GCaMP6s with the pan-neuronal driver *nSyb-LexA* were perfused with low osmolality solution. Again, strong calcium responses were observed only in the SEZ neurons labeled by *nan-Gal4* (Figure 4D; Figure S3). Moreover, Nan is necessary for the osmolality responses, as calcium responses to low osmolality were significantly decreased in a *nan<sup>36a</sup>* mutant background and were rescued with a *UAS-nan* transgene (Kim et al., 2003) (Figure 4E, F).

These data indicate that *nan* SEZ neurons are uniquely capable of responding to low extracellular osmolality with calcium increases. Because GCaMP6s preferentially reports calcium increases over decreases, we used the genetically encoded voltage sensor ArcLight (Cao et al., 2013) to test whether *nan* SEZ neuron activity is bidirectionally regulated by osmolality changes. Consistent with GCaMP6s imaging, decreasing extracellular osmolality decreased ArcLight fluorescence, indicating depolarization. In addition, we found that increasing extracellular osmolality significantly increased ArcLight fluorescence in these neurons, indicating hyperpolarization (Figure 4G, H). Taken together, these data suggest that *nan* SEZ neurons report bidirectional extracellular osmolality changes in the brain.

### The ISNs express *nanchung* and *AKHR* and respond to osmolality and AKH

During the course of these studies, we noted that the osmosensitive *nan* SEZ neurons had a similar spatial location and morphology as the ISNs that express *AKHR*, suggesting that they might be the same neurons. We tested this by examining overlap between *nan-Gal4* neurons and the ISNs, marked by *R34G02-LexA*, and indeed found *nan-Gal4* labels the ISNs as well as two additional SEZ neurons per hemisphere (Figure 5A). Consistent with this overlap, the ISNs responded to low osmolality and to AKH (Figure 5B, C). Responses were detected in neurites and cell bodies (Figure S3C, D). The *AKHR*-negative, *Nan*-positive neurons did not respond to osmolality or contribute to consumption behavior (Figure S4), arguing that the ISNs are a unique class of neurons that respond to osmolality and AKH. Thus, two screens for neurons regulating different homeostatic behaviors independently identified the ISNs.



ISNs are responsive to both osmolality and the hormone AKH. To examine how ISNs might integrate these two signals, we first asked whether Nan might be required for ISNs to sense AKH, and whether AKHR might be required for ISNs to sense osmolality. We therefore monitored osmolality responses in *AKHR* mutants and AKH responses in *nan* mutants (Figure 5C). By calcium imaging, loss of AKHR did not affect the ability of ISNs to respond to osmolality, nor did loss of Nan affect the AKH response. All imaging experiments were performed under controlled osmolality and AKH conditions, as monitoring GCaMP activity by necessity requires removing cuticle and exposing the brain to artificial hemolymph. Thus, these experiments indicate that each input increases ISNs activity via an independent molecular mechanism, but do not directly monitor interactions between ISN inputs.

We therefore asked whether ISN activity resulting from changes in one input, osmolality, might affect the ability of ISNs to respond to another input, AKH. To test this, we monitored calcium responses to AKH in ISNs of brains perfused with high or low extracellular osmolality. We found that high extracellular osmolality significantly reduced AKH responses by an average of 70% (Figure 5D, E). These results argue that ISNs sense extracellular AKH and osmolality via independent molecular mechanisms, AKHR and Nanchung, but that these two inputs regulate a common output, ISN activity.

### Starvation reduces *Drosophila* hemolymph osmolality

AKH levels increase with starvation and activate ISNs, consistent with its action as a hunger signal that drives feeding. Low extracellular osmolality also activates ISNs, suggesting that low osmolality might also act as an internal signal of nutrient deprivation. Flies starved for one day have approximately 75% lower hemolymph sugar levels (Na et al., 2013), and we hypothesized that this reduction in sugar levels might reduce hemolymph osmolality. To test this, we measured the hemolymph osmolality of single fed or starved flies with a temperature gradient osmometer (Arav and Rubinsky, 1994) (Figure 6A).

To confirm that we could detect physiological changes in osmolality, we measured hemolymph osmolality of single flies placed in dry (<20% RH) or humid (>80% RH) environments for 6-8 hours. Consistent with previous studies (Albers and Bradley, 2004), desiccation increased hemolymph osmolality by an average of 55 mOsm/kg. In addition, hemolymph osmolality of desiccated flies returned to control levels five minutes after water consumption (Figure 6B).

To test the effect of starvation on hemolymph osmolality, hemolymph was collected from single flies that were either fed or starved for 24 hours with access to water. We found that hemolymph osmolality of starved flies was lower than that of well-fed animals by ~30 mOsm/kg (Figure 6C). Importantly, physiological changes in osmolality of the magnitude observed following starvation were sufficient to elicit responses from ISNs in imaging preparations (Figure 6D, E). Thus, both low extracellular osmolality and increased AKH abundance may be starvation signals that increase ISN activity and promote feeding.

### ISNs oppositely regulate sugar and water consumption

Taken together, our data suggest a model in which ISNs are sensitive to internal signals for both water and sugar abundance, and are sufficient to modulate both water and sugar

consumption. To directly ask how ISN activity impacts sugar and water consumption, we drove *dTRPA1* expression with *ISN-Gal4*, which exclusively labels ISNs. Consistent with our previous observations, *ISN-Gal4, UAS-dTRPA1* flies avidly consumed sucrose upon TRP activation at 32°C, but not at 23°C. In contrast, these flies exhibited markedly reduced water consumption at 32°C but not at 23°C (Figure 7A).

To test the effect of inhibiting ISN activity on water and sugar consumption, we used *ISN-Gal4* to selectively drive expression of RNAi against *nSynaptobrevin (nSyb)*, which is required for synaptic transmission. Flies expressing *nSyb* RNAi in ISNs increased water consumption relative to controls by almost twofold. In contrast, expression of *nSyb* RNAi in ISNs decreased sucrose intake by an average of 44% (Figure 7B). Thus, neural activity in ISNs oppositely regulates sugar and water consumption behaviors: increased ISN activity both promotes sugar and restricts water consumption, whereas decreased activity promotes water and restricts sugar consumption.

To ask whether *nanchung* and *AKHR* are important for the ability of ISNs to regulate food and water intake, we expressed *nan* RNAi or *AKHR* RNAi selectively in ISNs and examined the effect on water and sucrose consumption. Flies expressing either of two independent *nan* RNAi constructs in ISNs increased water consumption and decreased sucrose consumption relative to controls, consistent with phenotypes observed when silencing ISNs (Figure 7C, S5A-C). Flies expressing *nan* RNAi under the control of *nSyb-Gal4* and *nan-Gal4* drivers produced the same reciprocal effects on water and sucrose consumption (Figure S5D, E). Like flies expressing *nan* RNAi, flies expressing *AKHR* RNAi in ISNs also increased water intake and decreased sucrose intake relative to controls (Figure 7D). The *nan* and *AKHR* RNAi data is consistent with the notion that both AKHR and osmolality contribute to the activation of ISNs. The loss of either signal decreased activity in ISNs, leading to loss-of-function phenotypes similar to those observed when silencing ISNs with *nSyb* RNAi.

The finding that the ISNs sense signals of hunger and thirst, AKH and osmolality, and oppositely regulate sugar and water consumption, suggests that hunger and thirst may be competing drives under some conditions. To test this, we examined whether there were consumption differences in flies with competing needs. Under conditions when flies were not thirsty, there was no drive to consume water and this was not affected by starvation state (Figure 7E). However, under conditions when flies were mildly starved (2 hours), we found that sucrose consumption was reduced in thirsty flies compared to water-sated flies (Figure 7E). These data argue that under conditions of competing needs, there is a balance between water and sucrose consumption. As the relative weight of these needs changes, the balance would be predicted to change as well.

## Discussion

In this study, we uncover a neural mechanism that coordinates two essential homeostatic behaviors: sugar and water consumption. This coordination is achieved by two neurons per SEZ hemisphere of the *Drosophila* brain, the ISNs, which are sensitive to internal signals for both hunger and thirst and whose activity oppositely regulates sugar and water consumption (Figure 7F). The antagonistic manner in which ISNs couple these behaviors suggests a

regulatory principle by which animal nervous systems might promote internal osmotic and metabolic homeostasis.

#### **Four neurons oppositely regulate sugar and water consumption in *Drosophila***

Low internal osmolality and high AKH are signals of water satiety and hunger, respectively. ISN activity increases both in the presence of low extracellular osmolality and AKH. We find that increasing ISN activity promotes sugar consumption and reduces water consumption. Conversely, high internal osmolality and low AKH are signals of thirst and food satiety. ISN activity decreases and AKH responses are reduced in the presence of high extracellular osmolality or insulin. Decreasing ISN activity increases water consumption and reduces sugar consumption.

How do ISNs achieve opposite regulation of a single behavior, consumption, in a manner that depends on the substance being consumed? One possibility is that the downstream targets of ISNs include interneurons involved in the behavioral response to water and sugar taste. This model predicts that increased ISN activity promotes the ability of sugar taste interneurons to drive consumption while inhibiting the ability of water taste interneurons to do so. It may be possible to test hypotheses about the neural circuits in which ISNs participate through the use of large scale calcium imaging.

#### **Molecules for sensing internal hunger and thirst cues**

ISNs regulate sugar and water consumption in a manner that appropriately reflects internal hunger and thirst states. Here, we show that two genes, *AKHR* and *nanchung*, are expressed in ISNs and function to confer sensitivity to these states.

AKHR is a G-protein coupled receptor expressed in the fat body and the brain that has been well characterized in the context of insect metabolic regulation (Bharucha et al., 2008; Candy, 2002). The ligand for this receptor, AKH, is secreted into the hemolymph by specialized neurosecretory cells in the corpus cardiacum (Kim and Rulifson, 2004), where it acts under conditions of food deprivation. Here, we identify a role for AKH in regulating the activity of four interneurons in the SEZ, the ISNs, and show that this activity promotes sugar consumption. AKH abundance in the hemolymph therefore promotes feeding via the ISNs. Manipulating AKHR exclusively in the ISNs provided a means to separate the metabolic and neural effects of AKH, uncovering a role for AKH in the nervous system.

Sensors for internal hemolymph osmolality have not previously been described. Here, we find that the non-selective cation channel Nanchung is expressed in ISNs and is required for their responses to low osmolality. Although we do not know if Nan is the direct osmosensor in ISNs, previous studies found that Nan confers low osmolality responses when expressed in heterologous cells (Kim et al., 2003), consistent with this notion. Nan family members of the TRPV4 family have been shown to participate in osmosensation in *C. elegans* and mammals (Colbert et al., 1997; Liedtke and Friedman 2003; Liedtke et al., 2003), suggesting an ancient and conserved function. Nanchung participates in sensory detection of mechanical stimuli in *Drosophila*, including proprioception, audition, and low humidity sensing (Gong et al., 2004; Kim et al., 2003; Liu et al., 2007; Zhang et al., 2013). It is interesting that the same molecule that is involved in external sensory detection of

mechanical stimuli also participates in internal detection of osmolality, a mechanical stimulus. Similar molecular re-tooling has recently been described for the GR43a gustatory receptor, which acts as a sensory receptor to monitor fructose in the environment and as an internal sensor monitoring circulating fructose levels in brain hemolymph (Miyamoto et al., 2012).

In the mammalian brain, osmosensitive neurons are generally found in areas which lack a blood-brain barrier. The blood-brain barrier of *Drosophila* expresses multiple aquaporins and may potentially regulate hemolymph osmolality (Limmer et al., 2014). Whether changes in hemolymph osmolality are regulated by the blood-brain barrier to impact ISN activity is an interesting question for future study.

### **Coupling of sugar and water consumption behaviors as a mechanism for homeostasis**

ISNs oppositely regulate the behavioral responses to hunger and thirst states. How might this type of coordination be adaptive? One possibility is suggested by the fact that sugar and water consumption perturb internal osmotic homeostasis in opposite directions. In *Drosophila* and mammals, sugar consumption leads to increased blood-sugar levels and increased blood osmolality. Conversely, water consumption leads to lowered blood osmolality. Our studies show that ISNs are sensitive to extracellular osmolality, and that they oppositely regulate sugar and water consumption. Under high osmotic conditions, decreased ISN activity promotes water consumption, reducing internal osmolality. Under low osmotic conditions, increased ISN activity promotes sucrose consumption, increasing internal osmolality. Thus, ISNs may monitor internal osmolality to reciprocally regulate sugar and water consumption to restore homeostasis.

Reciprocal regulation of food and water consumption has been reported in both classical and recent rodent studies. For example, increasing blood osmolality promotes water consumption and inhibits food consumption in rats, whereas decreasing osmolality has the opposite effect (Gutman and Krausz, 1969). In addition, ghrelin, a key internal signal for hunger in mammals, is sufficient not only to promote feeding but also to inhibit water consumption in rats (Mietlicki et al., 2009). Thus, vertebrates and invertebrates may share mechanisms for coupling water and sugar consumption in a manner that promotes homeostasis. In *Drosophila*, the convergence of internal signals onto the ISNs provides a mechanism to weigh homeostatic deviations and drive consumption to restore balance.

Other neurons in the *Drosophila* brain process homeostatic needs for water and sugar separately. For example, water reward and sugar reward are processed by different subsets of mushroom body input neurons, likely independent of gustatory sensory activation (Burke et al., 2012; Lin et al., 2014; Liu et al., 2012). Neuropeptide F, small Neuropeptide F, and dopamine are all signals of nutrient deprivation that promote nutrient intake (Hergarden et al., 2012; Inagaki et al., 2012; Inagaki et al., 2014; Lee et al., 2004; Marella et al., 2012). Circulating glucose and fructose in the hemolymph also report the nutritional state and alter feeding behavior by direct activation of a few central neurons (Miyamoto et al., 2012; Dus et al., 2015). The ISNs are unique in that they detect multiple internal state signals and use this information to weigh competing needs. In addition to parallel, independent pathways for

eating and drinking, this study demonstrates the existence of a pathway that couples these drives.

## Experimental Procedures

Additional details are available in Extended Experimental Procedures.

### Blue Dye Consumption experiments

Flies (15-20/vial) were transferred from food to filter paper soaked in 200mM sucrose and blue dye. After feeding animals were scored a 0 (no dye in abdomen), 1 (less than half of the abdomen was blue), or 2 (half or more of the abdomen contained dye). The scores of all flies in one vial were averaged and considered one trial.

### Proboscis Extension Response Assays

PER was performed as described (Marella et al., 2012), except that each animal was considered a data point, and was categorized as responding 0, 1, 2, or 3 times.

### Temporal Consumption Assays

Assays were performed as described (Pool et al., 2014). Animals were presented with a taste stimulus ten times and total consumption time was monitored. To generate thirsty flies, flies were placed in a sealed chamber with ~250g CaSO<sub>4</sub> (Drierite, stock# 23001) for 2 hours, unless otherwise noted.

### Immunohistochemistry

Immunohistochemistry was performed as described (Marella et al., 2012).

### Calcium and voltage imaging

GCaMP5G or GCaMP6s imaging of taste responses was performed as described (Harris et al., 2015). Stacks of 14-20 Z-slices were collected at approximately 0.3 Hz. For GCaMP5G or GCaMP6s imaging of AKH or osmolality responses, the brain was removed in ice-cold calcium and magnesium-free artificial hemolymph-like solution (AHL), transferred to a perfusion chamber with room temperature AHL, and immobilized with tungsten wire. Image analysis was performed in ImageJ.

### Hemolymph osmolality measurements

An osmometer was assembled and used as described (Arav and Rubinsky, 1994), with minor modifications detailed in Extended Experimental Procedures.

### Electrophysiology

Extracellular recordings were performed in live animals as described (Pool et al., 2014).

### Statistical Analyses

Student's t-test, with Sidak correction for multiple comparisons, was used to compare two groups. ANOVA followed by Tukey's post hoc test was used to compare three or more

groups. ANOVA followed by Dunnet's post hoc test was used to compare multiple responses of varying stimuli.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

The Scott lab provided comments on the manuscript. Dr. B. Rubinsky provided materials for osmolality measurements. This work was supported by a grant from the NIH (R01 DK09874; K.S.) and an NSF predoctoral fellowship (N.J.).

## References

- Akerboom J, Chen TW, Wardill TJ, Tian L, Marvin JS, Mutlu S, Calderon NC, Esposti F, Borghuis BG, Sun XR, et al. Optimization of a GCaMP calcium indicator for neural activity imaging. *J. Neurosci.* 2012; 32:13819–13840. [PubMed: 23035093]
- Albers MA, Bradley TJ. Osmotic regulation in adult *Drosophila melanogaster* during dehydration and rehydration. *J. Exper. Biol.* 2004; 207:2313–2321. [PubMed: 15159436]
- Arav A, Rubinsky B. Temperature gradient osmometer and anomalies in freezing temperatures. *Am. J. Physiol.* 1994; 267:1646–1652.
- Bharucha KN, Tarr P, Zipursky SL. A glucagon-like endocrine pathway in *Drosophila* modulates both lipid and carbohydrate homeostasis. *J. Exper. Biol.* 2008; 211:3103–3110. [PubMed: 18805809]
- Bjordal M, Arquier N, Kniazeff J, Pin JP, Leopold P. Sensing of amino acids in a dopaminergic circuitry promotes rejection of an incomplete diet in *Drosophila*. *Cell.* 2014; 156:510–521. [PubMed: 24485457]
- Bourque CW. Central mechanisms of osmosensation and systemic osmoregulation. *Nature Reviews.* 2008; 9:519–531.
- Burke CJ, Huetteroth W, Oswald D, Perisse E, Krashes MJ, Das G, Gohl D, Silies M, Certel S, Waddell S. Layered reward signalling through octopamine and dopamine in *Drosophila*. *Nature.* 2012; 492:433–437. [PubMed: 23103875]
- Cameron P, Hiroi M, Ngai J, Scott K. The molecular basis for water taste in *Drosophila*. *Nature.* 2010; 465:91–95. [PubMed: 20364123]
- Candy DJ. Adipokinetic hormones concentrations in the haemolymph of *Schistocerca gregaria*, measured by radioimmunoassay. *Insect Biochem. and Mol. Biol.* 2002; 32:1361–1367. [PubMed: 12530204]
- Cao G, Platasa J, Pieribone VA, Raccuglia D, Knust M, Nitabach M. Genetically targetted optical electrophysiology in intact neural circuits. *Cell.* 2013; 144:904–913. [PubMed: 23932121]
- Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr RA, Orger MB, Jayaraman V, et al. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature.* 2013; 499:295–300. [PubMed: 23868258]
- Chen Z, Wang Q, Wang Z. The amiloride-sensitive epithelial Na<sup>+</sup> channel PPK28 is essential for *Drosophila* gustatory water reception. *J. Neurosci.* 2010; 30:6247–6252. [PubMed: 20445050]
- Clyne JD, Miesenbock G. Sex-specific control and tuning of the pattern generator for courtship song in *Drosophila*. *Cell.* 2008; 133:354–363. [PubMed: 18423205]
- Colbert HA, Smith TL, Bargmann CI. OSM-9, a novel protein with structural similarity to channels, is required for olfaction, mechanosensation, and olfactory adaptation in *Caenorhabditis elegans*. *J. Neurosci.* 1997; 17:8259–8269. [PubMed: 9334401]
- Dethier, VG. *The Hungry Fly*. Harvard University Press; Cambridge, MA: 1976.
- Dus M, Lai JS, Gunapala KM, Min S, Tayler TD, Hergarden AC, Geraud E, Joseph CM, Suh GS. Nutrient Sensor in the Brain Directs the Action of the Brain-Gut Axis in *Drosophila*. *Neuron.* 2015; 87:139–151. [PubMed: 26074004]


- Gohl DM, Silies MA, Gao XJ, Bhalaria S, Luongo FJ, Lin CC, Potter CJ, Clandinin TR. A versatile in vivo system for directed dissection of gene expression patterns. *Nature Methods*. 2011; 8:231–237. [PubMed: 21473015]
- Gong Z, Son W, Chung YD, Kim J, Shin DW, McClung CA, Lee Y, Lee HW, Chang DJ, Kaang BK, et al. Two interdependent TRPV channel subunits, inactive and Nanchung, mediate hearing in *Drosophila*. *J. Neurosci*. 2004; 24:9059–9066. [PubMed: 15483124]
- Gutman Y, Krausz M. Regulation of food and water intake in rats as related to plasma osmolarity and volume. *Physiol. and Behav*. 1969; 4:311–313.
- Hamada FN, Rosenzweig M, Kang K, Pulver SR, Ghezzi A, Jegla TJ, Garrity PA. An internal thermal sensor controlling temperature preference in *Drosophila*. *Nature*. 2008; 454:217–220. [PubMed: 18548007]
- Harris DT, Kallman BR, Mullaney BC, Scott K. Representations of Taste Modality in the *Drosophila* Brain. *Neuron*. 2015; 86:1449–1460. [PubMed: 26051423]
- Hergarden AC, Taylor TD, Anderson DJ. Allatostatin-A neurons inhibit feeding behavior in adult *Drosophila*. *Proc. Nat. Acad. Sci. USA*. 2012; 109:3967–3972. [PubMed: 22345563]
- Inagaki HK, Jagadish S, Barne G, Ishimoto H, Ben-Tabou de-Leon S, Wong AM, Kitamoto T, Axel R, Anderson DJ. Visualizing neuromodulation in vivo: TANGO-mapping of dopamine signaling reveals appetite control of sugar sensing. *Cell*. 2012; 148:583–595. [PubMed: 22304923]
- Inagaki HK, Panse KM, Anderson DJ. Independent, reciprocal neuromodulatory control of sweet and bitter taste sensitivity during starvation in *Drosophila*. *Neuron*. 2014; 84:806–820. [PubMed: 25451195]
- Jenett A, Rubin GM, Ngo TT, Shepherd D, Murphy C, Dionne H, Pfeiffer BD, Cavallaro A, Hall D, Jeter J, et al. A GAL4-driver line resource for *Drosophila* neurobiology. *Cell Rep*. 2012; 2:991–1001. [PubMed: 23063364]
- Kim J, Chung YD, Park DY, Choi S, Shin DW, Soh H, Lee HW, Son W, Yim J, Park CS, et al. A TRPV family ion channel required for hearing in *Drosophila*. *Nature*. 2003; 424:81–84. [PubMed: 12819662]
- Kim SK, Rulifson EJ. Conserved mechanisms of glucose sensing and regulation by *Drosophila* corpora cardiaca cells. *Nature*. 2004; 431:316–320. [PubMed: 15372035]
- Lee G, Park JH. Hemolymph sugar homeostasis and starvation-induced hyperactivity affected by genetic manipulations of the adipokinetic hormone-encoding gene in *Drosophila melanogaster*. *Genetics*. 2004; 167:311–323. [PubMed: 15166157]
- Lee KS, You KH, Choo JK, Han YM, Yu K. *Drosophila* short neuropeptide F regulates food intake and body size. *J Biol Chem*. 2004; 279:50781–50789. [PubMed: 15385546]
- Liedtke W, Friedman JM. Abnormal osmotic regulation in *trpv4*<sup>-/-</sup> mice. *Proc. Nat. Acad. Sci. USA*. 2003; 100:13698–13703. [PubMed: 14581612]
- Liedtke W, Tobin DM, Bargmann CI, Friedman JM. Mammalian TRPV4 (VR-OAC) directs behavioral responses to osmotic and mechanical stimuli in *Caenorhabditis elegans*. *Proc. Nat. Acad. Sci. USA*. 2003; 100(Suppl 2):14531–14536. [PubMed: 14581619]
- Limmer S, Weiler A, Volkenhoff A, Babatz F, Klämbt C. The *Drosophila* blood-brain barrier: Development and function of a glial epithelium. *Front Neurosci*. 2014; 14:365. [PubMed: 25452710]
- Lin S, David O, Chandra V, Talbot C, Huetteroth W, Waddell S. Neural correlates of water reward in thirsty *Drosophila*. *Nat. Neurosci*. 2014; 17:1536–1542. [PubMed: 25262493]
- Liu C, Plaçais P-Y, Yamagata N, Pfeiffer BD, Aso Y, Friedrich AB, Siwanowicz I, Rubin GM, Preat T, Tanimoto H. A subset of dopamine neurons signals reward for odour memory in *Drosophila*. *Nature*. 2012; 488:512–516. [PubMed: 22810589]
- Liu L, Li Y, Wang R, Yin C, Dong Q, Hing H, Kim C, Welsh MJ. *Drosophila* hygrosensation requires the TRP channels water witch and nanchung. *Nature*. 2007; 450:294–298. [PubMed: 17994098]
- Liu Y, Luo J, Carlsson MA, Nässel DR. Serotonin and insulin-like peptides modulate leucokinin-producing neurons that affect feeding and water homeostasis in *Drosophila*. *J. Comp. Neurol*. 2015; 523:1840–1863. [PubMed: 25732325]
- Marella S, Mann K, Scott K. Dopaminergic modulation of sucrose acceptance behavior in *Drosophila*. *Neuron*. 2012; 73:941–950. [PubMed: 22405204]


- Mietlicki EG, Nowak EL, Daniels D. The effect of ghrelin on water intake during dipsogenic conditions. *Physiol. and Behav.* 2009; 96:37–43. [PubMed: 18761029]
- Miyamoto T, Slone J, Song X, Amrein H. A fructose receptor functions as a nutrient sensor in the *Drosophila* brain. *Cell.* 2012; 151:1113–1125. [PubMed: 23178127]
- Na J, Musselman LP, Pendse J, Baranski TJ, Bodmer R, Ocorr K, Cagan R. A *Drosophila* model of high sugar diet-induced cardiomyopathy. *PLoS Genetics.* 2013; 9:e1003175. [PubMed: 23326243]
- Paluzzi JP, Vanderveken M, O'Donnell MJ. The heterodimeric glycoprotein hormone, GPA2/GPB5, regulates ion transport across the hindgut of the adult mosquito, *Aedes aegypti*. *PLoS ONE.* 2014; 9:e86386. [PubMed: 24466069]
- Pfeiffer BD, Ngo TT, Hibbard KL, Murphy C, Jenett A, Truman JW, Rubin GM. Refinement of tools for targeted gene expression in *Drosophila*. *Genetics.* 2010; 186:735–755. [PubMed: 20697123]
- Pfeiffer BD, Truman JW, Rubin GM. Using translational enhancers to increase transgene expression in *Drosophila*. *Proc. Natl. Acad. Sci. USA.* 2012; 109:6626–6631. [PubMed: 22493255]
- Pool AH, Kvello P, Mann K, Cheung SK, Gordon MD, Wang L, Scott K. Four GABAergic interneurons impose feeding restraint in *Drosophila*. *Neuron.* 2014; 83:164–77. [PubMed: 24991960]
- Sellami A, Agricola HJ, Veenstra JA. Neuroendocrine cells in *Drosophila melanogaster* producing GPA2/GPB5, a hormone with homology to LH, FSH and TSH. *Gen. Comp. Endocrinol.* 2011; 170:582–588. [PubMed: 21118692]
- Sternson SM. Hypothalamic survival circuits: blueprints for purposive behaviors. *Neuron.* 2013; 77:810–824. [PubMed: 23473313]
- Zhang W, Yan Z, Jan LY, Jan YN. Sound response mediated by the TRP channels NOMPC, NANCHUNG, and INACTIVE in chordotonal organs of *Drosophila larvae*. *Proc. Nat. Acad. Sci. USA.* 2013; 110:13612–13617. [PubMed: 23898199]



### Highlights

- Four interoceptive neurons oppositely regulate sugar and waterconsumption
- These neurons detect a peptide hormone that signals nutrient deprivation
- These neurons directly sense changes in osmolality through an ion channel
- These neurons report homeostatic changes and drive consumption to restore balance





**Figure 2. ISNs respond to AKH and are inhibited by insulin**

A. (left) Expression of *AKHR-Gal4*, *UAS-mCD8::GFP* in brain. (second) *AKHR-Gal4*, *UAS-mCD8::RFP* in SEZ, (third) *R34G02-LexA*, *lexAop-mCD8::GFP* in SEZ, (right) overlay. Scale 50  $\mu$ m.

B. (left) GCaMP5G increase in ISNs in response to AKH (31.6 pM, 100 pM, 316 pM, 1 nM, 3.16 nM) applied at the blue lines. n=7, mean (black line)  $\pm$  SEM (grey). (right) Maximum GCaMP5G increase (% F/F) to AKH, mean  $\pm$  SEM, same data as trace. Repeated-measures one way ANOVA followed by holm-sidak post hoc test, comparing to 31.6 pM response. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001.

C. (left) GCaMP5G increase in ISNs to 1 nM AKH (blue line) or 30 mM KCl (red line) in the *AKHR* mutant. n=5, mean  $\pm$  SEM. (middle) GCaMP5G increase in ISNs to 316 pM AKH (blue line) pre- or post-TTX (100 nM) application. n=8, mean  $\pm$  SEM. (right) GCaMP5G increase (% F/F) to AKH in the *AKHR* mutant (t-test, \*p<0.05) or pre- or post-TTX application (t-test, ns). Mean  $\pm$  SEM, same data as traces.

D. (left) GCaMP5G increase in ISNs to 316 pM AKH (blue line) pre- or post-insulin (100 nM) application. n=11, mean  $\pm$  SEM. (middle) GCaMP5G increase in ISNs to 316 pM AKH (blue line) pre- or post-insulin (100 nM) and TTX (100 nM) application. n=10, mean  $\pm$  SEM. (right) GCaMP5G change (% F). Mean  $\pm$  SEM, insulin concentrations in the absence of TTX: One-way ANOVA with Dunnett's post hoc test comparing to 0 nM Insulin, \*\*\*p<0.001. Effect of TTX: t-test (100nM Insulin vs. 100nM insulin + TTX), \*\*\*p<0.001, same data as traces plus additional insulin concentrations.

E. (left) Example cell-attached recordings from *AKHR*<sup>+</sup> SEZ neurons in *AKHR* mutant or wild type flies, fed (f) or starved (s) 24H, and (right) summary graph. n = 6-11, mean  $\pm$  SEM.


SEM, one-way ANOVA, Tukey Post Hoc, \*\* $p < 0.01$ , \* $p < 0.05$ . See Figure S2, showing that ISNs are not taste-responsive.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




**Figure 3. Identification of *nanchung* and its role in water consumption**

A. Behavioral screen for genes regulating water consumption. *UAS-RNAi* lines were crossed to *nSyb-Gal4*. 10 RNAi and 10 sibling control flies were assayed/RNAi line. Average water consumption time in RNAi group versus sibling control was used to calculate % control consumption. Gray marks 2 standard deviations from mean. Two *nan* RNAi lines (BSC# 31674 and 31925) are highlighted in red. Behavioral data for *nan* RNAi BSC# 31925 are shown unless noted.

B. Water consumption in *nan* RNAi and *nan* mutant backgrounds. For all box plots, whiskers = 10<sup>th</sup> to 90<sup>th</sup> percentile, box = 25<sup>th</sup> to 75<sup>th</sup> percentile, and line in box = median. n=28-60 flies, one-way ANOVA, Tukey Post Hoc: for RNAi experiments (\*\*p<0.001), and for mutant experiments (\*\*p<0.001).

C. Activating *nan-Gal4* neurons affects water consumption. n=20-63 flies, t-test, Holm-Sidak correction, (23°C vs 32°C per genotype), \*\*\*p<0.001.

D. *nan-Gal4*, *UAS-mCD8::GFP* expression in the brain (top) and VNC (bottom). Scale 100 μm.



**Figure 4. Nanchung SEZ neurons respond to osmolality**

A. *nan-Gal4*, *UAS-GCaMP6s* expression in brain (left) and example  $\Delta F/F$  heat-map of the same brain (right) upon an extracellular osmolality decrease of 200mOsm/kg. Scale 50  $\mu\text{m}$ .

B.  $\Delta F/F$  traces (mean  $\pm$  SEM) for osmolality decreases of 25, 50, 100, and 200 mOsm/Kg (n=9-12 brains/condition).

C. Maximum  $\Delta F/F$  responses (mean  $\pm$  SEM) for osmolality changes. n=7-12 brains/condition, One way ANOVA, Dunnet's Post Hoc to mock (0 mOsm change) \*\*p<0.01, \*\*\*p<0.001, data from B plus additional osmolalities.

D. (top left) *nSyb-LexA*, *lexAop-GCaMP6s* expression in brain. (top right)  $\Delta F/F$  heat-map to an osmolality decrease of 200mOsm/kg in the same brain. (bottom left)  $\Delta F/F$  response (green) overlaid on *nSyb-LexA*, *lexAop-GCaMP6s* expression (grey). (bottom right)  $\Delta F/F$  response (green) overlaid on *nSyb-LexA*, *lexAop-GCaMP6s* (grey) and *nan-Gal4* expression (red) in same brain. Scale 50  $\mu\text{m}$ .


E.  $\Delta F/F$  traces upon osmolality decreases of 200 mOsm/Kg in *nan<sup>36a</sup>* mutant or wild type (WT) flies. n=28-33. E-H: data are mean  $\pm$  SEM.

F.  $\Delta F/F$  graphs for WT, mutant, and rescue flies (*UAS-nan*; *nan<sup>36a</sup>*). n=33, 28, and 12. one-way ANOVA, Tukey Post Hoc, \*p<0.05, \*\*p<0.01, data from E plus rescue.

G.  $\Delta F/F$  traces to 200 mOsm/Kg osmolality decreases in flies expressing *UAS-GCaMP6s* (green, n=9 brains) or *UAS-ArcLight* (magenta, n=11 brains) in ISNs.

H.  $\Delta F/F$  graphs for GCaMP6s and ArcLight from F. One sample t-tests for difference from theoretical mean of 0.0, \*p<0.05, \*\*\*p<0.001.

See also Figure S3, showing quantification of whole brain responses.



**Figure 5. Nanchung SEZ neurons are ISNs**

A. Co-expression of *R34G02-LexA* and *nan-Gal4* in ISNs. Zoom shows one *nan-Gal4* cluster. Scale 20  $\mu$ m.


B. (left) Low osmolality response and (right) AKH response in the same cell.

C. maximum  $\Delta F/F$  graphs (mean  $\pm$  SEM) to 200 mOsm/kg osmolality decreases (Osmo) or 1 nM AKH in control or mutant backgrounds. n=5-10 brains, t-test Holm-Sidak correction, ns.

D.  $\Delta F/F$  traces (mean  $\pm$  SEM) to 1 nM AKH in 440 mOsm/kg (high) or 240 mOsm/Kg (low) extracellular osmolality and graph of  $\Delta F/F$  responses. Top: n=9 brains. Bottom: n = 11 brains.

E. Quantification of  $\Delta F/F$  graphs from D, mean  $\pm$  SEM. brains, paired t-test, \*p<0.05.

See also Figure S4, showing that the AKHR-, Nan+ neurons do not respond to osmolality.



**Figure 6. *Drosophila* hemolymph osmolality decreases during starvation**

A. Top: Temperature (Temp.) gradient osmometer. Distance between sample and standard ice/liquid interfaces was used to calculate sample osmolality. Bottom: Freezing interfaces (white dots) from standards and single fly hemolymph. Scale 0.05 mm.


B. Desiccation affects hemolymph osmolality. n=14-17, one way ANOVA, Tukey's Post Hoc, \*\*p<0.01. Dehydrated: 6 hours at <20% RH. Hydrated: 6 hours in a humid box. Rehydrated: water was administered as described in methods.

C. Starvation affects hemolymph osmolality. n=14, t-test, \*\*\*p<0.001.

D. F/F traces (mean ± SEM) to 30 mOsm/Kg osmolality decreases or increases in flies expressing *UAS-GCaMP6s* (green, n=10 brains) or *UAS-ArcLight* (magenta, n=10 brains) in ISNs.

E. F/F graphs (mean ± SEM) for GCaMP6s and ArcLight from D. Wilcoxon matched-pairs signed rank test, \*p<0.05.





**Figure 7. ISNs promote sucrose consumption and inhibit water consumption**

A. ISN activation with *dTRPA1* alters water and 1M sucrose consumption. Desiccated (des.), hydrated (hyd.). n=30-76 flies. Water consumption: \*\*\*p<0.001, t-tests (23°C vs 32°C) with Holm-Sidak correction.

B. ISN silencing with *nSyb* RNAi affects water and sugar consumption. n=38-57 flies. B-D: one-way ANOVA, Tukey’s Post Hoc, \*\*\*p<0.001.

C. ISN-specific *nan* RNAi affects water and sugar consumption. n=37-67 flies.

D. ISN-specific *AKHR* RNAi affects water and sugar consumption. n=22-70 flies.

E. Desiccation affects sugar consumption in wild type flies. n = 84-101, \*\*\*p<.001, t-test (hyd. vs. des.).

F. Low osmolality (osmo) and high AKH increase ISN activity, which promotes sugar consumption and inhibits water consumption. Insulin indirectly inhibits ISNs. High osmo and low AKH reduce ISN activity to promote water consumption and inhibit sugar consumption. See Figure S5 for additional behavioral experiments.