

UNIVERSITY OF CALIFORNIA  
SANTA CRUZ

**The Role of Land Use in *Aedes* abundance and Insecticide Resistance in Central Africa and an Analysis of Arbovirus Transmission Potentials**

A dissertation submitted in partial satisfaction  
of the requirements for the degree of

DOCTOR OF PHILOSOPHY

in

ECOLOGY AND EVOLUTIONARY BIOLOGY

by

**Matthew J. Montgomery**

March 2023

The Dissertation of Matthew  
Montgomery is approved:

---

Professor Auston M. Kilpatrick, chair

---

Professor Chris Barker

---

Professor Bruce Lyon

---

Professor Jean Tsao

---

Peter Biehl  
Vice Provost and Dean of Graduate Studies



## Table of Contents

List of Tables.....	iv
List of Figures.....	v
Abstract.....	vii
Acknowledgements.....	ix
Introduction.....	1
Chapter 1: The effects of urbanization on <i>Aedes aegypti</i> and <i>Aedes albopictus</i> abundance across a broad latitudinal gradient in Central Africa.....	7
Chapter 2: Spatial distribution of insecticide resistant populations of <i>Aedes aegypti</i> and <i>Ae. albopictus</i> and first detection of V410L mutation in <i>Ae. aegypti</i> from Cameroon.....	34
Chapter 3: Comparing relative transmission potential to population seroprevalence for seven arboviruses: yellow fever, chikungunya, Zika, and four dengue serotypes.....	47
Conclusion.....	66
Appendix 1: Chapter 1 Supplemental Materials.....	71
Appendix 2: Chapter 2 Published Material Co-Author Approval.....	86
Appendix 3: Chapter 3 Supplemental Materials.....	87
Bibliography.....	113

## List of Tables

Table 1 Primer sequences for the evaluation of the level of expression of metabolic resistance genes by RT-qPCR and their origins in <i>Aedes albopictus</i> .....	38
Table 2 Primer sequences for the evaluation of the level of expression of metabolic resistance genes by RT-qPCR and their origins in <i>Aedes aegypti</i> .....	38
Table 3 1016 genotype numbers and the allelic frequency of the Isoleucine (I) mutation of <i>Aedes aegypti</i> .....	43
Table 4 1534 genotype numbers and the allelic frequency of the Cysteine (C) mutation of <i>Aedes aegypti</i> .....	43
Table 5 410 genotype numbers and the allelic frequency of the Leucine (L) mutation of <i>Aedes aegypti</i> .....	44
Table 3.1 Analysis of global seroprevalence data using a generalized linear model with a binomial distribution and a logit link with virus type as a fixed effect and study ID as a random effect.....	59

## List of Figures

Figure 1.1 Cities surveyed for adult mosquitoes in Cameroon.....	26
Figure 1.2 Larval habitat plotted against UI at 100m for five of the study cities.....	27
Figure 1.3 <i>A. aegypti</i> abundance on a log scale plotted against the urbanization index in a 1km radius around the study site for ten sites in each of six cities.....	28
Figure 1.4 <i>A. aegypti</i> abundance on a log scale plotted against temperature at the time of collection.....	29
Figure 1.5 <i>A. albopictus</i> (95% CI) abundance on a log scale plotted against the urbanization index in a 2km radius around the study site in each of three cities where <i>A. albopictus</i> was found.....	30
Figure 1.6 <i>A. albopictus</i> abundance on a log scale plotted against relative humidity at the time of collection.....	31
Figure 1.7 Total <i>A. aegypti</i> and <i>A. albopictus</i> abundance in response to prior month's rainfall across all sites.....	32
Figure 1.8 Locations of studies of <i>A. albopictus</i> abundance in response to urbanization.....	33
Figure 1 Map of study sites in six cities in Cameroon.....	37
Figure 2 Mortality rates of adult <i>Aedes aegypti</i> from urban and suburban habitats in six cities in Cameroon 24 h after exposure to insecticides alone or with 1h pre-exposure to synergist.....	39
Figure 3 <i>Aedes aegypti</i> insecticide mortality ( $\pm 1$ SE) between study cities and habitats.....	40
Figure 4 Mortality rates of adult <i>Aedes albopictus</i> from urban and suburban habitats in three cities in Cameroon 24 h after exposure to insecticides alone or with 1 h pre-exposure to synergist. ....	41
Figure 5 Expression of the CYP9M6F88/87, CYP9J28a, CYP9J10, CYP9J32 genes in samples of <i>Aedes aegypti</i> from urban Yaounde and urban and suburban Douala...	42

Figure 6 Expression of the CYP6P12 gene in samples of <i>Aedes albopictus</i> from urban Yaounde, suburban Kribi, and urban Douala.....	42
Figure 7 Frequency of the 1534C allele kdr mutation in <i>Aedes aegypti</i> across study cities by year.....	44
Figure 3.1 Human viremia (PFU/mL) by date of symptom onset.....	60
Figure 3.2 Predicted fraction of <i>A. aegypti</i> transmitting virus by blood meal viremia.....	61
Figure 3.3 Daily estimates of human infectiousness ( $\pm 1$ SE) for seven arboviruses for <i>A. aegypti</i> and <i>A. albopictus</i> .....	62
Figure 3.4 Relative seroprevalence for studies simultaneously comparing CHIKV, YFV, and ZIKV, CHIKV and DENV1-2, DENV1-4, and all remaining studies with studies ranked by average seroprevalence.....	63
Figure 3.5 Fitted coefficients of relative seroprevalence plotted against relative $R_0$ estimates for <i>A. aegypti</i> .....	64
Figure 3.6 Estimated seroprevalence for each arbovirus for an intermediate value of average seroprevalence.....	65

**Abstract**  
**The Role of Land Use in *Aedes* abundance and Insecticide Resistance in Central Africa and an Analysis of Arbovirus Transmission Potentials**  
**Matthew J. Montgomery**

Globally, mosquito-borne diseases are a major source of mortality, with the burden focused primarily in the poorest nations. Central to reducing mosquito borne disease is an understanding of the biology of those species responsible for transmitting disease to humans. Accordingly, this work focuses on the ecology and evolutionary biology of two vectors of global importance: *Aedes aegypti* and *Aedes albopictus*.

The first chapter examines the role of urbanization and species interactions in determining *Aedes* abundance across a 900km latitudinal range representing vast climatic gradients and the major ecotypes of Central Africa. Additionally, we performed pathogen screening on over 9,000 female mosquitoes for dengue, chikungunya, and Zika viruses.

The second chapter focuses on the distribution and potential causes of insecticide resistance in *Aedes* across the same broad climatic and ecotype gradient in Cameroon. In this chapter we detected significant variation in insecticide resistance between cities, habitat types, species, and insecticide class. We applied land use data to show the significant effect that urbanization can have on determining insecticide efficacy across an area as small as a single city and further revealed the underlying genetic and metabolic mechanisms for insecticide resistance.

The final chapter investigates viral fitness among the primary arboviruses transmitted by *A. aegypti* and *A. albopictus*: chikungunya, dengue (serotypes 1-4),

yellow fever, and Zika. Our innovative approach to comparing relative viral fitness is multi-facted, combining field observations of human infection rates with laboratory studies of host and vector competence.



## Acknowledgements

Firstly, I would like to thank my advisor Dr. Marm Kilpatrick for the tremendous help he has given me throughout this process and for allowing me the privilege to be a part of the Kilpatrick Lab. He is one of the most brilliant minds I have ever encountered and I have learned so much from him over the course of my graduate studies. As part of the Kilpatrick Lab, I also had the benefit of working with fellow graduate students Christa Seidl, Billy Gardner, and Nikka Malakooti who provided constructive input and advice throughout our time together. I also must thank my committee members, Dr. Bruce Lyon, Dr. Chris Barker, and Dr. Jean Tsao for not only serving on my committee, but for the expertise they provided in shaping my research projects at their early stages and guiding me throughout the process.

The Center for Research in Infectious Disease (CRID) in Yaounde, Cameroon performed the vast majority of actual field and laboratory work that composed the first two chapters. Their scientific expertise was invaluable to the completion of the research.

My second chapter, Spatial distribution of insecticide resistant populations of *Aedes aegypti* and *Ae. albopictus* and first detection of V410L mutation in *Ae. aegypti* from Cameroon, utilizes previously published materials with the permission of my co-authors (see Appendix) and my advisor who is an additional co-author on the published article. This work was based on a Department of Defense funded grant for a study which I designed and was the Principal Investigator. All field and lab work was performed by CRID via a contract funded by the aforementioned grant. I

performed the statistical analysis and wrote and created the figures for the majority of the manuscript.

On a personal note, I could not have completed this dissertation without the help of my family. My brother Tim selflessly offered his skills in GIS for calculating the urbanization indices in Chapter 2. I also must thank the two most important women in my life. My late mother, Lynda, always instilled in me the value of higher education and she would be so happy to know that I have earned my doctoral degree. Lastly my wife, Lynné provided invaluable support and encouragement throughout these years.

## Introduction

Globally, mosquito-borne diseases are a major source of mortality, with the burden focused primarily in the poorest nations. In the early half of the 20th century major progress was made towards reducing mosquito-borne disease burden through improved vector control, vaccines, and the development of new medications. However, in recent years emerging pathogens have swept across the globe in explosive epidemics, as demonstrated by West Nile Virus in 1999, the 2013 chikungunya outbreak, Zika virus in 2015, and dengue's rapidly increasing global burden (Lanciotti et al. 1999, Shragai et al. 2017, Wahid et al. 2017, WHO 2020). Central to reducing mosquito borne disease is an understanding of the biology of those species responsible for transmitting disease to humans. Accordingly, my dissertation work focuses on the ecology and evolutionary biology of two vectors of global importance: *Aedes aegypti* and *Aedes albopictus*. Both of these species are found in tropical to temperate regions on six continents, and they are the primary vectors of chikungunya, dengue, Yellow Fever, and Zika viruses as well as other human pathogens.

The first two chapters of my dissertation are focused on the biology of *A. aegypti* and *A. albopictus* in Cameroon, in central Africa. While these studies take place in a single nation, they were conducted over a large longitudinal gradient and across a range of different ecotypes and climates representative of a wide area of the African continent, with habitats ranging from tropical equatorial forests to the dry Sahel. The final chapter takes a global approach to the major viruses spread primarily

by these species and investigates how our understanding of the factors which predict a mosquito-borne disease epidemic correlate with empirical observations of human infections across the globe.

The first chapter examines the role of urbanization in determining *Aedes* abundance. Urbanization, defined as converting land from natural or agricultural land uses to areas built upon by humans, is among the most drastic and consequential form of human land use and plays a pivotal role in shaping the biotic and abiotic conditions faced by mosquitoes (Gubler et al. 2011). Previous research has shown that urbanization can influence mosquito abundance and distributions, and pathogen transmission to humans (Alirol et al. 2011). Consequently, it is imperative that the ecological factors leading to increased mosquito populations and thus vector borne disease risk be better understood.

The chapter begins with a literature review of the effects of urbanization on *Aedes aegypti* and *A. albopictus* abundance from studies outside of Africa. These studies show conflicting patterns in the response of *A. albopictus* to urbanization (and consistent increases in abundance for *A. aegypti*). We sought to investigate this highly varied response to urbanization by *A. albopictus* on the African continent. Additionally, we wanted to determine if *A. aegypti*'s abundance in response to urbanization would vary based on climatic factors or the presence of a competitor species. To address these questions, we obtained data on mosquito abundance from 231 collections across six cities spanning nearly 1,000km of latitudinal extent over a one year period. The six surveilled cities varied vastly in climate and ecotype, ranging

from coastal equatorial forests to the Sahel. Our results showed that urbanization increases abundance for *A. aegypti* throughout Cameroon, consistent with global patterns. Surprisingly, the slope of *A. aegypti* abundance in response to urbanization remained consistent across all cities even though the intercepts varied. Yet for *A. albopictus* there was heterogenous response to urbanization even within a single country: *A. albopictus* abundance increased with urbanization in one city, decreased with urbanization in another and showed no trend in a third city. To better understand the mechanisms of urbanizations effect on adult mosquito abundance we performed further analysis on climatic factors, larval habitat, structures, host abundance, and available host community composition.

Chapter 1 also included a wide scale effort to detect and identify the major viruses transmitted by *Aedes* mosquitoes (chikungunya, dengue, and Zika) across their global range in field caught populations in Cameroon and how urbanization could effect mosquito viral infection rates. Our extensive surveillance resulted in only a single detection of Zika virus among hundreds of pools containing many thousands of individual specimens. As such, our data were unable to address any hypotheses on how urbanization could influence disease distribution. This study supports numerous prior instances showing that field-caught mosquitoes generally have exceedingly low infection rates, even during periods where surveillance was concurrent with active human outbreaks. Reassuringly, our study shows that infection rates for chikungunya, dengue, and Zika were exceedingly low even in the absence of active public health vector control measures.

The second chapter focuses on the distribution and potential causes of insecticide resistance in *Aedes* across the same broad climatic and ecotype gradient in Cameroon. Public health efforts to mitigate vector borne disease often include insecticide use to reduce vector populations. However, mosquitoes can rapidly develop resistance to insecticides (Hamdan et al. 2005), which reduces their effectiveness (Ranson et al. 2011). Furthermore, there are multiple physiological and genetic mechanisms to insecticide resistance and these mechanisms can vary both between species and within geographically dispersed populations of a single species (Kamgang et al. 2011; Dusfour et al. 2019). Lastly selective pressure against resistance may occur through reduced mating performance, fecundity, or survival (Dusfour et al. 2019; Rigby et al. 2021). This may lead to fluctuations in insecticide resistance depending on insecticide usage in a region (Dusfour et al. 2019). Understanding the regional and species-specific susceptibility of mosquitoes to insecticides of public health importance is vital for converting knowledge of disease risk into effective mitigation strategies (Ranson et al. 2011; World Health Organization 2012; Lima et al. 2015; Bouzid et al. 2016; Alvarado-Castro et al. 2017).

In this chapter we detected significant variation in insecticide resistance between cities, habitat types, species, and insecticide class. We applied land use data to show the significant effect that urbanization can have on determining insecticide efficacy across an area as small as a single city and further revealed the underlying genetic and metabolic mechanisms for insecticide resistance. Our analysis also

includes temporal changes of allele frequency for a particular resistance mutation which shows increase in resistance allele frequency and distribution over rapid time scales across Cameroon, showing the high selective pressure for insecticide resistance in *A. aegypti*. We also detected the first instance of a particular resistance mutation, V410L, in central Africa. This mutation is found on other continents, but was only detected in Africa in distant (>1,400km) Angola in 2016, which indicates the high degree of gene flow between populations or far-reaching human-facilitated introduction of populations on the African continent (Ayres et al. 2020).

The final chapter investigates viral fitness among the primary arboviruses transmitted by *A. aegypti* and *A. albopictus*: chikungunya, dengue (serotypes 1-4), yellow fever, and Zika. Our innovative approach to comparing relative viral fitness is multi-faceted, combining field observations of human infection rates with laboratory studies of host and vector competence.

We first determined the relative transmission intensity of co-occurring arboviruses in human populations through the presence of arbovirus antibodies in human serosurveys. To obtain this data, we conducted a literature review and compiled data from historical serosurveys, which are studies where human blood samples are tested for specific disease antibodies. After identifying 213 unique serosurveys that simultaneously tested for at least two of our study pathogens among over 95,000 individuals across 5 continents, we were able to compare relative infection rates among seven arboviruses. Our results show that there are significant

differences in transmission intensity of these co-occurring viruses within the same human populations.

Next, we compiled estimates of relative viral transmission efficacy calculations generated from mathematical models using a vast trove of host and vector competence data for simultaneous calculations via the calculation of the relative reproductive number  $R_0$  for each pathogen, which describes the number of secondary cases arising from an initial primary case in a fully susceptible population (Dye 1992).

We hypothesized that among these co-occurring pathogens, those with higher relative  $R_0$  values should infect a larger fraction of a given host population and should have higher relative seroprevalence. Surprisingly, our results showed no significant correlation between relative  $R_0$  and transmission intensity across the global dataset, and within continents relative  $R_0$  was only significantly correlated with greater seroprevalence in Asia. These results indicate a stark disconnect between estimates of pathogen fitness and field observations of pathogen infection rates.

In this body of research I hope to show how human modifications of the environment through urbanization and insecticide use shapes the disease vector landscape that we face as humans and to calculate and compare arbovirus fitness while simultaneously presenting a new way to evaluate predictions of viral fitness using real world observations of human infection rates.



**Chapter 1: The effects of urbanization on *Aedes aegypti* and *A. albopictus*  
abundance across a broad latitudinal gradient in Central Africa**

**Abstract**

Urbanization shapes the biotic and abiotic conditions faced by many species. The effects of urbanization on disease vectors may increase or decrease their abundance and disease risk by altering larval habitat, microclimates, and host abundance. We investigated the effect of urbanization and species interactions on the abundance and disease risk posed by two cosmopolitan mosquito vectors of global health concern, *A. aegypti* and *A. albopictus*, in Cameroon, Central Africa. We collected adult mosquitoes during both the rainy and dry seasons at 63 sites on a rural to urban gradient from six cities spanning a 900 km latitudinal range from equatorial forest to the Sahel. We also measured larval habitat, host abundance, multiple measures of urbanization, and climate data. Urbanization increased larval habitat, the number of structures, and host availability, but a GIS urbanization index was the best predictor of mosquito abundance. *A. aegypti* abundance increased with rainfall and decreasing temperatures and increased 2.7% with each 1% of additional urbanization in all six cities. In contrast, *A. albopictus* abundance increased, decreased or showed no influence of urbanization in the three cities where this species was present; abundance also increased with rainfall and humidity. We found no evidence of interspecific competition among these species using adult abundance data, but analyses were limited spatially. We screened over 9,000 female mosquitoes for dengue,

chikungunya, and Zika viruses but only detected one virus (Zika) in a single mosquito pool. Our results show that urbanization consistently increases *A. aegypti* abundance across a broad range of habitats, while urbanization effects on *A. albopictus* abundance varied markedly over a relatively small geographic area. Urbanization is thought to increase mosquito abundance through mediating availability of larval habitat, temperature, humidity, number of structures, and host availability, yet our results showed temperature and humidity at the time of collection were not significantly correlated with urbanization and larval habitat, host availability, and number of structures were either not significantly correlated with abundance or were less effective predictors of abundance than the urbanization index. These results indicate that the precise mechanisms of urbanization effect on adult *A. aegypti* and *A. albopictus* abundance are not fully understood.

## **Introduction**

Urbanization, the conversion of natural and agricultural habitats to urban areas, alters the biotic and abiotic conditions faced by organisms and is increasing globally (Grimm et al. 2008; Alig et al. 2004; Chen et al. 2022). Urbanization often reshapes community composition, with introduced and human commensal species replacing native species (McKinney 2006). Urbanization also alters microclimates, including increasing temperatures through the Heat Island effect (Chapman et al. 2017, Deilami et al. 2018), and reducing humidity (McCluney et al. 2017). These changes in the biotic and abiotic environment also influence the transmission of pathogens (Bradely & Altizer 2007, Gottdenker et al. 2014, Hassel et al. 2017), and

especially vector-borne pathogens (Kilpatrick & Randolph 2012). However, the effect of urbanization on vector-borne disease depends on both the response of the vectors to urbanization and the microclimate effects on pathogen development in the vector, and both these factors can increase or decrease with urbanization depending on the vectors, the pathogen, and the local climate (Roger & Randolph 2000, Mordecai et al. 2013, Kraemer et al. 2015, Mordecai et al. 2017). Africa is a key area for understanding the effect of urbanization on vector borne disease because it has the world's highest rates of urban growth (OECD/SWAC 2020). *Aedes*-borne arboviruses, including dengue, chikungunya, and zika viruses, are an especially important group of pathogens because of recent increases in incidence and outbreaks (Bhargavi & Moa 2020, ECDC 2022, WHO 2022).

Urbanization can increase larval habitat, especially for key anthropogenic species like *Aedes aegypti* and *Aedes albopictus*, either through more impervious surfaces or through increased refuse, containers, and other objects which retain water (Braks et al. 2003; Morrison et al. 2004; Higa et al. 2010; Gubler et al. 2011; Li et al. 2014; Arduino et al. 2020). Furthermore, urbanization is often associated with storing water use throughout the year, which can provide mosquito larval habitat in periods with low rainfall, especially in arid climates (Faeth et al. 2005, Gubler et al. 2011; McCluney et al. 2017). Urbanization in poorer countries may generate more larval habitat compared to more developed nations due to poor sanitation, increased refuse, and more personal water storage, thus increasing mosquito abundance and disease risk (Service 1992; Gubler et al. 2011; Obenauer et al. 2017).

Temperature increases associated with urbanization can alter mosquito community composition and pathogen transmission through multiple mechanisms. Temperature tolerance (both minimum and maximum, as well as daily variation) vary considerably between mosquito species (Reinhold et al. 2018) and temperature influences survival (Bar-Zeev 1957; Couret et al. 2014), biting rate (Christophers 1960; Scott et al. 2000), development rate (Bar-Zeev 1958; Delatte et al. 2009), fecundity (Delatte et al. 2009; Carrington et al. 2013), vector competence (Lambrechts et al. 2011; Reinhold et al. 2018) and the extrinsic incubation period (Watts et al. 1987; Kilpatrick et al. 2008; Rohani et al. 2009; Reinhold et al. 2018). Increased temperature caused by urbanization can decrease larval development time and adult survival which can have opposing effects on transmission (Costa et al. 2010, Kilpatrick & Randolph 2012, Mordecai et al. 2013). In contrast, decreased humidity generally reduces mosquito survival without any positive effects on transmission (McCluney et al. 2017). Urbanization can also alter interspecific competition among mosquito larvae. For example, larval habitat that entirely dries out favors *A. aegypti* over *A. albopictus*, whereas the inverse is true in larval habitat that never entirely dries out (Costanzo et al. 2005).

The effect of urbanization on mosquito abundance can vary among species. *A. aegypti* abundance consistently increases with urbanization across its global range, with very few exceptions (Table S1). In contrast, *A. albopictus* abundance increased with urbanization in half of 15 previous studies, decreased in six, and had no effect in one (Table S2). Specifically, *A. albopictus* abundance was higher in urban areas than

rural areas in China (Li et al. 2014), and Mayotte (Bagny et al. 2012), but abundance was highest in suburban/transitional areas in Virginia (Barker et al. 2003) and Thailand (Tsuda et al. 2006) and rural areas in Brazil (Honorio et al. 2009) and Florida (Hornby et al. 1994). A key area where the effect of urbanization on *A. albopictus* has not been studied is the continent of Africa.

Our goal was to examine the effect of urbanization on the abundance and arbovirus infection rates of *A. aegypti* and *A. albopictus* mosquitoes across a broad latitudinal gradient in central Africa. We examined adult mosquito abundance, microclimate, and larval habitat at ten sites in each of six cities spanning the 900km latitudinal extent of Cameroon and tested *Aedes* mosquitoes for dengue, chikungunya, and Zika viruses which are important arboviruses and all circulate in this region (Kuniholm et al. 2006). We studied adult mosquitoes rather than mosquito larvae, which have been the focus of some earlier work (Bagny et al. 2012; Higa et al. 2010; Honorio et al. 2009; Tsuda et al. 2006), so we could detect pathogens and because adult mosquito abundance is often a better predictor for outbreaks than larval indices (Leandro et al. 2022). We hypothesized that *A. aegypti* would increase with urbanization, but we predicted the relationship would vary among cities because the benefits of urbanization might be counterbalanced in areas that were very hot and dry. We hypothesized that *A. aegypti* abundance might also be influenced by *A. albopictus*, which itself might show increasing or decreasing relationships with urbanization. While *A. aegypti* is native to Cameroon, *A. albopictus* is an introduced species that was first detected in 2000 (Simard et al. 2005). In some regions of the

world *A. albopictus* competes with and can displace *A. aegypti* populations through larval competition (Shagrai et al. 2017) and reproductive interference (Tripet et al. 2011). We examined multiple scales of urbanization on mosquito abundance and hypothesized that small-scale measures of urbanization (100-200m rather than >1km) would most closely correlate with abundance, based on the short-distance dispersal distances of *A. aegypti* and *A. albopictus* in mark-recapture studies (Medeiros et al. 2017, Juarez et al. 2020) .

## **Methods**

**Study area.** We collected mosquitoes in six cities spanning a 900 km latitudinal gradient in Cameroon (Figure 1) representing five African ecoregions: Atlantic equatorial coastal forest (Douala and Kribi), Congolian lowland forest (Yaounde), Northern Congolian Forest Savannah mosaic (Ngaoundere), East Sudanese Savannah (Garoua), and Sahelian (Maroua) (World Data Base on Protected Areas 2022). Average annual temperatures and relative humidity varied between 18°C to 28°C and 85% to 45%, respectively for our study cities. Cities range in altitude from 2m to 1200 m (Olivry, 1986). Within each city we surveyed at a minimum of ten sites across an urbanization gradient from forest to agrarian to dense urban centers.

**Land use analysis.** For each site we quantified urbanization at 100m, 200m, 500m, 1km, and 2km radii using an Urbanization Index (UI) (Gomez et al. 2008), where  $UI = (100\% - \% \text{ vegetation cover} + \% \text{ impervious surface})/2$ .

We performed spatial analysis using ArcGIS Pro version 2.8.7. We imported GPS field coordinates into ArcGIS Pro and spatial landcover data was obtained from the European Space Agency's (ESA) Climate Change Initiative (CCI) – S2 Prototype Land Cover 20M map of Africa which included 10 classes of land surface data at a 20 meter resolution and were classified as follows: "trees cover areas", "shrubs cover areas", "grassland", "cropland", "vegetation aquatic or regularly flooded", "lichen and mosses / sparse vegetation", "bare areas", "built up areas", "snow and/or ice" and "open water" (Figure S1). We used satellite imagery (Google Earth Pro version 7.3.4.8642) to visually quantify the number of structures at the 100m radius for each surveillance site.

**Larval habitat survey.** At each of the ten surveillance sites (with three exceptions) in five of the six cities (all except Yaounde) we recorded potential larval habitat within 50m of adult collection sites. We categorized and sampled any container with mosquito larvae present, and any larvae collected were reared to adulthood and identified to species for *A. aegypti* and *A. albopictus*.

**Adult mosquito collection.** We collected adult mosquitoes outdoors using two stationary human collectors using Modified CDC Backpack Aspirators (John Hock W. Hock Company, Gainesville, Florida). We used human landing catches, rather than passive traps, to increase the number of adult *Aedes* captured and thereby to increase our chances of detecting and accurately measuring arbovirus infection rates

(Schoeler et al. 2004; Scott & Morrison 2009). We conducted collections for 3 hours during daytime at consistent three hour periods (8am-11am and 3pm-6pm) and we recorded temperature and humidity at the starting time of each collection. We chose collection times based off of previous work in Africa showing both species were diurnal and that *A. aegypti* and *A. albopictus* activity peaked in both the morning and early evening (McClelland 1959; Trpis et al. 1973; Delatte et al. 2010). Collections took place over a one week period in each city once during both the dry and rainy seasons from September 2020-August 2021. When possible, we repeated collections a second time at each site during the same collection week. In Yaounde, we visited sites more frequently (ten times between September 2020 and October 2021). At the end of each collection period, we knocked down adult mosquitoes at  $-20^{\circ}\text{C}$  or using a chloroform solution and identified using them morphological identification keys (Edwards, 1941; Jupp, 1996) under a microscope. We grouped *Aedes* females by species and pooled in groups of up to 30 individuals and stored at  $-80^{\circ}\text{C}$  in RNAlater following the manufacturer's instructions (LifeTechnologies, 2011) for viral detection. The rest were stored at  $-20^{\circ}\text{C}$  in silicagel.

We analyzed mosquito abundance using generalized linear models (GLM) with a negative binomial distribution and log link using the *lme4* package in R version 3.6.1. We compared models with different urbanization radii, closely correlated climatic parameters (e.g. monthly rainfall versus number of monthly rainy days), and interactive versus additive effects of variables using Akaike's Information Criterion (AIC).



**Pathogen detection.** We pooled female mosquitoes in Eppendorf tubes and first ground them in 200  $\mu$ L of Leibovitz L15 medium equilibrated at room temperature. Then, we centrifuged the sample for 15 min at 15,000 rpm and transferred the supernatant to a new Eppendorf tube. We extracted potential total viral RNA present in each pool using the Qiagen extraction kit (QIAamp-viral RNA mini kit, Qiagen) following the manufacturer's instructions and stored at  $-80^{\circ}\text{C}$ .

We retrotranscribed extracted RNAs into cDNAs using the High-Capacity cDNA Reverse Transcription kit, 1000 reactions (Applied Biosystems, Foster city, California, USA). We prepared a mixture of 50  $\mu$ L final volume, including 25  $\mu$ L of RNA sample, 5  $\mu$ L of 10X reverse transcription buffer, 2 $\mu$ L of 100 mM dNTPs, 5 $\mu$ L of 10X hexa random primers 10.5 $\mu$ L, 2.5  $\mu$ L of reverse transcriptase, and 10.5  $\mu$ L of RNase-free  $\text{H}_2\text{O}$ . We incubated the samples for 10 min at  $25^{\circ}\text{C}$  and 1 h at  $37^{\circ}\text{C}$ .

We performed quantitative real-time PCR using TaqMan Universal PCR Master Mix reagents (Applied Biosystems, Foster city, California, USA) to amplify UTR genes for Zika (Conceição *et al.*, 2010; Grard *et al.*, 2014) and Dengue viruses (Leparac-Goffart *et al.*, 2009), and the E1 envelope protein for chikungunya virus (Ngoagouni *et al.*, 2017) (Table S3). Each PCR was performed in a 25  $\mu$ L reaction mixture containing 12.5 $\mu$ L of 2x PCR TaqMan Universal PCR Master Mix, 1 $\mu$ L of each primer, 1  $\mu$ L of each TaqMan probe (Applied Biosystems, Foster city, California, USA), 4.5  $\mu$ L of RNase-free  $\text{H}_2\text{O}$ , and 5  $\mu$ L template of cDNA. The amplification program consisted of a pre-activation heat step of 2 min at  $50^{\circ}\text{C}$  followed by 10 min at

95°C, and the amplification and fluorescence quantification steps of 45 cycles of 15 sec at 95°C and 1 min at 60°C. All assays were done on the Stratagene MX3005P qPCR machine (Agilent Technologies, Santa Clara, California, USA).

**Host availability.** In September of 2021 in Douala, Kribi, and Yaounde we counted potential hosts using two stationary observers between 8am-11am and 3pm-6pm on two separate days at each established surveillance site. We recorded all humans, chickens, goats, cows, and dogs visible over a 5 minute period. We counted human hosts to a maximum of 100 due to extremely large crowds in some sites rendering more accurate counts unfeasible.

**Indoor vs outdoor mosquito abundance surveys.** We conducted paired surveillance of an outdoor site with simultaneous mosquito collections inside 10 nearby homes at 10 sites in Yaounde in October of 2021 to investigate whether outdoor collections correlated with indoor household vector abundance. Outdoor mosquito collection took place for 3 hours as described above, while we surveyed the interior of willing households within 50 meters of the outdoor sites using the same Modified CDC Backpack Aspirators.

**Competition between *A. aegypti* and *A. albopictus*.** We calculated monthly population growth rates, ( $\lambda = N_{t+1}/N_t$ ), for each species for the two sites in Yaounde from which adequate numbers of both species were simultaneously collected. We then examined the potential effects of interspecific competition between *A. aegypti*

and *A. albopictus* by examining the effect of the abundance of *A. albopictus* on the population growth rate  $\log_{10}(\lambda)$  over the following month for *A. aegypti*, and vice versa. We analyzed the effect of mosquito abundance on population growth rates using generalized linear models (GLM) with a gaussian distribution and log link using the *lme4* package in R version 3.6.1.

## **Results**

**Larval Habitat Survey.** The number of containers increased with urbanization and the urbanization index at 100m (UI\_100m) was the best correlate of both containers ( $\text{Log}_{10}(\text{containers})=3.04+0.0067 (\pm 0.0021) * \text{UI}_{100\text{m}}$ ,  $P = 0.0020$ ; Figure 2) and larvae-positive containers ( $\text{Log}_{10}(\text{positive\_containers})=2.34+0.0078 (\pm 0.0024) * \text{UI}_{100\text{m}}$ ,  $P = 0.0012$ ). On average a site with UI of 0 would be expected to have 21 larval containers while a site with a UI of 100 would have 41. There was no significant relationship between the Container Index (larvae occupied containers/available containers) and UI ( $\text{logit}(\text{CI})=-0.05+0.0028 (\pm 0.0051) * \text{UI}_{100\text{m}}$ ,  $P = 0.57$ ).

**Host Availability.** Human abundance increased significantly with UI and 100m was the best correlate ( $\text{Log}_{10}(\text{Humans})=2.83+0.0097 (\pm 0.00085)*\text{UI}_{100\text{m}}$ ,  $P \leq 0.001$ ), while animal abundance was negatively correlated with UI ( $\text{Log}_{10}(\text{Animals})=1.7-0.025 (\pm 0.0035)*\text{UI}_{100\text{m}}$ ,  $P \leq 0.001$ ). On average a site with UI of 0 would be

expected to have 17 human hosts present while a site with a UI of 100 would have 45 (Figure S2).

**Mosquito Abundance.** We collected 25,853 adult mosquitoes from 231 two-person collection sessions from 63 sites across 6 cities (Table S4). Of these mosquitoes, 15,008 were *A. albopictus* and 2,633 were *A. aegypti*. No *A. albopictus* were found in our three study cities above 6° latitude (Ngaoundere, Garoua, and Maroua) (Table S5).

*A. aegypti* abundance increased with the UI in all cities (Figure 3), increased with rainfall, and decreased with temperature (Figure 4); humidity had no significant effect (Table S6). The best fitting model used the UI at 1km (UI\_2km:  $\Delta\text{AIC} = 4.080$ ; UI\_500m:  $\Delta\text{AIC} = 10.30$ ; UI\_200m:  $\Delta\text{AIC} = 23.13$ ; UI\_100m:  $\Delta\text{AIC} = 18.082$ ). The slope of the response to urbanization was consistent across cities (an additive model was preferred over a city \* urbanization interactive model;  $\Delta\text{AIC} 2.90$ ), with an average 2.7% increase in *A. aegypti* females for each corresponding 1% increase in urbanization. The intercepts for average *A. aegypti* abundance varied considerably between cities, with the greatest average abundance in Douala and the lowest in Yaounde. Abundance also increased with rainfall in the month prior ( $\text{Log}_{10}(\text{Abundance})=2.69+0.0019 (\pm 0.00071) * \text{Rainfall}$ ,  $P = 0.0056$ ) (Figure 5) and in the city of Yaounde, *A. aegypti* abundance peaked during the month of highest rainfall (Figure S3).

*A. aegypti* abundance was significantly correlated with the number of containers ( $\text{Log}_{10}(\textit{A. aegypti} \text{ Abundance})=0.43+0.017 (\pm 0.0059)*\text{Containers}$ ,  $P = 0.0032$ ) and larvae-positive containers ( $\text{Log}_{10}(\textit{A. aegypti} \text{ Abundance})=1.13+0.041 (\pm 0.010)*+\text{Containers}$ ,  $P \leq 0.001$ ), but not the number of structures in a 100m radius ( $\text{Log}_{10}(\textit{A. aegypti} \text{ Abundance})=4.85-0.0057 (\pm 0.0035)*\text{Structures}$ ,  $P = 0.10$ ). *A. aegypti* abundance also increased with human abundance ( $\text{Log}_{10}(\textit{A. aegypti} \text{ Abundance})=0.91+0.021(\pm 0.007) * \text{Humans}$ ,  $P = 0.018$ ). The time of collection had no significant effect ( $\text{Log}_{10}(\textit{A. aegypti} \text{ Abundance})=1.17 -0.58 (\pm 0.36) * \text{Time}$ ,  $P = 0.10$ )

*A. albopictus* showed heterogeneity in its response to urbanization in the three cities where it was found (Figure 6). The 2km UI was the best performing measure of urbanization (UI\_1km:  $\Delta\text{AIC} = 4.86$ ; UI\_500m:  $\Delta\text{AIC} = 10.40$ ; UI\_200m:  $\Delta\text{AIC} = 8.24$ ; UI\_100m:  $\Delta\text{AIC} = 4.13$ ). In Yaounde, *A. albopictus* increased with urbanization, but it decreased with urbanization in Douala and there was no significant pattern in either direction in Kribi. *A. albopictus* abundance also increased with rainfall, and humidity (Figure 7); temperature had no significant effect (Table S7).

*A. albopictus* abundance was not significantly correlated with the number of larval containers ( $\text{Log}_{10}(\textit{A. albopictus} \text{ abundance})=-6.74+0.0027 (\pm 0.0040)*\text{Containers}$ ,  $P = 0.49$ ), larvae positive containers ( $\text{Log}_{10}(\textit{A. albopictus} \text{ abundance})=-6.55+0.0038 (\pm 0.0074)*+\text{Containers}$ ,  $P = 0.60$ ) or the number of structures in a 100m radius ( $\text{Log}_{10}(\textit{A. albopictus} \text{ abundance})=3.76-0.0025 (\pm$

0.0024)\*Structures, P=0.29). *A. albopictus* abundance increased with human density ( $\text{Log}_{10}(A. albopictus \text{ abundance})=3.34+0.014 (\pm 0.0027)*\text{Humans}$ ,  $P \leq 0.001$ ). The time of collection had no significant effect ( $\text{Log}_{10}(A. albopictus \text{ Abundance})=3.022+0.059 (\pm 0.22) * \text{Time}$ ,  $P = 0.79$ ).

Combined abundance of *A. albopictus* and *A. aegypti* positively correlated with increased urbanization at the 2km radius in the three cities where both species coexist ( $\text{Log}_{10}(Aedes \text{ abundance})=-2.99+0.021 (\pm 0.0035)*\text{UI}_{2\text{km}}$ ,  $P \leq 0.001$ ). UI at 2km was the most parsimonious measurement for predicting the response of both species' abundance with urbanization (UI\_1km:  $\Delta\text{AIC} = 2.25$ ; UI\_500m:  $\Delta\text{AIC} = 4.015$ ; UI\_200m:  $\Delta\text{AIC} = 3.65$ ; UI\_100m:  $\Delta\text{AIC} = 1.89$ ).

The Urbanization Index at any radii was not significantly correlated with temperature (Table S8) or humidity (Table S9) in any of our study cities.

**Pathogen Detection.** We tested 1,660 *A. aegypti* and 7,771 *A. albopictus* females in 402 pools for chikungunya, dengue, and Zika viruses. Zika virus was found in one pool of 30 *A. albopictus* from Yaounde from June of 2021. No chikungunya or dengue virus was detected.

**Indoor/Outdoor Abundance.** No *A. aegypti* or *A. albopictus* were caught during indoor collections, while simultaneous outdoor collections caught a total of 28 and 397 respectively across all 10 outdoor surveillance sites.

**Competition between species.** There was no significant effect of *A. albopictus* abundance on the population growth rate  $\log_{10}(\lambda)$  of *A. aegypti* nor was there a significant effect of *A. aegypti*'s abundance on  $\log_{10}(\lambda)$  for *A. albopictus* over the next month (Figure S4). These results remained qualitatively consistent when season and rainfall were added as predictors.

## **Discussion**

We found that the effect of urbanization on mosquito abundance varied depending on the species of mosquito, but urbanization had a positive effect on combined *A. aegypti* and *A. albopictus* abundance. For *A. aegypti*, urbanization increased adult mosquito abundance consistently across the extent of Cameroon, which is broadly representative of the major ecotypes of sub-Saharan Africa and has vast climatic differences between the different cities in our study. We had predicted that the increase with urbanization would be smaller in the hottest, most arid environments, this hypothesis was not supported. We had based our hypotheses that *A. aegypti* abundance would differ in response to urbanization across cities based off of our expectation of finding a heat-island effect which would limit the benefits of urbanization in more northern cities, however we found no relationship between UI and temperature at the time of collection. We also found no support for competition among *A. aegypti* and *A. albopictus*, although our analysis was limited to just two sites in the city of Yaounde.

We investigated several mechanisms by which urbanization could influence mosquito abundance, including larval habitat, host abundance, and microclimates. Urbanization increased both larval habitat and host abundance, but the GIS urbanization index was more strongly correlated with *A. aegypti* and *A. albopictus* abundance than all of these predictors, indicating that urbanization influenced abundance through additional pathways beyond those we measured. Furthermore, adult *A. albopictus* abundance was not significantly correlated with either metric of larval habitat measured in our study. Thus the variability of *A. albopictus*'s varied response to urbanization across cities cannot be explained by differences in larval habitat or host availability between cities, nor can climatic factors explain this variation given that Kribi and Douala share a comparable coastal climate. In fact we have no mechanistic explanation for this pattern despite our study's robust collection of parameters thought to be associated with urbanization or mosquito abundance.

The spatial scale of urbanization that best correlated with mosquito abundance was relatively large, and varied between species (1km for *A. aegypti* and 2km for *A. albopictus*). This ran counter to our expectation that the 100m UI would correlate most strongly with abundance. These results indicate that urbanization affects populations on a scale larger than the typical dispersal distance of an individual mosquito. The spatial scales best correlated with other correlates of mosquito abundance varied as well, with containers and larvae positive containers best predicted by urbanization at 200m, while host availability was best correlated with urbanization at 500m.



During our study no Chikungunya or dengue virus was detected in either mosquito species, despite 41 documented concurrent human dengue cases in at least one city, Douala, at the time our study was conducted (Tchetgna et al. 2021). Zika virus was detected in a single pool from an urban area of Yaounde. A single detection precluded us from conducting any analysis on species infection rates or the possible effects of urbanization on disease positivity rates. Cameroon does not have active surveillance for human Zika cases. The most recent data on human incidence of Zika was a nationwide serological survey of 1,084 Cameroonian blood donors conducted in 2015 which identified Zika antibodies in 5% of the total population, with the city of Douala having the highest seroprevalence of 10% (Grake et al. 2015). Zika infection rates from our study were compared with Minimum Infection Rates (MIR) ( $[\text{number of positive pools} / \text{total specimens tested}] \times 1,000$ ) from prior entomological Zika surveillance studies of *A. aegypti* and *A. albopictus* (Table S10). Our results show an exceedingly low prevalence among Cameroonian *Aedes* compared to studies which detected Zika virus circulation in other regions.

Low rates of detection of entomological infection are typical for these pathogens and make studies of human disease risk difficult (Kuno et al. 1997; Scott & Morrison 2004). We tested a sizable sample of over 9,000 mosquitoes in 465 pools, which was a considerably greater number of mosquitoes tested compared with other studies which have used ~280-5,000 samples to detect dengue or chikungunya virus in *A. aegypti* and *A. albopictus* (Chow et al. 1998, Chung et al. 2002, Mendez et al. 2006, Edina et al. 2015, Vikram et al. 2015, Cevallos et al. 2018). Our large sample size and

no virus detection for Chikungunya or dengue and only one pool positive for Zika suggests that these viruses were either absent or circulating at low rates during the times and places we surveyed.

In Yaounde, we found no *A. aegypti* and *A. albopictus* during household surveys while simultaneous outdoor surveys captured numerous specimens. This supports prior research describing both species as exophilic (Kamgang et al. 2012). This calls for public health vector control measures that reduce human outdoor exposure to these mosquitoes in order to prevent arbovirus transmission. Based off of our findings, indoor measures to reduce mosquitoes feeding on humans, such as ITNs, are unlikely to have a significant effect on reducing human infections from *Aedes* transmitted arboviruses.

Africa is one of the most rapidly urbanizing regions on earth (Saghir & Santoro 2018). Our results, reflective of a broad range of habitats and climates across the African continent, suggest that *A. aegypti* populations will increase, while *A. albopictus* abundance will likely increase in some places and decrease in others. Furthermore, our results provide a more detailed investigation of the possible mechanisms underpinning urbanizations relationship to abundance. We found that while temperature and humidity could be important predictors of abundance, they were not significantly correlated with urbanization. Although increased larval habitat, host abundance, and the number of structures were significantly correlated with greater urbanization, they were less powerful predictors of adult mosquito abundance than the GIS based urbanization index. As such our results lead us to conclude that

urbanization likely influences adult mosquito through mechanisms that have yet to be fully determined.

### **Financial Support**

This study was funded by the United States Department of Defense's Defense Health Agency's Global Emerging Infections Surveillance (GEIS) program.







Figure 1.1. Cities surveyed for adult mosquitoes in Cameroon. Above approximately 6° latitude *A. albopictus* is not found.




**Figure 1.2. Larval habitat plotted against UI at 100m for five of the study cities. Line and ribbon shows fitted models with UI at 100m as a predictor and 95% CI, respectively.**



**Figure 1.3. *A. aegypti* abundance on a log scale plotted against the urbanization index in a 1km radius around the study site for ten sites in each of six cities. Collections resulting in zero specimens were plotted by adding 0.1. Lines and ribbons show fitted models with UI at 1km as a predictor and 95% CI, respectively.**




**Figure 1.4. *A. aegypti* abundance on a log scale plotted against temperature at the time of collection. Collections resulting in zero specimens were plotted by adding 0.1. Lines and ribbons show fitted models with temperature as predictor and 95% CI, respectively.**




**Figure 1.5. *A. albopictus* (95% CI) abundance on a log scale plotted against the urbanization index in a 2km radius around the study site for ten sites in each of three cities where *A. albopictus* was found. Lines and ribbons show fitted model with UI as predictor and 95% CI, respectively. The slope of urbanization in Kribi was not significantly different from zero. Collections resulting in zero specimens were plotted by adding 0.1.**






**Figure 1.6. *A. albopictus* abundance on a log scale plotted against relative humidity at the time of collection. Collections resulting in zero specimens were plotted by adding 0.1. Lines and ribbons show fitted models with humidity as predictor and 95% CI, respectively.**



**Figure 1.7. Total *A. aegypti* and *A. albopictus* abundance in response to prior month’s rainfall across all sites. Collections resulting in zero specimens were plotted by adding 0.1. Lines and ribbons show fitted models with prior month rain interacting with species as predictor and 95% CI, respectively.**



**Figure 1.8. Locations of studies of *A. albopictus* abundance in response to urbanization. Numbers correspond to study referenced in Table S2.**

RESEARCH ARTICLE

Open Access



# Spatial distribution of insecticide resistant populations of *Aedes aegypti* and *Ae. albopictus* and first detection of V410L mutation in *Ae. aegypti* from Cameroon

Matthew Montgomery<sup>1</sup>, James F. Harwood<sup>1</sup>, Aurelie P. Yougang<sup>2,3</sup>, Theodel A. Wilson-Bahun<sup>2,4</sup>, Armel N. Tedjou<sup>2,3</sup>, Christophe Rostand Keumeni<sup>2,3</sup>, Auston Marm Kilpatrick<sup>5</sup>, Charles S. Wondji<sup>2,6</sup> and Basile Kamgang<sup>2\*</sup>

## Abstract

**Background:** Dengue (DENV), chikungunya (CHIKV) and Zika virus (ZIKV), are mosquito-borne viruses of medical importance in most tropical and subtropical regions. Vector control, primarily through insecticides, remains the primary method to prevent their transmission. Here, we evaluated insecticide resistance profiles and identified important underlying resistance mechanisms in populations of *Aedes aegypti* and *Ae. albopictus* from six different regions in Cameroon to pesticides commonly used during military and civilian public health vector control operations.

**Methods:** *Aedes* mosquitoes were sampled as larvae or pupae between August 2020 and July 2021 in six locations across Cameroon and reared until the next generation, G1. *Ae. aegypti* and *Ae. albopictus* adults from G1 were tested following World Health Organization (WHO) recommendations and *Ae. aegypti* G0 adults screened with real time melting curve qPCR analyses to genotype the F1534C, V1016I and V410L *Aedes kdr* mutations. Piperonyl butoxide (PBO) assays and real time qPCR were carried out from some cytochrome p450 genes known to be involved in metabolic resistance. Statistical analyses were performed using Chi-square test and generalized linear models.

**Results:** Loss of susceptibility was observed to all insecticides tested. Mortality rates from tests with 0.25% permethrin varied from 24.27 to 85.89% in *Ae. aegypti* and from 17.35% to 68.08% in *Ae. albopictus*. Mortality rates for 0.03% deltamethrin were between 23.30% and 88.20% in *Ae. aegypti* and between 69.47 and 84.11% in *Ae. albopictus*. We found a moderate level of resistance against bendiocarb, with mortality rates ranging from 69.31% to 90.26% in *Ae. aegypti* and from 86.75 to 98.95% in *Ae. albopictus*. With PBO pre-exposure, we found partial or fully restored susceptibility to pyrethroids and bendiocarb. The genes *Cyp9M6F88/87* and *Cyp9J10* were overexpressed in *Ae. aegypti* populations from Douala sites resistant to permethrin and deltamethrin. *Cyp6P12* was highly expressed in alphacypermethrin and permethrin resistant *Ae. albopictus* samples. F1534C and V1016I mutations were detected in *A. aegypti* mosquitoes and for the first time V410L was reported in Cameroon.

\*Correspondence: [basile.kamgang@crid-cam.net](mailto:basile.kamgang@crid-cam.net); [kamgang\\_d@yahoo.fr](mailto:kamgang_d@yahoo.fr)

<sup>2</sup> Centre for Research in Infectious Diseases, P.O. Box 13591, Yaoundé, Cameroon

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Conclusions:** This study revealed that *Ae. aegypti* and *Ae. albopictus* are resistant to multiple insecticide classes with multiple resistance mechanisms implicated. These findings could guide insecticide use to control arbovirus vectors in Cameroon.

**Keywords:** Arbovirus, *Aedes aegypti*, *Aedes albopictus*, Insecticide resistance, Knock down resistance, Cytochrome P450 monooxygenases, Cameroon

## Background

*Aedes*-borne viral diseases such as dengue, chikungunya and Zika are increasingly reported in different regions across the world, including Africa [1]. For dengue, approximately half the world's population is at risk with an estimated 100–400 million new infections reported each year [1]. Vector control remains the cornerstone to prevent and fight against transmission. Use of insecticide based-control strategies is a common approach in response to arboviral disease outbreaks in order to quickly lower vector density and ongoing transmission [2], and is often the primary strategy to mitigate the threat of dengue, chikungunya, and Zika during military operations [3]. However, the increased use of insecticides may result in the selection of mosquitoes that carry genetic traits associated with insecticide resistance. Indeed, resistance to multiple insecticides used in public health has been documented in *Aedes aegypti* and *Ae. albopictus* in different locations across the world [4].

Insecticide resistance is usually achieved through two mechanisms: increased detoxification (metabolic resistance) and target insensitivity (target site resistance). Metabolic resistance through upregulation of detoxification genes is a common resistance mechanism in *Ae. albopictus* and *Ae. aegypti*, caused primarily by three main enzyme families, the monooxygenases (cytochrome P450s), glutathione S-transferases (GSTs) and carboxylesterases (COEs) [5, 6]. Target site resistance is caused by mutations in target genes such as acetylcholinesterase (*Ace-1*), the GABA receptor, and the voltage-gated sodium channel (VGSC) causing knockdown resistance (*kdr*) [5, 6].

Knockdown resistance is one of the main target site adaptations for both pyrethroids and dichlorodiphenyltrichloroethane (DDT). In *Ae. albopictus*, the *kdr* mutation is less prevalent with only four voltage-gated sodium channel (VGSC) mutations detected affecting two codons (1532 and 1534). In *Ae. aegypti*, many *kdr* mutations have been reported [4, 7, 8]. Among these mutations, the V1016I, F1534C and V410L mutations have been extensively investigated in pyrethroid-resistant *Ae. aegypti* populations from Asia, South America and, to a lesser extent, Africa. In Cameroon F1534C and V1016I mutations have been previously reported in *Ae.*

*aegypti* [9, 10]. V410L is a novel *kdr* mutation, located in domain I of segment 6 of the VGSC. It was described for the first time in a pyrethroid-resistant *Ae. aegypti* laboratory strain originating from Rio de Janeiro, Brazil [7] and several years later in Africa [11]. Considering that this mutation has shown a potential to reduce the sensitivity of sodium channels to type I and II pyrethroids and to increase resistance when associated with F1534C [12], it is imperative to determine the current distribution of this mutation in natural *Ae. aegypti* populations in Cameroon. This is due to the fact that its presence can greatly impair the usefulness of a wide variety of pyrethroid insecticides, which currently constitute the major class of insecticides used in *Aedes* control [13, 14]. In addition, several cytochrome P450 genes have been found over-expressed in *Aedes* spp. populations tested in the field and demonstrating pyrethroid resistance, with evidence indicating a role in pyrethroids metabolism by *CYP9J28*, *CYP9J10*, *CYP9J26*, *Cyp9M6F88/87* and *CYP6P12* [4, 15–19].

Our goal was to quantify the insecticide resistance profile and mechanisms involved in the resistance of populations of *Ae. aegypti* and *Ae. albopictus* from six cities that span most of the 1000 km north–south extent of Cameroon. In Cameroon, *Ae. aegypti* is present across the country, while the distribution of *Ae. albopictus* is limited to the south, under 6°N [20, 21]. We selected insecticides commonly used for public health and military force health protection applications to mitigate disease threats.

## Methods

### Mosquito sampling and rearing

*Aedes* mosquitoes were sampled as larvae or pupae between August 2020 and July 2021 in six locations across Cameroon: Douala (N 04° 02' 729", E 09° 42' 142"), Maroua N 10° 37' 284", E 14° 18' 381"), Garoua (N 09° 17' 7776", E 13° 23' 288"), Ngaoundéré N 07° 35' 414", E 13° 57' 365", Yaoundé (N 03° 51' 993", E 011° 27' 688"), and Kribi (N 02° 56' 862", E 09° 54' 003") (Fig. 1). Immature stages (field generation, G0) were collected from different potential larval habitats: domestic (e.g., jars, tanks), peri-domestic (e.g., used tires, discarded tanks), and natural (e.g., tree holes). In each location, larvae, or pupae from 20 positive larval habitats were collected, stored in plastic boxes, and transferred to insectary then pooled

according to the location and land use category (urban and suburban) and reared to adult stage for identification using taxonomic keys [22, 23]. Mosquitoes identified as *Ae. albopictus* or *Ae. aegypti* were blood-fed and allowed to reproduce to generation G1 for adult bioassays. Mosquito populations were maintained at insectary conditions (temperature  $27 \pm 2$  °C; relative humidity  $80\% \pm 10\%$ ), and females were blood fed using hemotek membranes to complete their gonotrophic cycle. Two susceptible lab strains were used for this study: Vector Control Research Unit (VCRU) strain for *Ae. albopictus* and Benin lab stain for *Ae. aegypti*.

#### Insecticide resistance bioassays

Bioassays were carried out according to the World Health Organization (WHO) protocol using 3–5 days old G1 generation mosquitoes. Four replicates of 20–25 females per tube were exposed to 0.03% and 0.05% deltamethrin, 0.05% alphacypermethrin, 0.1% bendiocarb, and 0.25% or 0.75% permethrin for 1 h. Mortality was recorded 24 h later; mosquitoes, alive or dead, after exposure were stored in RNA later or silica gel, respectively. Mortality rates were interpreted according to WHO recommendations: mortality rate of 98% or above indicates susceptibility, mortality rate between 90 and 97% suggests a possibility for resistance that should be confirmed with further bioassays, and mortality rate less than 90% indicates confirmed resistance [14].

#### Adult synergist assay with PBO

To evaluate the effectiveness of insecticide detoxification by the mosquitoes, the specific enzyme inhibitor 4% piperonyl butoxide (PBO) was evaluated to address the potential role of P450s in insecticide resistance. Female adults, 3–5 days old, were pre-exposed for one hour to PBO-impregnated papers and then immediately exposed to the selected insecticide. Mortality was scored 24 h later and compared to the results obtained with each insecticide without synergist according to the WHO standards [14].

#### Knockdown resistance (*kdr*) genotyping

Approximately 30 mosquitoes per city were genotyped for three different *kdr* mutations: V1016I, V410L and F1534C. Genomic DNA of 30 individual mosquitoes per population was extracted using the Livak protocol [24]. Three *kdr* mutations (F1534C, V1016I and V410L) were chosen for this study because these mutations have been described as involved in the pyrethroids resistance of *Ae. aegypti* mosquito [4, 25]. These mutations have been also reported in Africa [11, 15, 26]. Based on Moyes et al., review, F1534C and V410L are associated to insecticide resistance and V1016I is associated to insecticide

resistance when combined to other *kdr* mutations. Genotyping of the V1016I, V410L and F1534C mutations was performed by real-time melting curve quantitative PCR [27]. Each PCR reaction was performed in a 21.5 µl volume PCR tube containing 2 µl of DNA sample, 10 µl of SYBR® Green (SuperMix), and 1.25 µl of each primer. The thermocycle parameters were: 95 °C for 3 min, followed by 40 cycles of (95 °C for 20 s, 60 °C for 1 min and 72 °C for 30 s) and then a final step of 72 °C for 5 min, 95 °C for 1 min, 55 °C for 30 s and 95 °C for 30 s.

#### Gene expression


##### RNA extraction and cDNA synthesis

For this experiment three groups of mosquitoes were used: unexposed (control) to insecticide individuals from G1, exposed G1 individuals that survived resistance assays (resistant), and susceptible (laboratory susceptible strains). For each group three replicates of 10 mosquitoes per species were performed. RNA was extracted using the PicoPure RNA Isolation Kit (Arcturus® Picopure RNA Extraction Kit Life Technologies, California, USA), following the manufacturer's recommendations. Quality and quantity of RNA obtained were assessed using a "NanoDrop Lite" spectrophotometer (Thermo Scientific Inc., Wilmington, USA) and stored at  $-80$  °C.

Extracted RNA was used to synthesize complementary DNA (cDNA) using the Superscript III kit (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The resulting cDNA was purified using a QIAquick spin column (QIAquick PCR Purification Kit, Qiagen) and diluted 2-fold to accommodate the volumes of the reaction.

##### Quantitative-reverse transcriptase PCR

Candidate genes for expression analysis were chosen based on previous implications of involvement in metabolic resistance [4, 17]. Standard curve analyses were performed for each primer pair to check the specificity and efficiency of amplifications. Four cytochrome P450 candidate genes were chosen for analysis in *A. aegypti* (*Cyp9M6F88/87*, *Cyp9J28a*, *Cyp9J10*, and *Cyp9J32*) and only one in *A. albopictus* (*Cyp6P12*). The reactions were performed in a volume of 20 µl with 10 µl sybrGreen (Applied Biosystems, Texas, USA), 0.6 µl of each primer (10 µM), 7.8 µl of ddH<sub>2</sub>O, and 1 µl of cDNA, under the following conditions: 95 °C for 3 min, followed by 40 cycles of 95 °C for 10 s and 60 °C for 10 s. The relative expression level and fold change (FC) of each candidate gene compared to susceptible strains were calculated using the  $2^{-\Delta\Delta CT}$  method, integrating the efficiency of the PCR [17] after normalization with housekeeping genes: Aaeg60sL8, RPF7, RSP7, and tubulin. All primer



**Fig. 1** Map of study sites in six cities in Cameroon

sequences and their origins are shown for *A. albopictus* and *A. aegypti* in Tables 1 and 2, respectively. The Mx Pro software integrated into the Agilent brand TaqMan machine was used (MxPro-Mx3005P v4.10, Stratagene, California, USA).

#### Data analysis

Comparisons of mortality rates between species and land use category were conducted using generalized linear models (GLM) with a binomial distribution and logit link function using the *lme4* package in R version 3.6.1 (R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria).

The comparison of mortality rate after pre-exposure of mosquitoes to synergist and without pre-exposure to synergist was performed using a Chi-square test. The relative expression level and fold change (FC) of each target gene in field samples relative to the susceptible Benin (*Ae. aegypti*) or VCRU (*Ae. albopictus*) were calculated according to the  $2^{-\Delta\Delta^{CT}}$  method incorporating the PCR efficiency after normalization with the housekeeping genes. This analysis was performed using GraphPad Prism 8.0.2 (GraphPad Software, San Diego, California, USA). *P*-value < 0.05 was considered as statistically different.

## Results

### Insecticide resistance profile for *Aedes aegypti*

In total we tested 10 populations of *Ae. aegypti* with four insecticides (Fig. 2). Susceptibility for three pyrethroid varied among locations. To 0.25% permethrin, high resistance was observed with mortality rates ranging from 24.27% in urban Kribi to 85.89% in urban Ngaoundéré, probable resistance in urban Maroua (mortality rate of 95.43%); susceptibility in suburban Ngaoundéré, urban Garoua, suburban Garoua, and suburban Maroua (mortality rate from 97.82 to 98.80%). Remarkably, even when using 0.75% permethrin, which is triple the dose recommended for *Aedes* control, high levels of resistance were still observed. For example, the mortality rates to the 0.75% permethrin were 5.83% in urban Douala, and 40.48% in suburban Douala. For 0.03% deltamethrin, resistance was found in urban Yaoundé, urban Kribi, urban Ngaoundéré, urban and suburban Douala (mortality rates varying from 23.30% in urban Douala to 88.20% in urban Yaoundé); probable resistance in urban Maroua and susceptibility in suburban Ngaoundéré, urban Garoua, suburban Garoua, and suburban Maroua (mortality rate varying from 98.80% in suburban Maroua to 100% in suburban Ngaoundéré). Six of 10 populations tested were resistant to alphacypermethrin (urban Yaoundé, urban Kribi, urban Ngaoundéré, urban Maroua, and urban and suburban Douala); three others exhibited probable resistant (suburban Ngaoundéré, urban Garoua, and suburban Maroua) and the last one was fully susceptible (suburban Garoua). A moderate level of resistance was reported against bendiocarb with mortality rates varying from

69.31% in urban Yaoundé to 90.26% in suburban Maroua. In addition, probable resistance was detected in six populations with mortality rates ranging from 91.79% in Douala suburban to 95.50% in Ngaoundéré urban. Only the population from urban Maroua was susceptible to bendiocarb (mortality rate of 98.91%).

The best fitting model for predicting insecticide mortality in *Ae. aegypti* was a triple interaction of environment (urban or suburban), city, and insecticide (Additional file 1). On average across all cities and insecticides *A. aegypti* from urban populations had lower mortality to insecticides than those from suburban habitat ( $P < 0.01$ ) (Fig. 3). Across all cities and environments bendiocarb caused significantly higher levels of mortality compared to alphacypermethrin ( $P < 0.01$ ), deltamethrin ( $P < 0.01$ ), and 0.75% permethrin ( $P < 0.01$ ). However, these simple patterns had exceptions; in Yaoundé Bendiocarb had the lowest mortality rates and insecticide mortality was lower in suburban Garoua versus urban Garoua (Fig. 3).

Overall mortality to insecticides showed a strong latitudinal gradient, with higher mortality rates in northern cities (Fig. 3).

### Insecticide resistance profile for *Aedes albopictus*

Five *Ae. albopictus* populations were tested (urban Yaoundé, suburban Yaoundé, urban Kribi, suburban Douala, and suburban Kribi). The results are presented in Fig. 4. The analysis showed that resistance or probable resistance was observed to the three pyrethroids used. All populations tested with 0.25% permethrin showed high resistance with mortality rates between 17.35% (urban


**Table 1** Primer sequences for the evaluation of the level of expression of metabolic resistance genes by RT-qPCR and their origins in *Aedes albopictus*

Genes	Forward primer	Reverse primer	References
<i>Cyp6P12</i>	CGTGCGCTTTGGGATTGAG	ATCGTCGGTCCCAATCCTT	[17]
<i>RSP7</i>	AAGTTCGACACCTTCACGTC	CGCGCGCTCACTTATTAGAT	[17]
<i>qTubulin</i>	CCGCACTCGAGAAGGATTAC	GTGGTTCGGTTGACTTCGT	[17]

**Table 2** Primer sequences for the evaluation of the level of expression of metabolic resistance genes by RT-qPCR and their origins in *Aedes aegypti*

Genes	Forward primer	Reverse primer	References
<i>Cyp9J10</i>	ATCGGTGTTGGTAAAGTCTGT	CATGTCGTTGCGCATTATCCC	[46]
<i>Cyp9J28</i>	CCACTGACGTACGATGCGA	GCCGATCAGTGGACGGAGC	[17]
<i>Cyp9M6</i>	TCGGTGACAAATCCAACAAC	GTCCGGTACGACCAACGAAA	[18]
<i>Cyp9J32</i>	CGGTCCGCTTATGACGAAGAG	TTTGTTGCTCCGAAGAGTGG	[47]
<i>RPS3</i>	AGCGTGCCAAGTCGATGAA	GTGGCCGTGTCGACGTACT	[18]
<i>Ae60sL8</i>	CTGAAGGGAACCGTCAAGCAA	TCGGCGCAATGAACAAC	[17]





**Fig. 2** Mortality rates of adult *Aedes aegypti* from urban and suburban habitats in six cities in Cameroon 24 h after exposure to insecticides alone or with 1 h preexposure to synergist. Error bars represent standard error of the mean. PBO: Piperonyl butoxide

Yaoundé) and 81.15% (suburban Douala). When the dose of permethrin was increased to 0.75%, some recovery of susceptibility was restored. In urban Yaoundé, the 0.25% permethrin mortality rate was 17.35%, but when 0.75% permethrin was used the mortality rate increased to 91.59%. Two of five populations were found resistant to deltamethrin; urban Yaoundé with mortality rate of 69.47%, and suburban Douala with mortality rate of 84.11%, whereas the three others were shown to exhibit probable resistance (suburban Yaoundé, urban Kribi, and suburban Kribi). To alphacypermethrin, resistance was observed in suburban Douala with mortality rate of 71.21%, urban Yaoundé with mortality rate of 78.63%, and suburban Yaoundé with mortality rate of 89.40%; probable resistance was found in urban Kribi with mortality rate of 92.70%, and suburban Kribi with mortality rate of 95.17%. Moderate level of resistance was reported to bendiocarb with mortality rates varying from 86.75% in suburban Kribi to 98.95% in urban Yaoundé. Environment was not a statistically significant predictor for resistance in *Ae. albopictus*.


In cities where both *Ae. albopictus* and *Ae. aegypti* occur sympatrically (Douala, Kribi, and Yaoundé) *Ae. aegypti* had significantly higher levels of resistance than *Ae. albopictus*. The overall mean mortality to the insecticides standard doses tested were 54.7% and 80.1% for *Ae. aegypti* and for *Ae. albopictus* from sympatric cities ( $P < 0.02$ ,  $\chi^2 = 6.61$ ,  $df = 1$ ).

#### Tests with synergist PBO

A partial or full recovery of susceptibility to insecticides (permethrin, alphacypermethrin, deltamethrin and bendiocarb) was reported after PBO pre-exposure for both species (Figs. 2, 4).

For *Ae. albopictus* samples, a partial recovery of susceptibility was reported to permethrin from populations in urban Yaoundé (17.35% mortality without PBO and 82.54% after PBO exposure  $P < 0.001$ ), suburban Douala (81.15% mortality without PBO and 99.1% after PBO exposure  $P > 0.05$ ), and urban Kribi (68.08% mortality without PBO and 83.21% after PBO exposure  $P > 0.1$ ). Susceptibility was partially recovered to alphacypermethrin in *Ae. albopictus* populations from urban Yaoundé (78.63% mortality without PBO and 96.36% after PBO exposure  $P > 0.05$ ), suburban Douala (71.21% mortality without PBO and 95.94% after PBO exposure  $P > 0.01$ ), and to bendiocarb in populations from suburban Yaoundé (88.18% mortality without PBO and 98.9% after PBO exposure  $P > 0.25$ ). Full susceptibility recovery to deltamethrin was reported in urban Yaoundé (69.47% mortality without PBO and 100% after PBO exposure  $P < 0.005$ ) and to alphacypermethrin in suburban Yaoundé (89.40% mortality without PBO and 100% after PBO exposure  $P > 0.25$ ).

For *Ae. aegypti* samples, a partial recovery of susceptibility was reported to permethrin in populations from urban Yaoundé (78.32% mortality without PBO and



**Fig. 3** *Aedes aegypti* insecticide mortality ( $\pm 1$  SE) between study cities and habitats. Cities on the x-axis are arranged along a northward gradient

92.18% after PBO exposure  $P > 0.1$ ), suburban Douala (40.48% mortality without PBO and 62.13% after PBO exposure  $P < 0.01$ ) and urban Ngaoundéré (85.89% mortality without PBO and 93.25% after PBO exposure  $P > 0.5$ ). Also, susceptibility was partially recovered to alphacypermethrin in *Ae. aegypti* populations from urban Yaoundé (70.71% mortality without PBO and 95.45% after PBO exposure  $P > 0.01$ ), suburban Douala (52.27% mortality without PBO and 54.27% after PBO exposure  $P > 0.75$ ) and urban Ngaoundéré (88.52% mortality without PBO and 98.80% after PBO exposure  $P > 0.25$ ), to deltamethrin in populations from urban Yaoundé (88.20% mortality without PBO and 98.03% after PBO exposure  $P > 0.25$ ), suburban Douala (62.52% mortality without PBO and 68.83% after PBO exposure  $P > 0.25$ ) and urban Ngaoundéré (83.77% mortality without PBO and 98.86% after PBO exposure  $P > 0.1$ ), and to bendiocarb in populations from suburban Douala (91.61% mortality without PBO and 97.79% after PBO exposure  $P > 0.5$ ).

#### Gene expression

Using qPCR, expression of four cytP450 genes were quantified in *Ae. aegypti* (*Cyp9M6F88/87*, *Cyp9J28a*, *Cyp9J10*, and *Cyp9J32*). Among them, two were significantly overexpressed in field populations compared to a susceptible lab strain (Fig. 5). *Cyp9M6F88/87* was overexpressed in urban Douala sample resistant to permethrin [(fold change (FC)) =  $2.54 \pm 0.90$ ,  $P = 0.016$ ] and suburban Douala samples resistant to deltamethrin

(FC =  $5.49 \pm 1.64$ ,  $P = 0.003$ ); and *Cyp9J10* was overexpressed in suburban Douala (FC =  $3.16 \pm 0.40$ ,  $P = 0.013$ ).


Only one gene (*Cyp6P12*) was assessed in *Ae. albopictus* populations. This gene was significantly more expressed in urban Douala samples resistant to permethrin, compared to a lab susceptible strain (FC =  $5.54 \pm 0.73$ ,  $P = 0.001$ ) and in urban Yaoundé samples resistant to alphacypermethrin (FC =  $2.48 \pm 0.57$ ,  $P = 0.034$ ). However, in urban Douala samples resistant to deltamethrin, *Cyp6P12* expression was not significantly different compared to susceptible strain (Fig. 6).

#### Knockdown resistance (*kdr*) genotyping in *Ae. aegypti*

In this study, three *kdr* mutations were genotyped: V1016I, F1534C and V410L. The results are given in Tables 3, 4 and 5.

For V1016I *kdr* genotyping, 296 samples were examined in total. Among them, 227 (76.69%) were susceptible (1016 V/V), 59 (19.93%) were heterozygote resistant (1016 V/I), and 10 (3.38%) were homozygote resistant (1016I/I). The allele frequencies were 86.66% and 13.34% for alleles V and I respectively.

We examined 296 samples for F1534C genotyping. Among them, 170 (57.43%) were susceptible (1534F/F), 18 (6.08%) were heterozygote resistant (1534F/C), and 108 (36.49.05%) were homozygote resistant (1534C/C). The allele frequencies were 60.47% and 39.53% for alleles



**Fig. 4** Mortality rates of adult *Aedes albopictus* from urban and suburban habitats in three cities in Cameroon 24 h after exposure to insecticides alone or with 1 h preexposure to synergist. Error bars represent standard error of the mean. PBO Piperonyl butoxide



F and C, respectively. C allele frequency had increased from levels first found in 2017 in 5 of the studied cities (Fig. 7).

We genotyped 299 samples for V410L *ldr* and 212 of samples (70.91%) were susceptible (410 V/V), 59 (19.73%) were heterozygote resistant (410 V/I), and 28 (9.36%) were homozygote resistant (410I/I). The allele frequencies were 80.77% and 19.23% for alleles V and L, respectively.

### Discussion

The aim of this study was to evaluate insecticides resistance profiles of *Ae. albopictus* and *Ae. aegypti* collected in several locations in Cameroon and explore the potential mechanisms. Analyses revealed resistance to all four insecticides tested (deltamethrin, alphacypermethrin, permethrin, and bendiocarb) for both *Aedes* species. While the loss of susceptibility to these insecticides in *Aedes* was previously reported in some locations in

Cameroon [10, 28, 29] and outside Africa [30–33], this study expanded the spatial scale of insecticide resistance assessment in Cameroon while simultaneously comparing two species. The cause of resistance to insecticides in both *Ae. aegypti* and *Ae. albopictus* remains unclear as the use of insecticides against *Aedes* is scarce in African countries [10, 15, 30]. Nonetheless alleles, such as F1534C, that confer insecticide resistance continue to increase in frequency over time indicating selective pressure towards greater insecticide resistance. Given that the source of selective pressure for resistance is unknown, a return to susceptibility seems improbable [15] and could have operational consequences. The reduced susceptibility to the pyrethroids tested may pose a serious threat to future vector control programs, because pyrethroids are recommended for the control of adult *Aedes* mosquitoes, especially in case of disease outbreaks [34], and are commonly used in insecticide treated nets and uniforms.



*Aedes aegypti* populations from the northern part of Cameroon were more susceptible to pyrethroids than those from the southern part. This suggests that resistance to this insecticide family has not yet spread throughout the entire country and these insecticides are still effective in controlling *Aedes* in some locations of Cameroon. Previous studies on *Anopheles* in Cameroon showed the same latitudinal gradient to resistance distribution [35]. Climate is a potential contributing

mechanism; while the relationship between insecticide resistance and climate for *Aedes* mosquitoes is largely unexplored, solar radiation and humidity were the highest-ranked predictor variable for resistance levels in Cameroonian *Anopheles* [35]. As with our study, *Anopheles* mosquitoes from wet tropical areas generally had higher levels of resistance than those from dry arid climates.

**Table 3** 1016 genotype numbers and the allelic frequency of the Isoleucine (I) mutation of *Aedes aegypti*

Location	V1016 genotypes			VV + VI + II	Allelic frequencies	
	VV	VI	II		V	I
Urban Douala	21	6	0	27	0.89	0.1
Suburban Douala	22	8	0	30	0.87	0.13
Urban Yaoundé	27	2	1	30	0.93	0.07
Urban Maroua	13	14	2	29	0.69	0.31
Suburban Maroua	30	0	0	30	1	0
Urban Garoua	22	8	0	30	0.87	0.13
Suburban Garoua	27	2	1	30	0.93	0.07
Urban Ngaoundéré	21	8	1	30	0.83	0.17
Suburban Ngaoundéré	30	0	0	30	1	0
Urban Kribi	14	11	5	30	0.65	0.35
Total	227	59	10	296	0.87	0.13

V: Valine; I: Isoleucine; VV: absence of the V1016I mutation; VI: presence of the V1016I mutation with 2 alleles: one resistant, allele I and another susceptible V allele; II: presence of the V1016I mutation with the 2 resistant alleles

**Table 4** 1534 genotype numbers and the allelic frequency of the Cysteine (C) mutation of *Aedes aegypti*

Location	F1534 genotypes			FF + FC + CC	Allelic frequencies	
	FF	FC	CC		F	C
Urban Douala	0	3	26	29	0.05	0.95
Suburban Douala	1	5	24	30	0.12	0.88
Urban Yaoundé	27	2	1	30	0.93	0.07
Urban Maroua	8	1	19	28	0.30	0.70
Suburban Maroua	30	0	0	30	1	0
Urban Garoua	30	0	0	30	1	0
Suburban Garoua	26	2	1	29	0.93	0.07
Urban Ngaoundéré	18	5	7	30	0.68	0.32
Suburban Ngaoundéré	30	0	0	30	1	0
Urban Kribi	0	0	30	30	0	1
Total	170	18	108	296	0.60	0.40

F: phenylalanine; C: cysteine; FF: absence of the F1534C mutation; FC: presence of the F1534C mutation with 2 alleles: one resistant, allele C and another susceptible F allele; CC: presence of the F1534C mutation with the 2 resistant alleles

Pyrethroid resistance was more frequent in urban settings, which mirrors patterns in *Anopheles* mosquitoes in Cameroon [36]. This may be the result of a more intensive use of insecticide treated nets (ITN) and household insecticide (e.g., mosquito coils) usage in urban settings. However, both *Ae. aegypti* and *Ae. albopictus* feed primarily outdoors and during daylight in Cameroon and are thus unlikely to encounter ITNs [37, 38]. Land use could also affect resistance by determining the availability of larval habitat types since *Ae. aegypti* insecticide resistance levels can vary between larvae that developed in tires versus water containers [15].

With pre-exposure to PBO, partial or full recovery of susceptibility to all insecticides tested in *Ae. aegypti* and *Ae. albopictus* was observed. Similar results have been


seen in several other African countries [10, 15, 28, 29, 33], as well as outside Africa [16, 17]. These findings suggested the important role played by P450 genes in the resistance to pyrethroids (deltamethrin, alphacypermethrin, and permethrin) and carbamates (bendiocarb). These observations were confirmed by the overexpression of genes such as *Cyp9M6F88/87* and *Cyp9J10* in some *Ae. aegypti* populations and *Cyp6P12* in *Ae. albopictus* populations in Cameroon. The implication of these genes in metabolic resistance were previously demonstrated [16–19, 39].

Three possible *kdr* mutations were genotyped in this study: F1534C, V1016I, and V410L, which are involved in pyrethroids resistance in *Aedes* mosquito [11, 40–42]. These three *kdr* mutations have been previously

**Table 5** 410 genotype numbers and the allelic frequency of the Leucine (L) mutation of *Aedes aegypti*

Location	V410 genotypes			VV+VL+LL	Allelic frequencies	
	VV	VL	LL		V	L
Urban Douala	22	7	1	30	0.85	0.15
Suburban Douala	23	7	0	30	0.88	0.12
Urban Yaoundé	27	2	1	30	0.93	0.07
Urban Maroua	14	15	1	30	0.72	0.28
Suburban Maroua	30	0	0	30	1	0
Urban Garoua	22	8	0	30	0.87	0.13
Suburban Garoua	26	2	1	29	0.93	0.07
Urban Ngaoundéré	1	9	20	30	0.18	0.82
Suburban Ngaoundéré	30	0	0	30	1	0
Urban Kribi	17	9	4	30	0.72	0.28
Total	212	59	28	299	0.81	0.19

V: valine; L: leucine; VV: absence of the V410L mutation; VL: presence of the V410L mutation with 2 alleles: one resistant, allele L and another susceptible V allele; LL: presence of the V410L mutation with the 2 resistant alleles



**Fig. 7** Frequency of the 1534C allele *kdr* mutation in *Aedes aegypti* across study cities by year. Black circles represent mean pyrethroid survival ( $\pm 1$  SE) from studied cities for 2021

reported in Africa [15, 42–44], but only F1534C and V1016I were found previously in Cameroon [9, 10]. This study revealed for the first time the presence of a V410L mutation in *Ae. aegypti* in Cameroon. This mutation was first reported in 2017 in Brazil [7], and its first evidence in Africa was in 2020 in Angola [11]. V410L, alone confers low levels of resistance to insecticides, but when it co-occurs with V1016I or F1534C it yields higher levels

of resistance [7, 11]. The frequency of V410L was moderate compared to frequencies in Angola, supporting the hypothesis of a novel introduction of this mutation. The F1534C and V1016I *kdr* mutations are common in *Ae. aegypti* and have a worldwide distribution [4]. V1016I and V410L have a moderate frequency, while F1534C has a high frequency and has increased in *Ae. aegypti* populations across Cameroon since the mutation was first

observed there in 2017. This result supports the observation made previously in Cameroon suggesting an introduction and a gradual spread of the F1534C mutation in the country [10]. Broadly, the detection of *kdr* mutations are more frequent in Africa than was observed in the past [45]. Faced with their involvement in pyrethroid resistance, these three *kdr* mutations could have an impact on vector control measures.

The main limitation of the study is that we did not assess whether there is an association between the phenotypic resistance observed and the *kdr* mutations detected.

### Conclusions

Our study revealed the loss of susceptibility to four insecticides: deltamethrin, alphacypermethrin, permethrin, and bendiocarb, for both *Ae. aegypti* and *Ae. albopictus*. These insecticides are most likely to be used for mosquito control in outbreak responses and for reducing disease risk to deployed military personnel, and this study suggests these mitigation strategies will have limited efficacy. The level of loss of susceptibility varied according to the city, land use class, and species. Our results showed that several mechanisms are involved in resistance, and this can impact the strategies of *Aedes* control in Cameroon including the first detection of the V410L mutation in Cameroon. Further investigations including testing novel insecticides are needed to help to put in place effective strategies to control arbovirus vectors in Cameroon.

### Abbreviations

DNA: Deoxyribonucleic acid; *kdr*: Knockdown resistance; RNA: Ribonucleic acid; PBO: Piperonyl butoxide; PCR: Polymerase chain reaction; VGSC: Voltage-gated sodium channel; WHO: World Health Organization.

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40249-022-01013-8>.

**Additional file 1.** Logistic regression of *Aedes aegypti* mortality to insecticides across cities and habitat types. Suburban habitat, city of Douala, and Bendiocarb were reference levels.

### Acknowledgements

We would like to thank the residents living around the different collection sites and the garage owners for their cooperation during the field work. Some authors are military service members of the U.S. Government. This work was prepared as part of their official duties. Title 17, U.S.C. §105 provides that copyright protection under this title is not available for any work of the U.S. Government. Title 17, U.S.C. §101 defines a U.S. Government work as work prepared by a military service member or employee of the U.S. Government as part of that person's official duties. The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government.

### Author contributions

BK, CSW, JH, and MM conceived and designed the experiments. APY, BK, TAWB, CRK and ANT participated in mosquito collections. APY, CRK and TAWB performed the bioassays. APY, CRK, MM, and TAWB carried out the data analyses. APY, CRK, TAWB conducted the molecular analyses. APY, MM, JFH, AMK, BK and CSW wrote the paper. All authors read and approved the final manuscript.

### Funding

This work was supported by a Grant provided by the Armed Forces Health Surveillance Division, Global Emerging Infections Surveillance Branch (GEIS), ProMIS ID P0076\_20\_EC funded in FY 2020 and ProMIS ID P0042\_21\_N3 funded in FY 2021.

### Availability of data and materials

All the relevant data generated during this study are included in the manuscript.

### Declarations

#### Ethics approval and consent to participate

An oral consent form was obtained from the head or representative of each household or garage owner prior to collect the mosquito larvae.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup>U.S. Naval Medical Research Unit No. 3, Naval Air Station Sigonella, Italy. <sup>2</sup>Centre for Research in Infectious Diseases, P.O. Box 13591, Yaoundé, Cameroon. <sup>3</sup>Parasitology and Ecology Laboratory, Department of Animal Biology and Physiology, Faculty of Science, University of Yaoundé 1, P.O. Box 812, Yaoundé, Cameroon. <sup>4</sup>Laboratory of Vertebrate and Invertebrate Bioecology, Faculty of Science and Technology, Marien-Ngouabi University, Brazzaville, Congo. <sup>5</sup>Ecology and Evolutionary Biology Department, University of California, Santa Cruz, USA. <sup>6</sup>Vector Biology Department, Liverpool School of Tropical Medicine, London, UK.

Received: 3 June 2022 Accepted: 4 August 2022

Published online: 17 August 2022

### References

- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and burden of dengue. *Nature*. 2013;496:504–7.
- Wilson AL, Courtenay O, Kelly-Hope LA, Scott TW, Takken W, Torr SJ, et al. The importance of vector control for the control and elimination of vector-borne diseases. *PLoS Negl Trop Dis*. 2020;14:e0007831.
- Harwood JF, Farooq M, Richardson AG, Doud CW, Putnam JL, Szumlas DE, et al. Exploring new thermal fog and ultra-low volume technologies to improve indoor control of the dengue vector, *Aedes aegypti* (Diptera: Culicidae). *J Med Entomol*. 2014;51:845–54.
- Moyes CL, Vontas J, Martins AJ, Ng LC, Kooou SY, Dusfour I, et al. Contemporary status of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans. *PLoS Negl Trop Dis*. 2017;11:e0005625.
- Hemingway J, Hawkes NJ, McCarroll L, Ranson H. The molecular basis of insecticide resistance in mosquitoes. *Ins Biochem Mol Biol*. 2004;34:653–65.
- Perry T, Batterham P, Daborn PJ. The biology of insecticidal activity and resistance. *Ins Biochem Mol Biol*. 2011;41:411–22.
- Haddi K, Tomé HV, Du Y, Valbon WR, Nomura Y, Martins GF, et al. Detection of a new pyrethroid resistance mutation (V410L) in the sodium channel of *Aedes aegypti*: a potential challenge for mosquito control. *Sci Rep*. 2017;7:46549.
- Kosinski K, Lee Y, Romero-Weaver A, Chen T-Y, Collier T, Wang X, et al. Two novel single nucleotide polymorphisms in the voltage-gated sodium

- channel gene identified in *Aedes aegypti* mosquitoes from Florida. *J Fl Mosq Control Assoc.* 2022;69:8.
- Djiappi-Tchamen B, Nana-Ndjangwo MS, Mavridis K, Talipouo A, Nchoutpouen E, Makoudjou I, et al. Analyses of Insecticide Resistance Genes in *Aedes aegypti* and *Aedes albopictus* Mosquito Populations from Cameroon. *Genes.* 2021;12:828.
  - Youngang AP, Kamgang B, Bahun TAW, Tedjou AN, Nguiffo-Nguete D, Njikou F, et al. First detection of F1534C knockdown resistance mutation in *Aedes aegypti* (Diptera: Culicidae) from Cameroon. *Inf Dis Poverty.* 2020;9:1–12.
  - Ayres CF, Seixas G, Borrego S, Marques C, Monteiro I, Marques CS, et al. The V410L knockdown resistance mutation occurs in island and continental populations of *Aedes aegypti* in West and Central Africa. *PLoS Negl Trop Dis.* 2020;14:e0008216.
  - Saavedra-Rodríguez K, Maloof FV, Campbell CL, Garcia-Rejon J, Lenhart A, Penilla P, et al. Parallel evolution of vgsc mutations at domains I56, II56 and III56 in pyrethroid resistant *Aedes aegypti* from Mexico. *Sci Rep.* 2018;8:6747.
  - Macoris Mde L, Andrighetti MT, Wanderley DM, Ribolla PE. Impact of insecticide resistance on the field control of *Aedes aegypti* in the State of Sao Paulo. *Rev Soc Bras Med Trop.* 2014;47:573–8.
  - WHO: Entomological surveillance for *Aedes* spp. in the context of Zika virus: interim guidance for entomologists. World Health Organization. 2016. <https://apps.who.int/iris/handle/10665/204624>. Accessed 21 Sept 2021.
  - Badolo A, Sombié A, Pignatelli PM, Sanon A, Yaméogo F, Wangrawa DW, et al. Insecticide resistance levels and mechanisms in *Aedes aegypti* populations in and around Ouagadougou. *Burkina Faso PLoS Negl Trop Dis.* 2019;13:e0007439.
  - Ishak IH, Kamgang B, Ibrahim SS, Riveron JM, Irving H, Wondji CS. Pyrethroid resistance in Malaysian populations of dengue vector *Aedes aegypti* is mediated by CYP9 family of cytochrome P450 genes. *PLoS Negl Trop Dis.* 2017;11:e0005302.
  - Ishak IH, Riveron JM, Ibrahim SS, Stott R, Longbottom J, Irving H, et al. The Cytochrome P450 gene CYP6P12 confers pyrethroid resistance in *kdr*-free Malaysian populations of the dengue vector *Aedes albopictus*. *Sci Rep.* 2016;6:24707.
  - Kasai S, Komagata O, Itokawa K, Shono T, Ng LC, Kobayashi M, et al. Mechanisms of pyrethroid resistance in the dengue mosquito vector, *Aedes aegypti*: target site insensitivity, penetration, and metabolism. *PLoS Negl Trop Dis.* 2014;8:e2948.
  - Smith LB, Tyagi R, Kasai S, Scott JG. CYP-mediated permethrin resistance in *Aedes aegypti* and evidence for trans-regulation. *PLoS Negl Trop Dis.* 2018;12:e0006933.
  - Kamgang B, Happi JY, Boisier P, Njikou F, Herve JP, Simard F, Paupy C. Geographic and ecological distribution of the dengue and chikungunya virus vectors *Aedes aegypti* and *Aedes albopictus* in three major Cameroonian towns. *Med Vet Entomol.* 2010;24:132–41.
  - Tedjou AN, Kamgang B, Youngang AP, Njikou F, Wondji CS. Update on the geographical distribution and prevalence of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae), two major arbovirus vectors in Cameroon. *PLoS Negl Trop Dis.* 2019;13:e0007137.
  - Edwards FW. Mosquitoes of the Ethiopian Region. Hl-Culicine Adults and Pupae. London, British Museum (Natural History); 1941.
  - Jupp PG. Mosquitoes of southern Africa: Culicidae and Toxorhynchitinae. Hartheespoort: Ekogilde Publishers; 1996.
  - Livak KJ. Organization and mapping of a sequence on the *Drosophila melanogaster* X and Y chromosomes that is transcribed during spermatogenesis. *Genetics.* 1984;107:611–34.
  - Saavedra-Rodríguez K, Urdaneta-Marquez L, Rajatileka S, Moulton M, Flores A, Fernandez-Salas I, et al. A mutation in the voltage-gated sodium channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*. *Ins Mol Biol.* 2007;16:785–98.
  - Smith LB, Kasai S, Scott JG. Pyrethroid resistance in *Aedes aegypti* and *Aedes albopictus*: Important mosquito vectors of human diseases. *Pestic Biochem Physiol.* 2016;133:1–12.
  - Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C<sub>T</sub> method. *Nat Protoc.* 2008;3:1101.
  - Kamgang B, Youngang AP, Tchoupo M, Riveron JM, Wondji C. Temporal distribution and insecticide resistance profile of two major arbovirus vectors *Aedes aegypti* and *Aedes albopictus* in Yaoundé, the capital city of Cameroon. *Parasit Vectors.* 2017;10:469.
  - Youngang AP, Kamgang B, Tedjou AN, Wilson-Bahun TA, Njikou F, Wondji CS. Nationwide profiling of insecticide resistance in *Aedes albopictus* (Diptera: Culicidae) in Cameroon. *PLoS ONE.* 2020;15:e0234572.
  - Das M, Dutta P. Status of insecticide resistance and detoxifying enzyme activity of *Aedes albopictus* population in Sonitpur district of Assam. *India Int J Mosq Res.* 2014;1:35–41.
  - Ishak IH, Jaal Z, Ranson H, Wondji CS. Contrasting patterns of insecticide resistance and knockdown resistance (*kdr*) in the dengue vectors *Aedes aegypti* and *Aedes albopictus* from Malaysia. *Parasit Vectors.* 2015;8:181.
  - Marcombe S, Fustec B, Cattel J, Chonephetsarath S, Thammavong P, Phommavanh N, et al. Distribution of insecticide resistance and mechanisms involved in the arbovirus vector *Aedes aegypti* in Laos and implication for vector control. *PLoS Negl Trop Dis.* 2019;13:12.
  - Kamgang B, Wilson-Bahun TA, Youngang AP, Lenga A, Wondji CS. Contrasting resistance patterns to type I and II pyrethroids in two major arbovirus vectors *Aedes aegypti* and *Aedes albopictus* in the Republic of the Congo. *Central Africa Inf Dis Poverty.* 2020;9:1–10.
  - WHO: Pesticides and their application: for the control of vectors and pests of public health importance. 6th ed World Health Organization. 2006. <https://apps.who.int/iris/handle/10665/69223>. Accessed 18 Sept 2020.
  - Hancock PA, Lynd A, Wiebe A, Devine M, Essandoh J, Watsenga F, et al. Modelling spatiotemporal trends in the frequency of genetic mutations conferring insecticide target-site resistance in African mosquito malaria vector species. *BMC Biol.* 2022;20:1–17.
  - Mandeng SE, Awono-Ambene HP, Bigogo JD, Ekoko WE, Binyang J, Piamou M, et al. Spatial and temporal development of deltamethrin resistance in malaria vectors of the *Anopheles gambiae* complex from North Cameroon. *PLoS ONE.* 2019;14:e0212024.
  - Egid BR, Coulibaly M, Dadzie SK, Kamgang B, McCall PJ, Sedda L, et al. Review of the ecology and behaviour of *Aedes aegypti* and *Aedes albopictus* in Western Africa and implications for vector control. *Curr Res Parasitol Vector Borne Dis.* 2022;2:100074.
  - Kamgang B, Nchoutpouen E, Simard F, Paupy C. Notes on the blood-feeding behavior of *Aedes albopictus* (Diptera: Culicidae) in Cameroon. *Parasit Vectors.* 2012;5:57.
  - Rault LC, O'Neal ST, Johnson EJ, Anderson TD. Association of age, sex, and pyrethroid resistance status on survival and cytochrome P450 gene expression in *Aedes aegypti* (L.). *Pestic Biochem Physiol.* 2019;156:96–104.
  - Kasai S, Ng LC, Lam-Phua SG, Tang CS, Itokawa K, Komagata O, et al. First detection of a putative knockdown resistance gene in major mosquito vector. *Aedes albopictus* Japanese J Infect Dis. 2011;64:217–21.
  - Kawada H, Higa Y, Futami K, Muranami Y, Kawashima E, Osei JH, et al. Discovery of point mutations in the voltage-gated sodium channel from African *Aedes aegypti* populations: potential phylogenetic reasons for gene introgression. *PLoS Negl Trop Dis.* 2016;10:e0004780.
  - Sombié A, Saïki E, Yaméogo F, Sakurai T, Shirozu T, Fukumoto S, et al. High frequencies of F1534C and V1016I *kdr* mutations and association with pyrethroid resistance in *Aedes aegypti* from Somgandé (Ouagadougou). *Burkina Faso Trop Med Health.* 2019;47:2.
  - Konan LY, Oumbouke WA, Silué UG, Coulibaly IZ, Ziogba J-CT, Guessan RK, et al. Insecticide Resistance Patterns and Mechanisms in *Aedes aegypti* (Diptera: Culicidae) Populations Across Abidjan, Côte d'Ivoire Reveal Emergent Pyrethroid Resistance. *J Med Entomol.* 2021;58:1808–1816.
  - Kudom AA. Entomological surveillance to assess potential outbreak of *Aedes*-borne arboviruses and insecticide resistance status of *Aedes aegypti* from Cape Coast. *Ghana Acta Trop.* 2020;202:105257.
  - Weetman D, Kamgang B, Badolo A, Moyes CL, Shearer FM, Coulibaly M, et al. *Aedes* mosquitoes and *Aedes*-borne arboviruses in Africa: current and future threats. *Int J Environ Res Public Health.* 2018;15:220.
  - Bariami V, Jones CM, Poupardin R, Vontas J, Ranson H. Gene amplification, ABC transporters and cytochrome P450s: unraveling the molecular basis of pyrethroid resistance in the dengue vector. *Aedes aegypti* *PLoS Negl Trop Dis.* 2012;6:e1692.
  - Seixas G, Grigoraki L, Weetman D, Vicente JL, Silva AC, Pinto J, et al. Insecticide resistance is mediated by multiple mechanisms in recently introduced *Aedes aegypti* from Madeira Island (Portugal). *PLoS Negl Trop Dis.* 2017;11:e0005799.



### **Chapter 3: Comparing relative transmission potential to population seroprevalence for seven arboviruses: yellow fever, chikungunya, zika, and four dengue serotypes**

#### Abstract

Dengue (DENV), chikungunya (CHIKV), yellow fever (YFV), and Zika virus (ZIKV) are mosquito-borne viruses which account for millions of human infections annually, primarily in tropical regions. Despite the presence of multiple arboviruses in many locations, little comparative work has been done to examine relative transmission intensity through seroprevalence, and to what extent this is determined by differences in host and vector in competence between viruses. We calculated relative seroprevalence estimates for DENV (serotypes 1-4), CHIKV, YFV, and ZIKV from 204 seroprevalence studies with a total of 95,000 individuals on five continents. We then calculated the relative epidemic potential,  $R_0$ , for the same seven arboviruses for both *A. aegypti* and *A. albopictus* using prior clinical and laboratory studies on vector and host competence. We hypothesized relative seroprevalence of a virus would be positively correlated to its  $R_0$  estimates. Globally, CHIKV and YFV had the highest relative seroprevalence, followed by ZIKV and DENV-2, DENV-3, DENV-1, and DENV-4, whereas in Asia YFV seroprevalence was zero and in the Americas ZIKV had the highest seroprevalence and CHIKV was the third lowest. Relative  $R_0$  values were highest for CHIKV followed by the four DENV and ZIKV, with YFV being much lower. Relative  $R_0$  values were higher for *A. aegypti* than *A. albopictus*. Relative  $R_0$

for *A. aegypti* was significantly correlated with higher seroprevalence for the seven arboviruses in Asia, but not in other continents or globally. These results demonstrate substantial variation in seroprevalence and relative  $R_0$  among these arboviruses, but also regional variation in relative seroprevalence differences. This suggests that predicting relative exposure of different pathogens may require more detailed local spatio-temporal estimates of host and vector competence.

## Introduction

Viruses transmitted by arthropods (arboviruses) are of increasing global concern and account for millions of annual cases of human illness and tens of thousands of fatalities (Kilpatrick & Randolph 2012, Puntasecca et al. 2021, Yang et al. 2021). Increased global trade and travel have led to regular introductions of novel pathogens including many arboviruses such as West Nile, Zika, and Chikungunya (Dufft et al. 2009, Kilpatrick & Randolph 2012; Possas et al. 2017, WHO 2020a). The total fraction of the population infected over the course of an epidemic is determined, in part, by the reproductive number of the pathogen,  $R_0$ , which also influences long term steady state seroprevalence (Dye 1992). Despite the presence of multiple arboviruses in human populations, little comparative work has been done to determine relative transmission intensity through seroprevalence, and to what extent this is determined by differences in host and vector in competence between viruses.

Arboviruses spread by *Aedes* mosquitoes, including CHIKV, DENV, yellow fever (YFV), and Zika virus (ZIKV) account for the majority of human arbovirus

infections (World Health Organization 2020). Most transmission of these arboviruses occurs in urban areas by two anthropophilic vectors, *A. aegypti* and *A. albopictus* (Weaver & Reisen 2010). The urban cycles of these viruses share common vectors and a single host, humans. As a result, in most urban areas, relative transmission intensity will depend only on differences in host and vector competence among the viruses which are key components of  $R_0$  (Dye 1992). Host competence depends on both the levels of viremia (the concentration of virus in blood) reached within human hosts and the duration of infectious viremia. Vector competence, the probability of a mosquito transmitting virus after feeding on an infected host, depends on host viremia, temperature, and time since feeding, and often differs between vector species and populations (Azar & Weaver 2019, Souza-Neto et al. 2019).

Our goal was to estimate the relative seroprevalence of these seven arboviruses across human populations and compare these to estimates of relative  $R_0$  based on host and vector competence. Antibodies against these viruses persist for up to 60 years following exposure (Imrie et al. 2007; Wang & Sekaran 2010), and thus provide an estimate of previous exposure of a population (Imrie et al. 2007; Wang & Sekaran 2010). We collected a large number of seroprevalence estimates to try to average across sources of variation, including movement of people among locations and episodic transmission. We then estimated relative  $R_0$  across all seven arboviruses using measures of human viremia and mosquito vector competence from previous studies. Numerous prior studies have estimated  $R_0$  for individual viruses or pairs of arboviruses in one or two vectors (Chitnis et al 2006, Gardner & Ryman 2010,

Lambrechts et al. 2010, Weaver et al. 2015, Manore et al. 2014, Vasilakis & Weaver 2017, Mordecai et al. 2017, Liu et al. 2020), but none of these studies has estimated relative  $R_0$  for all seven of the arboviruses transmitted primarily by *A. aegypti* and *A. albopictus*. We compared relative seroprevalence to relative  $R_0$  for the seven arboviruses for both *A. aegypti* and *A. albopictus* and hypothesized that these would be positively correlated.

## Methods

### **Relative transmission potential, $R_0$**

We compiled data on host and vector competence for CHIKV, YFV, ZIKV, and the four serotypes of DENV for both *A. aegypti* and *A. albopictus* (Table S1). Human viremia data were reported relative to time since symptom onset or resolution and measured in either PFU/mL or viral copies/mL. DENV viremia was converted from viral copies/mL to PFU/mL using the equation  $\text{Log}_{10} \text{PFU/mL} = [0.974 * \text{Log}_{10} \text{viral copies/mL}] - 2.807$  (Fernandes-Monteiro et al. 2015). We analyzed six days of viremia for each virus after which infectiousness of most viruses decreased to zero or near zero (see below). We fit a linear model to compare viremia between viruses for each of the six days in R and performed a post-hoc comparison of viremia between viruses using the emmeans package (Lenth 2022).

For vector competence, we collected data on the fraction of *A. aegypti* and *A. albopictus* mosquitoes that transmitted virus after feeding on infected blood with known viremia. The majority of vector competence studies provided infectious dose

blood-meal viremia in PFU/mL. For studies which measured viremia in Tissue Culture Infectious Dose (TCID50) we converted those data to PFU/mL using the equation  $\text{Log}_{10} \text{PFU mL}^{-1} = -0.567306 + 0.987227 \text{Log}_{10} \text{CID50 mL}^{-1}$  (Kilpatrick et al. 2007) and for studies measuring viremia in Mouse Lethal Dose (MLD50) we used  $1.395 \text{Log}_{10} \text{PFU mL}^{-1} / \text{MLD50}$  (Freire et al. 2007).

We fit models to estimate the fraction of mosquitoes transmitting virus as a function of time since feeding and host viremia. We fit empirical data on the fraction transmitting on a given day to the cumulative distribution of a gamma distribution representing the extrinsic incubation period, in days, using a Bayesian framework via the R package *rethinking* (Ferguson et al. 2016, McElreath 2016). We fit all data for three viruses (CHIKV, YFV, and ZIKV) and both mosquitoes simultaneously with an additive model of vector species and virus to leverage information for virus-vector pairs with limited information. We used vague priors for all parameters (gaussian distributions with mean = 0, SD = 20); alternative priors produced nearly identical posterior estimates. We performed similar analyses for the four serotypes of DENV. We fit separate models for DENV because data for all four viruses for both viremia and vector competence came from a single location and set of studies (Nguyen et al. 2013, Whitehorn et al. 2015) and the units that viremia were measured in differed between the DENV studies (copies/ml) and the other three viruses (PFU/ml).

We calculated epidemic potential by estimating the infectiousness of human viremias for each virus on each day using the vector competence models (estimated 14 days after feeding) and then summing across six days of viremia for each virus.

We estimated uncertainty in relative  $R_0$  estimates using both variation in daily viremia among people and the uncertainty in the vector-competence-viremia relationships.

**Seroprevalence estimates.** We collected seroprevalence data from the literature by searching Google Scholar and Web of Science. Our search terms consisted of a pathogen (CHIKV, DENV 1-4, YFV, and ZIKV) or the term “arbovirus”, seroprevalence or serosurvey or antibodies, and a location including countries and continents. We included articles that simultaneously measured antibodies for at least two of the seven arboviruses in the same population so that we could estimate the relative differences in seroprevalence among two or more viruses. We excluded studies done in response to outbreaks of specific viruses and studies of DENV seroprevalence which did not specify serotype. We categorized seroprevalence data by continent for Asia, Africa, and the Americas.

**Calculating comparative seroprevalence.** We compared seroprevalence among viruses using a generalized linear model with a binomial distribution with virus as a fixed effect and study ID as a random effect using the *lme4* package in R version 4.2.1 (Bates et al. 2015). We fit models both to all serosurvey data and to data for each continent. For Asia, where all YFV studies had a seroprevalence of zero, we fit a similar model using both study ID and virus as fixed effects using the *brglm2* package in R to better estimate the YFV coefficient (Kosmidis 2017). We then used the fitted

coefficients for each arbovirus as the response variable and estimates of relative  $R_0$  for either *A. aegypti* or *A. albopictus* as the predictor variable using a linear model. We performed these analysis for both global and continental seroprevalence data with a single estimate for relative  $R_0$  for each virus because we had insufficient host and vector competence data to estimate continent-specific relative  $R_0$  values.

## Results

**Viremia.** Viremia among human patients differed significantly between arboviruses, with CHIKV having significantly higher viremia than any other virus. None of the other viruses differed significantly from one another in viremia (Figure 1, Table S2).

**Vector Competence.** Vector competence varied significantly among viruses and between mosquito species (Figure 2; Figures S1 and S2, Table S3). Vector competence was higher for *A. aegypti* for all seven arboviruses, with the exception of DENV-1 and DENV-4 at titers above 8 PFU/mL (Figure 2).

**Relative  $R_0$ .** Daily human infectiousness to mosquitoes and thus relative  $R_0$  differed between arboviruses (Figure 3).  $R_0$  was greatest for CHIKV, followed by DENV-2, DENV-1, DENV-3, DENV-4, ZIKV, and YFV with *A. aegypti* as the vector. For *A. albopictus* the ranked order of  $R_0$  from greatest to least was CHIKV, DENV-1, DENV-3, DENV-4, ZIKV, DENV-2, and YFV.

**Seroprevalence for seven arboviruses.** We reviewed 611 articles and identified 80 which contained 202 unique serosurveys that matched our criteria (Supplemental References). Serosurvey data represented 95,000 individual blood samples from 46 countries (Figure S3). In the full dataset, relative seroprevalence differed significantly among arboviruses, with CHIKV and YFV being highest, followed by ZIKV and DENV-2, DENV-3, DENV-1, and DENV-4 (Figure 4, Table 1, Figure 5A). However, there were substantial differences among the three continents (Tables S4-S6). Patterns in Africa were somewhat similar to the global dataset, in part because Africa made up a substantial fraction of all studies (Figure S3), but DENV-4 and DENV-1 switched places in relative seroprevalence, and ZIKV was relatively lower (Figure 5 A,B). In the Americas, ZIKV and DENV-2 seroprevalence was the highest and DENV-4 was the lowest. Finally, in Asia, YFV was absent and all YFV seroprevalence estimates were 0%, whereas CHIKV and DENV-3 had the highest seroprevalence. The differences in seroprevalence among arboviruses were much larger in the Americas and Asia than in global and especially African datasets (Figure 6, Tables S4-S6). In areas in the Americas with average transmission, ZIKV and CHIKV had very high estimated seroprevalence (>80%), whereas the four dengue viruses were much lower (<50%) (Figure 6C). In Asia, CHIKV, DENV-3 and DENV-2 had the highest estimated seroprevalence whereas YFV seroprevalence was zero in all studies (Figure 6D).



**Comparison of relative  $R_0$  and relative seroprevalence.** Relative seroprevalence was significantly positively correlated with relative  $R_0$  for *A. aegypti* in Asia, but no other correlations were significant with either species as the vector (Figure 5; Figure S4). CHIKV had very high relative  $R_0$  and seroprevalence in all regions, and DENV-4 had a low seroprevalence in all regions and a moderately low relative  $R_0$ . However, ZIKV had the 2<sup>nd</sup> lowest relative  $R_0$  and the highest seroprevalence in the Americas and was moderately high in Asia and the global dataset. The four serotypes of DENV also showed spatial variation in relative seroprevalence and differences among these four viruses was variable and not correlated with relative  $R_0$  in most regions (Figure 5).

### Discussion

We found large differences among arboviruses in seroprevalence, but differences varied substantially among continents, and differences in seroprevalence were only correlated with relative  $R_0$  for one vector, *A. aegypti*, in one of three regions and not in the global analysis. We also found large differences in relative  $R_0$  among viruses due to both significant differences in viremia as well as vector competence. The lack of correlation between relative  $R_0$  and seroprevalence for the global dataset and for Africa was surprising, because we had a large dataset of seroprevalence studies and a relatively large dataset to estimate vector competence. However, the analysis only included seven arboviruses and two viruses with low relative  $R_0$  values played a key role in the lack of a correlation with relative

seroprevalence: YFV and ZIKV. YFV had the lowest estimated relative  $R_0$ , by far, but had the second highest relative seroprevalence in Africa and in the global dataset. ZIKV had the second lowest relative  $R_0$ , but had the highest seroprevalence in the Americas, and intermediate seroprevalence in Asia and globally. Clearly the relative  $R_0$  values didn't match the observed relative seroprevalence for these viruses.

The low estimate of relative  $R_0$  for YFV was especially puzzling. We had a robust dataset for calculating YFV vector competence, but these data suggested that it was significantly lower than any other virus for both vectors. Human viremia data were limited for YFV, but no more so than for CHIKV or ZIKV, all of which had no more than 11 viremia values for a single day post symptom onset. In any case, YFV had the lowest viremia of the seven viruses on 3 of the 6 days and overall were in the lowest group with DENV-1, DENV-2 which had much higher relative vector competence. The low estimated relative  $R_0$  of YFV is surprising in light of its very high seroprevalence in Africa. Human YFV infections were detected in 46 countries between 1970-2016 (Shearer 2018), and in 10 African countries and 2 South American countries in 2022 (WHO 2022, PAHO 2022). Given the recent and ongoing outbreaks of YFV in Africa (WHO 2022) and potential for expansion, more research is needed on human viremia data for YFV and to understand the variation in YFV vector competence among mosquito populations (Figure 2).

The mismatch between relative seroprevalence and relative  $R_0$  may arise from spatial or temporal variation among viruses, mosquitoes, or human populations and their contributions to relative epidemic potential. For example, there are substantial

differences between the Asian and Reunion strains of CHIKV in vector competence (Tsetsarkin et al. 2007). Similarly, there are clear spatial and temporal differences in vector competence among mosquito populations (Gubler et al. 1979, Bosio et al. 1998, Bosio et al. 2000, Goncalves et al. 2014). Innate differences in viremia among human populations may also contribute to different epidemic potential in different populations, but they have not been studied. However, evidence suggest that host competence is mediated by prior exposure to related viruses. For example, prior exposure to DENV may lead to partial protection against YFV, and been posited as one of the reasons for YFVs absence from Asia (Kuno 2020), and patients with secondary infections with DENV clear the virus more quickly than primary infections (Ben-Shachar et al 2016). Lastly, the lack of correlation between relative  $R_0$  and seroprevalence could be due to factors influencing seroprevalence, including episodic transmission, human migration, or differential waning of antibodies. The variation in relative seroprevalence rankings among studies (Figure 4) indicate that there is substantial uncertainty in which virus will lead to the highest exposure a given population at a point in time.

Despite these uncertainties, our results provide predictions about the relative likelihood of pathogen establishment among arboviruses. If the pathogen with both the highest seroprevalence and highest relative  $R_0$ , CHIKV, is unable to become established in a given location (e.g. the United States), then our results suggest it is less likely that the other arboviruses in this study would be able to become established in the same population. There have been no locally acquired cases of CHIKV in the

US since 2015, despite numerous travel-associated cases (CDC 2022), suggesting that ZIKV, YFV, and the four serotypes of DENV are also unlikely to become established unless conditions (e.g. climate, larval habitat, poverty) change.

There appear to be substantial differences among the seven arboviruses we examined based both on relative seroprevalence and estimates of relative  $R_0$ . However, the lack of a correlation between relative seroprevalence and relative  $R_0$  for most regions suggests that finer scale data on viruses, vectors and hosts may be needed to predict the relative transmission potential of different viruses in a given population. Measurements of vector competence and viremia for all seven arboviruses in replicated population of mosquitoes and humans would help illuminate the differences in relative  $R_0$ , and may uncover cryptic coevolution between viruses, vectors and hosts.

**Acknowledgements.** We thank Angela Brierly for assistance compiling seroprevalence estimates. Funding was provided by NSF grants DEB-1717498, DEB 1911853, DEB-1051342, CNH-1115069, and EF-0914866.

## **Tables and Figures**

**Table 3.1. Analysis of global seroprevalence data using a generalized linear model with a binomial distribution and a logit link with virus type as a fixed effect and study ID as a random effect. CHIKV was the reference level.**

	<b>Estimate</b>	<b>Std. Error</b>	<b>z value</b>	<b>P-value</b>
<b>Intercept (CHIKV)</b>	-1.27	0.15	-8.61	< 0.001
<b>DEN-1</b>	-1.33	0.034	-39.41	< 0.001
<b>DEN-2</b>	-0.64	0.034	-18.67	< 0.001
<b>DEN-3</b>	-0.82	0.038	-21.79	< 0.001
<b>DEN-4</b>	-2.21	0.041	-53.29	< 0.001
<b>YFV</b>	-0.047	0.022	-2.16	0.030
<b>ZIKV</b>	-0.65	0.022	-29.51	< 0.001





Figure 3.1. Human viremia (PFU/mL) by date of symptom onset.



**Figure 3.2.** Predicted fraction of *A. aegypti* transmitting virus by blood meal viremia. Circles represent vector competence study data with circle size corresponding to *A. aegypti* sample size. Data for CHIKV, YFV, and ZIKV are plotted for days 7 and 14 post-feeding. DENV1-4 was plotted for day 14 since source study data only measured vector competence on day 14.






Figure 3.3. Daily estimates of human infectiousness ( $\pm 1$  SE) for seven arboviruses for *A. aegypti* (top) and *A. albopictus* (bottom).






**Figure 3.4. Relative seroprevalence for studies simultaneously comparing CHIKV, YFV, and ZIKV (A), CHIKV and DENV1-2 (B), DENV1-4 (C), and all remaining studies (D) with studies ranked by average seroprevalence along the x-axis. Lines represent predicted seroprevalence values based on fitted model with virus type and study ID as predictor variables.**



**Figure 3.5. Fitted coefficients of relative seroprevalence plotted against relative  $R_0$  estimates for *A. aegypti* for all data (A), Africa (B), the Americas (C), and Asia (D). Seroprevalence coefficients were significantly positively correlated with relative  $R_0$  in Asia (Panel D; Seroprevalence Fitted Coefficients =  $-5.82 + 2.62 (\pm 0.81) * R_0$ ,  $P = 0.023$ ), but not for the global dataset (Panel A; Seroprevalence Fitted Coefficients =  $-0.72 - 0.062 (\pm 0.48) * R_0$ ,  $P = 0.90$ ), in Africa (Panel B; Seroprevalence Fitted Coefficients =  $-0.51 - 0.080 (\pm 0.29) * R_0$ ,  $P = 0.80$ ) or the Americas (Panel C; Seroprevalence Fitted Coefficients =  $-0.10 + 0.045 (\pm 0.69) * R_0$ ,  $P = 0.95$ ). YFV was not included in the Americas due to a lack of serological data.**



**Figure 3.6. Estimated seroprevalence for each arbovirus for all data (A), Africa (B), the Americas (C), and Asia (D) for an intermediate value of average seroprevalence. Statistically significant differences between viruses are indicated with compact letter display. YFV was not included in the Americas due to a lack of serological data.**

## **Conclusion:**

The unifying goal of this research is an improved understanding of the ecological factors that affect human disease risk whether through how land use influences mosquito abundance, the efficacy of insecticides used to control mosquito populations, and how variations in host and vector transmission capabilities of viruses determines the scope of epidemics. Better understanding the ecological determinants of human disease risk can then be used to devise better control strategies which reduce human disease burden. The ongoing urbanization of our planet is one of the most drastic changes effecting the natural world and as such a pressing area of study in ecology. Urbanization, broadly defined as the process by which land is covered by impervious surfaces by humans, is among the most drastic and consequential form of human land use and plays a pivotal role in shaping the biotic and abiotic conditions faced by many organisms (Gubler et al. 2011). This research provides new insights into species abundance in response to urbanization and thus how the process shapes disease risk across a grand temporal and spatial scale from two globally important mosquito species. While urbanization was strongly correlated with mosquito abundance and insecticide resistance, the precise mechanisms of this relationship remains largely unresolved.

Urbanization positively increases the abundance of one major global vector at a constant rate despite vastly different climates and ecotypes, ranging from the arid Sahel to lush Equatorial Forests. While simultaneously, urbanization decreases, has no effect, and increases the abundance of another species even in cities sharing near

identical climates. These results show that while one species' abundance in response to urbanization can be a linear relationship, other species can have completely divergent relationships on a city-by-city basis.

The correlates of urbanization one would expect to determine mosquito abundance, such as host abundance, larval habitat availability, and number of human structures, were indeed significantly correlated with abundance, but none were better predictors than the broader vague categorization of our urbanization index. Other factors important to mosquito abundance which we expected to be mediated by urbanization, temperature and humidity, were not significantly correlated with urbanization.

Furthermore, urbanization shapes disease risk by decreasing the efficacy of insecticides, the primary public health tool for controlling arbovirus disease outbreaks. The mechanisms of the positive relationship between insecticide resistance (i.e. increased survival to insecticide exposure) are another large gap in our understanding. Generally urbanization takes place at the expense of agricultural land where the vast majority of insecticide use occurs. In the absence of public health vector control programs (as is the case in Cameroon) what mechanisms then would potentially lead to greater resistance with increased urbanization? Does this phenomenon hold true for other vectors, or arthropods in general, as urbanization increases? Further elucidating the mechanisms of urbanization's effects on abundance and insecticide resistance is clearly an area in need of more research.

Considering our results in the context of climate change, another area of urgent focus in the field of ecology, the potential effects of climate change on disease risk are unclear as a result of this research. Increased temperatures led to lower abundance of *A. aegypti* across all habitats and increased humidity was significantly correlated with greater *A. albopictus* abundance. Increased rainfall was correlated with greater abundance of both species as well. Furthermore, insecticide resistance decreased along a south to north gradient as the climate became hotter and drier. As such, as hotter drier climates arise this could likely reduce disease risk by limiting vector populations.

Our estimate of arbovirus transmission potential (relative  $R_0$ ) provide further hope that increased burden and spread of vector borne disease to all corners of the world is not inexorable. If the pathogen with both the highest seroprevalence and highest relative  $R_0$ , CHIKV, is unable to become established in a given location (e.g. the United States), then our results suggest it is less likely that the other arboviruses in this study would be able to become established in the same population. These calculations provide a plausible explanation for why the ranges of our studied arboviruses are considerably smaller than their vector populations.

Comparing these calculations of relative  $R_0$  to human seroprevalence as a measure of transmission intensity shows a disconnect between calculated predictions of pathogen fitness and the actual transmission intensity of these pathogens in field settings. Our results show that a pathogen, such as yellow fever, may have relatively poor host and vector competence in some populations and thus have lower predicted


fitness, but simultaneously have higher relative levels of host antibodies in field observations. Essentially, pathogens which by our measure should have the lowest fitness and thus lower infection rates somehow show among the highest level of fitness in infecting humans. The potential explanations for this mismatch are a target rich area for future research. Our novel methodology of comparing calculations of relative transmission potential from laboratory derived data on host and vector competences with seroprevalence (or some other measure of prior infection) can be applied to other host-pathogen systems to test if calculations of pathogen fitness match field observations.

In our study system, the mismatch between relative seroprevalence and relative  $R_0$  may arise from spatial or temporal variation among viruses, mosquitoes, or human populations and their contributions to relative epidemic potential. In both *A. aegypti* and *A. albopictus*: vector competence can vary enormously within the same mosquito populations over relatively short spatial and time scales (<1 year) (Gubler et al. 1979, Kilpatrick et al. 2010, Goncalvez et al. 2014). Host competence and seroprevalence can also be mediated by prior exposure to other arboviruses, possibly different rates of waning in antibodies, or perhaps by variations between human population in viremia. The extent to which any of these variations determine transmission intensity within a population is largely unexplored. Further studies on the factors underpinning host and vector competence are needed to understand if our estimates of host and vector competence can be extrapolated beyond any one particular place and time.


The ranges of both *A. aegypti* and *A. albopictus* are expanding and our increasingly interconnected world allows for pathogens to spread quickly (Shragai et al. 2017). Understanding and reducing disease risk relies on accurate assessments of host and vector competence, knowledge of the environmental factors determining vector abundance, and availability of effective means of controlling vector populations. Unlike most fields of ecology, we seek to actively disrupt and destroy our study organisms of disease-spreading vectors and hope the knowledge gained from this research will be useful in doing so.




## Appendix 1



**Figure S1: Example of land use data buffers (100m-2km) generated for collection sites in the city of Kribi.**




**Figure S2. Available human hosts observed at collection sites in Douala, Kribi, and Yaounde. Lines and ribbons show fitted models with urbanization index as predictor and 95% CI, respectively.**



**Figure S3. Monthly rainfall and mosquito abundance for the city of Yaounde.**

**Dark circles represent the mean catch rate per hour across all sites (+/- 1SE).**



**Figure S4. Monthly population growth rate of *A. albopictus* plotted against abundance of *A. aegypti* in the preceding month (A) and vice versa (B). The relationship was not significant for an effect of *A. aegypti* abundance on *A. albopictus* population growth rates the following month ( $\text{Log}(\text{ALBPopGrowth})=0.22-0.011 (\pm 0.0089)*\text{AEG}$ ,  $P = 0.27$ ) nor for an effect of *A. albopictus* abundance on the following month *A. aegypti* population growth rates ( $\text{Log}(\text{AEGPopGrowth})=0.14-0.0026 (\pm 0.0059)*\text{ALB}$ ,  $P = 0.66$ ).**

**Table S1. Previous studies on *Ae. aegypti* abundance and urbanization.**

<b>Study</b>	<b>Location</b>	<b>Urbanization Metric</b>	<b>Urbanization Correlation w/<i>Ae. aegypti</i> Abundance</b>
<b>Braks et al. 2003</b>	Rio de Janeiro & Nova Iguacu, Brazil; Boca Raton & West Palm Beach, FL, USA	Qualitative (Urban, suburban, rural)	Positive
<b>Carbajo et al. 2006</b>	Buenos Aires, Argentina	Quantitative GIS Land Cover Analysis (100m and 300m)	Positive
<b>Rey et al. 2006</b>	Manatee, Miami-Dade, & Palm Beach Co., FL, USA	Quantitative GIS Land Cover Analysis (100m)	Positive
<b>Tsuda et al. 2006</b>	Chiangmai Province, Thailand	Qualitative (Urban, Transition, Rural)	Positive
<b>Honorio et al. 2009</b>	Rio de Janeiro, Brazil	Qualitative (Urban, Low Vegetation, Medium Vegetation, High Vegetation)	Positive
<b>Higa et al. 2010</b>	Vietnam	Qualitative (Urban, Transition, Rural)	Mixed (regional variation)
<b>Fatima et al. 2016</b>	Pakistan	Quantitative GIS Land Cover Analysis (30m)	Positive
<b>Zahouli et al. 2016</b>	Cote D'Ivoire	Qualitative (Urban, Suburban, Rural)	Positive
<b>Ndenga et al. 2017</b>	Kenya	Qualitative (Urban, rural)	Positive
<b>Overgaard et al. 2017</b>	Colombia	Qualitative (Urban, rural)	Positive
<b>Dalpadado et al. 2018</b>	Gampaha District, Sri Lanka	Qualitative (Urban, Suburban, Rural)	Positive
<b>Estallo et al. 2018</b>	Cordoba, Argentina	Quantitative GIS Land Cover Analysis (10m)	Positive
<b>Talaga et al. 2020</b>	Kourou, French Guiana	Quantitative GIS Land Cover Analysis (70m)	Moderately Urbanized site had highest density

**Table S2. Previous studies on *Ae. albopictus* abundance and urbanization. Study number indicates study location and result in Figure 8.**

#	Study	Location	Urbanization Metric	Urbanization Correlation w/ <i>Ae. albopictus</i> Abundance
1	<b>Hornby et al. 1994</b>	Lee County, FL, USA	Qualitative (Urban, Suburban)	Negative
2	<b>Barker et al. 2003</b>	Virginia, USA	Qualitative (Forest, Yard Bordering Forest, Yard)	Positive (Unforested sites had highest abundance)
3	<b>Braks et al. 2003</b>	Rio de Janeiro & Nova Iguacu, Brazil; Boca Raton & West Palm Beach, FL, USA	Qualitative (Urban, suburban, rural)	Negative
4	<b>Rey et al. 2006</b>	Manatee, Miami-Dade, & Palm Beach Co., FL, USA	Quantitative GIS Land Cover Analysis (100m)	Negative
5	<b>Tsuda et al. 2006</b>	Chiangmai Province, Thailand	Qualitative (Urban, Transition, Rural)	Negative
6	<b>Honorio et al. 2009</b>	Rio de Janeiro, Brazil	Qualitative (Urban, Low Vegetation, Medium Vegetation, High Vegetation)	Negative
7	<b>Higa et al. 2010</b>	Vietnam	Qualitative (Urban,	None

			Transition, Rural)	
8	<b>Bagny et al. 2012</b>	Mayotte, France	Quantitative GIS Land Cover Analysis (25m)	Positive
9	<b>Li et al. 2014</b>	Guangzho, China	Qualitative (Urban, suburban, rural)	Positive
1 0	<b>Samson et al. 2015</b>	Cap-Haitien, Haiti	Quantitative GIS Land Cover Analysis (6.5m)	Positive
1 1	<b>Baldacchino et al. 2017</b>	Belluno & Trento, Italy	Quantitative GIS Land Cover Analysis (250m)	Positive
1 2	<b>Dalpadado et al. 2018</b>	Gampaha District, Sri Lanka	Qualitative (Urban, Suburban, Rural)	Negative
1 3	<b>McClure et al. 2018</b>	Big Island, Hawaii, USA	Quantitative GIS Land Cover Analysis	Positive
1 4	<b>Arduino et al. 2020</b>	Sao Paulo, Brazil	Qualitative (Urban, Forest, Grass- Shrubs)	Positive
1 5	<b>Westby et al. 2021</b>	St. Louis, Missouri, USA	Qualitative (Urban, Suburban, Rural)	Positive (Urban & Suburban equal, but both higher than rural)

Table S3: Sequence of primers used for viral detection.

Primers	Sequence (5'-3')	References
ZIKV-forward	nt9271-AARTACACATACCARAACAAAgTggT9297	(Lanciotti <i>et al.</i> , 2008)
ZIKV-reverse	nt9352-TCCRCTCCCYCTYTggTCTTg-9373	
ZIKV-probe	nt9304-FAM-CTYAgACCAgCTgAAR-BBQ-9320	
CHIKV-forward	AAGCTYCGCGTCCTTTACCAAG	(Pastorino <i>et al.</i> , 2005)
CHIKV-reverse	CCAAATTGTCCYGGTCTTCCT	
CHIKV-probe	FAM-CCAATGTCYTCMGCCTGGACACCTTT-TAMRA	
DENV-forward	AGGACYAGAGGTTAGAGGAGA	(Leparc-Goffart <i>et al.</i> , 2009)
DENV-reverse	CGYTCTGTTGCCTGGAWTGAT	
DENV-probe	FAM-ACAGCATATTGACGCTGGGARAGACC-TAMRA	

Table S4: Average hourly mosquito collection rate by city and species.

City	<i>A. aegypti</i> females/hour	SD of <i>A. aegypti</i> females/hour	<i>A. albopictus</i> females/hour	SD of <i>A. albopictus</i> females/hour
Douala	18.2	29.7	8.9	10.3
Garoua	2.3	3.1	0.0	0.0



Kribi	4.4	11.1	13.2	11.3
Maroua	1.4	2.1	0.0	0.0
Ngaoundere	1.2	1.3	0.0	0.0
Yaounde	0.7	2.4	21.7	30.6

Table S5: Hourly collection rate of mosquitoes from each collection site. Urbanization indices are included for each location.

Site	City	UI 100M	UI 200M	UI 500M	UI 1KM	UI 2KM	<i>A. aegypti</i> females/ hr	<i>A. albopictus</i> females/hr
<b>Akwa</b>	Douala	96	90	90	85	81	5.3	9.7
<b>Bepanda</b>	Douala	85	85	87	88	92	7.2	0.0
<b>Bonaberi</b>	Douala	97	93	91	76	64	21.5	17.0
<b>Brazzaville</b>	Douala	100	100	98	92	89	10.0	0.0
<b>Deido</b>	Douala	100	100	98	91	78	67.0	2.3
<b>Kotto</b>	Douala	100	99	98	93	79	2.7	19.3
<b>Logbessou</b>	Douala	75	86	88	59	58	15.7	10.5
<b>New Bell</b>	Douala	19	30	70	80	88	23.7	0.7
<b>Ngodji</b>	Douala	0	0	0	0	5	0.0	5.3
<b>Yassa</b>	Douala	24	25	23	25	22	3.0	29.3
<b>Bockle-Siha</b>	Garoua	0	0	0	1	1	0.0	0.0
<b>Camp-Chinois</b>	Garoua	97	97	74	63	42	1.3	0.0
<b>Foulberre</b>	Garoua	100	98	98	83	54	6.5	0.0
<b>Lainde</b>	Garoua	56	51	40	50	36	3.5	0.0
<b>Nassarao</b>	Garoua	67	54	43	21	14	1.7	0.0
<b>Poumpoumre</b>	Garoua	35	32	56	61	50	5.2	0.0
<b>Rocade</b>	Garoua	43	17	7	16	22	3.3	0.0

<b>Round e Adja</b>	Garoua	100	96	63	58	61	0.0	0.0
<b>Round e-Adja</b>	Garoua	100	96	63	58	61	0.0	0.0
<b>Sangue re-Paul</b>	Garoua	0	0	0	2	1	0.0	0.0
<b>Yelwa</b>	Garoua	100	98	85	68	54	0.8	0.0
<b>Zoological Garden</b>	Garoua	53	43	63	70	57	2.7	0.0
<b>Bebwambé</b>	Kribi	0	5	13	17	20	0.6	11.6
<b>Bikondo</b>	Kribi	0	0	0	1	2	0.0	2.0
<b>Dombe</b>	Kribi	85	77	68	55	32	1.0	11.7
<b>Kingué 1</b>	Kribi	1	14	29	37	32	0.0	0.0
<b>Kingué 2</b>	Kribi	96	89	75	69	47	43.7	2.0
<b>Londji</b>	Kribi	0	0	3	21	25	1.9	21.5
<b>Mboamanga</b>	Kribi	91	83	60	52	42	17.0	22.5
<b>Mokolo</b>	Kribi	63	58	61	56	43	1.0	13.3
<b>Ngoyé</b>	Kribi	74	63	57	58	51	1.2	10.5
<b>Polongwe</b>	Kribi	34	12	2	2	6	1.1	13.9
<b>Zaire</b>	Kribi	100	94	73	57	35	2.2	10.7
<b>Comice</b>	Maroua	88	93	76	55	43	0.7	0.0
<b>Djarengol</b>	Maroua	64	56	49	51	43	3.0	0.0
<b>Domayo</b>	Maroua	74	55	58	71	52	0.5	0.0
<b>Dougoi</b>	Maroua	93	88	79	79	62	1.6	0.0
<b>Florina</b>	Maroua	0	0	2	1	8	0.5	0.0
<b>Louggo 1</b>	Maroua	0	0	0	0	0	0.2	0.0
<b>Makabaye</b>	Maroua	41	34	19	12	13	0.0	0.0
<b>Salak</b>	Maroua	16	22	13	10	7	4.3	0.0
<b>Zaika</b>	Maroua	1	0	0	1	8	0.3	0.0
<b>Bamnyanga</b>	Ngaoundere	46	31	24	14	18	1.3	0.0

<b>Beka Hosere</b>	Ngaoundere	0	0	2	3	12	1.0	0.0
<b>Burkina</b>	Ngaoundere	100	98	82	54	31	0.3	0.0
<b>Djakboul</b>	Ngaoundere	60	62	72	61	59	1.2	0.0
<b>Gadamabanga</b>	Ngaoundere	0	0	1	2	12	0.8	0.0
<b>Joli Soir</b>	Ngaoundere	100	100	98	87	65	0.8	0.0
<b>Maborno</b>	Ngaoundere	50	32	14	10	6	0.3	0.0
<b>Madagascar</b>	Ngaoundere	96	98	93	79	54	0.7	0.0
<b>Mboundjere</b>	Ngaoundere	100	100	88	79	65	4.3	0.0
<b>Sabongari</b>	Ngaoundere	93	83	57	53	41	2.0	0.0
<b>Afanoya I</b>	Yaounde	0	0	0	1	1	0.0	5.1
<b>Afanoya II</b>	Yaounde	25	10	2	1	1	0.3	8.9
<b>Essos</b>	Yaounde	100	100	98	96	91	0.3	28.2
<b>Mending</b>	Yaounde	89	86	85	84	68	0.0	24.8
<b>Messa</b>	Yaounde	79	66	81	86	86	5.5	16.3
<b>Mvog-Ada</b>	Yaounde	100	100	93	95	86	1.1	24.8
<b>Nkolbisson</b>	Yaounde	69	62	43	38	48	0.1	18.2
<b>Omnisport</b>	Yaounde	99	89	85	93	92	0.1	66.4
<b>Tsinga</b>	Yaounde	63	72	58	54	43	0.2	16.1
<b>Zamassi</b>	Yaounde	0	4	1	0	0	0.1	0.9

**Table S6: Analysis of *A. aegypti* abundance with urbanization index (UI), temperature, humidity, and prior month's rain in six cities as predictors using a generalized linear model with negative binomial distribution and a log link. The city of Douala was the reference level.**

Predictor	Estimate	SE	Z-Value	P
(Intercept)	6.075	2.80	2.17	0.030
UI_1KM	0.026	0.0044	5.80	≤0.001
Monthly_Rain (mm)	0.0027	0.0011	2.52	0.012
City of Garoua	-0.39	0.75	-0.52	0.61
City of Kribi	-0.98	0.71	-1.39	0.17
City of Maroua	-1.49	0.73	-2.03	0.042
City of Ngaoundere	-2.40	0.82	-2.92	0.0035
City of Yaounde	-3.91	0.64	-6.12	≤0.001
Temperature	-0.18	0.070	-2.62	0.0088
Humidity	-0.0068	0.011	-0.60	0.55

**Table S7: Analysis of *A. albopictus* abundance with urbanization index (UI) interacting with city and temperature, humidity, and prior month's rain in three cities as predictors with a generalized linear model with a negative binomial distribution and a log link. The city of Douala was the reference level.**

Predictor	Estimate	SE	Z- value	P
(Intercept)	-0.314	2.45	-0.13	0.90
UI_2KM	-0.023	0.012	-1.96	0.049
City of Kribi	-1.076	0.96	-1.12	0.26
City of Yaounde	-1.924	0.86	-2.23	0.026
Monthly_Rain (mm)	0.0013	0.00067	2.0080	0.045
Temperature	0.072	0.057	1.25	0.21
Humidity	0.021	0.010	2.015	0.044
UI_2KM:Kribi	0.021	0.019	1.10	0.27
UI_2KM:Yaounde	0.047	0.012	3.93	≤0.001

**Table S8. Analysis of temperature in response to urbanization index (UI) at 100m interacting with city as predictors with a generalized linear model with a negative binomial distribution and a log link. The city of Kribi was the reference level.**

Predictor	Estimate	SE	Z- value	P
(Intercept)	3.33	0.059	56.41	≤0.001
UI_100M	-0.00026	0.0010	-0.27	0.79
City of Douala	0.045	0.14	0.33	0.74
City of Yaounde	-0.094	0.070	-1.33	0.18
City of Ngaoundere	-0.11	0.11	-1.06	0.29
City of Garoua	0.15	0.095	1.55	0.12
City of Maroua	0.010	0.089	0.12	0.91
UI_100M:Douala	0.000062	0.0018	0.034	0.97
UI_100M:Yaounde	0.00050	0.0011	0.45	0.65
UI_100M:Ngaoundere	0.000046	0.0015	0.030	0.98
UI_100M:Garoua	-0.00024	0.0014	-0.17	0.87
UI_100M:Maroua	0.00011	0.0015	0.074	0.94

**Table S9. Analysis of humidity in response to urbanization index (UI) at 100m interacting with city as predictors with a generalized linear model with a negative binomial distribution and a log link. The city of Kribi was the reference level.**

Predictor	Estimate	SE	Z- value	P
(Intercept)	4.41	0.076	58.00	≤0.001
UI_100M	-0.00034	0.0013	-0.27	0.79
City of Douala	-0.089	0.18	-0.49	0.62
City of Yaounde	-0.052	0.090	-0.58	0.56
City of Ngaoundere	-0.16	0.13	-1.16	0.25
City of Garoua	-0.25	0.13	-1.92	0.055
City of Maroua	-0.23	0.12	-1.96	0.050
UI_100M:Douala	0.0011	0.0024	0.48	0.63
UI_100M:Yaounde	0.000024	0.0014	0.017	0.99
UI_100M:Ngaoundere	-0.00077	0.0019	-0.40	0.69
UI_100M:Garoua	0.00097	0.0020	0.50	0.62
UI_100M:Maroua	0.0017	0.0019	0.886	0.38

**Table S10: Compiled Minimum Infection Rates (MIR) for studies which detected Zika virus in field caught specimens of *A. aegypti* and *A. albopictus*.**

Study	Country/ Location	Species	Zika+ Pools	Total Pools	Specimens Tested	Max Pool Size	MIR
<b>Marchette 1969</b>	Malaysia	<i>A. aegypti</i>	1	58	1277	80	0.8
<b>Grard 2014</b>	Gabon	<i>A. albopictus</i>	2	91	2130	25	0.9
<b>Ferreira-de-Brito 2016</b>	Brazil	<i>A. aegypti</i>	3	198	550	10	5.5
<b>Guerbois 2016</b>	Mexico	<i>A. aegypti</i>	15	55	279	NA	53.8
<b>Ho 2017</b>	Singapore	<i>A. aegypti</i> and <i>A. albopictus</i> **	9	517	1375	5	6.5
<b>Cevallos 2018</b>	Ecuador	<i>A. aegypti</i>	2	14	193	10	10.4
<b>Correa-Morales 2019</b>	Mexico	<i>A. aegypti</i>	260	3120	14,145	Data Unavailable	18.4
<b>Correa-Morales 2019</b>	Mexico	<i>A. albopictus</i>	7	52	78	Data Unavailable	89.7
<b>Singh 2019</b>	India	<i>A. aegypti</i>	3	55	203	10	14.8
<b>Ali 2020</b>	Malaysia	<i>A. albopictus</i>	6	NA	186	25	32.3
<b>Calle-Tobon 2020</b>	Colombia	<i>A. aegypti</i>	98	Data Unavailable	6585	10	14.9
<b>Campos 2020</b>	Cape Verde	<i>A. aegypti</i>	2	Not Pooled	816	1	2.5
<b>Kosoltanapawit 2020</b>	Thailand	<i>A. aegypti</i>	2	Not Pooled	130	1	15.4
<b>Phumee 2020*</b>	Thailand	<i>A. aegypti</i>	Data Unavailable	Data Unavailable	Data Unavailable	10	22.4
<b>Parra 2022</b>	Sao Paulo, Brazil	<i>A. aegypti</i>	55	607	1026	10	53.6
<b>Parra 2022</b>	Sao Paulo, Brazil	<i>A. albopictus</i>	1	11	12	10	83.3

<b>This Study</b>	Cameroon	<i>A. albopictus</i>	1	289	7490	30	0.1
-------------------	----------	----------------------	---	-----	------	----	-----


- Study provided an MIR without number of specimens or pools tested.
- \* Study data did not allow for differentiation of *A. aegypti* and *A. albopictus* so results were combined.

## Appendix 2

DocuSign Envelope ID: 9D79BC25-CE1E-44BF-BF9E-895BAC74C7A0

The text of this dissertation includes a reprint of the following previously published material: Montgomery, M., Harwood, J.F., Yougang, A.P. *et al.* Spatial distribution of insecticide resistant populations of *Aedes aegypti* and *Ae. albopictus* and first detection of V410L mutation in *Ae. aegypti* from Cameroon. *Infect Dis Poverty* **11**, 90 (2022). The co-authors listed in this publication have approved of its inclusion.


DocuSigned by:  
  
DC9A13EB1B15443...  
James F. Harwood

DocuSigned by:  
  
FE03B432FB334B8...  
Aurelie P. Yougang

DocuSigned by:  
  
7CBF4C3228C64AA...  
Armel N. Tedjou

DocuSigned by:  
  
08E94CA78E0A4D6...  
Christophe Rostand Keumeni

DocuSigned by:  
  
4D4FFB8BE589489...  
Theodel A. Wilson-Bahun

DocuSigned by:  
  
E92CADF9116849A...  
Charles S. Wondji

DocuSigned by:  
  
7B6F28C07BF540B...  
Basile Kamgang



### Appendix 3

**Table S1. Human viremia data. CHIKV (Appassakij et al 2013, Riswari et al. 2015), DENV1-4 (Nguyen et al. 2013), YFV (Nassar et al. 1995, Monath et al. 2012), ZIKV (Mansuy et al. 2017).**

Day	Viremia	SE	SD	N	Virus	
1	5.35		0.15	0.21	2	YFV
2	4.06		0.61	1.84	9	YFV
3	4.24		0.50	1.41	8	YFV
4	2.44		0.36	0.80	5	YFV
5	1.90		0.40	0.57	2	YFV
6	2.50		0.40*	0.96	1	YFV
1	8.20		0.35	NA	2	CHIKV
2	7.81		0.61	NA	11	CHIKV
3	5.57		1.49	NA	6	CHIKV
4	4.46		1.40	NA	8	CHIKV
5	3.15		0.55	NA	3	CHIKV
6	2.45		0.46	NA	5	CHIKV
7	2.63	NA		NA	1	CHIKV
11	1.08	NA		NA	1	CHIKV
13	2.98	NA		NA	1	CHIKV
17	1.86	NA		NA	1	CHIKV
-2	5.80		0.63	1.26	1	ZIKV
0	4.53		0.57	1.13	4	ZIKV
1	4.91		0.40	0.89	5	ZIKV
2	4.26		0.35	1.09	10	ZIKV
3	3.74		0.28	0.69	6	ZIKV
4	3.65		1.10	1.91	3	ZIKV
5	3.61		1.08	1.87	3	ZIKV
7	1.70	NA		NA	1	ZIKV

10	1.85	NA	NA	1	ZIKV
1	8.56	NA	NA	1	DENV3
2	7.81	0.30	1.25	18	DENV1
2	8.01	0.25	1.13	20	DENV2
2	8.74	0.25	0.71	8	DENV3
2	6.86	0.74	1.28	3	DENV4
3	8.17	0.15	1.24	65	DENV1
3	7.59	0.14	1.24	74	DENV2
3	7.99	0.26	1.04	16	DENV3
3	6.75	0.46	1.66	13	DENV4
4	7.79	0.17	1.46	78	DENV1
4	6.85	0.16	1.41	83	DENV2
4	7.52	0.33	1.65	25	DENV3
4	6.06	0.42	1.96	22	DENV4
5	6.67	0.18	1.57	78	DENV1
5	5.44	0.16	1.41	83	DENV2
5	6.33	0.31	1.53	25	DENV3
5	4.98	0.40	1.87	22	DENV4
6	5.44	0.14	1.19	77	DENV1

6	3.95	0.14	1.24	82	DENV2
6	4.77	0.36	1.75	24	DENV3
6	3.47	0.24	1.12	21	DENV4
7	4.50	0.11	0.87	60	DENV1
7	3.01	0.14	1.13	61	DENV2
7	3.62	0.34	1.39	17	DENV3
7	2.86	0.00	0.00	19	DENV4
8	3.77	0.31	0.96	10	DENV1
8	2.59	0.40	1.14	8	DENV2
8	2.93	0.37	0.99	7	DENV3
8	2.86	NA	NA	9	DENV4

\*Average SE for prior 5 days of YFV.

**Table S2. Analysis of viremia with day and virus as predictors. Day 1 and CHIKV were the reference levels.**

	Estimate	Std. Error	t-Value	P-Value
<b>(Intercept)</b>	7.2730	0.4176	17.416	< 2e-16
<b>VirusDENV1</b>	-1.8500	0.4511	-4.101	0.000289
<b>VirusDENV2</b>	-1.9128	0.4511	-4.241	0.000196
<b>VirusDENV3</b>	-1.5284	0.4511	-3.388	0.001983
<b>VirusDENV4</b>	-1.5085	0.4511	-3.344	0.002227
<b>VirusYFV</b>	-1.8596	0.4511	-4.123	0.000272
<b>VirusZIKV</b>	-1.4701	0.4511	-3.259	0.002779

	Estimate	Std. Error	t-Value	P-Value
Day2	-0.5155	0.4176	-1.235	0.226592
Day3	-1.3048	0.4176	-3.124	0.003931
Day4	-2.2862	0.4176	-5.474	6.1E-06
Day5	-3.1883	0.4176	-7.635	1.63E-08
Day6	-4.7024	0.4176	11.260	2.69E-12

**Table S3. Results from a Bayesian model fit to vector competence data. Coefficients were fit for the mean extrinsic incubation period (EIP) in days, including virus, mosquito species, and viremia, with *A. aegypti* and ZIKV as the reference level. The model also included two additional study ID intercept parameters (ID1 for *A. albopictus* transmitting CHIV, and ID2 for *A. aegypti* transmitting ZIKV) to account for substantial differences among these two individual studies and all others in the fraction transmitting.**

	Mean EIP	SE	95% CI
Intercept ( <i>A. aegypti</i> , Zika virus)	2.64	0.12	2.41 - 2.87
<i>A. albopictus</i>	0.11	0.052	0.0080 - 0.21
CHIKV	-0.33	0.066	-0.46 - -0.21
YFV	0.26	0.055	0.15 - 0.37
Viremia (PFU/ml)	0.18	0.021	0.14 - 0.23

	Mean EIP	SE	95% CI
ID1 ( <i>A. albopictus</i> CHIKV)	-2.24	0.28	-2.78 - -1.70
ID2 ( <i>A. aegypti</i> ZIKA)	-0.89	0.16	-1.20 - -0.58

**Table S4. Analysis of seroprevalence data for Africa using a generalized linear model with a binomial distribution and a logit link with virus type as a fixed effect and study ID as a random effect. CHIKV was the reference level.**

	Estimate	Std. Error	z value	P-value
<b>Intercept (CHIKV)</b>	-1.47	0.20	-7.24	< 0.001
<b>DEN-1</b>	-1.40	0.046	30.68	< 0.001
<b>DEN-2</b>	-0.71	0.053	13.39	< 0.001
<b>DEN-3</b>	-0.58	0.10	-6.08	< 0.001
<b>DEN-4</b>	-0.86	0.10	-8.33	< 0.001
<b>YFV</b>	-0.10	0.022	-4.60	< 0.001
<b>ZIKV</b>	-0.81	0.024	33.87	< 0.001


**Table S5. Analysis of seroprevalence data for the Americas using a generalized linear model with a binomial distribution and a logit link with virus type as a fixed effect and study ID as a random effect. CHIKV was the reference level..**

	Estimate	Std. Error	z value	P-value
<b>Intercept (CHIKV)</b>	-1.76	0.61	-2.90	0.0038
<b>DEN-1</b>	0.031	0.71	0.043	0.97


	Estimate	Std. Error	z value	P-value
<b>DEN-2</b>	0.57	0.71	0.80	0.42
<b>DEN-3</b>	-0.15	0.71	-0.21	0.83
<b>DEN-4</b>	-1.28	0.71	-1.81	0.07
<b>ZIKV</b>	0.70	0.068	10.44	< 0.001

**Table S6. Analysis of seroprevalence data for Asia using a generalized linear model with a binomial distribution and a logit link with virus type as a fixed effect and study ID as a random effect. CHIKV was the reference level.**

	Estimate	Std. Error	z value	P-value
<b>Intercept (CHIKV)</b>	-1.13	0.05	-22.12	< 0.001
<b>DEN-1</b>	-2.072	0.08	-24.74	< 0.001
<b>DEN-2</b>	-0.74	0.08	-9.68	< 0.001
<b>DEN-3</b>	-0.20	0.08	-2.44	0.014
<b>DEN-4</b>	-2.093	0.09	-23.18	< 0.001
<b>YFV</b>	-6.64	1.31	-5.078	< 0.001
<b>ZIKV</b>	-1.087	0.31	-3.55	< 0.001



**Figure S1. Predicted fraction of *A. albopictus* transmitting virus by blood meal viremia. Circles represent vector competence study data with circle size corresponding to *A. aegypti* sample size. Data for CHIKV, YFV, and ZIKV are plotted for days 7 and 14 post-feeding. DENV1-4 was plotted for day 14 since source study data only measured vector competence on day 14.**




**Figure S2. Predicted fraction of mosquitoes transmitting virus by blood meal viremia on day 14.**





**Figure S3. Number of serosurveys and total sample size by country.**



**Figure S4. Fitted coefficient of arbovirus seroprevalence plotted against relative  $R_0$  estimates for *A. albopictus*. The relationship was not statistically significant in any location: global (Fitted Coefficient =  $-0.61 - 0.21 (\pm 0.59) * R_0$ , ( $P = 0.74$ ), Africa (Fitted Coefficient =  $-0.55 - 0.067 (\pm 0.31) * R_0$ , ( $P = 0.84$ ), the Americas (Fitted Coefficient =  $-2.91 + 0.71 (\pm 1.59) * R_0$ , ( $P = 0.68$ ), or Asia (Fitted Coefficient =  $-4.25 + 2.14 (\pm 1.15) * R_0$ , ( $P = 0.12$ ). YFV was not included in the Americas due to a lack of serological data.**

#### Supplemental References

Aitken, T. H., Downs, W. G., & Shope, R. E. (1977). *Aedes aegypti* strain fitness for yellow fever virus transmission. The American journal of tropical medicine and hygiene, 26(5 Pt 1), 985-989.

Amaya-Larios, I. Y., Martínez-Vega, R. A., Mayer, S. V., Galeana-Hernández, M., Comas-García, A., Sepúlveda-Salinas, K. J., Falcón-Lezama, J. A., Vasilakis, N., & Ramos-Castañeda, J. (2014). Seroprevalence of Neutralizing Antibodies Against

Dengue Virus in Two Localities in the State of Morelos, Mexico. *The American Journal of Tropical Medicine and Hygiene*, 91(5), 1057–1065.

Andayi, F., Charrel, R. N., Kieffer, A., Richet, H., Pastorino, B., Leparc-Goffart, I., Ahmed, A. A., Carrat, F., Flahault, A., & Lamballerie, X. de. (2014). A Sero-epidemiological Study of Arboviral Fevers in Djibouti, Horn of Africa. *PLOS Neglected Tropical Diseases*, 8(12), e3299.

Ang, L. W., Thein, T. L., Ng, Y., Boudville, I. C., Chia, P. Y., Lee, V. J. M., & Leo, Y. S. (2019). A 15-year review of dengue hospitalizations in Singapore: Reducing admissions without adverse consequences, 2003 to 2017. *PLoS neglected tropical diseases*, 13(5), e0007389.

Appassakij, H., Khuntikij, P., Kemapunmanus, M., Wutthanarungsan, R., & Silpapojakul, K. (2013). Viremic profiles in asymptomatic and symptomatic chikungunya fever: A blood transfusion threat? *Transfusion*, 53(10pt2), 2567–2574.

Ayorinde, A. F., Oyeyiga, A. M., Nosegbe, N. O., & Folarin, O. A. (2016). A survey of malaria and some arboviral infections among suspected febrile patients visiting a health centre in Simawa, Ogun State, Nigeria. *Journal of Infection and Public Health*, 9(1), 52–59.

Mogadishu, Somalia 2011: Co-circulation of three dengue virus serotypes.

*International Journal of Infectious Diseases*, 21, 3.

Botros, B. A., Watts, D. M., Soliman, A. K., Salib, A. W., Moussa, M. I., Mursal, H., Douglas, C., & Farah, M. (1989). Serological evidence of dengue fever among refugees, Hargeysa, Somalia. *Journal of medical virology*, 29(2), 79–81.

Braga, J. U., Bressan, C., Dalvi, A. P. R., Calvet, G. A., Daumas, R. P., Rodrigues, N., Wakimoto, M., Nogueira, R. M. R., Nielsen-Saines, K., Brito, C., Filippis, A. M. B. de, & Brasil, P. (2017). Accuracy of Zika virus disease case definition during simultaneous Dengue and Chikungunya epidemics. *PLOS ONE*, 12(6), e0179725.

Brès, P. (1970). [Recent data from serological surveys on the prevalence of arbovirus infections in Africa, with special reference to yellow fever]. *Bulletin of the World Health Organization*, 43(2), 223–267.

Calvo, Eliana Patricia, Coronel-Ruiz, Carolina, Velazco, Syrley, Velandia-Romero, Myriam, & Castellanos, Jaime E.. (2016). Diagnóstico diferencial de dengue y chikungunya en pacientes pediátricos. *Biomédica*, 36(Suppl. 2), 35-43.

Carrillo-Hernández, M. Y., Ruiz-Saenz, J., Villamizar, L. J., Gómez-Rangel, S. Y., & Martínez-Gutierrez, M. (2018). Co-circulation and simultaneous co-infection of dengue, chikungunya, and zika viruses in patients with febrile syndrome at the Colombian-Venezuelan border. *BMC Infectious Diseases*, 18(1), 61.

Chouin-Carneiro, T., Vega-Rua, A., Vazeille, M., Yebakima, A., Girod, R., Goindin, D., ... & Failloux, A. B. (2016). Differential susceptibilities of *Aedes aegypti* and *Aedes albopictus* from the Americas to Zika virus. *PLoS neglected tropical diseases*, 10(3), e0004543.

Comach, G., Blair, P. J., Sierra, G., Guzman, D., Soler, M., Quintana, M. C. de, Bracho-Labadie, M., Camacho, D., Russell, K. L., Olson, J. G., & Kochel, T. J. (2009). Dengue Virus Infections in a Cohort of Schoolchildren from Maracay,

Venezuela: A 2-Year Prospective Study. *Vector-Borne and Zoonotic Diseases*, 9(1), 87–92.

Duffy, M. R., Chen, T. H., Hancock, W. T., Powers, A. M., Kool, J. L., Lanciotti, R. S., Pretrick, M., Marfel, M., Holzbauer, S., Dubray, C., Guillaumot, L., Griggs, A., Bel, M., Lambert, A. J., Laven, J., Kosoy, O., Panella, A., Biggerstaff, B. J., Fischer, M., & Hayes, E. B. (2009). Zika virus outbreak on Yap Island, Federated States of Micronesia. *The New England journal of medicine*, 360(24), 2536–2543.

Eligio-García, L., Crisóstomo-Vázquez, M. del P., Caballero-García, M. de L., Soria-Guerrero, M., Méndez-Galván, J. F., López-Cancino, S. A., & Jiménez-Cardoso, E. (2020). Co-infection of Dengue, Zika and Chikungunya in a group of pregnant women from Tuxtla Gutiérrez, Chiapas: Preliminary data. 2019. *PLOS Neglected Tropical Diseases*, 14(12), e0008880.

Espino, C. (2009). "Active Surveillance and Incidence Rate of Dengue Infection in a Cohort of High Risk Population in Maracay, Venezuela." USF Tampa Graduate Theses and Dissertations.

Fagbami, A. H. (1979). Zika virus infections in Nigeria: Virological and seroepidemiological investigations in Oyo State. *The Journal of Hygiene*, 83(2), 213–219.

Fontenille, D. (1988). *Etude des circuits de vection d'arbovirus, a madagascar* [These de doctorat, Paris, EPHE].

FUKUMI, Hideo, Kaoru HAYASHI, Kumato MIFUNE, Atsuo UJIIYE, Osamu SUENAGA, Koichi FUTATSUKI, Sachiko MATSUO, and Ichiro MIYAGI.

"Epidemiological Studies on Viral and Parasitic Diseases, and Vector Insects in East Africa." 熱帯医学 9, no. 3 (1967): 127-135.

Gomez-Govea, M. A., Zamudio-Osuna, M. de J., Murillo, K. del C. T., Ponce, G., Cavazos, M. E. de la O., Tavitas-Aguilar, M. I., Flores-Suárez, A. E., Villarreal-Perez, J. Z., & Rodriguez-Sanchez, I. P. (2017). Chikungunya Fever in Patients from Northeastern Mexico. *Southwestern Entomologist*, 42(1), 143–152.

Guzmán, M. G., Kourí, G., Valdés, L., Bravo, J., Vázquez, S., & Halstead, S. B. (2002). Enhanced severity of secondary dengue-2 infections: death rates in 1981 and 1997 Cuban outbreaks. *Revista Panamericana de Salud Pública*, 11(4), 223-227.

Henderson, B. E., Metselaar, D., Cahill, K., Timms, G. L., Tukei, P. M., & Williams, M. C. (1968). Yellow fever immunity surveys in northern Uganda and Kenya and eastern Somalia, 1966-67. *Bulletin of the World Health Organization*, 38(2), 229–237.

Huy, R., Buchy, P., Conan, A., Ngan, C., Ong, S., Ali, R., ... & Vong, S. (2010). National dengue surveillance in Cambodia 1980-2008: epidemiological and virological trends and the impact of vector control. *Bulletin of the World Health Organization*, 88, 650-657.

Hyams, K. C., Oldfield, E. C., Scott, R. M., Bourgeois, A. L., Gardiner, H., Pazzaglia, G., Moussa, M., Saleh, A. S., Dawi, O. E., & Daniell, F. D. (1986). Evaluation of febrile patients in Port Sudan, Sudan: Isolation of dengue virus. *The American Journal of Tropical Medicine and Hygiene*, 35(4), 860–865.

Ido, E., Ahuka, S., Karhemere, S., Ibuki, K., Shibata, M., Kameoka, M., & Muyembe, J. J. (2016). Dengue virus infection during an outbreak of chikungunya virus in Democratic Republic of Congo. *Annals of African Medicine*, 10(1).

Johnson, B.W., Chambers, T.V, Crabtree, M.B, et al. (2002). Vector competence of Brazilian *Aedes aegypti* and *Ae. albopictus* for a Brazilian yellow fever virus isolate, *Transactions of The Royal Society of Tropical Medicine and Hygiene*, 96 (6), 611–613

Kajeguka, D. C., Kaaya, R. D., Mwakalinga, S., Ndossi, R., Ndaro, A., Chilongola, J. O., Mosha, F. W., Schiøler, K. L., Kavishe, R. A., & Alifrangis, M. (2016). Prevalence of dengue and chikungunya virus infections in north-eastern Tanzania: A cross sectional study among participants presenting with malaria-like symptoms. *BMC Infectious Diseases*, 16(1), 1–9.

Kama, M., Aubry, M., Naivalu, T., Vanhomwegen, J., Mariteragi-Helle, T., Teissier, A., Paoaafaite, T., Hué, S., Hibberd, M. L., Manuguerra, J.-C., Christi, K., Watson, C. H., Nilles, E. J., Aaskov, J., Lau, C. L., Musso, D., Kucharski, A. J., & Cao-Lormeau, V.-M. (2019). Sustained Low-Level Transmission of Zika and Chikungunya Viruses after Emergence in the Fiji Islands. *Emerging Infectious Diseases*, 25(8), 1535–1538.

Kikuti, M., Cunha, G. M., Paploski, I. A. D., Kasper, A. M., Silva, M. M. O., Tavares, A. S., Cruz, J. S., Queiroz, T. L., Rodrigues, M. S., Santana, P. M., Lima, H. C. A. V., Calcagno, J., Takahashi, D., Gonçalves, A. H. O., Araújo, J. M. G., Gauthier, K., Diuk-Wasser, M. A., Kitron, U., Ko, A. I., Ribeiro, G. S. (2015). Spatial

Distribution of Dengue in a Brazilian Urban Slum Setting: Role of Socioeconomic Gradient in Disease Risk. *PLOS Neglected Tropical Diseases*, 9(7), e0003937.

Kokernot, R. H., Casaca, V. M. R., Weinbren, M. P., & McIntosh, B. M. (1965). Survey for antibodies against arthropod-borne viruses in the sera of indigenous residents of Angola. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 59(5), 563–570.

Kokernot, R. H., Szlamp, E. L., Levitt, J., & McIntosh, B. M. (1965). Survey for antibodies against arthropod-borne viruses in the sera of indigenous residents of the caprivi strip and bechuanaland protectorate\*. *Transactions of The Royal Society of Tropical Medicine and Hygiene*, 59(5), 553–562.

Kularatne SAM, Gihan MC, Weerasinghe SC, et al. (2009). Concurrent outbreaks of Chikungunya and Dengue fever in Kandy, Sri Lanka, 2006–07: a comparative analysis of clinical and laboratory features *Postgraduate Medical Journal* 85:342-346.

Kuniholm, M. H., Wolfe, N. D., Huang, C. Y.-H., Mpoudi-Ngole, E., Tamoufe, U., Burke, D. S., & Gubler, D. J. (2006). SEROPREVALENCE D DISTRIBUTION OF FLAVIVIRIDAE, TOGAVIRIDAE, AND BUNYAVIRIDAE ARBOVIRAL INFECTIONS IN RURAL CAMEROONIAN ADULTS. *The American Journal of Tropical Medicine and Hygiene*, 74(6), 1078–1083.

Leslie T, Martin NJ, Jack-Roosberg C, Odongo G, Beausoleil E, Tuck J, Raviprakash K, Kochel TJ. (2014). Dengue serosurvey in Sint Eustatius. *PLoS One*. 10;9(6):e95002.



- Li, M. I., Wong, P. S. J., Ng, L. C., & Tan, C. H. (2012). Oral Susceptibility of Singapore *Aedes (Stegomyia) aegypti* (Linnaeus) to Zika Virus. *PLOS Neglected Tropical Diseases*, 6(8), e1792.
- Macnamara, F. N., Horn, D. W., & Porterfield, J. S. (1959). Yellow fever and other arthropod-borne viruses: A consideration of two serological surveys made in South Western Nigeria. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 53(2), 202–212.
- Luca, M. D., Severini, F., Toma, L., Boccolini, D., Romi, R., Remoli, M. E., Sabbatucci, M., Rizzo, C., Venturi, G., Rezza, G., & Fortuna, C. (2016). Experimental studies of susceptibility of Italian *Aedes albopictus* to Zika virus. *Eurosurveillance*, 21(18), 30223.
- Mansuy, J. M., Mengelle, C., Pasquier, C., Chapuy-Regaud, S., Delobel, P., Martin-Blondel, G., & Izopet, J. (2017). Zika Virus Infection and Prolonged Viremia in Whole-Blood Specimens. *Emerging Infectious Diseases*, 23(5), 863–865.  
<https://doi.org/10.3201/eid2305.161631>
- Manu, S. K. (2017). Detection of Dengue and Chikungunya Viruses in Febrile Patients at Greater Accra Regional Hospital [Thesis, University of Ghana].
- McCarthy, M. C., Haberberger, R. L., Salib, A. W., Soliman, B. A., El-Tigani, A., Khalid, I. O., & Watts, D. M. (1996). Evaluation of arthropod-borne viruses and other infectious disease pathogens as the causes of febrile illnesses in the Khartoum Province of Sudan. *Journal of Medical Virology*, 48(2), 141–146.

- Mcintosh, B. M., Harwin, R. M., Paterson, H. E., & Westwater, M. L. (1963). An epidemic of chikungunya in south-eastern Southern Rhodesia. *The Central African Journal of Medicine*, 9, 351–359.
- Miller, B. R., & Mitchell, C. J. (1986). Passage of yellow fever virus: its effect on infection and transmission rates in *Aedes aegypti*. *The American journal of tropical medicine and hygiene*, 35(6), 1302-1309.
- Ming, C. K., Thein, S., Thaug, U., Tin, U., Myint, K. S., Swe, T., Halstead, S. B., & Diwan, A. R. (1974). Clinical and laboratory studies on haemorrhagic fever in Burma, 1970-72. *Bulletin of the World Health Organization*, 51(3), 227–235.
- Mitchell, C. J., Miller, B. R., & Gubler, D. J. (1987). Vector competence of *Aedes albopictus* from Houston, Texas, for dengue serotypes 1 to 4, yellow fever and Ross River viruses. *Journal of the American Mosquito Control Association*, 3(3), 460-465.
- Monath, T.P. (2012). *Vaccines*, Plotkin, S., Orenstein, W., Offitt, P.A.,.. 6th Edition.
- Moore, D. L., Causey, O. R., Carey, D. E., Reddy, S., Cooke, A. R., Akinkugbe, F. M., David-West, T. S., & Kemp, G. E. (1975). Arthropod-borne viral infections of man in Nigeria, 1964-1970. *Annals of Tropical Medicine and Parasitology*, 69(1), 49–64.
- Morrill, J. C., Johnson, B. K., Hyams, C., Okoth, F., Tukei, P. M., Mugambi, M., & Woody, J. (1991). Serological evidence of arboviral infections among humans of coastal Kenya. *The Journal of Tropical Medicine and Hygiene*, 94(3), 166–168.
- Morrison, A. C., Minnick, S. L., Rocha, C., Forshey, B. M., Stoddard, S. T., Getis, A., Focks, D. A., Russell, K. L., Olson, J. G., Blair, P. J., Watts, D. M., Sihuincha,

- M., Scott, T. W., & Kochel, T. J. (2010). Epidemiology of Dengue Virus in Iquitos, Peru 1999 to 2005: Interepidemic and Epidemic Patterns of Transmission. *PLOS Neglected Tropical Diseases*, 4(5), e670.
- Nassar, E. D. S., Chamelet, E. L., Coimbra, T. L., Souza, L., Suzuki, A., Ferreira, I. B., ... & Rosa, A. (1995). Jungle yellow fever: clinical and laboratorial studies emphasizing viremia on a human case. *Revista do Instituto de Medicina Tropical de São Paulo*, 37, 337-341.
- Nogueira, R. M. R., Miagostovich, M. P., Lampe, E., Souza, R. W., Zagne, S. M. O., & Schatzmayr, H. G. (1993). Dengue epidemic in the state of Rio de Janeiro, Brazil, 1990–1: Co-circulation of dengue 1 and dengue 2 serotypes. *Epidemiology & Infection*, 111(1), 163–170.
- Nur, Y. A., Groen, J., Heuvelmans, H., Tuynman, W., Copra, C., & Osterhaus, A. D. (1999). An outbreak of West Nile fever among migrants in Kisangani, Democratic Republic of Congo. *The American Journal of Tropical Medicine and Hygiene*, 61(6), 885–888.
- Ochieng, C., Ahenda, P., Vittor, A. Y., Nyoka, R., Gikunju, S., Wachira, C., Waiboci, L., Umuro, M., Kim, A. A., Nderitu, L., Juma, B., Montgomery, J. M., Breiman, R. F., & Fields, B. (2015). Seroprevalence of Infections with Dengue, Rift Valley Fever and Chikungunya Viruses in Kenya, 2007. *PLoS ONE*, 10(7), e0132645.
- Oliveira, J. V., Duarte, A., Santos, C., Pessoa, L., Filho, C. M., Lima, J., Carvalho, D., Xavier, T., Figueiredo, E., Giovanetti, M., Almeida, B., Goes, J., Lima, F., Alcantara, L. C., & Siqueira, I. (2018). Prevalence of Zika, dengue and Chikungunya

virus infection in pregnant women and surveillance of congenital Zika infection in Salvador, Brazil. *International Journal of Infectious Diseases*, 73, 182.

Omarjee, R., Prat, C. M., Flusin, O., Boucau, S., Tenebray, B., Merle, O., Huc-Anais, P., Cassadou, S., & Leparc-Goffart, I. (2014). Importance of case definition to monitor ongoing outbreak of chikungunya virus on a background of actively circulating dengue virus, St Martin, December 2013 to January 2014.

*Eurosurveillance*, 19(13), 20753.

Osterrieth, P., Rathe, E., & Deleplanque-Liegeois, P. (1961). [Simultaneous isolation of the yellow fever and Chikungunya viruses at Bili (Haut-Uele, Belgian Congo)].

*Annales De La Societe Belge De Medecine Tropicale* (1920), 41, 207–212.

Pesko, K., Westbrook, C. J., Mores, C. N., Lounibos, L. P., & Reiskind, M. H. (2009). Effects of infectious virus dose and bloodmeal delivery method on susceptibility of *Aedes aegypti* and *Aedes albopictus* to chikungunya virus. *Journal of Medical Entomology*, 46(2), 395–399.

Pinto, M. R. (1967). Survey for antibodies to arboviruses in the sera of children in Portuguese Guinea. *Bulletin of the World Health Organization*, 37(1), 101–108

Proesmans, S., Katshongo, F., Milambu, J., Fungula, B., Mavoko, H. M., Ahuka-Mundeke, S., Luz, R. I. da, Esbroeck, M. V., Ariën, K. K., Cnops, L., Smet, B. D.,

Lutumba, P., Geertruyden, J.-P. V., & Vanlerberghe, V. (2018). Dengue and Chikungunya among Febrile Outpatients in Kinshasa, Democratic Republic of Congo: A cross-sectional study. *BioRxiv*, 486407.

- Pruszyński, C. A., Stenn, T., Acevedo, C., Leal, A. L., & Burkett-Cadena, N. D. (2020). Human Blood Feeding by *Aedes aegypti* (Diptera: Culicidae) in the Florida Keys and a Review of the Literature. *Journal of Medical Entomology*, 57(5), 1640–1647.
- Restrepo, B. N., Piedrahita, L. D., Agudelo, I. Y., Parra-Henao, G., & Osorio, J. E. (2012). Frequency and clinical features of dengue infection in a schoolchildren cohort from medellin, Colombia. *Journal of tropical medicine*, 2012, 120496.
- Richard, V., Paoaafaite, T., & Cao-Lormeau, V.-M. (2016). Vector Competence of *Aedes aegypti* and *Aedes polynesiensis* Populations from French Polynesia for Chikungunya Virus. *PLOS Neglected Tropical Diseases*, 10(5), e0004694.
- Riswari, S. F., Ma'roef, C. N., Djauhari, H., Kosasih, H., Perkasa, A., Yudhaputri, F. A., Artika, I. M., Williams, M., van der Ven, A., Myint, K. S., Alisjahbana, B., Ledermann, J. P., Powers, A. M., & Jaya, U. A. (2016). Study of viremic profile in febrile specimens of chikungunya in Bandung, Indonesia. *Journal of Clinical Virology*, 74, 61–65.
- Robin, Y. "Rapport de la 7ème conference technique de l'Organisation de Coordination et de Coopération pour la lutte contre les Grandes Endémies. OCCGE, Bobo-Dioulasso, Burkina Faso." Bobo-Dioulasso: OCCGE (1967).
- Robin, Y., Brès, P., Lartigue, J. J., Gidel, R., Lefèvre, M., Athawet, B., & Hery, G. (1968). Les arbovirus en Côte-d'Ivoire: enquête sérologique dans la population humaine. *Bull Soc Pathol Exot*, 61, 833-45.

Rodhain, F., Gonzalez, J. P., Mercier, E., Helynck, B., Larouze, B., & Hannoun, C. (1989). Arbovirus infections and viral haemorrhagic fevers in Uganda: A serological survey in Karamoja district, 1984. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 83(6), 851–854.

Rojas, D. P., Barrera-Fuentes, G. A., Pavia-Ruz, N., Salgado-Rodriguez, M., Chendendoza, A., Manrique-Saide, P., Vazquez-Prokopec, G. M., Halloran, M. E., Longini, I. M., & Gomez-Dantes, H. (2018). Epidemiology of dengue and other arboviruses in a cohort of school children and their families in Yucatan, Mexico: Baseline and first year follow-up. *PLOS Neglected Tropical Diseases*, 12(11), e0006847.

Sánchez-Carbonel, J., Tantaléan-Yépez, D., Aguilar-Luis, M. A., Silva-Caso, W., Weilg, P., Vásquez-Achaya, F., Costa, L., Martins-Luna, J., Sandoval, I., & del Valle-Mendoza, J. (2018). Identification of infection by Chikungunya, Zika, and Dengue in an area of the Peruvian coast. Molecular diagnosis and clinical characteristics. *BMC Research Notes*, 11(1), 175.

Santana, V. dos S., Lavezzo, L. C., Mondini, A., Terzian, A. C. B., Bronzoni, R. V. de M., Rossit, A. R. B., Machado, R. L. D., Rahal, P., Nogueira, M. C. L., & Nogueira, M. L. (2010). Concurrent dengue and malaria in the Amazon region. *Revista Da Sociedade Brasileira de Medicina Tropical*, 43(5), 508–511.

Schwarz, N. G., Girmann, M., Randriamampionona, N., Bialonski, A., Maus, D., Krefis, A. C., Njarasoa, C., Rajanalison, J. F., Ramandrisoa, H. D., Randriarison, M. L., May, J., Schmidt-Chanasit, J., & Rakotozandrindrainy, R. (2012). Seroprevalence

of antibodies against Chikungunya, Dengue, and Rift Valley fever viruses after febrile illness outbreak, Madagascar. *Emerging infectious diseases*, 18(11), 1780–1786.

Smithburn, K. C. (1952). Neutralizing antibodies against certain recently isolated viruses in the sera of human beings residing in East Africa. *Journal of immunology* (Baltimore, Md. : 1950), 69(2), 223–234.

Smithburn, K. C., Taylor, R. M., Rizk, F., & Kader, A. (1954). Immunity to certain arthropod-borne viruses among indigenous residents of Egypt. *The American Journal of Tropical Medicine and Hygiene*, 3(1), 9–18.

Smithburn, K. C., Kerr, J. A., & Gatne, P. B. (1954). Neutralizing antibodies against certain viruses in the sera of residents of India. *Journal of immunology* (Baltimore, Md. : 1950), 72(4), 248–257.

Soghaier, M. A., Abdelgadir, D. M., Abdelkhalig, S. M., Kafi, H., Zarroug, I. M. A., Sall, A. A., Eldegai, M. H., Elageb, R. M., Osman, M. M., & Khogali, H. (2018). Evidence of pre-existing active Zika virus circulation in Sudan prior to 2012. *BMC Research Notes*, 11(1), 906.

Stewart-Ibarra, A. M., Ryan, S. J., Kenneson, A., King, C. A., Abbott, M., Barbachano-Guerrero, A., Beltrán-Ayala, E., Borbor-Cordova, M. J., Cárdenas, W. B., Cueva, C., Finkelstein, J. L., Lupone, C. D., Jarman, R. G., Maljkovic Berry, I., Mehta, S., Polhemus, M., Silva, M., & Endy, T. P. (2018). The Burden of Dengue Fever and Chikungunya in Southern Coastal Ecuador: *Epidemiology, Clinical*

Presentation, and Phylogenetics from the First Two Years of a Prospective Study. *The American Journal of Tropical Medicine and Hygiene*, 98(5), 1444–1459.

Stoddard, S. T., Wearing, H. J., Jr, R. C. R., Morrison, A. C., Astete, H., Vilcarrromero, S., Alvarez, C., Ramal-Asayag, C., Sihuinchu, M., Rocha, C., Halsey, E. S., Scott, T. W., Kochel, T. J., & Forshey, B. M. (2014). Long-Term and Seasonal Dynamics of Dengue in Iquitos, Peru. *PLOS Neglected Tropical Diseases*, 8(7), e3003.

Sutherland, L. J., Cash, A. A., Huang, Y.-J. S., Sang, R. C., Malhotra, I., Moormann, A. M., King, C. L., Weaver, S. C., King, C. H., & LaBeaud, A. D. (2011). Serologic Evidence of Arboviral Infections among Humans in Kenya. *The American Journal of Tropical Medicine and Hygiene*, 85(1), 158–161.

Thaung, U., Ming, C. K., Swe, T. H. A. N., & Thein, S. (1975). Epidemiological features of dengue and chikungunya infections in Burma. *Southeast Asian J Trop Med Public Health*, 6(2), 276-83.

Theamboonlers, A., Rianthavorn, P., Praianantathavorn, K., Wuttirattanakowit, N., & Poovorawan, Y. (2009). Clinical and molecular characterization of chikungunya virus in South Thailand. *Jpn J Infect Dis*, 62(4), 303-305

Ushijima, Y., Abe, H., Nguema Ondo, G., Bikangui, R., Massinga Loembé, M., Zadeh, V. R., Essimengane, J. G. E., Mbouna, A. V. N., Bache, E. B., Agnandji, S. T., Lell, B., & Yasuda, J. (2021). Surveillance of the major pathogenic arboviruses of public health concern in Gabon, Central Africa: Increased risk of West Nile virus and dengue virus infections. *BMC Infectious Diseases*, 21(1), 265.



- Van Den Hurk, A. F., McElroy, K., Pyke, A. T., McGee, C. E., Hall-Mendelin, S., Day, A., ... & Higgs, S. (2011). Vector competence of Australian mosquitoes for yellow fever virus. *The American journal of tropical medicine and hygiene*, 85(3), 446.
- Vega-Rúa, A., Zouache, K., Girod, R., Failloux, A.-B., & Lourenço-de-Oliveira, R. (2014). High level of vector competence of *Aedes aegypti* and *Aedes albopictus* from ten American countries as a crucial factor in the spread of Chikungunya virus. *Journal of Virology*, 88(11), 6294–6306.
- Villarroel, P. M. S., Nurtop, E., Pastorino, B., Roca, Y., Drexler, J. F., Gallian, P., Jaenisch, T., Leparc-Goffart, I., Priet, S., Ninove, L., & Lamballerie, X. de. (2018). Zika virus epidemiology in Bolivia: A seroprevalence study in volunteer blood donors. *PLOS Neglected Tropical Diseases*, 12(3), e0006239.
- Watts, D. M., el-Tigani, A., Botros, B. A., Salib, A. W., Olson, J. G., McCarthy, M., & Ksiazek, T. G. (1994). Arthropod-borne viral infections associated with a fever outbreak in the northern province of Sudan. *The Journal of Tropical Medicine and Hygiene*, 97(4), 228–230.
- Wong, P.-S. J., Li, M. I., Chong, C.-S., Ng, L.-C., & Tan, C.-H. (2013). *Aedes* (*Stegomyia*) *albopictus* (Skuse): A potential vector of Zika virus in Singapore. *PLoS Neglected Tropical Diseases*, 7(8), e2348.
- Yamashiro, T., Disla, M., Petit, A., Taveras, D., Castro-Bello, M., Lora-Orste, M., ... & Nishizono, A. (2004). Seroprevalence of IgG specific for dengue virus among

adults and children in Santo Domingo, Dominican Republic. *The American journal of tropical medicine and hygiene*.

Yergolkar, P. N., Tandale, B. V., Arankalle, V. A., Sathe, P. S., Gandhe, S. S., Gokhle, M. D., Jacob, G. P., Hundekar, S. L., & Mishra, A. C. (2006). Chikungunya Outbreaks Caused by African Genotype, India. *Emerging Infectious Diseases*, 12(10), 1580–1583.

Yoosuf AA, Shiham I, Mohamed AJ, Ali G, Luna JM, Pandav R, Gongal GN, Nisaluk A, Jarman RG, Gibbons RV. (2009). First report of chikungunya from the Maldives. *Trans R Soc Trop Med Hyg.*;103(2):192-6.

Zaidi, M. B., Cedillo-Barron, L., González y Almeida, M. E., Garcia-Cordero, J., Campos, F. D., Namorado-Tonix, K., & Perez, F. (2020). Serological tests reveal significant cross-reactive human antibody responses to Zika and Dengue viruses in the Mexican population. *Acta Tropica*, 201, 105201.

## Bibliography

- Ali, R., Azmi, R. A., Wasi Ahmad, N., Abd Hadi, A., Muhamed, K. A., Rasli, R., Yoon Ling, C., Anak Chua, H., Lian Wan, K., & Lee, H. L. (2020). Entomological Surveillance Associated with Human Zika Cases in Miri Sarawak, Malaysia. *The American Journal of Tropical Medicine and Hygiene*, 102(5), 964–970.
- Aliota, M. T., Peinado, S. A., Velez, I. D., & Osorio, J. E. (2016). The wMel strain of *Wolbachia* Reduces Transmission of Zika virus by *Aedes aegypti*. *Scientific Reports*, 6(1), 28792.
- Alva-Urcia, C., Aguilar-Luis, M. A., Palomares-Reyes, C., Silva-Caso, W., Suarez-Ognio, L., Weigl, P., Manrique, C., Vasquez-Achaya, F., Valle, L. J. del, & Valle-Mendoza, J. del. (2017). Emerging and reemerging arboviruses: A new threat in Eastern Peru. *PLOS ONE*, 12(11), e0187897.
- Arduino, M. de B., Mucci, L. F., dos Santos, L. M., & Soares, M. F. de S. (n.d.). Importance of microenvironment to arbovirus vector distribution in an urban area, São Paulo, Brazil. *Revista Da Sociedade Brasileira de Medicina Tropical*, 53.
- Ayres, C.F., Seixas, G., Borrego, S., Marques, C., Monteiro, I., Marques, C.S., Gouveia, B., Leal, S., Troco, A.D., Fortes, F., 2020. The V410L knockdown resistance mutation occurs in island and continental populations of *Aedes aegypti* in West and Central Africa. *PLOS Neglected Tropical Diseases* 14, e0008216.
- Azar, S. R., & Weaver, S. C. (2019). Vector Competence: What Has Zika Virus Taught Us? *Viruses*, 11(9), 867.
- Baba, M., Logue, C. H., Oderinde, B., Abdulmaleek, H., Williams, J., Lewis, J., Laws, T. R., Hewson, R., Marcello, A., & Agaro, P. D. (2013). Evidence of arbovirus co-infection in suspected febrile malaria and typhoid patients in Nigeria. *The Journal of Infection in Developing Countries*, 7(01), 051–059.
- Badolo, A., Sombié, A., Pignatelli, P.M., Sanon, A., Yaméogo, F., Wangrawa, D.W., Sanon, A., Kanuka, H., McCall, P.J., Weetman, D., 2019. Insecticide resistance levels and mechanisms in *Aedes aegypti* populations in and around Ouagadougou, Burkina Faso. *PLoS neglected tropical diseases* 13, e0007439.
- Badolo, A., Sombié, A., Yaméogo, F., Wangrawa, D. W., Sanon, A., Pignatelli, P. M., Sanon, A., Viana, M., Kanuka, H., Weetman, D., & McCall, P. J. (2022). First comprehensive analysis of *Aedes aegypti* bionomics during an arbovirus outbreak in west Africa: Dengue in Ouagadougou, Burkina Faso, 2016–2017. *PLOS Neglected Tropical Diseases*, 16(7), e0010059.

- Bagny Beilhe, L., S. Arnoux, H. Delatte, et al. (2012). Spread of invasive *Aedes albopictus* and decline of resident *Aedes aegypti* in urban areas of Mayotte 2007–2010. *Biol. Invasions* 14: 1623–1633.
- Baldacchino, F., Marcantonio, M., Manica, M., Marini, G., Zorer, R., Delucchi, L., Arnoldi, D., Montarsi, F., Capelli, G., Rizzoli, A., & Rosà, R. (2017). Mapping of *Aedes albopictus* Abundance at a Local Scale in Italy. *Remote Sensing*, 9(7), 749.
- Barker, C.M., S.L. Paulson, S. Cantrell, et al. (2003). Habitat preferences and phenology of *Ochlerotatus triseriatus* and *Aedes albopictus* (Diptera: Culicidae) in southwestern Virginia. *J. Med. Entomol.* 40: 403–410.
- Ben-Shachar R, Schmidler S, Koelle K. (2016). Drivers of Inter-individual Variation in Dengue Viral Load Dynamics. *PLoS Comput Biol* 12(11): e1005194.
- Bhargavi, B. S., & Moa, A. (2020). Global outbreaks of zika infection by epidemic observatory (EpiWATCH), 2016-2019. *Global Biosecurity*, 2(1), Article 1.
- Bhatt, S., Gething, P.W., Brady, O.J., Messina, J.P., Farlow, A.W., Moyes, C.L., Drake, J.M., Brownstein, J.S., Hoen, A.G., Sankoh, O., 2013. The global distribution and burden of dengue. *Nature* 496, 504-507.
- Bosa, H. K., Montgomery, J. M., Kimuli, I., & Lutwama, J. J. (2014). Dengue fever outbreak in Bosio, C. F., B. J. Beaty, and W. C. Black IV. 1998. Quantitative genetics of vector competence for dengue-2 virus in *Aedes aegypti*. *Am. J. Trop. Med. Hyg.* 59: 965-970.
- Bosio, C. F., R. E. Fulton, M. L. Salasek, B. J. Beaty, and W. C. Black IV. 2000. Quantitative trait loci that control vector competence for dengue- 2 virus in the mosquito *Aedes aegypti*. *Genetics* 156: 687-698.
- Bradley, C. A., & Altizer, S. (2007). Urbanization and the ecology of wildlife diseases. *Trends in Ecology & Evolution*, 22(2), 95–102.
- Brady, O. J., Johansson, M. A., Guerra, C. A., Bhatt, S., Golding, N., Pigott, D. M., Delatte, H., Grech, M. G., Leisnham, P. T., Maciel-de-Freitas, R., Styer, L. M., Smith, D. L., Scott, T. W., Gething, P. W., & Hay, S. I. (2013). Modelling adult *Aedes aegypti* and *Aedes albopictus* survival at different temperatures in laboratory and field settings. *Parasites & Vectors*, 6(1), 351.
- Braks, M. a. H., Honório, N. A., Lounibos, L. P., Lourenço-De-Oliveira, R., & Juliano, S. A. (2004). Interspecific Competition Between Two Invasive Species of Container Mosquitoes, *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae), in Brazil. *Annals of the Entomological Society of America*, 97(1), 130–139.

- Calle-Tobón, A., Pérez-Pérez, J., Rojo, R., Rojas-Montoya, W., Triana-Chavez, O., Rúa-Uribe, G., & Gómez-Palacio, A. (2020). Surveillance of Zika virus in field-caught *Aedes aegypti* and *Aedes albopictus* suggests important role of male mosquitoes in viral populations maintenance in Medellín, Colombia. *Infection, Genetics and Evolution*, *85*, 104434.
- Campos, M., Ward, D., Morales, R. F., Gomes, A. R., Silva, K., Sepúlveda, N., Gomez, L. F., Clark, T. G., & Campino, S. (2020). Surveillance of *Aedes aegypti* populations in the city of Praia, Cape Verde: Zika virus infection, insecticide resistance and genetic diversity. *Parasites & Vectors*, *13*(1), 481.
- Carbajo, A. E., Curto, S. I., & Schweigmann, N. J. (2006). Spatial distribution pattern of oviposition in the mosquito *Aedes aegypti* in relation to urbanization in Buenos Aires: Southern fringe bionomics of an introduced vector. *Medical and Veterinary Entomology*, *20*(2), 209–218.
- Cebrián-Camisón, S., Martínez-de la Puente, J., & Figuerola, J. (2020). A Literature Review of Host Feeding Patterns of Invasive *Aedes* Mosquitoes in Europe. *Insects*, *11*(12), 848.
- Cevallos, V., Ponce, P., Waggoner, J. J., Pinsky, B. A., Coloma, J., Quiroga, C., Morales, D., & Cárdenas, M. J. (2018). Zika and Chikungunya virus detection in naturally infected *Aedes aegypti* in Ecuador. *Acta Tropica*, *177*, 74–80.
- Chan, M., & Johansson, M. A. (2012). The Incubation Periods of Dengue Viruses. *PLoS ONE*, *7*(11), e50972.
- Chapman, S., Watson, J., Salazar, A., Thatcher, M., & Mcalpine, C. (2017). The impact of urbanization and climate change on urban temperatures: A systematic review. *Landscape Ecology*, *32*.
- Chen, S., Huang, Q., Muttarak, R., Fang, J., Liu, T., He, C., Liu, Z., & Zhu, L. (2022). Updating global urbanization projections under the Shared Socioeconomic Pathways. *Scientific Data*, *9*(1).
- Conceicao, T.M., Da Poian, A.T., Sorgine, M.H., 2010. A real-time PCR procedure for detection of dengue virus serotypes 1, 2, and 3, and their quantitation in clinical and laboratory samples. *Journal of virological methods* *163*, 1-9.
- Correa-Morales, F., González-Acosta, C., Mejía-Zúñiga, D., Huerta, H., Pérez-Rentería, C., Vazquez-Pichardo, M., Ortega-Morales, A. I., Hernández-Triana, L. M., Salazar-Bueyes, V. M., & Moreno-García, M. (2019). Surveillance for Zika in Mexico: Naturally infected mosquitoes in urban and semi-urban areas. *Pathogens and Global Health*, *113*(7), 309–314.

- Costa, E. Santos, de M., Correia, J. et al. (2010). Impact of small variations in temperature and humidity on the reproductive activity and survival of *Aedes aegypti* (Diptera, Culicidae). *Revista Brasileira de Entomologia* 54, 488–493.
- Costanzo, K. S., Kesavaraju, B., & Juliano, S. A. (2005). Condition-Specific Competition in Container Mosquitoes: The Role of Noncompeting Life-History Stages. *Ecology*, 86(12), 3289–3295.
- Cox, J., Grillet, M.E., Ramos, O.M., Amador, M. & Barrera, R. (2007) Habitat segregation of dengue vectors along an urban environmental gradient. *American Journal of Tropical Medicine and Hygiene*, 76, 820–826.
- Dalpadado, C. P. R. D., & Amarasinghe, L. D. (2018). *Abundance and distribution pattern of Aedes aegypti and Aedes albopictus in selected urban, sub-urban and rural areas of Gampaha District, Sri Lanka*. Proceedings of the 3rd International Research Symposium on Pure and Applied Sciences, 26th October 2018 - Faculty of Science, University of Kelaniya, Sri Lanka
- Das, M., Dutta, P., 2014. Status of insecticide resistance and detoxifying enzyme activity of *Aedes albopictus* population in Sonitpur district of Assam, India. *International Journal of Mosquito Research* 1, 35-41.
- Deilami, K., Kamruzzaman, Md., & Liu, Y. (2018). Urban heat island effect: A systematic review of spatio-temporal factors, data, methods, and mitigation measures. *International Journal of Applied Earth Observation and Geoinformation*, 67, 30–42.
- Delatte, H., Desvars, A., Bouétard, A., Bord, S., Gimonneau, G., Vourc'h, G., & Fontenille, D. (2010). Blood-Feeding Behavior of *Aedes albopictus*, a Vector of Chikungunya on La Réunion. *Vector-Borne and Zoonotic Diseases*, 10(3), 249–258.
- Delgado, F. G., Torres, K. I., Castellanos, J. E., Romero-Sánchez, C., Simon-Lorière, E., Sakuntabhai, A., & Roth, C. (2018). Improved Immune Responses Against Zika Virus After Sequential Dengue and Zika Virus Infection in Humans. *Viruses*, 10(9), 480.
- Djiappi-Tchamen, B., Nana-Ndjangwo, M.S., Mavridis, K., Talipouo, A., Nchoutpouen, E., Makoudjou, I., Bamou, R., Mayi, A.M.P., Awono-Ambene, P., Tchuinkam, T., 2021. Analyses of Insecticide Resistance Genes in *Aedes aegypti* and *Aedes albopictus* Mosquito Populations from Cameroon. *Genes* 12, 828.
- Edwards, J., (1941). *Mosquitoes of the Ethiopian region*. Oxford University Press, 513 p.
- Egid, B.R., Coulibaly, M., Dadzie, S.K., Kamgang, B., McCall, P.J., Sedda, L., Toe, H.K., Wilson, A.L., 2021. Review of the ecology and behaviour of *Aedes aegypti* and

*Aedes albopictus* in Western Africa and implications for vector control. *Current Research in Parasitology & Vector-Borne Diseases*, 100074.

Estallo, E. L., Sangermano, F., Grech, M., Ludueña-Almeida, F., Frías-Céspedes, M., Ainete, M., Almirón, W., & Livdahl, T. (2018). Modelling the distribution of the vector *Aedes aegypti* in a central Argentine city. *Medical and Veterinary Entomology*, 32(4), 451–461.

European Centre for Disease Prevention and Control (ECDC) (2022). *Dengue worldwide overview*. Retrieved October 12, 2022, from <https://www.ecdc.europa.eu/en/dengue-monthly>

Faeth, S. H., Warren, P. S., Shochat, E., & Marussich, W. A. (2005). Trophic Dynamics in Urban Communities. *BioScience*, 55(5), 399–407.

Faraji, A., Egizi, A., Fonseca, D. M., Unlu, I., Crepeau, T., Healy, S. P., & Gaugler, R. (2014). Comparative host feeding patterns of the Asian tiger mosquito, *Aedes albopictus*, in urban and suburban Northeastern USA and implications for disease transmission. *PLoS neglected tropical diseases*, 8(8), e3037.

Fatima, S. H., Atif, S., Rasheed, S. B., Zaidi, F., & Hussain, E. (2016). Species Distribution Modelling of *Aedes aegypti* in two dengue-endemic regions of Pakistan. *Tropical Medicine & International Health*, 21(3), 427–436.

Ferguson, N. M., Cucunubá, Z. M., Dorigatti, I., Nedjati-Gilani, G. L., Donnelly, C. A., Basáñez, M. G., Nouvellet, P., & Lessler, J. (2016). EPIDEMIOLOGY. Countering the Zika epidemic in Latin America. *Science (New York, N.Y.)*, 353(6297), 353–354.

Fernandes-Monteiro, A. G., Trindade, G. F., Yamamura, A. M., Moreira, O. C., de Paula, V. S., Duarte, A. C., Britto, C., & Lima, S. M. (2015). New approaches for the standardization and validation of a real-time qPCR assay using TaqMan probes for quantification of yellow fever virus on clinical samples with high quality parameters. *Human vaccines & immunotherapeutics*, 11(7), 1865–1871.

Ferreira-de-Brito, A., Ribeiro, I. P., Miranda, R. M. de, Fernandes, R. S., Campos, S. S., Silva, K. A. B. da, Castro, M. G. de, Bonaldo, M. C., Brasil, P., & Lourenço-de-Oliveira, R. (2016). First detection of natural infection of *Aedes aegypti* with Zika virus in Brazil and throughout South America. *Memorias Do Instituto Oswaldo Cruz*, 111(10), 655–658.

Freire, M. S., Mann, G. F., Marchevsky, R. S., Yamamura, A. M. Y., Almeida, L. F. C., Jabor, A. V., Malachias, J. M. N., Coutinho, E. S. F., & Galler, R. (2005). Production of yellow fever 17DD vaccine virus in primary culture of chicken embryo

- fibroblasts: Yields, thermo and genetic stability, attenuation and immunogenicity. *Vaccine*, 23(19), 2501–2512.
- Gake, B., Vernet, M. A., Leparc-Goffart, I., Drexler, J. F., Gould, E. A., Gallian, P., & Lamballerie, X. de. (2017). Low seroprevalence of Zika virus in Cameroonian blood donors. *Brazilian Journal of Infectious Diseases*, 21, 481–483.
- Gervasi, S. S., Civitello, D. J., Kilvitis, H. J., & Martin, L. B. (2015). The context of host competence: A role for plasticity in host–parasite dynamics. *Trends in Parasitology*, 31(9), 419–425.
- Gonçalves, C.M., Melo, F.F., Bezerra, J.M. et al. (2014), Distinct variation in vector competence among nine field populations of *Aedes aegypti* from a Brazilian dengue-endemic risk city. *Parasites Vectors* 7, 320.
- Gottdenker, N. L., Streicker, D. G., Faust, C. L., & Carroll, C. R. (2014). Anthropogenic land use change and infectious diseases: A review of the evidence. *EcoHealth*, 11(4), 619–632.
- Grard, G., Caron, M., Mombo, I.M., Nkoghe, D., Mboui Ondo, S., Jiolle, D., Fontenille, D., Paupy, C., Leroy, E.M., 2014. Zika virus in Gabon (Central Africa)-2007: a new threat from *Aedes albopictus*? *PLOS Neglected Tropical Diseases* 8, 1-6.
- Grimm, N. B., S. H. Faeth, N. E. Golubiewski, C. L. Redman, J. G. Wu, X. M. Bai, and J. M. Briggs. 2008. Global change and the ecology of cities. *Science* 319:756–760.
- Gubler, D. J. (2011). Dengue, Urbanization and Globalization: The Unholy Trinity of the 21st Century. *Tropical Medicine and Health*, 39(4 Suppl), 3–11.
- Gubler, D. J., S. Nalim, R. Tau, H. Saipen, and J. Sulianti Saroso. 1979. Variation in susceptibility to oral infection with dengue viruses among geographic strains of *Aedes aegypti*. *Am. J. Trop. Med. Hyg.* 28: 1045-1052.
- Guerbois, M., Fernandez-Salas, I., Azar, S. R., Danis-Lozano, R., Alpuche-Aranda, C. M., Leal, G., Garcia-Malo, I. R., Diaz-Gonzalez, E. E., Casas-Martinez, M., Rossi, S. L., Del Río-Galván, S. L., Sanchez-Casas, R. M., Roundy, C. M., Wood, T. G., Widen, S. G., Vasilakis, N., & Weaver, S. C. (2016). Outbreak of Zika Virus Infection, Chiapas State, Mexico, 2015, and First Confirmed Transmission by *Aedes aegypti* Mosquitoes in the Americas. *The Journal of Infectious Diseases*, 214(9), 1349–1356.
- Haddi, K., Tomé, H.V., Du, Y., Valbon, W.R., Nomura, Y., Martins, G.F., Dong, K., Oliveira, E.E., 2017. Detection of a new pyrethroid resistance mutation (V410L) in



the sodium channel of *Aedes aegypti*: a potential challenge for mosquito control. *Scientific reports* 7, 46549.

Hancock, P.A., Lynd, A., Wiebe, A., Devine, M., Essandoh, J., Wat'senga, F., Manzambi, E.Z., Agossa, F., Donnelly, M.J., Weetman, D., 2022. Modelling spatiotemporal trends in the frequency of genetic mutations conferring insecticide target-site resistance in African mosquito malaria vector species. *BMC biology* 20, 1-17.

Harwood, J.F., Helmey, W.L., Turnwall, B., Justice, K.D., Farooq M., Richardson, A.G. 2016. Controlling *Aedes aegypti* in Cryptic Environments with Manually Carried Ultra-Low Volume and Mist Blower Pesticide Applications. *Journal of the American Mosquito Control Association*. 32:217-223

Hassell, J. M., Begon, M., Ward, M. J., & Fèvre, E. M. (2017). Urbanization and Disease Emergence: Dynamics at the Wildlife-Livestock-Human Interface. *Trends in Ecology & Evolution*, 32(1), 55–67.

Hemingway, J., Hawkes, N.J., McCarroll, L., Ranson, H., 2004. The molecular basis of insecticide resistance in mosquitoes. *Insect biochemistry and molecular biology* 34, 653-665.

Higa, Y., N.T. Yen, H. Kawada, et al. (2010). Geographic distribution of *Aedes aegypti* and *Aedes albopictus* collected from used tires in Vietnam. *J. Am. Mosq. Control Assoc.* 26: 1–9.

Ho, Z. J. M., Hapuarachchi, H. C., Barkham, T., Chow, A., Ng, L. C., Lee, J. M. V., Leo, Y. S., Prem, K., Lim, Y. H. G., Sessions, P. F. de, Rabaa, M. A., Chong, C. S., Tan, C. H., Rajarethinam, J., Tan, J., Anderson, D. E., Ong, X., Cook, A. R., Chong, C. Y., ... Sessions, O. M. (2017). Outbreak of Zika virus infection in Singapore: An epidemiological, entomological, virological, and clinical analysis. *The Lancet Infectious Diseases*, 17(8), 813–821.

Hock, J.W., 2016. Improved CDC Backpack Aspirator—Model 1412 Instructions. John W. Hock company, 4 p.

Honorio, N., M. Castro, F. de Barros, et al. (2009). The spatial distribution of *Aedes aegypti* and *Aedes albopictus* in a transition zone, Rio de Janeiro, Brazil. *Cad. Saude Publica* ' 25: 1203–1214.

Hornby, J.A., D.E. Moore & T.W. Miller. (1994). *Aedes albopictus* distribution, abundance, and colonization in Lee County, Florida, and its effect on *Aedes aegypti*. *J. Am. Mosq. Control Assoc.* 10: 397–402.

Imrie, A., Meeks, J., Gurary, A., et al. (2007). Antibody to Dengue 1 Detected More Than 60 Years after Infection. *Viral Immunology*, 20(4), 672–675.

Integrated *Aedes aegypti* Mosquito Surveillance as Early Warning System for Arbovirus Transmission, Brazil. *Emerg Infect Dis.* 2022 Apr;28(4):701-706.

Ishak, I.H., Jaal, Z., Ranson, H., Wondji, C.S., 2015. Contrasting patterns of insecticide resistance and knockdown resistance (kdr) in the dengue vectors *Aedes aegypti* and *Aedes albopictus* from Malaysia. *Parasit Vectors* 8, 181.

Ishak, I.H., Kamgang, B., Ibrahim, S.S., Riveron, J.M., Irving, H., Wondji, C.S., 2017. Pyrethroid resistance in Malaysian populations of dengue vector *Aedes aegypti* is mediated by CYP9 family of cytochrome P450 genes. *PLOS Neglected Tropical Diseases* 11, e0005302.

Ishak, I.H., Riveron, J.M., Ibrahim, S.S., Stott, R., Longbottom, J., Irving, H., Wondji, C.S., 2016. The Cytochrome P450 gene CYP6P12 confers pyrethroid resistance in kdr-free Malaysian populations of the dengue vector *Aedes albopictus*. *Scientific reports* 6, 24707.

Juarez, J. G., Garcia-Luna, S., Chaves, L. F., Carbajal, E., Valdez, E., Avila, C., Tang, W., Martin, E., Barrera, R., Hemme, R. R., Mutebi, J.-P., Vuong, N., Roark, E. B., Maupin, C. R., Badillo-Vargas, I. E., & Hamer, G. L. (2020). Dispersal of female and male *Aedes aegypti* from discarded container habitats using a stable isotope mark-capture study design in South Texas. *Scientific Reports*, 10(1), 6803.

Jupp, P.G., 1996. Mosquitoes of southern Africa: Culicinae and Toxorhynchitinae. Ekogilde Publishers, Hartebeespoort, 156 pp. p.

Kamgang, B., Happi, J. Y., Boisier, P., Njiokou, F., Hervé, J.-P., Simard, F., & Paupy, C. (2010). Geographic and ecological distribution of the dengue and chikungunya virus vectors *Aedes aegypti* and *Aedes albopictus* in three major Cameroonian towns. *Medical and Veterinary Entomology*, 24(2), 132–141.

Kamgang, B., Nchoutpouen, E., Simard, F., & Paupy, C. (2012). Notes on the blood-feeding behavior of *Aedes albopictus* (Diptera: Culicidae) in Cameroon. *Parasites & Vectors*, 5(1), 57.

Kamgang, B., Yougang, A.P., Tchoupo, M., Riveron, J.M., Wondji, C., 2017. Temporal distribution and insecticide resistance profile of two major arbovirus vectors *Aedes aegypti* and *Aedes albopictus* in Yaoundé, the capital city of Cameroon. *Parasites & vectors* 10, 469.

Kamgang B, Vazeille M, Tedjou A, et al (2020). “Different Populations of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) from Central Africa Are Susceptible to Zika Virus Infection.” PLOS Neglected Tropical Diseases 14, no. 3 (March 23, 2020).

Kamgang, B., Wilson-Bahun, T.A., Yougang, A.P., Lenga, A., Wondji, C.S., 2020. Contrasting resistance patterns to type I and II pyrethroids in two major arbovirus vectors *Aedes aegypti* and *Aedes albopictus* in the Republic of the Congo, Central Africa. *Infectious Diseases of Poverty* 9, 1-10.

Kasai, S., Komagata, O., Itokawa, K., Shono, T., Ng, L.C., Kobayashi, M., Tomita, T., 2014. Mechanisms of pyrethroid resistance in the dengue mosquito vector, *Aedes aegypti*: target site insensitivity, penetration, and metabolism. *PLoS Negl Trop Dis* 8, e2948.

Kasai, S., Ng, L.C., Lam-Phua, S.G., Tang, C.S., Itokawa, K., Komagata, O., Kobayashi, M., Tomita, T., 2011. First detection of a putative knockdown resistance gene in major mosquito vector, *Aedes albopictus*. *Japanese journal of infectious diseases* 64, 217-221.

Kawada, H., Higa, Y., Futami, K., Muranami, Y., Kawashima, E., Osei, J.H., Sakyi, K.Y., Dadzie, S., de Souza, D.K., Appawu, M., 2016. Discovery of point mutations in the voltage-gated sodium channel from African *Aedes aegypti* populations: potential phylogenetic reasons for gene introgression. *PLoS Negl Trop Dis* 10, e0004780.

Kilpatrick, A. M., LaDeau, S. L., & Marra, P. P. (2007). Ecology Of West Nile Virus Transmission and its Impact on Birds in the Western Hemisphere. *The Auk*, 124(4), 1121–1136.

Kilpatrick, A. M., & Randolph, S. E. (2012). Drivers, dynamics, and control of emerging vector-borne zoonotic diseases. *Lancet*, 380(9857), 1946–1955.

Konan, L.Y., Oumbouke, W.A., Silué, U.G., Coulibaly, I.Z., Ziogba, J.-C.T., N’Guessan, R.K., Coulibaly, D., Béné, J.B.V., Lenhart, A., 2021. Insecticide Resistance Patterns and Mechanisms in *Aedes aegypti* (Diptera: Culicidae) Populations Across Abidjan, Côte d’Ivoire Reveal Emergent Pyrethroid Resistance. *Journal of Medical Entomology*.

Kosmidis I (2017). *brglm2*: Bias reduction in generalized linear models. R package version 0.1.5.

Kosoltanapiwat, N., Tongshoob, J., Singkhaimuk, P., Nitatsukprasert, C., Davidson, S. A., & Ponlawat, A. (2020). Entomological Surveillance for Zika and Dengue Virus in *Aedes* Mosquitoes: Implications for Vector Control in Thailand. *Pathogens*, 9(6), 442.

- Kraemer, L., Moritz, U., Sinka, M., et al. (2015). The Global Distribution of the Arbovirus Vectors *Aedes Aegypti* and *Ae. Albopictus*.” Edited by Mark Jit. *ELife* 4.
- Kraemer, M. U. G., Reiner, R. C., Brady, O. J. et al. (2019). Past and future spread of the arbovirus vectors *Aedes aegypti* and *Aedes albopictus*. *Nature Microbiology*, 4(5), 854–863.
- Kudom, A.A., 2020. Entomological surveillance to assess potential outbreak of Aedes-borne arboviruses and insecticide resistance status of *Aedes aegypti* from Cape Coast, Ghana. *Acta tropica* 202, 105257.
- Kuniholm M, Wolfe D, Huang C, et al. (2006). Seroprevalence and distribution of Flaviviridae, Togaviridae, and Bunyaviridae arboviral infections in rural Cameroonian adults. *Am J Trop Med Hyg* 74: 1078–1083.
- Kuno, G. (2020). The Absence of Yellow Fever in Asia: History, Hypotheses, Vector Dispersal, Possibility of YF in Asia, and Other Enigmas. *Viruses*, 12(12), Article 12.
- Lambrechts, L., Paaijmans, K. P., Fansiri, T., et al. (2011). Impact of daily temperature fluctuations on dengue virus transmission by *Aedes aegypti*. *Proceedings of the National Academy of Sciences of the United States of America*, 108(18), 7460–7465.
- Lanciotti, R.S., Kosoy, O.L., Laven, J.J., Velez, J.O., Lambert, A.J., Johnson, A.J., Stanfield, S.M., Duffy, M.R. (2008). Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerging Infectious Diseases* 14, 1232-1239.
- Leandro AS, de Castro WAC, Lopes RD, Delai RM, Villela DAM, de-Freitas RM. Citywide
- Lenth, R. (2022). emmeans: Estimated Marginal Means, aka Least-Squares Means\_. R package version 1.8.2.
- Leparc-Goffart, I., Baragatti, M., Temmam, S., Tuiskunen, A., Moureau, G., Charrel, R., De Lamballerie, X., 2009. Development and validation of real-time one-step reverse transcription-PCR for the detection and typing of dengue viruses. *Journal of Clinical Virology* 45, 61-66.
- Li, Y., Kamara, F., Zhou, G., Puthiyakunnon, S. et al. (2014). Urbanization Increases *Aedes albopictus* Larval Habitats and Accelerates Mosquito Development and Survivorship. *PLOS Neglected Tropical Diseases* 8.

Li Y, Dou Q, Lu Y, Xiang H, Yu X, Liu S. (2020) Effects of ambient temperature and precipitation on the risk of dengue fever: A systematic review and updated meta-analysis. *Environ Res*; 191:110043.

Lifetechnologies, 2011. RNAlater® Tissue Collection: RNA Stabilization Solution. Applied Biosystems, 12 p.

Livak, K.J., 1984. Organization and mapping of a sequence on the *Drosophila melanogaster* X and Y chromosomes that is transcribed during spermatogenesis. *Genetics* 107, 611-634.

Macoris Mde, L., Andrighetti, M.T., Wanderley, D.M., Ribolla, P.E., 2014. Impact of insecticide resistance on the field control of *Aedes aegypti* in the State of Sao Paulo. *Rev Soc Bras Med Trop* 47, 573-578.

Magalhaes, T., Robison, A., Young, M. C., Black, W. C., Foy, B. D., Ebel, G. D., & Rückert, C. (2018). Sequential Infection of *Aedes aegypti* Mosquitoes with Chikungunya Virus and Zika Virus Enhances Early Zika Virus Transmission. *Insects*, 9(4), 177.

Mandeng, S.E., Awono-Ambene, H.P., Bigoga, J.D., Ekoko, W.E., Binyang, J., Piamou, M., Mbakop, L.R., Fesuh, B.N., Mvondo, N., Tabue, R., 2019. Spatial and temporal development of deltamethrin resistance in malaria vectors of the *Anopheles gambiae* complex from North Cameroon. *PloS one* 14, e0212024.

Marchette NJ, Garcia R, Rudnick A. Isolation of Zika virus from *Aedes aegypti* mosquitoes in Malaysia. *Am J Trop Med Hyg.* 1969 May;18(3):411-5.

Marcombe, S., Fustec, B., Cattel, J., Chonephetsarath, S., Thammavong, P., Phommavanh, N., David, J.-P., Corbel, V., Sutherland, I.W., Hertz, J.C., 2019. Distribution of insecticide resistance and mechanisms involved in the arbovirus vector *Aedes aegypti* in Laos and implication for vector control. *PLoS Neglected Tropical Diseases* 13.

McClelland, G. a. H. (1959). Observations on the Mosquito, *Aedes* (*Stegomyia*) *aegypti* (L.), in East Africa. I.—The Biting Cycle in an Outdoor Population at Entebbe, Uganda. *Bulletin of Entomological Research*, 50(2), 227–235.

McCluney, K. E., Burdine, J. D., & Frank, S. D. (2017). Variation in arthropod hydration across US cities with distinct climate. *Journal of Urban Ecology*, 3.

McClure, K. M., Lawrence, C., & Kilpatrick, A. M. (2018). Land Use and Larval Habitat Increase *Aedes albopictus* (Diptera: Culicidae) and *Culex quinquefasciatus* (Diptera: Culicidae) Abundance in Lowland Hawaii. *Journal of Medical Entomology*, 55(6), 1509–1516.

McElreath R (2016). rethinking: Statistical Rethinking Course and Book Package. R package version 1.59, URL <https://github.com/rmcelreath/rethinking>.

McKinney, M. L. (2006) 'Urbanization as a Major Cause of Biotic Homogenization', *Biological Conservation*, 127: 247–60

McLean DM, Clarke AM, Coleman JC, Montalbetti CA, Skidmore AG, et al. (1974) Vector capability of *Aedes aegypti* mosquitoes for California encephalitis and dengue viruses at various temperatures. *Can J Microbiol* 20: 255–262.

Medeiros, M.C.I., Boothe E.C., Roark E.B., Hamer G.L. (2017). Dispersal of male and female *Culex quinquefasciatus* and *Aedes albopictus* mosquitoes using stable isotope enrichment. *PLoS Negl Trop Dis* 11(1).

Mordecai, E. A., Paaijmans, K. P., Johnson, L. R., Balzer, C., Ben-Horin, T., de Moor, E., McNally, A., Pawar, S., Ryan, S. J., Smith, T. C., & Lafferty, K. D. (2013). Optimal temperature for malaria transmission is dramatically lower than previously predicted. *Ecology Letters*, 16(1), 22–30.

Mordecai, E. A., Cohen, J. M., Evans, M. V., Gudapati, P., Johnson, L. R., Lippi, C. A., Miazgowicz, K., Murdock, C. C., Rohr, J. R., Ryan, S. J., Savage, V., Shocket, M. S., Ibarra, A. S., Thomas, M. B., & Weikel, D. P. (2017). Detecting the impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic models. *PLOS Neglected Tropical Diseases*, 11(4), e0005568.

Moyes, C.L., Vontas, J., Martins, A.J., Ng, L.C., Koou, S.Y., Dusfour, I., Raghavendra, K., Pinto, J., Corbel, V., David, J.-P., 2017. Contemporary status of insecticide resistance in the major *Aedes* vectors of arboviruses infecting humans. *PLOS Neglected Tropical Diseases* 11, e0005625.

Ndenga, B. A., Mutuku, F. M., Ngugi, H. N., Mbakaya, J. O., Aswani, P., Musunzaji, P. S., Vulule, J., Mukoko, D., Kitron, U., & LaBeaud, A. D. (2017). Characteristics of *Aedes aegypti* adult mosquitoes in rural and urban areas of western and coastal Kenya. *PLOS ONE*, 12(12), e0189971.

Ngoagouni, C., Kamgang, B., Kazanji, M., Paupy, C., Nakoune, E., 2017. Potential of *Aedes aegypti* and *Aedes albopictus* populations in the Central African Republic to transmit enzootic chikungunya virus strains. *Parasites and Vectors* 10, 1-5.

Nguyen, N. M., Kien, D. T. H., Tuan, T. V., Quyen, N. T. H., et al. (2013). Host and viral features of human dengue cases shape the population of infected and infectious *Aedes aegypti* mosquitoes. *Proceedings of the National Academy of Sciences*, 110(22), 9072–9077.

- Obenauer, J. F., Andrew Joyner, T., & Harris, J. B. (2017). The importance of human population characteristics in modeling *Aedes aegypti* distributions and assessing risk of mosquito-borne infectious diseases. *Tropical Medicine and Health*, 45(1), 38.
- OECD/SWAC (2020), Africa's Urbanisation Dynamics 2020: Africapolis, Mapping a New Urban Geography, West African Studies, OECD Publishing, Paris.
- Olivry, J.C., 1986. Fleuves et rivières du Cameroun. Vol 9, France Edition. ORSTOM, Paris, 781 p.
- Overgaard, H. J., Olano, V. A., Jaramillo, J. F., Matiz, M. I., Sarmiento, D., Stenström, T. A., & Alexander, N. (2017). A cross-sectional survey of *Aedes aegypti* immature abundance in urban and rural household containers in central Colombia. *Parasites & Vectors*, 10(1), 356.
- Pan American Health Organization (PAHO) (2022). Epidemiological Alert Yellow Fever 31 August 2022
- Parra, M. C. P., Lorenz, C., de Aguiar Milhim, B. H. G., Dibo, M. R., Guirado, M. M., Chiaravalloti-Neto, F., & Nogueira, M. L. (2022). Detection of Zika RNA virus in *Aedes aegypti* and *Aedes albopictus* mosquitoes, São Paulo, Brazil. *Infection, Genetics and Evolution*, 98, 105226.
- Pastorino, B., Bessaud, M., Grandadam, M., Murri, S., Tolou, H.J., Peyrefitte, C.N., 2005. Development of a TaqMan® RT-PCR assay without RNA extraction step for the detection and quantification of African Chikungunya viruses. *Journal of Virological Methods* 124, 65-71.
- Pereira-dos-Santos, T., Roiz, D., Lourenço-de-Oliveira, R., & Paupy, C. (2020). A Systematic Review: Is *Aedes albopictus* an Efficient Bridge Vector for Zoonotic Arboviruses? *Pathogens*, 9(4), 266.
- Perry, T., Batterham, P., Daborn, P.J., 2011. The biology of insecticidal activity and resistance. *Insect biochemistry and molecular biology* 41, 411-422.
- Phumee, A., Buathong, R., Boonserm, R., Intayot, P., Aungsananta, N., Jittmittraphap, A., Joyjinda, Y., Wacharapluesadee, S., & Siriyasatien, P. (2019). Molecular Epidemiology and Genetic Diversity of Zika Virus from Field-Caught Mosquitoes in Various Regions of Thailand. *Pathogens*, 8, 30.
- Plourde, A. R., & Bloch, E. M. (2016). A literature review of Zika virus. *Emerging infectious diseases*, 22(7), 1185.

- Possas, C., Brasil, P., Marzochi, M. C., Tanuri, A., Martins, R. M., Marques, E. T., Homma, A. (2017). Zika puzzle in Brazil: peculiar conditions of viral introduction and dissemination-A Review. *Memórias do Instituto Oswaldo Cruz*, 112, 319-327.
- Puntasecca, C. J., King, C. H., & LaBeaud, A. D. (2021). Measuring the global burden of chikungunya and Zika viruses: A systematic review. *PLoS neglected tropical diseases*, 15(3), e0009055.
- Rault, L.C., O'Neal, S.T., Johnson, E.J., Anderson, T.D., 2019. Association of age, sex, and pyrethroid resistance status on survival and cytochrome P450 gene expression in *Aedes aegypti* (L.). *Pesticide biochemistry and physiology* 156, 96-104.
- Rey, J. R., Nishimura, N., Wagner, B., Braks, M. A. H., O'Connell, S. M., & Lounibos, L. P. (2006). Habitat Segregation of Mosquito Arbovirus Vectors in South Florida. *Journal of Medical Entomology*, 43(6), 1134–1141.
- Rogers, D. J., & Randolph, S.E. (2000). The global spread of malaria in a future, warmer world. *Science* 289:1763-1766.
- Rohani, A., Wong, Y.C., Zamre, I., et al. (2009) The effect of extrinsic incubation temperature on development of dengue serotype 2 and 4 viruses in *Aedes aegypti* (L.). *SE Asian J. Trop. Med.*, 40, 942.
- Saavedra-Rodríguez, K., Urdaneta-Marquez, L., Rajatileka, S., Moulton, M., Flores, A., Fernández-Salas, I., Bisset, J., Rodríguez, M., McCall, P., Donnelly, M., 2007. A mutation in the voltage-gated sodium channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*. *Insect molecular biology* 16, 785-798.
- Saavedra-Rodríguez, K., Maloof, F.V., Campbell, C.L., Garcia-Rejon, J., Lenhart, A., Penilla, P., Rodríguez, A., Sandoval, A.A., Flores, A.E., Ponce, G., 2018. Parallel evolution of vgsc mutations at domains IS6, IIS6 and IIS6 in pyrethroid resistant *Aedes aegypti* from Mexico. *Scientific reports* 8, 6747.
- Saghir, J., & Santoro, J. (2018). Urbanization in Sub-Saharan Africa. In *Meeting Challenges by Bridging Stakeholders*. Washington, DC, USA: Center for Strategic & International Studies.
- Samson, D. M., Archer, R. S., Alimi, T. O., Arheart, K. L., Impoinvil, D. E., Oscar, R., Fuller, D. O., & Qualls, W. A. (2015). New baseline environmental assessment of mosquito ecology in northern Haiti during increased urbanization. *Journal of Vector Ecology: Journal of the Society for Vector Ecology*, 40(1), 46–58.
- Schmittgen, T.D., Livak, K.J., 2008. Analyzing real-time PCR data by the comparative C T method. *Nature protocols* 3, 1101.



Schneider, J.R., Morrison, A.C., Astete, et al. (2004). Adult Size and Distribution of *Aedes aegypti* (Diptera: Culicidae) Associated with Larval Habitats in Iquitos, Peru. *J Med Entomol* 41, 634–642.

Schoeler, G. B., Schleich, S. S., Manweiler, S. A., & Sifuentes, V. L. (2004). Evaluation of surveillance devices for monitoring *Aedes aegypti* in an urban area of northeastern Peru. *Journal of the American Mosquito Control Association*, 20(1), 6–11.

Scott, T., & Morrison, A. (2009). Vector dynamics and transmission of dengue virus: Implications for dengue surveillance and prevention strategies: Vector dynamics and dengue prevention. *Dengue Virus*, 338, 115–128.

Service, M. W. (1970). Studies on the biology and taxonomy of *Aedes* (*Stegomyia*) *vittatus* (Bigot) (Diptera: Culicidae) in northern Nigeria. *Trans R Soc Entomol Soc Lond* 122, 101–43.

Simard F, Nchoutpouen E, Toto JC, Fontenille D (2005). Geographic Distribution and Breeding Site Preference of *Aedes albopictus* and *Aedes aegypti* (Diptera: Culicidae) in Cameroon, Central Africa. *Journal of Medical Entomology* 42, 726–731.

Singh, H., Singh, O. P., Akhtar, N., Sharma, G., Sindhania, A., Gupta, N., & Valecha, N. (2019). First report on the transmission of Zika virus by *Aedes* (*Stegomyia*) *aegypti* (L.) (Diptera: Culicidae) during the 2018 Zika outbreak in India. *Acta Tropica*, 199, 105114.

Sivan, Arun et al. “Host-Feeding Pattern of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) in Heterogeneous Landscapes of South Andaman, Andaman and Nicobar Islands, India.” *Parasitology research (1987)* 114.9 (2015): 3539–3546.

Smith, L.B., Kasai, S., Scott, J.G., 2016. Pyrethroid resistance in *Aedes aegypti* and *Aedes albopictus*: Important mosquito vectors of human diseases. *Pestic Biochem Physiol* 133, 1-12.

Smith, L.B., Tyagi, R., Kasai, S., Scott, J.G., 2018. CYP-mediated permethrin resistance in *Aedes aegypti* and evidence for trans-regulation. *PLoS neglected tropical diseases* 12, e0006933.

Sombié, A., Saiki, E., Yaméogo, F., Sakurai, T., Shirozu, T., Fukumoto, S., Sanon, A., Weetman, D., McCall, P.J., Kanuka, H., 2019. High frequencies of F1534C and V1016I *kdr* mutations and association with pyrethroid resistance in *Aedes aegypti* from Somgandé (Ouagadougou), Burkina Faso. *Tropical medicine and health* 47, 2.

- Souza-Neto, J. A., Powell, J. R., & Bonizzoni, M. (2019). *Aedes aegypti* vector competence studies: A review. *Infection, Genetics and Evolution*, 67, 191–209.
- Tchetgna, H. S., Yousseu, F. S., Kamgang, B., Tedjou, A., McCall, P. J., & Wondji, C. S. (2021). Concurrent circulation of dengue serotype 1, 2 and 3 among acute febrile patients in Cameroon. *PLOS Neglected Tropical Diseases*, 15(10), e0009860.
- Tedjou A, Kamgang B, Yougang A, et al. (2019). “Update on the Geographical Distribution and Prevalence of *Aedes Aegypti* and *Aedes Albopictus* (Diptera: Culicidae), Two Major Arbovirus Vectors in Cameroon.” *PLOS Neglected Tropical Diseases* 13: 3.
- Thein, S., Aung, M. M., Shwe, T. N., Aye, M., Zaw, A., Aye, K., Aye, K. M., & Aaskov, J. (1997). Risk factors in dengue shock syndrome. *The American Journal of Tropical Medicine and Hygiene*, 56(5), 566–572.
- Trpis, M., McClelland, G. A. H., Gillett, J. D., Teesdale, C., & Rao, T. R. (1973). Diel periodicity in the landing of *Aedes aegypti* on man. *Bulletin of the World Health Organization*, 48(5), 623–629.
- Tsetsarkin, K. A., Vanlandingham, D. L., McGee, C. E., & Higgs, S. (2007). A single mutation in chikungunya virus affects vector specificity and epidemic potential. *PLoS pathogens*, 3(12), e201.
- Tsuda, Y., W. Suwonkerd, S. Chawprom, et al. (2006). Different spatial distribution of *Aedes aegypti* and *Aedes albopictus* along an urban–rural gradient and the relating environmental factors examined in three villages in northern Thailand. *J. Am. Mosq. Control Assoc.* 22: 222–228.
- Valentine, M. J., Murdock, C. C., & Kelly, P. J. (2019). Sylvatic cycles of arboviruses in non-human primates. *Parasites & Vectors*, 12(1), 463.
- Vikram, K., Nagpal, B. N., Pande, V., Srivastava, A., Saxena, R., Singh, H., ... & Valecha, N. (2015). Detection of dengue virus in individual *Aedes aegypti* mosquitoes in Delhi, India. *Journal of vector borne diseases*, 52(2), 129.
- Waggoner, J. J., Gresh, L., Vargas, M. J., Ballesteros, G., Tellez, Y., Soda, K. J., Sahoo, M. K., Nuñez, A., Balmaseda, A., Harris, E., & Pinsky, B. A. (2016). Viremia and Clinical Presentation in Nicaraguan Patients Infected With Zika Virus, Chikungunya Virus, and Dengue Virus. *Clinical Infectious Diseases*, 63(12), 1584–1590.
- Wang, S. M., & Sekaran, S. D. (2010). Early Diagnosis of Dengue Infection Using a Commercial Dengue Duo Rapid Test Kit for the Detection of NS1, IGM, and IGG. *The American Journal of Tropical Medicine and Hygiene*, 83(3), 690–695.

Weaver, S. C., & Reisen, W. K. (2010). Present and future arboviral threats. *Antiviral Research*, 85(2), 328–345.

Weetman, D., Kamgang, B., Badolo, A., Moyes, C.L., Shearer, F.M., Coulibaly, M., Pinto, J., Lambrechts, L., McCall, P.J., 2018. *Aedes* mosquitoes and *Aedes*-borne arboviruses in Africa: current and future threats. *International journal of environmental research and public health* 15, 220.

Westby, K. M., Adalsteinsson, S. A., Biro, E. G., Beckermann, A. J., & Medley, K. A. (2021). *Aedes albopictus* Populations and Larval Habitat Characteristics across the Landscape: Significant Differences Exist between Urban and Rural Land Use Types. *Insects*, 12(3), 196.

Whitehorn, J., Kien, D. T., Nguyen, N. M., Nguyen, H. L., Kyrylos, P. P., Carrington, L. B., Tran, C. N., Quyen, N. T., Thi, L. V., Le Thi, D., Truong, N. T., Luong, T. T., Nguyen, C. V., Wills, B., Wolbers, M., & Simmons, C. P. (2015). Comparative Susceptibility of *Aedes albopictus* and *Aedes aegypti* to Dengue Virus Infection After Feeding on Blood of Viremic Humans: Implications for Public Health. *The Journal of infectious diseases*, 212(8), 1182–1190.

WHO, 2006. Pesticides and their application: for the control of vectors and pests of public health importance.

WHO, 2016. Entomological surveillance for *Aedes* spp. in the context of Zika virus: interim guidance for entomologists.

Wilson, A.L., Courtenay, O., Kelly-Hope, L.A., Scott, T.W., Takken, W., Torr, S.J., Lindsay, S.W., 2020. The importance of vector control for the control and elimination of vector-borne diseases. *PLoS neglected tropical diseases* 14, e0007831.

World Data Base on Protected Areas (2022). <https://dopa-explorer.jrc.ec.europa.eu/ecoregion/30705OSM> Mapnik, January 2021 version of the World Database on Protected Areas (WDPA)

World Health Organization (2020a). Vector-borne diseases. (n.d.). Retrieved January 1, 2021, from <https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases>

World Health Organization (2022). Dengue and Severe Dengue. Retrieved October 12, 2022, from <https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue>

World Health Organization (WHO) (2022). Yellow fever - East, West, and Central Africa 2 September 2022, Disease Outbreak News.

Yang, X., Quam, M. B., Zhang, T., & Sang, S. (2021). Global burden for dengue and the evolving pattern in the past 30 years. *Journal of travel medicine*, 28(8).

Yougang, A.P., Kamgang, B., Bahun, T.A.W., Tedjou, A.N., Nguiffo-Nguete, D., Njiokou, F., Wondji, C.S., 2020a. First detection of F1534C knockdown resistance mutation in *Aedes aegypti* (Diptera: Culicidae) from Cameroon. *Infectious Diseases of Poverty* 9, 1-12.

Yougang, A.P., Kamgang, B., Tedjou, A.N., Wilson-Bahun, T.A., Njiokou, F., Wondji, C.S., 2020b. Nationwide profiling of insecticide resistance in *Aedes albopictus* (Diptera: Culicidae) in Cameroon. *PloS one* 15, e0234572.

Zahouli, J. B. Z., Koudou, B. G., Müller, P., Malone, D., Tano, Y., & Utzinger, J. (2017). Effect of land-use changes on the abundance, distribution, and host-seeking behavior of *Aedes* arbovirus vectors in oil palm-dominated landscapes, southeastern Côte d'Ivoire. *PLOS ONE*, 12(12), e0189082.