

UC San Diego

UC San Diego Previously Published Works

Title

The new paradigm of hepatitis C therapy: integration of oral therapies into best practices

Permalink

<https://escholarship.org/uc/item/75w5b426>

Journal

Journal of Viral Hepatitis, 20(11)

ISSN

1352-0504

Authors

Afdhal, NH
Zeuzem, S
Schooley, RT
[et al.](#)

Publication Date

2013-11-01

DOI

10.1111/jvh.12173

Peer reviewed

REVIEW

The new paradigm of hepatitis C therapy: integration of oral therapies into best practices

N. H. Afdhal,¹ S. Zeuzem,² R. T. Schooley,³ D. L. Thomas,⁴ J. W. Ward,⁵ A. H. Litwin,⁶ H. Razavi,⁷ L. Castera,⁸ T. Poynard,⁹ A. Muir,¹⁰ S. H. Mehta,¹¹ L. Dee,¹² C. Graham,¹³ D. R. Church,¹⁴ A. H. Talal,¹⁵ M. S. Sulkowski,¹⁶ and I. M. Jacobson¹⁷ for the New Paradigm of HCV Therapy Meeting Participants*

¹Division of Gastroenterology and Hepatology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA; ²Department of Medicine, J.W. Goethe University Hospital, Frankfurt, Germany; ³Division of Infectious Diseases, San Diego School of Medicine, University of California, La Jolla, CA, USA; ⁴Division of Infectious Diseases, Johns Hopkins University School of Medicine, Baltimore, MD, USA; ⁵Division of Viral Hepatitis, Centers for Disease Control and Prevention, Atlanta, GA, USA; ⁶Departments of Medicine and Psychiatry and Behavioral Sciences, Montefiore Medical Center, Albert Einstein College of Medicine, Bronx, NY, USA; ⁷Center for Disease Analysis, Louisville, CO, USA; ⁸Service d'Hépatologie, Hôpital Beaujon, Assistance Publique Hôpitaux de Paris, Clichy, France; ⁹Service d'Hépatologie, Groupe Hospitalier Pitie-Salpêtrière, Paris, France; ¹⁰Gastroenterology and Hepatology Research Group, Duke Clinical Research Institute, Durham, NC, USA; ¹¹Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA; ¹²Fair Pricing Coalition and AIDS Action Baltimore, Baltimore, MD, USA; ¹³Division of Infectious Disease, Beth Israel Deaconess Medical Center, Boston, MA, USA; ¹⁴Massachusetts Department of Public Health, Bureau of Infectious Disease, Boston, MA, USA; ¹⁵Division of Gastroenterology and Hepatology, Department of Internal Medicine, University at Buffalo, Buffalo, NY, USA; ¹⁶Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA; and ¹⁷Division of Gastroenterology and Hepatology, Weill Cornell Medical College, New York, NY, USA

Received August 2013; accepted for publication August 2013

SUMMARY. Emerging data indicate that all-oral antiviral treatments for chronic hepatitis C virus (HCV) will become a reality in the near future. In replacing interferon-based therapies, all-oral regimens are expected to be more tolerable, more effective, shorter in duration and simpler to administer. Coinciding with new treatment options are novel methodologies for disease screening and staging, which create the possibility of more timely care and treatment. Assessments of histologic damage typically are performed using liver biopsy, yet noninvasive assessments of histologic damage have become the norm in some European countries and are becoming more widespread in the United States. Also in place are new Centers for Disease Control and Prevention (CDC) initiatives to simplify testing, improve provider and patient awareness and expand recommendations for HCV screening beyond risk-based strategies. Issued in 2012, the CDC recommendations aim to

increase HCV testing among those with the greatest HCV burden in the United States by recommending one-time testing for all persons born during 1945–1965. In 2013, the United States Preventive Services Task Force adopted similar recommendations for risk-based and birth-cohort-based testing. Taken together, the developments in screening, diagnosis and treatment will likely increase demand for therapy and stimulate a shift in delivery of care related to chronic HCV, with increased involvement of primary care and infectious disease specialists. Yet even in this new era of therapy, barriers to curing patients of HCV will exist. Overcoming such barriers will require novel, integrative strategies and investment of resources at local, regional and national levels.

Keywords: diagnosis, directly acting antiviral agents, health services, hepatitis C, pharmacoeconomics.

Abbreviations: AASLD, American Association for the Study of Liver Diseases; ACA, Affordable Care Act; APRI, AST-to-Platelet Ratio Index; CDC, Centers for Disease Control and Prevention; FDA, US Food and Drug Administration; HCV, hepatitis C virus; *IL28B*, interleukin-28B; PEG-IFN, pegylated interferon-alfa; RBV, ribavirin; SVR, sustained virological response; USPSTF, United States Preventive Services Task Force.

Correspondence: Nezam H. Afdhal, MD, Division of Gastroenterology and Hepatology, Beth Israel Deaconess Medical Center and Harvard Medical School, 110 Francis Street, Boston, MA 02215, USA.

E-mail: nafdhal@caregroup.harvard.edu

*The participants and their affiliations are listed in the Appendix 1.

© 2013 The Authors. Journal of Viral Hepatitis published by John Wiley & Sons Ltd
This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

INTRODUCTION

The treatment landscape for hepatitis C is in flux. From 2002 to 2011, the standard of care treatment for chronic infection with hepatitis C virus (HCV) was 24 or 48 weeks of therapy with pegylated interferon-alfa (PEG-IFN) and ribavirin (RBV). For patients with genotype 1 virus, the likelihood of achieving a sustained virological response (SVR), defined as having undetectable serum HCV RNA at 24 weeks after cessation of treatment, was only 40–50% after 48 weeks of therapy. In 2011, the HCV protease inhibitors telaprevir and boceprevir entered the market to be used in combination with PEG-IFN and RBV for genotype 1 HCV infection. The protease inhibitors increase the likelihood of SVR to 67–75% in treatment-naïve patients with genotype 1 HCV [1–4]. However, adding a protease inhibitor to a PEG-IFN backbone, which is itself difficult to tolerate, has increased the potential for toxicity and has placed a resource-intensive burden on treating physicians. In addition, the triple therapy regimens have limited efficacy in treatment-experienced null responders [5].

Several directly acting antiviral agents are being evaluated for their potential use in combination with either RBV or other antivirals of different classes. In early 2013, a small study of the nucleotide sofosbuvir in combination with RBV was reported, and among 25 treatment-naïve, HCV genotype 1 patients, 21 (84%) had an SVR after 12 weeks of therapy [6]. In another small study reported in early 2013, a total of 31 of 33 (94%) previously treatment-naïve genotype 1 patients were HCV RNA negative 12 weeks after cessation of therapy with the NS3 protease inhibitor ABT-450, combined with low-dose ritonavir, the non-nucleoside NS5B polymerase inhibitor ABT-333 and ribavirin [7]. In phase 3 studies of sofosbuvir with RBV, SVR rates have been as high as 78% in HCV genotype 2 and 3 patients [8], and it is anticipated that in the United States, all-oral combination therapies will be available for HCV genotype 2 or 3 patients by 2014. By 2015, regimens including only directly acting antivirals are expected to be available for persons with any HCV genotype.

At this time of new treatment opportunities, novel changes have been made to improve the methods by which extent of liver disease is diagnosed. Assessment of the extent of histologic damage, an important component of patient evaluation, has been traditionally carried out by liver biopsy. Noninvasive assessments of histologic damage such as elastography have become the norm in several European countries and are becoming more frequently used in the United States. Strategies for HCV screening also have been revised. To improve the identification of persons living with chronic HCV infection, the U.S. Centers for Disease Control and Prevention (CDC) expanded its risk-based approach to HCV testing, publishing a recommendation in 2012 that all persons born during 1945–1965 receive one-time testing for HCV. In 2013, the United States Preventive Services

Task Force (USPSTF) adopted similar recommendations for risk-based and birth-cohort-based testing. Growing evidence suggests that in the United States, HCV infections are rapidly increasing among persons aged 15–24 primarily because of injection drug use [9]. This trend suggests that screening efforts should also ensure that young injection drug users are tested and engaged in care.

The 2010 Patient Protection and Affordable Care Act (ACA) will expand opportunities for persons to purchase health insurance and have access to hepatitis C testing, care and treatment. The ACA will facilitate implementation of HCV testing because it requires nongrandfathered private health plans to cover clinical preventive services given an A or B grade by the USPSTF without cost-sharing and provides incentives for Medicaid programmes to cover these services. By prohibiting insurance companies from declining to sell or renew policies because of pre-existing conditions such as hepatitis C, ACA will help more patients access HCV care and treatment services [10].

Improvements in therapies, diagnostic techniques and screening for HCV will create a new era for HCV treatment. Although the exact effects these changes will have on the future landscape of HCV care cannot be elucidated, certain outcomes are likely. For instance, as methods for diagnosis of liver disease and treatment of HCV become simpler, safer and more effective, primary healthcare providers may manage greater numbers of HCV-infected patients. This expansion into primary care may become necessary if the number of patients undergoing treatment increases because of screening efforts and improved prospects for treatment success for regimens containing all directly acting antivirals. Also subject to change are pricing and reimbursement models, as well as the pharmacoeconomics of curing HCV.

To discuss the new paradigm of HCV therapy, representatives from leading academic medical centres, government agencies, insurance providers and the pharmaceutical and biotechnology industries met in Boston, Massachusetts, USA, on March 22 and 23, 2013. The focus of the meeting, or Think Tank, was to predict how shifts in HCV screening, diagnosis and treatment will affect access to and delivery of care; identify barriers to treating HCV; discuss successful strategies for identifying and treating patients; and discuss the pharmacoeconomics of treatment for patients, providers, pharmaceutical companies and health-care payers. Here, we describe the current challenges and opportunities for curing HCV in the forthcoming era.

CURRENT EVOLUTION OF ALL-ORAL THERAPIES FOR HCV

Arrays of IFN-free regimens for treating HCV are currently in the later stages of clinical development. At scientific meetings, data have been presented from phase 2 and 3 studies of various all-oral regimens. The results of individ-

Table 1 Reported results for all-oral therapies for hepatitis C virus in clinical development

	No. patients	Duration, weeks	SVR rates	Reference
Treatment-naïve patients				
Genotype 1 (1a or 1b)				
ABT-450/r + ABT-333 + RBV	33	12	94% SVR12	Poordad <i>et al.</i> [7]
ABT-450/r + ABT-267 + ABT-333 + RBV	80	8	88% SVR12	Kowdley <i>et al.</i> [11]
ABT-450/r + ABT-333 + RBV	41	12	85%	Kowdley <i>et al.</i> [11]
ABT-450/r + ABT-267 + RBV	79	12	90%	Kowdley <i>et al.</i> [11]
ABT-450/r + ABT-267 + ABT-333	79	12	87%	Kowdley <i>et al.</i> [11]
ABT-450/r + ABT-267 + ABT-333 + RBV	79	12	98%	Kowdley <i>et al.</i> [11]
Daclatasvir + Asunaprevir + BMS-791325	16	24	88%	Everson <i>et al.</i> [98]
Daclatasvir + Asunaprevir + BMS-791325	16	12	94%	Everson <i>et al.</i> [98]
Faldaprevir + Deleobuvir	46	28	39% SVR12	Zeuzem <i>et al.</i> [15]
Faldaprevir + Deleobuvir + RBV	316	16, 28, or 40	52–69% SVR12	Zeuzem <i>et al.</i> [15]
Mericitabine + Danoprevir + RBV	64	24	71% SVR12	Gane <i>et al.</i> [99]
Sofosbuvir + RBV	25	12	84%	Gane <i>et al.</i> [6]
Sofosbuvir + Daclatasvir	55	12 or 24	98% SVR4	Sulkowski <i>et al.</i> [14]
Sofosbuvir + Daclatasvir + RBV	56	12 or 24	96% SVR4	Sulkowski <i>et al.</i> [14]
Sofosbuvir + Ledipasvir + RBV	25	12	100% SVR12	Gane <i>et al.</i> [100]
Genotype 2 or 3				
Sofosbuvir + RBV	10	12	100%	Gane <i>et al.</i> [6]
Sofosbuvir	10	12	60%	Gane <i>et al.</i> [6]
Sofosbuvir + RBV	253	12	67% SVR12	Gane <i>et al.</i> [101]
Sofosbuvir + Daclatasvir	14	24	100%	Sulkowski <i>et al.</i> [14]
Sofosbuvir + Daclatasvir + RBV	14	24	93%	Sulkowski <i>et al.</i> [14]
Prior nonresponse				
Genotype 1 (1a or 1b)				
ABT-450/r + ABT-333 + RBV	17	12	47% SVR12	Poordad <i>et al.</i> [7]
ABT-450/r + ABT-267 + RBV	45	12	89%	Kowdley <i>et al.</i> [11]
ABT-450/r + ABT-267 + ABT-333 + RBV	45	12	93%	Kowdley <i>et al.</i> [11]
Daclatasvir + Asunaprevir	11	24	36%	Lok <i>et al.</i> [102]
Sofosbuvir + RBV	10	12	10%	Gane <i>et al.</i> [6]
Sofosbuvir + Daclatasvir	21	12	100%	Sulkowski <i>et al.</i> [13]
Sofosbuvir + Daclatasvir + RBV	20	12	95%	Sulkowski <i>et al.</i> [13]
Sofosbuvir + Simeprevir + RBV	27	12	96% SVR8	Lawitz <i>et al.</i> [12]
Sofosbuvir + Simeprevir	14	12	93% SVR8	Lawitz <i>et al.</i> [12]
Sofosbuvir + Ledipasvir + RBV	10	12	100% SVR12	Gane <i>et al.</i> [100]
Genotype 1b				
Daclatasvir + Asunaprevir	21	24	91%	Suzuki <i>et al.</i> [103]
Genotype 2 or 3				
Sofosbuvir + RBV	201	12 or 16	SVR12 12 week: 50% 16 week: 73%	Jacobson <i>et al.</i> [8]
IFN-ineligible or intolerant				
Genotype 1b				
Daclatasvir + Asunaprevir	22	24	64%	Suzuki <i>et al.</i> [103]
Genotype 2 or 3				
Sofosbuvir + RBV	207	12	78% SVR12	Jacobson <i>et al.</i> [8]

ABT-450/r, ritonavir-boosted ABT-450; IFN, interferon; RBV, ribavirin; SVR, viral negativity 24 weeks post-therapy. SVR12, SVR8 and SVR4 refer to viral negativity at 12, 8 and 4 weeks post-therapy.

ual studies will not be described in great detail here but are summarized in Table 1. The more promising regimens have the following characteristics: a strong safety profile, SVR rates approaching or even exceeding 90%, minimal

pill burden and minimal potential for drug–drug interactions.

Several major conclusions and predictions regarding the future of all-oral therapies were discussed. A reasonable

anticipation is that a genotype-specific, all-oral therapy for HCV genotypes 2 and 3 with sofosbuvir and ribavirin will be available by 2014. By 2015, genotype-1-specific therapies should follow, and these will comprise any of three regimens currently under development by AbbVie Pharmaceuticals, Bristol-Myers Squibb and Gilead Sciences (Table 1). True pangenotypic regimens will probably not be available until 2016 or 2017 and will require development of pangenotypic NS5A inhibitors and protease inhibitors that can be combined with each other and with nucleotide polymerase inhibitors. Some of these combinations are in phase 1 or early phase 2 studies across multiple genotypes.

Another major area of discussion was whether pretreatment and on-treatment predictors of response, including those used for PEG-IFN, can help predict response to all-oral therapies. Although there is evidence that many of these factors still predict response to relatively weak interferon-sparing regimens, more potent regimens, with SVR rates >90%, readily overcome the traditional obstacles seen with PEG-IFN. For example, in phase 2 studies of the Abbott multidrug regimen [11] or sofosbuvir plus simeprevir [12] or daclatasvir [13], prior interferon response was not strongly related to response to all-oral treatment. In fact, interferon null responders did just as well as naïve patients and had SVR rates in the 90% range. It is increasingly apparent that regimens consisting of potent agents that individually or cumulatively impose a high barrier to resistance attenuate or eliminate factors such as 1a/1b subtype, *IL28B* status, viral load, race, metabolic syndrome, obesity and age as major determinants of response. In addition, with potent directly acting antiviral combinations, nearly all patients are negative within 4 weeks, which means the traditional strategy of using virologic response at week 4 or 12 to determine the duration of treatment may be moot. The presence of cirrhosis, which often excludes patients from early phase trials, may yet be a differentiating factor in SVR rates, but this remains to be further determined for genotype 1, and as with other factors, presence of cirrhosis can probably be overcome by a sufficiently potent regimen or longer treatment duration. In studies of sofosbuvir and RBV in patients with HCV genotypes 2 or 3, cirrhosis was a significant negative predictor of response for treatment-naïve patients with HCV genotype 3 and for prior treatment-failure patients with either genotype 2 or 3, but these studies only included 1 potent directly acting antiviral. The effect of portal hypertension and hepatocellular dysfunction (Child's class B and C) on SVR in patients with more advanced liver disease remains an area requiring additional investigation.

The final major questions for discussion encompassed the need for RBV and duration of therapy, which are in some ways connected. As with the pretreatment predictors, ribavirin use and treatment duration appear to matter with relatively weak regimens but may not with sufficiently potent combinations. In studies of the polymerase inhibitor

sofosbuvir with either the NS5A inhibitor daclatasvir [14] or the protease inhibitor simeprevir [12], SVR rates were independent of RBV use. However, in a study combining the protease inhibitor faldaprevir and the non-nucleoside polymerase inhibitor deleobuvir, omitting RBV resulted in a marked reduction in efficacy in genotype 1a patients [15]. And for HCV genotype 1a patients in the phase 2 AVIATOR trial [11], the removal of RBV from a regimen containing the ritonavir-boosted protease inhibitor ABT-450/r, the NS5A inhibitor ABT-267 and the non-nucleoside polymerase inhibitor ABT-333 resulted in a 10% loss of efficacy. The optimal duration of therapy remains unknown, but with potent regimens, 12 weeks is probably the maximum required for most patients (with the potential exception for patients with advanced cirrhosis). Eight-week treatment regimens can be explored, although this may result in a moderate (~10%) reduction in SVR [11] depending on the regimen.

For regimens containing only direct-acting antivirals, one could imagine a scenario where more potentially difficult-to-treat patients are distinguished from a potentially more easily treatable population. Difficult-to-treat patients may be best served by undergoing an individualized regimen under the care of a specialist. Individualized therapy could be based upon HCV genotype, fibrosis stage, comorbidities, concomitant medications or prior directly acting antiviral drug exposure. Populations of patients who may require individualized therapy but for whom evidence-based treatment data are limited include those with cirrhosis, including decompensated cirrhosis, HIV coinfection, renal failure, an organ transplant or other conditions resulting in being immunocompromised. In the future, it is possible that the population of HCV positive individuals with F0-2 histology will undergo treatment without further stratification such as via HCV genotype or *IL28B* polymorphisms, because SVR rates will likely be in the 90% range.

SCREENING STRATEGIES FOR HCV

In the United States, mortality associated with hepatitis C is on the rise and currently exceeds that for HIV [16]. On the basis of survey data from 1999 to 2002, it has been estimated that 3.2 (2.7–3.9) million persons in the United States have chronic HCV infection [17]. The strongest risk of HCV infection is a history of injection drug use [17]. Of persons with chronic infection, 74% were born during the years 1945 through 1965 [18].

In 1998, CDC recommended a risk-based approach to screening, with routine HCV testing for persons with risk factors including injection drug use, having received clotting factor concentrates produced before 1987, being on chronic haemodialysis, having persistently abnormal alanine aminotransferase levels, being a recipient of donated blood from a person who tested positive for HCV, or having received a blood transfusion or organ transplant before

July 1992 [19]; in 1999, CDC recommended HCV testing for persons with HIV. The 2009 guidelines from the American Association for the Study of Liver Diseases (AASLD) [20] and the 2006 guidelines from the American College of Gastroenterology [21] also recommend screening in high-risk patients. However, risk-based screening strategies can be limited either by clinician reluctance to ask about risk factors or by patient unawareness or reluctance to disclose risk behaviours. As a result, use of risk-based strategies alone has resulted in a large proportion of infected persons remaining undiagnosed; in the United States, various estimates indicate that 45–85% of persons with HCV are unaware of their infection status [22–25]. To augment risk-based screening, in 2012 CDC published a recommendation for one-time testing without prior ascertainment of HCV risk for persons born during 1945–1965 [25]. This birth-cohort approach was designed to both target persons with the highest prevalence of HCV infection and remove any behavioural stigma from screening. In the state of New York, legislation requiring all patients born within the birth-cohort period to be offered hepatitis C screening when they visit healthcare providers has passed the legislature and is awaiting the governor's approval.

It has been estimated that with implementation of the birth-cohort screening strategy, 121 000 deaths from HCV will be averted [26]. In recognition of these and other data [27,28], in June 2013 the USPSTF issued a final recommendation regarding HCV testing, assigning a Grade B to two recommendations: screening for HCV infection in persons at high risk for infection and one-time HCV screening for adults born between 1945 and 1965 [29]. A USPSTF Grade B designation expands access to clinical preventive services.

For HCV screening to become widely adopted in diverse clinical settings providing care for persons at risk for HCV infection, efforts are needed at local, regional and national levels. Approximately 79 million persons were born during 1945–1965 (the Baby Boom Generation), making birth-cohort based screening a daunting task. However, since the release of the CDC recommendations, multiple independent studies of HCV testing have shown birth-cohort-based approaches superior to risk-based strategies alone [30,31]. To increase the numbers of HCV-infected persons aware of their infection, CDC is implementing a national multimedia campaign, Know More Hepatitis, that includes education for consumers and healthcare providers (<http://www.cdc.gov/knowmorehepatitis/>). Specific initiatives include messaging on airport dioramas and billboards in cities such as Atlanta, Washington, DC, Salt Lake City, Orlando and Las Vegas; online medical education for health professionals (<http://depts.washington.edu/hepstudy/hepC/>); and the launch of an annual National Hepatitis Testing Day observed on May 19th.

Centers for Disease Control and Prevention is also conducting demonstration projects to evaluate the implemen-

tation of risk-based and birth-cohort strategies for HCV strategies in over 25 clinical settings. For example, the Hepatitis C Assessment and Testing Project in New York City evaluated community-based screening interventions in three urban primary care clinics [32]. Both risk-based and birth-cohort-based interventions were associated with an increased proportion of patients tested for HCV. Both risk-based and birth-cohort HCV screening approaches can be integrated within electronic medical records.

The new HCV screening recommendations are expected to increase demand for testing to detect current HCV infection. To meet this demand, CDC recently simplified the HCV testing sequence [33]. Patients should first be tested for HCV antibody. Patients who are reactive for HCV antibody should next be tested with an FDA-approved nucleic acid testing assay for the detection of HCV RNA indicative of current HCV infection. Rapid tests for HCV antibody allow access to HCV testing in settings lacking laboratory-based diagnostic services. Rapid tests for HCV antibody detection include OraQuick [34], which is approved by the United States Food and Drug Administration (FDA), as well as Chembio [34,35], MedMira [34,35] and mBio, which are under development.

HCV DIAGNOSTIC TESTING AND DISEASE STAGING

It is likely that the rapid improvements in treatment efficacy and tolerability anticipated with interferon-sparing regimens will also transform our approach to disease staging. The historic low efficacy and safety of interferon-based regimens led to the recommendation for liver staging to determine whether the benefits of treatment would outweigh the risks. For most patients, this required that there be more than just portal fibrosis. Liver biopsy was considered the best test for this purpose, but the procedure is costly, invasive and in a small minority of cases can result in complications such as significant bleeding, organ puncture, or death [36]. And the accuracy in staging disease is often compromised by variability in tissue sampling or in interobserver or intra-observer histopathological scoring.

As treatment efficacy for genotype 2 and 3 infections improved, biopsy was no longer routinely recommended to justify treatment necessity [20]. Likewise, continued improvements in treatment efficacy and safety for all genotypes will change the primary goal of staging from justification of treatment benefit to identification of persons with cirrhosis or bridging fibrosis because they may need longer treatment courses and require hepatocellular carcinoma screening and portal hypertension management. Accordingly, the most important characteristic of a staging test is the negative predictive value for detection of cirrhosis.

Noninvasive methods of assessing histology are becoming more widely used. These include measurement of

Table 2 FibroScan and FibroSure* for diagnosis of cirrhosis

	FibroScan	FibroSure
AUROC, mean (95% CI)	0.94 (0.93–0.95) [41]	0.86 (0.71–0.92) [38]
Sensitivity (95% CI)	0.83 (0.79–0.86) [104]	0.85 [38]
Specificity (95% CI)	0.89 (0.87–0.91) [104]	0.81 [38]
Advantages	Evaluates a genuine property of the liver High performance for cirrhosis User-friendly, point-of-contact test Good reproducibility	Good reproducibility High applicability (>95%)
Disadvantages	Decreased performance in obese patients Applicability lower than serum biomarkers: failure in 3% of cases and unreliable results in 16% (obesity, ascites, limited operator experience) Requires a dedicated device Inflammation, extra-hepatic cholestasis, and right heart failure can provide false positive results	Nonspecific of the liver

*Known as FibroTest in Europe.

serum biomarkers of fibrosis and measurement of liver stiffness through elastography (Table 2). The noninvasive methods have practically no complications and can be performed repeatedly to dynamically monitor progression of fibrosis. The rate of adoption of noninvasive diagnostic tests for liver fibrosis differs between international regions, and the United States lags behind Europe in this regard. The 2012 European Association for the Study of the Liver (EASL) guidelines for treating chronic hepatitis C suggest that while liver biopsy is still regarded as the reference method for grading inflammation and staging fibrosis, transient elastography can be used to assess liver fibrosis, and noninvasive serum markers are recommended for detecting significant fibrosis (METAVIR score F2–F4) [37].

In 2004, the biomarker assay FibroSure (named FibroTest in Europe) was launched in the United States for assessing fibrosis and necroinflammatory activity. The assay, which can only be performed in validated laboratories, predicts a histology score on the basis of patient age, sex and results for serum haptoglobin, α 2-macroglobulin, apolipoprotein A1, γ -glutamyltransferase and bilirubin analyses [38]. In a review of 25 studies in chronic HCV, FibroTest had an AUROC of 0.79 (0.70–0.89) for diagnosis of significant fibrosis ($F \geq 2$) and 0.86 (0.71–0.92) for liver cirrhosis [38].

If the birth-cohort screening to enhance diagnosis of HCV infection is fully implemented, it has been estimated that as many as 800 000 additional cases of HCV infection would be identified [26]. Should there be such a large-scale influx of newly diagnosed patients, tests of serum biomarkers will likely be a more practical approach to liver disease staging than one restricted to liver biopsy-based staging. In the United States, the FibroSure test costs approximately \$250, which is a fraction of the cost of biopsy. The biomarker assay AST-to-Platelet Ratio Index

(APRI) is not proprietary and costs no more than a routine blood draw and routine liver function tests. APRI is calculated as (AST/upper limit of normal range)/platelet count ($10^9/L$) $\times 100$. However, a recent large meta-analysis suggested that APRI can identify hepatitis C-related fibrosis with only a moderate degree of accuracy (AUROC of 0.77 for significant fibrosis and 0.80 for severe fibrosis) [39]. Alternate *in vitro* diagnostic testing for liver fibrosis was subsequently developed, including FibroIndex and Forns index [38]. For identifying cirrhosis, the age-platelet index, APRI and Hepascore have median AUROCs of 0.80 or greater (range 0.80–0.91) [38].

Transient elastography, using the FibroScan device (Echosens, Paris, France), is widely used in several European countries and has more recently been adopted in Asia and Canada. In April of 2013, FibroScan was approved by the FDA for use in the United States. The main limitation of FibroScan use in practice is its limited applicability (80%), mostly due to patient obesity or limited operator experience [40]. Results of a meta-analysis suggest FibroScan is a reliable method for diagnosing significant fibrosis (AUROC = 0.84), severe fibrosis (0.89) and particularly cirrhosis (0.94) [41]. However, for diagnosing significant fibrosis, a high variation of the AUROC was found depending on the type of underlying liver disease [41]. When compared and validated externally in a multicenter prospective study, FibroScan outperformed serum biomarkers of fibrosis for the prediction of cirrhosis (AUROCS 0.89–90 vs 0.77–0.86) but had similar performance for the diagnosis of significant fibrosis [42]. Both FibroScan and FibroTest have a prognostic value similar to liver biopsy for predicting complications and outcome of liver disease [43,44]. Combining FibroScan with Fibrotest may increase diagnostic performance for significant fibrosis [45,46], and this approach has been recommended in the 2012 EASL Guide-




Fig. 1 The hepatitis C care cascade. Among patients infected with hepatitis C virus, fewer than 10% are treated and cured. Barriers exist in screening methods, patient referral to appropriate providers, attending necessary appointments and initiating treatment [49,53,54,61,74,75,105–113].

lines as first line evaluation of liver fibrosis in patients with chronic hepatitis C [37].

The Think Tank recognized that the goals of liver staging are changing and that there is an urgency to revise guidelines accordingly. The accurate exclusion of cirrhosis has been recommended as the most important role for HCV staging in clinical practice, and an algorithm has been proposed on how to best utilize noninvasive tests to achieve this goal [47].

BARRIERS TO CARE AND STRATEGIES TO ADDRESS THEM

Among persons infected with HCV, a substantial portion fails to progress towards a cure at every step of treatment, from recognition of disease to viral clearance (Fig. 1) [48]. In the United States, at least half of those infected with HCV do not know their status [22–25]. Among patients who are recognized as being positive for HCV antibody, it is estimated that fewer than half are linked to care [49]. Failure to link to care represents both a lack of referral to a specialist for treatment and failure to attend the appointment. Even after being linked to proper caregivers, patients can fail to receive pretreatment work-up, meet eligibility criteria for treatment or agree to initiate treatment.

Reasons for these failures can be attributed to barriers at the level of patients, providers and the healthcare system itself. Patients can have limited access to health care, because of lack of or limited insurance [50], low health literacy or not having a usual source of medical care. They can also have competing health priorities, such as mental health issues [51] or comorbidities [52]. Issues related to patient behaviours or environment, such as substance abuse [53,54], lack of drug treatment, lack of social support [55] or unstable employment or housing [56], may also

limit uptake of treatment. Patients may also have limited knowledge of HCV and its treatment and may not perceive it as being something they need to worry about because the disease is largely asymptomatic [57]. Finally, many patients fear the side effects of IFN-based regimens [57].

Primary care providers can have misconceptions about whom to screen, risk of progression of liver disease or ztherapy itself [58,59]. Even specialists in liver disease may have limited experience treating HCV [60]. Providers also can be selective about which patients they consider as good candidates for therapy and fail to recommend treatment because of concerns about nonadherence, drug use [61] or risk of re-infection.

Governments and payers play key roles in delivering HCV services; surveying infection, testing and treatment rates; and educating the public and as well as healthcare providers. Unfortunately, the United States has poorly developed surveillance systems, inadequate educational initiatives and fragmented viral hepatitis services [62]. Also at issue are insufficient numbers of providers who can and are willing to treat HCV [63] and insufficient resources for case managers, navigators and social workers.

Training community-based healthcare providers to treat HCV may become a key method for broadening access to cure. Community-based health centres often have advantages of being culturally appropriate and accessible to patients in both urban and rural areas. In these settings, ongoing relationships with providers may establish trust and an avenue for communication. The Extension for Community Healthcare Outcomes model was developed as a means of using video-conferencing to train primary care providers through interactions with specialists to treat complex diseases, such as HCV. In New Mexico, the programme was successful in generating rates of SVR at 21 sites in rural areas and prisons that were similar to rates

of SVR at the University of New Mexico's HCV clinic [64]. Other strategies to improve rates of initiation and completion of therapy include having peer navigators and integrated care programmes.

The Think Tank believed that screening in conjunction with an all-oral treatment paradigm would reduce the barriers to care and allow treatment within primary care and community sites for many HCV-infected patients.

PHARMACOECONOMICS OF HEPATITIS C

The sequelae of hepatitis C impose a high economic burden. It has been estimated that in 2012, the healthcare cost of HCV was \$6.5 billion, and it has been predicted the cost will peak at \$9.1 billion in 2024 [65]. A retrospective analysis of data from a large, managed care organization claims database suggested that the annual all-cause medical costs of patients diagnosed with HCV were almost twice as high as enrollees without diagnosed HCV [66]. The health burden of HCV largely relates to the development of advanced liver disease, which can lead to liver transplant. In the United States, HCV is the leading cause of hepatocellular carcinoma [67], and it is likely that more cases of hepatocellular carcinoma, decompensated cirrhosis and liver transplants due to HCV will be observed in the coming years [68]. The medical cost of hepatocellular carcinoma has been estimated as \$23 755–44 200 per year per person, and the cost of liver transplant has been estimated as \$201 110 per year per person [69]. Additional disease burden and costs are generated by extrahepatic manifestations of HCV infection including cryoglobulinemic vasculitis, lymphoproliferative disorders, renal disease and rheumatoid-like polyarthritis [70].

The addition of telaprevir or boceprevir to PEG-IFN plus RBV has changed the pharmacoeconomics of treating HCV. Adding the directly acting antivirals to PEG-IFN and RBV can increase the cost of treatment up to \$50 000, depending upon individual regimens needed [71], yet the antivirals also increase the success rates of therapy. At present, economic evaluations of telaprevir or boceprevir with PEG-IFN and RBV are limited. A decision analysis of telaprevir and boceprevir indicated that triple therapy including telaprevir or boceprevir was cost-effective when compared with dual PEG-IFN and RBV therapy in patients with genotype 1 infection [72], although the results were dependent on the cost of protease inhibitors, treatment adherence rates and extent of fibrosis. More recently, a study from Mount Sinai in New York has estimated that the real cost of reaching end of therapy with triple therapy may be as high as \$147 000 when the cost of side effect management is included [73].

As new all-oral regimens enter the market, several factors will affect their cost-effectiveness: success rates in patients with advanced liver disease or difficult-to-treat HCV genotypes, costs related to monitoring and managing

treatment-related toxicities, extent of clinically relevant viral resistance and duration of therapy. The costs of new agents will also be considered against the costs of current IFN-based therapy, which is challenging to administer and has side effects requiring ongoing management. The dominant factor in cost assessments of treatment should be the efficacy of the treatment because all those who fail experience most or all of the cost and none of the benefit. But high projected costs of new directly acting antiviral treatments may result in lack of access for some patients. Industry-created assistance and co-pay programmes can be instrumental in making treatment more affordable and accessible. Public health programmes to support engaging HCV-infected persons in care should also be explored to provide infrastructure for wrap-around services that may not be reimbursable (e.g. coordination of care or peer support). Given the projected high costs of treatment, relatively minor investments in patient support mechanisms are easily justified but not often implemented because of the nature of the fee-for-service healthcare delivery system in the United States. Enhanced communication between physicians and third-party payers may increase the availability of new therapies to patients.

SPECIAL POPULATIONS

Patients with drug addiction

The majority of prevalent and incident infections of HCV occur among injection drug users. Surveillance data have provided evidence that among persons aged 15–24 years, injection drug use is causing a rapid increase in HCV infections [9]. The increase appears to be occurring predominantly in non-Hispanic white males and females. More effort is needed to better understand this trend and to ensure that young injection drug users are tested and engaged in care.

Fewer than 20% of drug users with HCV initiate antiviral therapy [49,54,74,75], principally because of lack of knowledge about HCV, an exaggerated concern about treatment-related side effects and a low perceived need for treatment. There has been reluctance among many healthcare providers to treat drug users because of concerns about adherence, potential reinfection even if SVR is attained, and overall lack of experience and consequent discomfort with the care of patients with addiction problems. Despite concerns regarding adherence to HCV treatment, results of a recent meta-analysis suggest that treatment completion rates among drug users who initiate therapy are over 80% [76]. Addiction treatment and support services (including peer support) increase HCV treatment completion rates [77–80]. Multidisciplinary models for the management of HCV among people who inject drugs have been described in community-based clinics, substance abuse treatment clinics and hospital-based clin-

ics [79,81–83]. For example, integrating internist-addiction medicine specialists from a methadone maintenance treatment programme into a hepatitis clinic improved adherence with HCV evaluation and treatment relative to standard referral practices in patients with prior or ongoing drug use [78]. Education of both patients and providers about the disease and close collaboration between HCV treaters and those who treat addiction are important elements to promote successful treatment of this patient population. Evidence-based international recommendations for treating hepatitis C in people who inject drugs were recently released [84].

As HCV treatment shifts to all-oral regimens, wider uptake among drug users is likely to occur because of decreased side effects, elimination of mental illness as an exclusion for therapy and elimination of needle exposure during therapy. As suggested by modelling data, even modest increases in the numbers of active injection drug users who receive treatment may interrupt HCV transmission enough to result in substantial declines in HCV prevalence [85,86]. If the modelling data are verified by field studies, timely HCV detection and treatment and their integration with other services for drug addiction will take on new urgency. The Think Tank emphasized the need for interventions that facilitate access to HCV therapy for drug users, such as promoting HCV treatment among addiction medicine specialists. In clinical studies of novel therapies, exclusion criteria often limit participation of patients with a history of injection drug use, even if patients have not been using for a long time. Broadening criteria to include such patients would better inform efficacy of treatment in this population.

Patients with cirrhosis

Diagnosis of cirrhosis in patients with HCV is important in part because these patients have a higher incidence of hepatocellular carcinoma and a potential for bleeding from oesophageal varices. Screening for each of these may result in reductions in morbidity and mortality. Although the presence of cirrhosis decreases the likelihood of response to current triple therapy regimens and increases the risk of side effects [87], its presence does not rule out the possibility of initiating therapy. Patients with compensated cirrhosis (Childs-Pugh A) may be candidates for triple therapy if they have well-maintained hepatic synthetic function and no complications of portal hypertension (as assessed by serum albumin and platelets). Indeed, treatment has traditionally been considered strongly indicated in well-compensated cirrhosis to prevent further disease progression or decompensation. In contrast, patients with decompensated cirrhosis (Childs-Pugh B or C) are no longer considered candidates for receiving current triple therapy regimens. Some of the newer regimens have demonstrated promising rates of efficacy for patients with cirrhosis [88,89]. The most extensively studied oral regimen, with data from a phase 3

programme, is sofosbuvir and ribavirin in patients with genotypes 2 and 3, in which cirrhosis had an impact in patients with genotype 3 that may be ameliorated with longer duration of therapy [8]. With the proliferation of novel drugs and regimens under investigation, studies are needed to address issues such as the pharmacokinetics and pharmacodynamics in the setting of cirrhosis; tolerability and efficacy across Childs-Pugh A, B and C patients; and impact of SVR on clinical outcomes. Drug–drug interactions, especially in post-transplant patients, must also be evaluated. There is an urgent need for these issues to be addressed as early in drug development programmes as possible.

Patients with HCV–HIV coinfection

Persons with HIV infection have a high prevalence of chronic HCV infection with a tendency towards more rapid progression to cirrhosis and potentially less access to liver transplantation. Some reports suggest that relative to HCV mono-infected patients, HIV–HCV coinfecting patients also have higher rates of comorbid conditions such as drug use, major depression and anaemia [90]. With HIV therapies, drug interactions may occur and may be difficult to predict; therefore, novel direct antiviral therapies for HCV will need to be evaluated for their potential for interaction with at least some HIV antiretrovirals. Coinfecting patients have reduced rates of response to therapy with PEG-IFN and RBV [91], but adding telaprevir or boceprevir increases efficacy of therapy [92,93]. Some of the newer regimens may have even greater efficacy [94,95]. Curing HCV in coinfecting patients is linked to improved clinical outcomes and longer survival [93,96]. Conducting studies of the newer regimens in coinfecting patients will be important for generating data needed to develop practice guidelines and justify third-party payment.

PRIORITIES FOR EDUCATION AND RESEARCH

Although all-oral therapies are likely to be simpler to administer than IFN-based therapies, educating patients and providers will remain a challenge. Both patients and providers need to receive clear messages on the natural history of HCV, with warning signs and an explanation of why diagnosis and treatment is important. Providers will need to be educated regarding best practices for screening, diagnosis and treatment. Partnerships between members of academia, community health centres, the HCV-affected community, the pharmaceutical industry, healthcare payers and federal, state, and local government entities are very useful for performing postmarketing studies and education (Table 3). One example is the CDC Foundation's Viral Hepatitis Action Coalition (<http://www.viralhepatitis-action.org/>).

Updated treatment guidelines serve as a valuable resource for providers and also influence payer policies. The most recent guidelines for diagnosing, managing and

Table 3 Delivering solutions for hepatitis C

Advance and simplify treatment
Improve efficacy, safety and tolerability
Eliminate interferon
Shorten treatment duration
Develop pan-genotypic all-oral regimens
Increase awareness and screening
Support efforts to enhance public awareness, education, and testing
Develop programmes to educate providers
Participate in public-private partnerships (e.g. Viral Hepatitis Action Coalition)
Ensure access to therapy
Provide patient assistance, co-pay programmes
Collaborate with international partners
Develop innovative access models (e.g. licensing agreements)

treating hepatitis C in the United States were published by the AASLD in 2009 [20], before FDA approval of telaprevir and boceprevir. A 2011 update revised treatment recommendations for patients with HCV genotype 1 [97], yet the approach to testing and staging was not reassessed. Expert opinion pieces can be helpful when guidelines are outdated and should be considered as a means to provide guidance in a rapidly changing field. In July 2013, the AASLD and Infectious Diseases Society of America announced a collaboration to develop clinical recommendations for managing hepatitis C. To serve the medical community in the next few years, one can anticipate a need for much more frequent revisions by the professional societies to keep pace with the evolution of a diverse group of therapies. The more nimble and rapid methods for updating guidelines in HIV could inform processes for updating guidelines in HCV.

Research in HCV should include evaluations of screening, care and therapy in community healthcare clinics, drug treatment programmes and other settings providing care to persons at risk for HCV infection. Improved and expanded disease surveillance throughout the country is indicated to better understand trends in transmission and diagnosis. Serum biomarker assays to identify patients likely to achieve a successful treatment outcome early on should be incorporated into ongoing clinical trials of novel therapies. As novel approaches towards screening and treatment are developed, especially in rural or underserved settings, it will be important that outcomes be reported so that successful strategies can be imparted to others.

To understand the effects of cure on long-term health outcomes, endpoints other than SVR should be evaluated in clinical studies, and this is particularly true for confirmation of a reduction in the risk of development of hepatocellular carcinoma, liver failure and liver-related and overall mortality in patients with cirrhosis. Registries carefully noting those who achieved viral eradication would be

useful for charting areas of success as well as ongoing need. Such registries may also be helpful in identifying less frequent side effects not noted in the registration trials as well as outcomes in specific patient subgroups.

CONCLUSIONS

We have outlined the many challenges that lie ahead for healthcare providers as we attempt to reverse the rising morbidity and mortality associated with HCV. The new opportunities afforded by screening and improved diagnostics, education and treatment have created great excitement both in the medical community and in our patients, and the opportunities raise the prospect of eradicating HCV-related liver disease and eventually transmission. In the United States, HCV has all the attributes of an eradicable disease except sufficient public investment. Delivering care effectively, safely and broadly to all patient populations in an economically acceptable fashion must be our goal now and over the next 5–10 years.

ACKNOWLEDGEMENTS

The costs of this meeting were sponsored by Boehringer Ingelheim, Ingelheim, Germany; Gilead Sciences, Inc., Foster City, CA; Idenix Pharmaceuticals, Cambridge, MA; Janssen Research & Development, Raritan, NJ; Janssen Therapeutics, Titusville, NJ; Laboratory Corporation of America, Burlington, NC; Liver Institute for Education and Research, NJ; Novartis Pharmaceuticals Corporation, East Hanover, NJ; Quest Diagnostics, Madison, NJ; Roche Molecular Diagnostics, Pleasanton, CA; and Vertex Pharmaceuticals, Inc., Cambridge, MA. Jennifer King, PhD, helped draft the article.

FINANCIAL DISCLOSURES

The following authors disclose financial relationships with one or more meeting sponsors. Nezam H. Afdhal: research support from Abbott Laboratories, Bristol-Myers Squibb, Gilead Sciences, Inc., GlaxoSmithKline, Merck & Co, Inc., and Vertex Pharmaceuticals, Inc.; consultant or advisory arrangements with Boehringer Ingelheim, Echosens, Gilead Sciences, Inc., GlaxoSmithKline, Kadmon Corporation, Ligand Pharmaceuticals, Inc., Medgenics Inc., Merck & Co., Inc., Novartis Pharmaceutical Corporation, Spring Bank Pharmaceuticals, Inc., Quest Diagnostics, and Vertex Pharmaceuticals, Inc.; and stock ownership in Medgenics, Inc. and Spring Bank Pharmaceuticals, Inc. Ira M. Jacobson: advisor for AbbVie Pharmaceuticals, Achillion Pharmaceuticals, Inc., Boehringer Ingelheim, Bristol-Myers Squibb, Enanta Pharmaceuticals, Inc., Gilead Sciences, Inc., GlaxoSmithKline, Idenix Pharmaceuticals, Kadmon Corporation, Merck & Co., Inc., Novartis Pharmaceuticals Corporation, Genentech, Inc., Janssen Therapeutics, and Vertex

Pharmaceuticals, Inc.; consultant for Abbott Laboratories, Achillion Pharmaceuticals, Inc., Boehringer Ingelheim, Bristol-Myers Squibb, Enanta Pharmaceuticals, Inc., Gilead Sciences, Inc., GlaxoSmithKline, Idenix Pharmaceuticals, Kadmon Corporation, Merck & Co., Inc., Novartis Pharmaceutical Corporation, Genentech, Inc., Janssen Therapeutics, and Vertex Pharmaceuticals, Inc.; grant support from Abbott Laboratories, Achillion Pharmaceuticals, Inc., Boehringer Ingelheim, Bristol-Myers Squibb, Gilead Sciences, Inc., Merck & Co., Novartis Pharmaceutical Corporation, Pfizer Inc., Genentech, Inc., Janssen Therapeutics, and Vertex Pharmaceuticals, Inc.; and speakers bureau for Bristol-Myers Squibb, Gilead Sciences, Inc., Merck & Co., Genentech, Inc., and Vertex Pharmaceuticals, Inc. Stefan Zeuzem: honoraria and consultant fees from Abbott Laboratories, Achillion Pharmaceuticals, Inc., Boehringer Ingelheim, Bristol-Myers Squibb, Gilead Sciences, Inc., Idenix Pharmaceuticals, Janssen Therapeutics, Merck & Co., Novartis Pharmaceutical Corporation, Roche, Presidio Pharmaceuticals, Inc., Santaris Pharma A/S, and Vertex Pharmaceuticals, Inc.; and speakers bureau for Abbott Laboratories, Boehringer Ingelheim, Bristol-Myers Squibb, Gilead Sciences, Inc., Janssen Therapeutics, Merck & Co., and Vertex Pharmaceuticals, Inc. Robert Schooley: grant support from Boehringer Ingelheim and Bristol-Myers Squibb; consultant for Johnson & Johnson, Merck & Co., and Santaris Pharma A/S; board member for Gilead Sciences, Inc., GlobeImmune Inc., and Monogram Biosciences; stock ownership in GlobeImmune Inc. and Achillion Pharmaceuticals. David L. Thomas: grant support from Merck & Co., Inc.; consultant for Gilead Sciences, Inc., Janssen Therapeutics, and Merck & Co.; provisions of medicines, equipment, or administrative support from Gilead Sciences, Inc.; medications for Nitt Study from Merck & Co., Inc. Alain Litwin: consultant for Boehringer Ingelheim, Janssen Therapeutics, Kadmon Corporation, and Gilead Sciences, Inc.; grant support from Vertex Pharmaceuticals, Inc. Homie

Razavi: consultant for Abbott Laboratories, Boehringer Ingelheim, Gilead Sciences, Inc., Merck & Co., Inc., Novartis Pharmaceutical Corporation, and Vertex Pharmaceuticals, Inc. Laurent Castera: speakers bureau for Gilead Sciences, Inc., Janssen Therapeutics, and Merck & Co. Inc.; board member for Merck & Co., Inc., Bristol-Myers Squibb, and Gilead Sciences, Inc. Thierry Poynard: speakers bureau and consultant for Merck & Co., Inc.; stock ownership in BioPredictive. Andrew Muir: grant support from Abbott Laboratories, Achillion Pharmaceuticals, Bristol-Myers Squibb, Gilead Sciences, Inc., GlaxoSmithKline, GlobeImmune Inc., Medtronic, Inc., Merck & Co., Inc., Pfizer, Inc., Roche, Scynexis, Inc., and Vertex Pharmaceuticals, Inc.; consultant for Achillion Pharmaceuticals, Bristol-Myers Squibb, Gilead Sciences, Inc., Merck & Co., Inc., Profectus Biosciences, Inc., Scynexis, Inc., and Vertex Pharmaceuticals, Inc. Camilla Graham, former employee of Vertex Pharmaceuticals (left 9/28/12), Andrew Talal: grant support from Abbott Laboratories, Bristol-Myers Squibb, Boehringer Ingelheim, Roche, Gilead Sciences, Inc., Merck & Co., Inc., Tibotec Pharmaceuticals, Ltd., and Vertex Pharmaceuticals, Inc.; advisor for Abbott Diagnostics, Boehringer Ingelheim, Merck & Co., and Pfizer, Inc.; developed educational presentations for Medscape and Merck & Co., Inc. Mark Sulkowski: advisor and consultant for AbbVie Pharmaceuticals, Boehringer Ingelheim, Bristol-Myers Squibb, Gilead Sciences, Inc., Idenix Pharmaceuticals, Janssen Therapeutics, Merck & Co., Inc., Roche, and Vertex Pharmaceuticals; grants from AbbVie Pharmaceuticals, Boehringer Ingelheim, Bristol-Myers Squibb, Gilead Sciences, Inc., Janssen Therapeutics, Merck & Co., Inc., Roche, and Vertex Pharmaceuticals. Daniel Church, Lynda Dee, and Shruti Mehta have no potential conflicts to disclose. John Ward has no potential conflicts of interest to disclose; the findings and conclusions in this report are those of the author(s) and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

REFERENCES

- 1 Kwo PY, Lawitz EJ, McCone J *et al*. Efficacy of boceprevir, an NS3 protease inhibitor, in combination with peginterferon alfa-2b and ribavirin in treatment-naïve patients with genotype 1 hepatitis C infection (SPRINT-1): an open-label, randomised, multicentre phase 2 trial. *Lancet* 2010; 376: 705–716.
- 2 Poordad F, McCone J Jr, Bacon BR *et al*. Boceprevir for untreated chronic HCV genotype 1 infection. *N Engl J Med* 2011; 364: 1195–1206.
- 3 Jacobson IM, McHutchison JG, Dusheiko G *et al*. Telaprevir for previously untreated chronic hepatitis C virus infection. *N Engl J Med* 2011; 364: 2405–2416.
- 4 McHutchison JG, Everson GT, Gordon SC *et al*. Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009; 360: 1827–1838.
- 5 Fontaine H, Pol S. Antiviral activity of telaprevir and boceprevir for the treatment of hepatitis C virus infection in treatment-experienced patients. *Clin Res Hepatol Gastroenterol* 2011; 35(Suppl. 2): S59–S63.
- 6 Gane EJ, Stedman CA, Hyland RH *et al*. Nucleotide polymerase inhibitor sofosbuvir plus ribavirin for hepatitis C. *N Engl J Med* 2013; 368: 34–44.
- 7 Poordad F, Lawitz E, Kowdley KV *et al*. Exploratory study of oral combination antiviral therapy for hepatitis C. *N Engl J Med* 2013; 368: 45–53.
- 8 Jacobson IM, Gordon SC, Kowdley KV *et al*. Sofosbuvir for hepatitis C genotype 2 or 3 in patients without treatment options. *N Engl J Med* 2013; 368: 1867–1877.
- 9 Centers for Disease Control Prevention (CDC). Hepatitis C virus infection among adolescents and young

- adults: Massachusetts, 2002–2009. *MMWR Morb Mortal Wkly Rep* 2011; 60: 537–541.
- 10 Ngo-Metzger Q, Ward JW, Valdiserri RO. Expanded hepatitis C virus screening recommendations promote opportunities for care and cure. *Ann Intern Med* 2013; 159: 364–365.
 - 11 Kowdley KV, Lawitz E, Poordad F *et al.* Safety and efficacy of interferon-free regimens of ABT-450/r, ABT-267, ABT-333 +/- ribavirin in patients with chronic HCV GT1 infection: results from the AVIATOR Study. *J Hepatol* 2013; 58 (Suppl. S1): S2.
 - 12 Lawitz E, Ghalib R, Rodriguez-Torres M *et al.* Suppression of viral load through 4 weeks post-treatment results of a once-daily regimen of simeprevir + sofosbuvir with or without ribavirin in hepatitis C virus GT 1 null responders. *Program and Abstracts of the 20th Conference on Retroviruses and Opportunistic Infections* 2013; Abstract 155LB.
 - 13 Sulkowski MS, Gardiner DF, Rodriguez-Torres M *et al.* Sustained virologic response with daclatasvir plus sofosbuvir +/- ribavirin (RBV) in chronic HCV genotype (GT) 1-infected patients who previously failed telaprevir (TVR) or boceprevir (BOC). *J Hepatol* 2013; 58(Suppl. 1): S570.
 - 14 Sulkowski MS, Gardiner DF, Rodriguez-Torres M *et al.* High rate of sustained virologic response with the all-oral combination of daclatasvir (NS5A inhibitor) plus sofosbuvir (nucleotide NS5B inhibitor), with or without ribavirin, in treatment-naive patients chronically infected with HCV genotype 1, 2, or 3. *Hepatology* 2012; 56(Suppl. S1): Abstract LB-2.
 - 15 Zeuzem S, Soriano V, Asselah T *et al.* Faldaprevir and deleobuvir for HCV genotype 1 infection. *N Engl J Med* 2013; 369: 630–639.
 - 16 Ly KN, Xing J, Klevens RM, Jiles RB, Ward JW, Holmberg SD. The increasing burden of mortality from viral hepatitis in the United States between 1999 and 2007. *Ann Intern Med* 2012; 156: 271–278.
 - 17 Armstrong GL, Wasley A, Simard EP, McQuillan GM, Kuhnert WL, Alter MJ. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann Intern Med* 2006; 144: 705–714.
 - 18 Smith BD, Patel N, Beckett GA, Jewett A, Ward JW. Hepatitis C virus antibody prevalence, correlates and predictors among persons born from 1945 through 1965, United States, 1999–2008. *Hepatology* 2011; 54(Suppl. S1): Abstract 241.
 - 19 Recommendations for prevention and control of hepatitis C virus (HCV) infection and HCV-related chronic disease. Centers for Disease Control and Prevention. *MMWR Recomm Rep* 1998; 47(RR-19): 1–39.
 - 20 Ghany MG, Strader DB, Thomas DL, Seeff LB. Diagnosis, management, and treatment of hepatitis C: an update. *Hepatology* 2009; 49: 1335–1374.
 - 21 Dienstag JL, McHutchison JG. American Gastroenterological Association medical position statement on the management of hepatitis C. *Gastroenterology* 2006; 130: 225–230.
 - 22 Roblin DW, Smith BD, Weinbaum CM, Sabin ME. HCV screening practices and prevalence in an MCO, 2000–2007. *Am J Manag Care* 2011; 17: 548–555.
 - 23 Southern WN, Drainoni M-L, Smith BD *et al.* Hepatitis C testing practices and prevalence in a high-risk urban ambulatory care setting. *J Viral Hepat* 2011; 18: 474–481.
 - 24 Volk ML, Tocco R, Saini S, Lok ASF. Public health impact of antiviral therapy for hepatitis C in the United States. *Hepatology* 2009; 50: 1750–1755.
 - 25 Smith BD, Morgan RL, Beckett GA *et al.* Recommendations for the identification of chronic hepatitis C virus infection among persons born during 1945–1965. *MMWR Recomm Rep* 2012; 61: 1–32.
 - 26 Rein DB, Smith BD, Wittenborn JS *et al.* The cost-effectiveness of birth-cohort screening for hepatitis C antibody in U.S. primary care settings. *Ann Intern Med* 2012; 156: 263–270.
 - 27 Morgan RL, Baack B, Smith BD, Yartel A, Pitasi M, Falck-Ytter Y. Eradication of hepatitis C virus infection and the development of hepatocellular carcinoma: a meta-analysis of observational studies. *Ann Intern Med* 2013; 158: 329–337.
 - 28 Van der Meer AJ, Veldt BJ, Feld JJ *et al.* Association between sustained virological response and all-cause mortality among patients with chronic hepatitis C and advanced hepatic fibrosis. *JAMA* 2012; 308: 2584–2593.
 - 29 Moyer VA, on behalf of the U.S. Preventive Services Task Force. Screening for hepatitis C virus infection in Adults: U.S. preventive services task force recommendation statement. *Ann Intern Med* 2013; 159: 349–357.
 - 30 Eckman MH, Talal AH, Gordon SC, Schiff E, Sherman KE. Cost-effectiveness of screening for chronic hepatitis C infection in the United States. *Clin Infect Dis* 2013; 56: 1382–1393.
 - 31 McEwan P, Ward T, Yuan Y, Kim R, L'italien G. The impact of timing and prioritization on the cost-effectiveness of birth cohort testing and treatment for hepatitis C virus in the United States. *Hepatology* 2013; 58: 54–64.
 - 32 Litwin AH, Smith BD, Drainoni M-L *et al.* Primary care-based interventions are associated with increases in hepatitis C virus testing for patients at risk. *Dig Liver Dis* 2012; 44: 497–503.
 - 33 Centers for Disease Control Prevention (CDC). Testing for HCV infection: an update of guidance for clinicians and laboratorians. *MMWR Morb Mortal Wkly Rep* 2013; 62: 362–365.
 - 34 Smith BD, Teshale E, Jewett A *et al.* Performance of premarket rapid hepatitis C virus antibody assays in 4 national human immunodeficiency virus behavioral surveillance system sites. *Clin Infect Dis* 2011; 53: 780–786.
 - 35 Jewett A, Smith BD, Garfein RS, Cuevas-Mota J, Teshale EH, Weinbaum CM. Field-based performance of three pre-market rapid hepatitis C virus antibody assays in STAHR

- (Study to Assess Hepatitis C Risk) among young adults who inject drugs in San Diego, CA. *J Clin Virol* 2012; 54: 213–217.
- 36 Baranova A, Lal P, Biringdinc A, Younossi ZM. Non-Invasive markers for hepatic fibrosis. *BMC Gastroenterol* 2011; 11: 91.
 - 37 European Association for the Study of the Liver. EASL clinical practice guidelines: management of hepatitis C virus infection. *J Hepatol* 2011; 55: 245–264.
 - 38 Chou R, Wasson N. Blood tests to diagnose fibrosis or cirrhosis in patients with chronic hepatitis C virus infection: a systematic review. *Ann Intern Med* 2013; 158: 807–820.
 - 39 Lin Z-H, Xin Y-N, Dong Q-J *et al.* Performance of the aspartate aminotransferase-to-platelet ratio index for the staging of hepatitis C-related fibrosis: an updated meta-analysis. *Hepatology* 2011; 53: 726–736.
 - 40 Castéra L, Foucher J, Bernard P-H *et al.* Pitfalls of liver stiffness measurement: a 5-year prospective study of 13,369 examinations. *Hepatology* 2010; 51: 828–835.
 - 41 Friedrich-Rust M, Ong M-F, Martens S *et al.* Performance of transient elastography for the staging of liver fibrosis: a meta-analysis. *Gastroenterology* 2008; 134: 960–974.
 - 42 Degos F, Perez P, Roche B *et al.* Diagnostic accuracy of FibroScan and comparison to liver fibrosis biomarkers in chronic viral hepatitis: a multicenter prospective study (the FIBROSTIC study). *J Hepatol* 2010; 53: 1013–1021.
 - 43 Klibansky DA, Mehta SH, Curry M, Nasser I, Challies T, Afdhal NH. Transient elastography for predicting clinical outcomes in patients with chronic liver disease. *J Viral Hepat* 2012; 19: e184–e193.
 - 44 Vergniol J, Foucher J, Terrebbonne E *et al.* Noninvasive tests for fibrosis and liver stiffness predict 5-year outcomes of patients with chronic hepatitis C. *Gastroenterology* 2011; 140: 1970–1979.
 - 45 Castéra L, Vergniol J, Foucher J *et al.* Prospective comparison of transient elastography, Fibrotest, APRI, and liver biopsy for the assessment of fibrosis in chronic hepatitis C. *Gastroenterology* 2005; 128: 343–350.
 - 46 Poynard T, Ingiliz P, Elkrif L *et al.* Concordance in a world without a gold standard: a new non-invasive methodology for improving accuracy of fibrosis markers. *PLoS ONE* 2008; 3: e3857.
 - 47 Bonder A, Afdhal NH. Biopsy no more; changing the screening and diagnostic algorithm for hepatitis C. *Clin Gastroenterol Hepatol* 2013; 11: 309–310.
 - 48 Holmberg SD, Spradling PR, Moorman AC, Denniston MM. Hepatitis C in the United States. *N Engl J Med* 2013; 368: 1859–1861.
 - 49 Fishbein DA, Lo Y, Reinus JF, Gourevitch MN, Klein RS. Factors associated with successful referral for clinical care of drug users with chronic hepatitis C who have or are at risk for HIV infection. *J Acquir Immune Defic Syndr* 2004; 37: 1367–1375.
 - 50 Ong JP, Collantes R, Pitts A, Martin L, Sheridan M, Younossi ZM. High rates of uninsured among HCV-positive individuals. *J Clin Gastroenterol* 2005; 39: 826–830.
 - 51 Schaefer M, Capuron L, Friebe A *et al.* Hepatitis C infection, antiviral treatment and mental health: a European expert consensus statement. *J Hepatol* 2012; 57: 1379–1390.
 - 52 Miller L, Fluker S-A, Osborn M, Liu X, Strawder A. Improving access to hepatitis C care for urban, underserved patients using a primary care-based hepatitis C clinic. *J Natl Med Assoc* 2012; 104: 244–250.
 - 53 Morrill JA, Shrestha M, Grant RW. Barriers to the treatment of hepatitis C. Patient, provider, and system factors. *J Gen Intern Med* 2005; 20: 754–758.
 - 54 Mehta SH, Genberg BL, Astemborski J *et al.* Limited uptake of hepatitis C treatment among injection drug users. *J Community Health* 2008; 33: 126–133.
 - 55 Blasiole JA, Shinkunas L, Labrecque DR, Arnold RM, Zickmund SL. Mental and physical symptoms associated with lower social support for patients with hepatitis C. *World J Gastroenterol* 2006; 12: 4665–4672.
 - 56 Treloar C, Newland J, Rance J, Hopwood M. Uptake and delivery of hepatitis C treatment in opiate substitution treatment: perceptions of clients and health professionals. *J Viral Hepat* 2010; 17: 839–844.
 - 57 Khokhar OS, Lewis JH. Reasons why patients infected with chronic hepatitis C virus choose to defer treatment: do they alter their decision with time? *Dig Dis Sci* 2007; 52: 1168–1176.
 - 58 Zickmund SL, Brown KE, Bielefeldt K. A systematic review of provider knowledge of hepatitis C: is it enough for a complex disease? *Dig Dis Sci* 2007; 52: 2550–2556.
 - 59 Shehab TM, Sonnad SS, Lok AS. Management of hepatitis C patients by primary care physicians in the USA: results of a national survey. *J Viral Hepat* 2001; 8: 377–383.
 - 60 Kanwal F, Hoang T, Spiegel BMR *et al.* Predictors of treatment in patients with chronic hepatitis C infection – role of patient versus nonpatient factors. *Hepatology* 2007; 46: 1741–1749.
 - 61 Stoové MA, Gifford SM, Dore GJ. The impact of injecting drug use status on hepatitis C-related referral and treatment. *Drug Alcohol Depend* 2005; 77: 81–86.
 - 62 Mitchell AE, Colvin HM, Palmer Beasley R. Institute of medicine recommendations for the prevention and control of hepatitis B and C. *Hepatology* 2010; 51: 729–733.
 - 63 Volk ML. Antiviral therapy for hepatitis C: why are so few patients being treated? *J Antimicrob Chemother* 2010; 65: 1327–1329.
 - 64 Arora S, Thornton K, Murata G *et al.* Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med* 2011; 364: 2199–2207.
 - 65 Razavi H, El Khoury A, Elbasha E *et al.* Chronic hepatitis C virus (HCV) disease burden and cost in the United States. *Hepatology* 2013; 57: 2164–2170.
 - 66 McAdam-Marx C, McGarry LJ, Hane CA, Biskupiak J, Deniz B, Brixner DI. All-cause and incremental per patient per year cost

- associated with chronic hepatitis C virus and associated liver complications in the United States: a managed care perspective. *J Manag Care Pharm* 2011; 17: 531–546.
- 67 Di Bisceglie AM, Lyra AC, Schwartz M *et al.* Hepatitis C-related hepatocellular carcinoma in the United States: influence of ethnic status. *Am J Gastroenterol* 2003; 98: 2060–2063.
- 68 Davis GL, Albright JE, Cook SF, Rosenberg DM. Projecting future complications of chronic hepatitis C in the United States. *Liver Transpl* 2003; 9: 331–338.
- 69 El Khoury AC, Klimack WK, Wallace C, Razavi H. Economic burden of hepatitis C-associated diseases in the United States. *J Viral Hepat* 2012; 19: 153–160.
- 70 Jacobson IM, Cacoub P, Dal Maso L, Harrison SA, Younossi ZM. Manifestations of chronic hepatitis C virus infection beyond the liver. *Clin Gastroenterol Hepatol* 2010; 8: 1017–1029.
- 71 Gellad ZF, Reed SD, Muir AJ. Economic evaluation of direct-acting antiviral therapy in chronic hepatitis C. *Antivir Ther* 2012; 17(6 Pt B): 1189–1199.
- 72 Liu S, Cipriano LE, Holodniy M, Owens DK, Goldhaber-Fiebert JD. New protease inhibitors for the treatment of chronic hepatitis C: a cost-effectiveness analysis. *Ann Intern Med* 2012; 156: 279–290.
- 73 Bichoupan K, Martel-Laferriere V, Ng M *et al.* Real world costs of telaprevir-based triple therapy, including costs of managing adverse events, at the Mount Sinai Medical Center, NY: \$147,000 per EOT. *J Hepatol* 2013; 58(Suppl. 1): S324–S325.
- 74 Hallinan R, Byrne A, Agho K, Dore GJ. Referral for chronic hepatitis C treatment from a drug dependency treatment setting. *Drug Alcohol Depend* 2007; 88: 49–53.
- 75 Jowett SL, Agarwal K, Smith BC *et al.* Managing chronic hepatitis C acquired through intravenous drug use. *QJM* 2001; 94: 153–158.
- 76 Dimova RB, Zeremski M, Jacobson IM, Hagan H, Des Jarlais DC, Talal AH. Determinants of hepatitis C virus treatment completion and efficacy in drug users assessed by meta-analysis. *Clin Infect Dis* 2013; 56: 806–816.
- 77 Harris KA Jr, Arnsten JH, Litwin AH. Successful integration of hepatitis C evaluation and treatment services with methadone maintenance. *J Addict Med* 2010; 4: 20–26.
- 78 Martinez AD, Dimova R, Marks KM *et al.* Integrated internist – addiction medicine – hepatology model for hepatitis C management for individuals on methadone maintenance. *J Viral Hepat* 2012; 19: 47–54.
- 79 Sylvestre DL, Zweben JE. Integrating HCV services for drug users: a model to improve engagement and outcomes. *Int J Drug Policy* 2007; 18: 406–410.
- 80 Alavi M, Grebely J, Micallef M *et al.* Assessment and treatment of hepatitis C virus infection among people who inject drugs in the opioid substitution setting: ETHOS study. *Clin Infect Dis* 2013; 57(Suppl. 2): S62–S69.
- 81 Bruggman P, Litwin AH. Models of care for the management of hepatitis C virus among people who inject drugs: one size does not fit all. *Clin Infect Dis* 2013; 57(Suppl. 2): S56–S61.
- 82 Evon DM, Simpson K, Kixmiller S *et al.* A randomized controlled trial of an integrated care intervention to increase eligibility for chronic hepatitis C treatment. *Am J Gastroenterol* 2011; 106: 1777–1786.
- 83 Moussalli J, Delaquaize H, Boubilley D *et al.* Factors to improve the management of hepatitis C in drug users: an observational study in an addiction centre. *Gastroenterol Res Pract* 2010; 2010: article ID 261472.
- 84 Robaey G, Grebely J, Mauss S *et al.* Recommendations for the management of hepatitis C virus infection among people who inject drugs. *Clin Infect Dis* 2013; 57(Suppl. 2): S129–S137.
- 85 Martin NK, Hickman M, Hutchinson SJ, Goldberg DJ, Vickerman P. Combination interventions to prevent HCV transmission among people who inject drugs: modeling the impact of antiviral treatment, needle and syringe programs, and opiate substitution therapy. *Clin Infect Dis* 2013; 57(Suppl. 2): S39–S45.
- 86 Hellard ME, Jenkinson R, Higgs P *et al.* Modelling antiviral treatment to prevent hepatitis C infection among people who inject drugs in Victoria, Australia. *Med J Aust* 2012; 196: 638–641.
- 87 Fontaine H, Hezode C, Dorival C *et al.* SVR12 rates and safety of triple therapy including telaprevir or boceprevir in 221 cirrhotic non responders treated in the French Early Access Program (ANRS CO20-CUPIC). *J Hepatol* 2013; 58(Suppl. 1): S27.
- 88 Lawitz E, Wyles D, Davis M *et al.* Sofosbuvir + peginterferon + ribavirin for 12 weeks achieves 90% SVR12 in genotype 1, 4, 5, or 6 HCV infected patients: The Neutrino Study. *J Hepatol* 2013; 58(Suppl. 1): S567.
- 89 Soriano V, Gane EJ, Angus PW *et al.* Efficacy and safety of the interferon (IFN)-free combination of BI 201335+ BI 207127+/- ribavirin (RBV) in treatment-naive patients with HCV genotype (GT) 1 infection and compensated liver cirrhosis: results from the SOUND-C2 study. *Hepatology* 2012; 56(Suppl. S1): 234A.
- 90 Butt AA, Justice AC, Skanderson M, Good C, Kwoh CK. Rates and predictors of hepatitis C virus treatment in HCV-HIV-coinfected subjects. *Aliment Pharmacol Ther* 2006; 24: 585–591.
- 91 Torriani FJ, Rodriguez-Torres M, Rockstroh JK *et al.* Peginterferon Alfa-2a plus ribavirin for chronic hepatitis C virus infection in HIV-infected patients. *N Engl J Med* 2004; 351: 438–450.
- 92 Sulkowski MS, Sherman KE, Dieterich DT *et al.* Combination therapy with telaprevir for chronic hepatitis C Virus genotype 1 infection in patients with HIV: a randomized trial. *Ann Intern Med* 2013; 159: 86–96.
- 93 Sulkowski M, Pol S, Mallolas J *et al.* Boceprevir versus placebo with pegylated interferon alfa-2b and ribavirin for treatment of hepatitis C virus genotype 1 in patients with HIV: a randomised, double-blind,

- controlled phase 2 trial. *Lancet Infect Dis* 2013; 13: 597–605.
- 94 Dieterich D, Soriano V, Nelson M *et al.* STARTVerso 4: High rates of early virologic response in hepatitis C virus genotype 1/HIV co-infected patients treated with fladaprevir + pegylated interferon and ribavirin. *Program and Abstracts of the 20th Conference on Retroviruses and Opportunistic Infections* 2013 Mar 3; Abstract 40LB.
- 95 Dieterich D, Rockstroh JK, Orkin C *et al.* Simeprevir with pegylated interferon/ribavirin in patients co-infected with chronic hepatitis C virus and HIV-1: Week-24 interim analysis of the TMC435-C212 Study. *Program and Abstracts of the 20th Conference on Retroviruses and Opportunistic Infections* 2013 Mar 3; Abstract 154LB.
- 96 Limketkai BN, Mehta SH, Sutcliffe CG *et al.* Relationship of liver disease stage and antiviral therapy with liver-related events and death in adults coinfecting with HIV/HCV. *JAMA* 2012; 308: 370–378.
- 97 Ghany MG, Nelson DR, Strader DB, Thomas DL, Seeff LB. An update on treatment of genotype 1 chronic hepatitis C virus infection: 2011 practice guideline by the American Association for the Study of Liver Diseases. *Hepatology* 2011; 54: 1433–1444.
- 98 Everson GT, Sims KD, Rodriguez-Torres M *et al.* An interferon-free, ribavirin-free 12-week regimen of daclatasvir (DCV), asunaprevir (ASV), and BMS-791325 yielded SVR4 of 94% in treatment-naive patients with genotype (GT) 1 chronic hepatitis C virus (HCV) infection. *Hepatology* 2012; 56 (Suppl. S1): Abstract LB-3.
- 99 Gane EJ, Pockros P, Zeuzem S *et al.* Interferon-free treatment with a combination of mericitabine and danoprevir/r with or without ribavirin in treatment-naive HCV genotype 1-infected patients. *J Hepatol* 2012; 56(Suppl. 2): S555–S556.
- 100 Gane EJ, Stedman CA, Hyland RH, Ding X, Pang PS, Symonds WT. ELECTRON: 100% SVR rates for once-daily sofosbuvir plus ledipasvir plus ribavirin given for 12 weeks in treatment-naive and previously treated patients with HCV genotype 1. *Program and Abstracts of the 20th Conference on Retroviruses and Opportunistic Infections* 2013 Mar 3; Abstract 41LB.
- 101 Gane E, Lawitz E, Rodriguez-Torres M *et al.* Phase 3 randomized controlled trial of all-oral treatment with sofosbuvir + ribavirin for 12 weeks compared to 24 weeks of PEG + ribavirin in treatment-naive GT2/3 HCV-infected patients. *J Hepatol* 2013; 58(Suppl. 1): S3.
- 102 Lok AS, Gardiner DF, Lawitz E *et al.* Preliminary study of two antiviral agents for hepatitis C genotype 1. *N Engl J Med* 2012; 366: 216–224.
- 103 Suzuki F, Ikeda K, Toyota J *et al.* Dual oral therapy with the NS5A inhibitor daclatasvir (BMS-790052) and NS3 protease inhibitor asunaprevir (BMS-650032) in HCV genotype 1B-infected null responders or ineligible/intolerant to peginterferon/ribavirin. *J Hepatol* 2012; 56(Suppl. 2): S7–S8.
- 104 Tsochatzis EA, Gurusamy KS, Ntaoula S, Cholongitas E, Davidson BR, Burroughs AK. Elastography for the diagnosis of severity of fibrosis in chronic liver disease: a meta-analysis of diagnostic accuracy. *J Hepatol* 2011; 54: 650–659.
- 105 Adeyemi OM, Jensen D, Attar B *et al.* Hepatitis C treatment eligibility in an urban population with and without HIV coinfection. *AIDS Patient Care STDS* 2004; 18: 239–245.
- 106 Falck-Ytter Y, Kale H, Mullen KD, Sarbah SA, Sorescu L, McCullough AJ. Surprisingly small effect of antiviral treatment in patients with hepatitis C. *Ann Intern Med* 2002; 136: 288–292.
- 107 Fleming CA, Craven DE, Thornton D, Tumilty S, Nunes D. Hepatitis C virus and human immunodeficiency virus coinfection in an urban population: low eligibility for interferon treatment. *Clin Infect Dis* 2003; 36: 97–100.
- 108 Grebely J, Raffa JD, Lai C *et al.* Low uptake of treatment for hepatitis C virus infection in a large community-based study of inner city residents. *J Viral Hepat* 2009; 16: 352–358.
- 109 Groom H, Dieperink E, Nelson DB *et al.* Outcomes of a Hepatitis C screening program at a large urban VA medical center. *J Clin Gastroenterol* 2008; 42: 97–106.
- 110 Hall CS, Charlebois ED, Hahn JA, Moss AR, Bangsberg DR. Hepatitis C virus infection in San Francisco's HIV-infected urban poor. *J Gen Intern Med* 2004; 19: 357–365.
- 111 Mehta SH, Lucas GM, Mirel LB *et al.* Limited effectiveness of antiviral treatment for hepatitis C in an urban HIV clinic. *AIDS* 2006; 20: 2361–2369.
- 112 Restrepo A, Johnson TC, Widjaja D *et al.* The rate of treatment of chronic hepatitis C in patients co-infected with HIV in an urban medical centre. *J Viral Hepat* 2005; 12: 86–90.
- 113 Rocca LG, Yawn BP, Wollan P, Kim WR. (2004) Management of patients with hepatitis C in a community population: diagnosis, discussions, and decisions to treat. *Ann Fam Med* 2: 116–124.

APPENDIX 1

MEETING PARTICIPANTS

Nezam H. Afdhal, MD, and Rachel Baden, MD, Harvard Medical School, Beth Israel Deaconess Medical Center, Boston, MA; Cynthia Beam, PhD, and Julie Krop, MD, Vertex Pharmaceuti-

icals, Inc., Cambridge, MA; Clifford Brass, MD, Novartis Pharmaceuticals Corporation, East Hannover, NJ; David Brumley, MD, MBA Tufts Health Plan Watertown, MA; Laurent Castera, MD, PhD, Hôpital Beaujon, Clichy, France; Daniel Church, MD, MPH, and Alfred DeMaria, Jr., MD,

Massachusetts Department of Public Health, Jamaica Plain, MA; Kevin Cranston, MDiv, Massachusetts Health and Human Services, Boston, MA; Lynda Dee, JD, Fair Pricing Coalition, Baltimore, MD; Arnold Doyle, Janssen Therapeutics, Titusville, NJ; Julia Engstrom-Melnyk, PhD, Roche

Diagnostic Corporation, Indianapolis, IN; Michael Fried, MD, University of North Carolina at Chapel Hill, Chapel Hill, NC; Camilla Graham, MD, MPH, and Michael Wong, MD, Beth Israel Deaconess Medical Center, Boston, MA; William Guyer, PharmD, John G. McHutchison, MD, Joe Steele, MS, and William Symonds, PharmD, Gilead Sciences, Inc., Foster City, CA; Ira M. Jacobson, MD, Weill Medical College of Cornell University, New York, NY; Alain Litwin, MD, MPH, Albert Einstein College of Medicine of Yeshiva University, Valley Cottage, NY; Anthony Martinez, MD, and Andrew Talal, MD, MPH, University at Buffalo, Buffalo, NY; Henry Masur, MD, National Institutes of Health, Bethesda, MD; Douglas Mayers, MD, and Ronald C. Renaud, Jr., Idenix Pharmaceuticals, Inc., Cambridge,

MA; Shruti H. Mehta, PhD, MPH, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD; Andrew Muir, MD, MHS, Duke Clinical Research Institute, Durham, NC; Christopher J. Myatt, PhD, MBio Diagnostics, Inc., Boulder, CO; Stanley J. Nades, MD, and Rick Pesano, MD, PhD, Quest Diagnostics, San Juan Capistrano, CA; William O'Brien, MD, MS, Boehringer Ingelheim Pharmaceuticals, Ridgefield, CT; Christos J. Petropoulos, BS, PhD, Monogram Biosciences, A Labcorp Specialty Testing Group, South San Francisco, CA; Gastón R. Picchio, PhD, Janssen of Johnson and Johnson, Titusville, NJ; Thierry Poinard, MD, PhD, Hospitalier Pitié-Salpêtrière, Paris, France; Avi Prabhakar, MD, Janssen Services, LLC, Titusville, NJ; Kellie A. Rademacher, PharmD,

Prime Therapeutics, LLC, Eagan, MN; Homie Razavi, PhD, MBA, Center for Disease Analysis, Louisville, CO; Gary Rice, Diplomat Specialty Pharmacy, Flint, MI; Robert Schooley, MD, and David Wyles, MD, UC San Diego School of Medicine, La Jolla, CA; Joseph Sebastian, PhD, LabCorp, Research Triangle Park, NC; Bryce D. Smith, PhD, and John Ward, MD, Centers for Disease Control and Prevention, Atlanta, GA; Mark Sulkowski, MD, and David Thomas, MD, MPH, The Johns Hopkins University School of Medicine, Baltimore, MD; Chris Taylor, National Alliance of State and Territorial AIDS Directors (NASTAD), Washington, DC; Stefan Zeuzem, MD, J.W. Goethe University Hospital, Frankfurt, Germany.