

UC Davis

UC Davis Previously Published Works

Title

A pharmacist-led community-based survey study: Determining the impact of the Covid-19 pandemic on actionable factors associated with worse cancer outcomes and cancer health disparities

Permalink

<https://escholarship.org/uc/item/77g5w1bq>

Authors

Titus-Lay, Erika

Nehira, Jeffrey

Courtney, Jennifer

et al.

Publication Date

2023-09-01

DOI

10.1016/j.rcsop.2023.100311


Peer reviewed



ELSEVIER

Contents lists available at ScienceDirect

Exploratory Research in Clinical and Social Pharmacy

journal homepage: www.elsevier.com/locate/rcsop

A pharmacist-led community-based survey study: Determining the impact of the Covid-19 pandemic on actionable factors associated with worse cancer outcomes and cancer health disparities

Erika Titus-Lay^a, Jeffrey Nehira^a, Jennifer Courtney^a, Jacquelyn Jee^a, Marissa Kumar^a, Jenny Tiet^a, Vivi Le^a, Blythe Durbin-Johnson^b, Moon S. Chen Jr^c, Ruth Vinall^{a,*}

^a California Northstate University College of Pharmacy (CNUCOP), Elk Grove, CA 95757, USA

^b Department of Public Health Sciences, School of Medicine, University of California Davis, One Shields Avenue, Davis, CA 95616, USA

^c Division of Hematology and Oncology, UC Davis School of Medicine, Sacramento, CA 95817, USA

ARTICLE INFO

Keywords:

Community health
Covid-19
Cancer outcomes
Cancer health disparities
Pharmacist-led study

ABSTRACT

Purpose: The goals of this cross-sectional community-based survey study were to assess the impact of the Covid-19 pandemic on actionable factors which are known to contribute to worse cancer outcomes, and to determine whether race and ethnicity-based differences exist.

Methods: A survey study which captured demographic information and changes in cancer outcomes-related factors since the start of the Covid-19 pandemic, was conducted at a public Covid-19 vaccination clinic over a period of 10 days during March 2021. Surveys were administered in multiple languages. Chi-square tests and ANOVA followed by post-hoc Dunnett testing assessed for race and ethnicity-based differences.

Results: A total of 949 people participated (61.6% participation rate). Ninety-three surveys were removed based on inclusion criteria giving a final participant number of 856. Many participants reported postponing cancer screenings (17.8%) and cancellation of medical appointments (22.8% and 25.8% reported cancelled appointments by providers or themselves, respectively) due to the pandemic. Participants also reported decreased physical activity (44.7%) and increased tobacco and/or marijuana usage (7.0%). Conversely, participants reported consuming more fruits and vegetables (21.4%) and decreasing alcohol consumption (21.4%). Several race-related differences but no ethnicity-related differences were observed.

Conclusion: Our data can be used to help guide pharmacist-led targeted outreach in our community which will help mitigate Covid-19 pandemic-driven changes in behaviors associated with worse cancer outcomes and exacerbation of cancer health disparities. To our knowledge, this is the first cancer outcomes-related study to be conducted at a public Covid-19 vaccination site and is the first pharmacist-led study in this area.

* Corresponding author.

E-mail address: rvinall@cnsu.edu (R. Vinall).

<https://doi.org/10.1016/j.rcsop.2023.100311>

Received 5 May 2023; Received in revised form 12 July 2023; Accepted 12 July 2023

Available online 13 July 2023

2667-2766/© 2023 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

There are multiple factors which are known to be associated with worse cancer outcomes and many of these are actionable, i.e. it is possible to put measures in place to help mitigate them.^{1,2} For example, it is well known that there is a strong association between cancer screening rates and cancer outcomes³⁻⁵ and that tobacco usage increases overall cancer risk.⁶⁻⁸ Tobacco usage is the number one actionable factor associated with cancer deaths and it has been associated with worse outcomes for oral, lung, liver, stomach, kidney, bladder, colorectal, uterine, pancreatic cancers.⁹ Decreased physical exercise, increased alcohol consumption, and poor diet have also been widely recognized as actionable factors which are associated with several cancer types.¹⁰⁻¹³ Many of these factors have been shown to be more prevalent in minority populations and this can contribute to cancer health disparities.¹⁴ For example, it has been reported that prevalence of physical inactivity outside of work is highest in Asian adults.¹⁵ Resulting disparities are reflected by differences in the following common measures of cancer outcomes within minority communities: cancer incidence, prevalence, stage at diagnosis, mortality, survival, morbidity, survivorship and quality of life after cancer treatment, financial burden.¹⁶

Data from several national studies indicates that the Covid-19 pandemic may have negatively impacted factors which contribute to worse cancer outcomes and that this negative impact may have been greater in minority populations.¹⁷⁻²⁵ However, there is a lack of community-based studies which have looked at contributing factors. Community-based studies are important because data from these can help support and inform subsequent community-based outreach efforts which can be targeted to specific populations. Providing effective outreach and education are essential components of any wellness program and have been shown to be successful in helping reduce cancer morbidity and mortality rates.^{26,27} For example, smoking cessation programs have helped support an over 70% reduction in cigarette smoking since 1965.²⁸

Pharmacists are well positioned to play an important role in community outreach efforts based on their training, experience, and penetrance and accessibility in all communities and there are several examples of successful outreach studies involving pharmacists.^{27,29-35} Pharmacists already play a key role in helping mitigate worse cancer outcomes by providing smoking cessation counseling and dispensing smoking cessation medications, and studies have shown this has been effective in reducing smoking rates³⁰⁻³². All licensed pharmacists can provide education and help patients choose from over-the-counter nicotine replacement therapy options available for smoking cessation. In many states, including California, pharmacists with advanced training and working under a collaborative practice agreement or state protocol can authorize prescription medications for smoking cessation. Similarly, pharmacists with advanced training can also authorize prescription medications for alcohol cessation. Pharmacists also have training in motivational interviewing which can be utilized to encourage healthier habits through behavior change.³⁶ It is important for pharmacists to have this training and these skillsets because based on their accessibility and penetrance within communities they often serve as the initial healthcare professional to community members to recognize signs of early cancer and/or cancer risk factors. This is particularly true within underserved communities where many minority populations live. The potential exists for pharmacists to serve a bigger role in cancer control through extending counseling services relating to other actionable factors and this is supported by several studies and opinion pieces.³⁷⁻³⁹ An expanded role in this area also aligns with the Surgeon General's 2018 call to action relating to community health and prosperity.⁴⁰

The primary goals of the current study were to determine the impact of the Covid-19 pandemic on actionable factors which are known to contribute to worse cancer outcomes and to determine whether race and ethnicity-based differences exist in our community. People attending a public Covid-19 vaccination clinic at California Northstate University in Elk Grove (Sacramento County, California) were recruited to participate in a paper-based survey study which captured zip code, race/ethnicity, gender, age, income and education metrics as well as actionable factors associated with worse cancer outcomes (postponement of healthcare visits and cancer screenings, tobacco/marijuana and alcohol usage, consumption of fruits/vegetables, level of physical activity). The study was pharmacist-led. Pharmacy students as well as students from other healthcare profession programs were engaged in the distribution and collection of surveys under faculty member supervision. Our long-term goal is for our pharmacy faculty and students, and others, to use this data to provide pharmacist-led targeted outreach within our community and beyond and to thereby help improve cancer outcomes. To our knowledge, this is the first cancer outcomes-related survey study to be conducted at a Covid-19 vaccination site and is the first reported pharmacist-led study relating to this subject area.

2. Methods

This is a cross-sectional research survey study. Data for this research survey study was obtained from people attending a public Covid-19 vaccination clinic at California Northstate University in Elk Grove (Sacramento County, California) over a period of 10 days during March 2021. Anyone eligible to receive the Covid-19 vaccine per CDC guidelines during this timeframe was able to register for the clinic, there were no restrictions based on healthcare membership or coverage. Clinic registration was made available through the CalVax system. Based on registration numbers for the clinic and an anticipated 70% participation rate, we estimated that we would be able to collect 1000 completed surveys during the given timeframe. Convenience sampling was employed for this study: all Covid-19 clinic attendees were provided with a survey and invited to participate in the survey study after they had registered at the clinic. An oral overview of the research study was provided to all attendees following clinic registration and it was made clear that participation in the study was voluntary: attendees were given the option to leave their blank survey at their table if they did not want to participate in the study. If they did want to participate, they were asked to provide written consent and leave their completed survey at their table. The following inclusion criterion was applied: age 18 or older. Participants who completed the survey did not receive any compensation. This study was approved by the CNU IRB committee (protocol #2103-01-20).

The survey used in this study is a modified version of the survey developed by Scarinici et al.⁴¹ (Appendix 1). The original survey

was developed as part of a collaboration between 17 National Institutes of Health National Cancer Institute-funded Comprehensive Cancer Centers. Unlike Scarinici et al., we did not capture perceived susceptibility to Covid-19 infection or perceptions of prevention measures. Areas that were assessed in this study included impact of the Covid-19 pandemic on cancer prevention behaviors (tobacco and/or marijuana usage, alcohol consumption, fruit and vegetable consumption, postponement of cancer screenings, access to healthcare, physical activity, access to cancer-specific healthcare, access to medications) and cancer survivor behaviors (access and adherence to treatment, including medications). Demographic data (age, race, ethnicity, income bracket, education level) were also captured.

Table 1
Participant characteristics, by race.

Characteristic	Participants, No. (%)						P-value (ANOVA or Chi Squared)
	Total (N = 856 [100%])	White (n = 411 [48.0%])	Black or African American (n = 43 [5.0%])	American Indian or Alaskan Native (n = 7 [0.7%])	Asian or Asian American (n = 254 [29.7%])	Native Hawaiian, other Pacific Islander (n = 16 [1.9%])	
Gender							0.667
Male	334 (39.0)	156 (38.0)	18 (41.9)	3 (42.9)	106 (41.7)	4 (25.0)	
Female	515 (60.2)	256 (61.6)	24 (55.8)	4 (57.1)	165 (57.1)	11 (68.8)	
Prefer not to answer	6 (0.7)	2 (0.5)	1 (2.3)	0 (0)	2 (0.8)	1 (6.3)	
Age, median (range)	46.8 (18.1, 95.5)	49.5 (18.1, 95.5)	51.3 (23.9, 83.9)	45.9 (29.0, 76.3)	45.2 (19.7, 90.3)	44.3 (20.6, 58.8)	
Diagnosed with cancer during their lifetime	55 (6.4)	35 (8.4)	3 (7.0)	0 (0)	12 (4.7)	0 (0)	0.327
General health							0.835
Excellent	208 (24.3)	100 (24.3)	7 (16.3)	2 (28.6)	66 (26.0)	4 (25.0)	
Very good	358 (41.8)	179 (43.6)	15 (34.9)	4 (57.1)	100 (39.4)	7 (43.8)	
Good	241 (28.2)	108 (26.3%)	18 (41.9)	1 (14.3)	73 (28.7)	5 (31.3)	
Fair	41 (4.8)	19 (4.6)	3 (7.0)	0 (0)	15 (5.9)	0 (0)	
Poor	3 (0.4)	1 (0.2)	0 (0)	0 (0)	0 (0)	0 (0)	
Annual income							<0.001
\$0 to \$9999	25 (2.9)	5 (1.2%)	1 (2.3%)	0 (0%)	16 (6.3%)	1 (6.3%)	
\$10,000 to \$14,999	15 (1.8)	4 (1.0%)	0 (0%)	0 (0%)	10 (3.9%)	0 (0%)	
\$15,000 to \$19,999	12 (1.4)	2 (0.5%)	1 (2.3%)	0 (0%)	8 (3.1%)	0 (0%)	
\$20,000 to \$34,999	66 (7.7)	23 (5.6%)	3 (7.0%)	0 (0%)	27 (10.6%)	3 (18.8%)	
\$35,000 to \$49,999	65 (7.6)	25 (6.1%)	3 (7.0%)	0 (0%)	26 (10.2%)	2 (12.5%)	
\$50,000 to \$74,999	117 (13.7)	51 (12.4%)	9 (20.9%)	3 (42.9%)	35 (13.8%)	0 (0%)	
\$75,000 to \$99,999	119 (13.9)	59 (14.4%)	8 (18.6%)	1 (14.3%)	29 (11.4%)	3 (18.8%)	
\$100,000 to \$199,999	235 (27.5)	138 (33.6%)	13 (30.2%)	2 (28.6%)	50 (19.7%)	3 (18.8%)	
\$200,000 or more	82 (9.6)	56 (13.6%)	5 (11.6%)	0 (0%)	11 (4.3%)	1 (6.3%)	
Education							<0.001
Less than high school	15 (1.8)	5 (1.2%)	0 (0%)	0 (0%)	8 (3.1%)	0 (0%)	
Some high school, no diploma	10 (1.2)	1 (0.2%)	0 (0%)	0 (0%)	8 (3.1%)	0 (0%)	
GED	2 (0.2)	0 (0%)	0 (0%)	2 (28.6%)	0 (0%)	0 (0%)	
High school graduate	56 (6.5)	17 (4.1%)	2 (4.7%)	0 (0%)	28 (11.0%)	1 (6.3%)	
Some college but no degree	126 (14.7)	56 (13.6%)	10 (23.3%)	1 (14.3%)	38 (15.0%)	3 (18.8%)	
Associate degree- occupational/ vocational	35 (4.1)	17 (4.1%)	1 (2.3%)	0 (0%)	11 (4.3%)	1 (6.3%)	
Associate degree- academic program	41 (4.8)	17 (4.1%)	2 (4.7%)	0 (0%)	12 (4.7%)	2 (12.5%)	
Bachelor's degree (e. g., BA, AB, BS)	319 (37.3)	159 (38.7%)	13 (30.2%)	2 (28.6%)	98 (38.6%)	5 (31.3%)	
Master's degree (e.g., MA, MS, Meng, Med, MSW)	175 (20.4)	108 (26.3%)	12 (27.9%)	1 (14.3%)	23 (9.1%)	3 (18.8%)	
Professional school degree (e.g., MD, DDS, DVM, JD)	22 (2.6)	10 (2.4%)	1 (2.3%)	0 (0%)	8 (3.1%)	0 (0%)	
Doctorate degree (e. g., PhD, EdD)	23 (2.7)	13 (3.2%)	1 (2.3%)	0 (0%)	6 (2.4%)	0 (0%)	

For our study, the survey was administered in paper format. Participants were offered paper questionnaires in the following languages: English, Chinese, Hmong, Vietnamese. Faculty members and pharmacy students who are native speakers for each language translated the surveys; each translated survey was cross-checked by at least one other native speaker. Data from paper surveys was manually entered into SurveyMonkey software. The data were extracted and then R software was used for all subsequent data analyses. Analyses were conducted using R version 4.2.1.⁴² Participant responses were compared between races or ethnicities using chi-square tests. Table cells (race/response combinations) with results differing from what would be expected under independence were identified using adjusted standardized residuals.⁴³ Age was compared between groups using ANOVA, followed by post-hoc Dunnett testing comparing White to other races. Other and unknown races/ethnicities were not included in statistical testing, nor were responses of “Other”, “Prefer not to answer”, “Don't know/not sure”, or not applicable responses.

3. Results

3.1. Participant demographics

A total of 949 people out of an estimated 1540 people attending a public Covid-19 vaccination clinic at California Northstate University in Elk Grove (Sacramento County, California) participated in this survey study over a period of 10 days during March 2021 (61.6% participation rate). Ninety-three completed surveys were removed based on our inclusion criteria (age = 18 or older) giving a final participant number of 856. Median age was 47.1 [18.1–95.5] years; 411 White participants (48.0%), 43 Black/African American participants (5.0%), 254 Asian/Asian American participants (29.7%), 16 Native Hawaiian/other Pacific Islander participants (1.9%), 7

Table 2
Participant characteristics, by ethnicity.

Characteristic	Participants, No. (%)			P value (ANOVA or Chi Squared)
	Total (N = 856)	Hispanic (n = 129 [15.1%])	Non-Hispanic (n = 720 [84.4%])	
Gender				0.0118
Male	334 (39.0)	37 (28.7)	294 (40.8)	
Female	515 (60.2)	91 (70.5)	420 (58.3)	
Prefer not to answer	6 (0.7)	1 (0.8)	5 (0.7)	
Age, median (range)	47.1 (18.1, 95.5)	42.6 (18.6, 95.5)	47.7 (18.1, 90.3)	0.0196
Diagnosed with cancer during their lifetime	55 (6.4)	3 (2.3)	52 (7.2)	0.0607
General health				0.899
Excellent	208 (24.3)	30 (23.3%)	177 (24.6%)	
Very good	358 (41.8)	53 (41.1%)	303 (42.1%)	
Good	241 (28.2)	35 (27.1%)	203 (28.2%)	
Fair	41 (4.8)	8 (6.2%)	33 (4.6%)	
Poor	3 (0.4)	0 (0%)	2 (0.3%)	
Annual income				0.563
\$0 to \$9999	25 (2.9)	2 (1.6%)	23 (3.2%)	
\$10,000 to \$14,999	15 (1.8)	1 (0.8%)	14 (1.9%)	
\$15,000 to \$19,999	12 (1.4)	1 (0.8%)	11 (1.5%)	
\$20,000 to \$34,999	66 (7.7)	10 (7.8%)	56 (7.8%)	
\$35,000 to \$49,999	65 (7.6)	13 (10.1%)	52 (7.2%)	
\$50,000 to \$74,999	117 (13.7)	21 (16.3%)	95 (13.2%)	
\$75,000 to \$99,999	119 (13.9)	14 (10.9%)	103 (14.3%)	
\$100,000 to \$199,999	235 (27.5)	35 (27.1%)	200 (27.8%)	
\$200,000 or more	82 (9.6)	11 (8.5%)	70 (9.7%)	
Education				0.625
Less than high school	15 (1.8)	4 (3.1%)	11 (1.5%)	
Some high school, no diploma	10 (1.2)	1 (0.8%)	9 (1.3%)	
GED	2 (0.2)	0 (0%)	2 (0.3%)	
High school graduate	56 (6.5)	9 (7.0%)	47 (6.5%)	
Some college but no degree	126 (14.7)	22 (17.1%)	103 (14.3%)	
Associate degree-occupational/vocational	35 (4.1)	4 (3.1%)	31 (4.3%)	
Associate degree-academic program	41 (4.8)	6 (4.7%)	34 (4.7%)	
Bachelor's degree (e.g., BA, AB, BS)	319 (37.3)	41 (31.8%)	276 (38.3%)	
Master's degree (e.g., MA, MS, Meng, Med, MSW)	175 (20.4)	33 (25.6%)	142 (19.7%)	
Professional school degree (e.g., MD, DDS, DVM, JD)	22 (2.6)	1 (0.8%)	20 (2.8%)	
Doctorate degree (e.g., PhD, EdD)	23 (2.7)	3 (2.3%)	20 (2.8%)	

American Indian/Alaskan native participants (0.7%), 129 Hispanic participants (15.1), and 720 non-Hispanic participants (84.1%) (Tables 1 and 2). There were 334 male and 515 female participants (39.0% and 60.2%, respectively). Only 3 participants requested a translated survey (Chinese). Our data indicate that significantly fewer male Hispanics participated compared to non-Hispanics (28.7% versus 40.8%, respectively, $p = 0.0118$) and Hispanic participants were significantly younger (median age of 42.6 compared to 47.7 for non-Hispanics, $p = 0.0196$). Race-related gender differences were not observed. Most of the percentages for race, ethnicity, and gender are similar to the reported demographics for Sacramento county (61.4% White, 10.9% Black/African American, Asian (17.8%), Native Hawaiian/other Pacific Islander participants (1.3%), American Indian/Alaskan native (1.6%), Hispanic (24.4%), 50.8% female)⁴⁰. There were more Asian participants than may have been expected based on the demographic data from Sacramento county, and this may be in part due to because of the location of the clinic in Sacramento county: the city of Elk Grove has a significantly higher Asian population compared to other areas in Sacramento county⁴¹. There were also fewer Black participants than may have been expected based on comparison with demographic data from both Sacramento county and the city of Elk Grove.^{44,45}

Statistically significant race-based differences were observed for annual income and education (Table 1, $p \leq 0.001$): fewer White participants had a combined income of \$0 - \$9999 and more Asian participants had incomes in this range and the \$10,000 - \$14,000 range than would be expected if wealth and race were independent in this cohort. Similarly, more White participants and fewer Asian participants had incomes in the \$100,000 - \$199,000 and \$200,000 or more range than would be expected under independence. Ethnicity-based differences were not observed for income or education (Table 2). Statistically significant race-based differences were also observed for education level (Table 1, $p \leq 0.001$): more White participants had a master's degree and fewer had a high school diploma as their highest level of education compared to other races. More American Indian/Alaskan Native participants had a GED than would be expected under independence. More Asian participants had some high school/no diploma or had a high school diploma as their highest level of education and fewer had a master's degree than would be expected under independence. Ethnicity-based differences were not observed for level of education (Table 2).

Table 3
Impact of Covid-19 on cancer screening and lifestyle factors relating to cancer risk, by race.

Characteristic	Participants, No. (%)						P-value (ANOVA or Chi Squared)
	Total (N = 856 [100%])	White (n = 411 [48.0%])	Black or African American (n = 43 [5.0%])	American Indian or Alaskan Native (n = 7 [0.7%])	Asian or Asian American (n = 254 [29.7%])	Native Hawaiian, other Pacific Islander (n = 16 [1.9%])	
Postponed a scheduled cancer screening*	48 (5.6)	35 (8.5)	5 (11.6)	0 (0)	3 (1.2)	0 (0)	0.0411
Tobacco and/or marijuana usage							<0.001
Increased usage	60 (7.0)	40 (9.7)	2 (4.7)	1 (14.3)	6 (2.4)	0 (0)	
Decreased usage	30 (3.5)	8 (1.9)	6 (14.0)	0 (0)	0 (0)	11 (11.7)	
No change	503 (58.8)	278 (67.6)	19 (44.2)	3 (42.9)	123 (48.4)	8 (50)	
Consumption of fruits and vegetables							<0.001
Decreased consumption	69 (8.1)	33 (8.0)	6 (14.0)	1 (14.3)	18 (7.1)	3 (18.8)	
Increased consumption	183 (21.4)	82 (20.0)	12 (27.9)	2 (28.6)	52 (20.5)	4 (25.0)	
No change	582 (68.0)	290 (70.6)	25 (58.1)	2 (28.6)	174 (68.5)	8 (50.0)	
Alcohol consumption							<0.001
Increased consumption	69 (8.1)	77 (18.7)	5 (11.6)	1 (14.3)	16 (6.3)	2 (12.5)	
Decreased consumption	183 (21.4)	52 (12.7)	9 (20.9)	0 (0)	37 (14.6)	4 (25.0)	
No change	582 (68.0)	261 (63.5)	26 (60.5)	4 (57.1)	141 (55.5)	7 (43.8)	
Physical activity							<0.001
Decreased activity	383 (44.7)	176 (42.8)	21 (48.8)	4 (57.1)	103 (40.6)	9 (56.3)	
Increased activity	155 (18.1)	90 (21.9)	4 (9.3)	1 (14.3)	38 (15.0)	2 (12.5)	
No change	294 (34.3)	140 (34.1)	18 (41.9)	1 (14.3)	99 (39.0)	5 (31.3)	

* NOTE: A total of 152 participants (17.8%) said 'yes' to they were scheduled to have a cancer screening, this included 96 (23.4%) White, 11 (25.6%) Black or African American, 0 (0%) American Indian or Alaskan Native, 28 (11%) Asian or Asian American, and 0 (0%) Native Hawaiian and other Pacific Islander participants.

3.2. Impact of the Covid-19 pandemic on cancer screening and lifestyle factors relating to cancer risk

Our survey data indicates that the Covid-19 pandemic negatively impacted cancer screening. A total of 152 participants (17.8%) had a scheduled cancer screening appointment and of these 48 participants (5.6%) reported having postponed it for reasons relating to Covid-19 (Tables 3 and 4). Race differences were observed ($p = 0.0411$) with fewer Asian or Asian American participants reporting having postponed a screening appointment than would be expected under independence, however, fewer Asian or Asian American participants had a scheduled cancer screening appointment. White participants were more likely than other races to have a scheduled cancer screening appointment. Ethnicity-based differences were not observed for scheduling of a screening appointment or postponement of an appointment.

Out of the lifestyle factors assessed (tobacco and/or marijuana usage, consumption of fruits and vegetables, alcohol consumption, and physical activity), our survey data indicates that the Covid-19 pandemic had the largest negative impact on physical activity levels of participants (Tables 3 and 4). A total of 383 participants (44.7%) reported a decrease in physical activity while 155 (18.1%) and 294 (34.3%) of participants reported an increase in physical exercise or no change, respectively. No race or ethnicity-related differences were observed.

A negative impact on tobacco and/or marijuana usage was also observed: 60 participants (7.0%) reported increased usage during the Covid-19 pandemic compared to only 30 participants (3.5%) reporting decreased usage and 503 (58.8%) reporting no change in usage (Tables 3 and 4). More white participants reported increased usage of tobacco and/or marijuana and more Black or African American participants reported decreased usage than would be expected under independence. No ethnicity-related differences were observed.

Our survey data indicate that many participants decreased their alcohol consumption during the Covid-19 pandemic: 183 participants (21.4%) reported decreasing alcohol consumption compared to 69 participants (8.1%) who increased consumption and 585 participants (68.0%) who didn't change alcohol consumption (Tables 3 and 4). Race differences were observed: significantly more White participants reported increasing alcohol consumption and significantly fewer Asian or Asian American participants reported increasing alcohol consumption than would be expected under independence. No ethnicity-related differences were observed.

Our survey data indicate that the pandemic had a positive impact on fruit and vegetable consumption: 183 participants (21.4%) reported increased consumption of fruit and vegetables during the Covid-19 pandemic compared to 69 participants (8.1%) and 582 participants (68.0%) who reported decreased or no change in consumption of fruit and vegetables, respectively (Tables 3 and 4). No race or ethnicity-related differences were observed.

Table 4
Impact of Covid-19 on cancer screening and lifestyle factors relating to cancer risk, by ethnicity.

Characteristic	Participants, No. (%)			P value (ANOVA or Chi Squared)
	Total (N = 856)	Hispanic (n = 129 [15.1%])	Non-Hispanic (n = 720 [84.4%])	
Postponed a scheduled cancer screening*	48 (5.6)	5 (3.9)	43 (6.0)	0.697
Tobacco and/or marijuana usage				0.520
Increased usage	60 (7.0)	9 (7.0)	43 (6.0)	
Decreased usage	30 (3.5)	8 (6.2)	22 (3.1)	
No change	503 (58.8)	71 (55.0)	428 (59.4)	
Consumption of fruits and vegetables				0.421
Decreased consumption	69 (8.1)	8 (6.2)	61 (8.5)	
Increased consumption	183 (21.4)	34 (26.4)	148 (20.6)	
No change	582 (68.0)	84 (65.1)	492 (68.3)	
Alcohol consumption				0.128
Increased consumption	120 (14%)	24 (18.6)	96 (13.3)	
Decreased consumption	127 (14.8)	24 (18.6)	102 (14.2)	
No change				
Physical activity				0.00599
Decreased activity	383 (44.7)	65 (50.4)	316 (43.9)	
Increased activity	155 (18.1)	30 (23.3)	124 (17.2)	
No change	294 (34.3)	28 (21.7)	262 (36.4)	

* NOTE: A total of 152 participants (17.8%) said 'yes' to they were scheduled to have a cancer screening, this included 21 (16.3%) Hispanics and 131 (18.2%) non-Hispanics.

3.3. Impact of the Covid-19 pandemic on access to healthcare, medications

Our survey data demonstrate that the Covid-19 pandemic negatively impacted access to healthcare, however, no race or ethnicity-based differences were observed for any of the metrics assessed (Tables 5 and 6). A large number of survey participants reported having had a medical appointment cancelled by their healthcare provider (22.8%) or having cancelled an appointment themselves (25.8%) due to concerns relating to Covid-19. A large number of survey participants also reported postponing routine vaccinations (8.8%, e.g. Shingles, flu, HPV vaccinations) for themselves or their children. The Covid-19 pandemic did not appear to have a big impact on the ability of participants to obtain prescription or over-the-counter medications (3.5% and 4.4% of participants reported being unable to obtain medications, respectively). A total of 57 participants (6%) reported having been diagnosed with cancer during their lifetime, only 2 of these participants stated that the Covid-19 pandemic had impacted their medical care (3.5%) and our survey data indicate that Covid-19 did not have a significant impact on their access to cancer-related healthcare and medications.

4. Discussion

This survey study was conducted over a 10-day period at a pharmacist-led public Covid-19 vaccination clinic held at California Northstate University (CNU) in Elk Grove (Sacramento County, California) in March 2021. To our knowledge, this is the first study to have conducted a cancer outcomes-related survey at a public Covid-19 vaccination site. It is also the first pharmacist-led study in this area. Our data indicates that the Covid-19 pandemic has had both negative and positive impacts on factors which are known to be associated with cancer outcomes. In regard to negative impacts, the data indicate that the Covid-19 pandemic resulted in postponement of cancer screenings and cancellation of medical appointments as well as decreased physical activity and increased tobacco and/or marijuana usage. Conversely, the data indicates that the Covid-19 pandemic increased consumption of fruits and vegetables and decreased consumption of alcohol. Several race-related differences but no ethnicity-related differences were observed.

Survey participation rates were high and for the most part representative of our diverse community. A slightly lower level of participation than might be expected was observed for Black and African Americans and fewer male Hispanics compared to male non-Hispanics participated. These findings align with other studies which demonstrated significantly higher levels of Covid-19 vaccine hesitancy in African American and Hispanic populations during the timeframe of our study.^{46–48} Vaccination rates subsequently increased, most likely due to targeted outreach to these populations.^{46,48,49} Significantly more females participated in our study compared to males. The reasons for this are unclear, however several studies have shown that women are generally more likely to participate in health research survey studies compared to men and this may at least partially help explain this observed difference.^{50–52} It is likely that our engagement of pharmacy students and other healthcare professions students and our emphasis on providing culturally appropriate patient counseling, and, when possible, linguistically appropriate counseling, helped increase overall survey participation rates. At the CNU Covid-19 vaccination clinic, where this study was held, pharmacists and pharmacy students took the lead on patient counseling and vaccinations while physicians and medical students took the lead for post-vaccination surveillance thereby making the clinic an interdisciplinary effort. Students also helped with survey administration and collection under the supervision of faculty members. Other studies have demonstrated that involving medical professionals, including pharmacists, in survey administration as well as focus on cultural and linguistic competency can increase survey participation.^{53,54} Pharmacy students receive extensive patient counseling and cultural competency training, and this makes pharmacists well positioned to help recruit participants

Table 5
Impact of Covid-19 on access to healthcare, access to medications.

Characteristic	Participants, No. (%)						P-value (ANOVA or Chi Squared)
	Total (N = 856 [100%])	White (n = 411 [48.0%])	Black or African American (n = 43 [5.0%])	American Indian or Alaskan Native (n = 7 [0.7%])	Asian or Asian American (n = 254 [29.7%])	Native Hawaiian, other Pacific Islander (n = 16 [1.9%])	
Participant cancelled a medical appointment	221 (25.8)	107 (26.0)	13 (30.2)	3 (42.9)	57 (22.4)	2 (12.5)	0.647
Participant's medical appointment was cancelled by facility	195 (22.8)	104 (25.3)	10 (23.3)	2 (28.6)	48 (18.9)	1 (6.3)	0.285
Participant postponed vaccinations for themselves or their children	75 (8.8)	42 (10.2)	3 (7.0)	1 (14.3)	17 (6.7)	2 (12.5)	0.673
Participant was unable to obtain one or more prescription medications	30 (3.5)	17 (4.1)	3 (7.0)	0 (0)	6 (2.4)	1 (20.0)	0.350
Participant was unable to obtain one or more over-the-counter medications	38 (4.4)	22 (5.3%)	3 (7.0%)	0 (0%)	9 (3.5%)	1 (6.3%)	0.883

Table 6
Impact of Covid-19 on access to healthcare, access to medications.

Characteristic	Participants, No. (%)			P value (ANOVA or Chi Squared)
	Total (N = 856)	Hispanic (n = 129 [15.1%])	Non-Hispanic (n = 720 [84.4%])	
Participant cancelled a medical appointment	221 (25.8)	39 (30.2)	181 (25.1)	0.127
Participant's medical appointment was cancelled by facility	195 (22.8)	32 (24.8)	159 (22.1)	0.564
Participant postponed vaccinations for themselves or their children	75 (8.8)	9 (7.0)	66 (9.2)	0.643
Participant was unable to obtain one or more prescription medications	30 (3.5)	5 (3.9)	25 (3.5)	0.956
Participant was unable to obtain one or more over-the-counter medications	38 (4.4)	7 (5.4)	31 (4.3)	0.659

for research studies including cancer control-related research studies.

A key finding from our study was that the Covid-19 pandemic resulted in many people postponing cancer screenings. This aligns with what others have shown.^{55,56} For example, Oakes et al. reported that breast cancer and cervical cancer screening rates dropped by as much as 40% at the start of the pandemic and have remained lower than pre-pandemic levels.¹⁸ While significantly fewer Asians or Asian Americans reported postponing scheduled cancer screening appointments, significantly fewer Asians or Asian Americans had cancer screening appointments scheduled compared to other races. White participants reported the highest levels of scheduled cancer screenings. The combined data supports the urgent need to put mechanisms in place to actively encourage cancer screening with an emphasis placed on increasing scheduling of screening appointments for minority populations: reduced screening rates in minority population are well known to contribute to cancer health disparities.⁵⁷ Pharmacists often serve as the initial health care educator within underserved communities and they are one of the most trusted healthcare professions.⁵⁸ They are provided with training not only on cancer treatments but also on cancer risks and screenings. These attributes, and their ability to hear and observe signs of early cancer or cancer risk, allow them to advise individuals visiting their pharmacies to seek immediate medical attention and/or to educate patients on how to reduce their cancer risk. Several opinion pieces have noted that the potential exists for pharmacists to play a bigger role in cancer control efforts, and this could include placing more focus on encouraging cancer screenings.³⁷⁻³⁹

Our finding that a high number of participants reported decreased levels of physical activity as a result of the Covid-19 pandemic also aligns with what others have reported. For example, a recent systematic review by Mehraeen et al. demonstrated the Covid-19 pandemic reduced physical activity in all age groups regardless of race and gender.¹⁹ Our data indicates that tobacco and marijuana usage also increased. Our survey captured tobacco and marijuana usage as a combined metric making it difficult to directly compare our data with other studies which typically assessed these separately, however, it is well known that a strong association between tobacco and marijuana usage exists⁵⁹ and therefore we believe an comparison with tobacco-specific studies is still meaningful. A report from the Federal Trade Commission, cigarette sales increased 0.4% in 2020 compared to 2019²⁰ and this aligns with our finding, however, a meta-analysis by Sarich et al. revealed mixed results with some studies reporting increased tobacco usage and others reporting decreased usage.⁶⁰ Results from studies which assessed marijuana usage during the pandemic are also mixed. Salles et al. reported an overall decrease in marijuana usage⁶¹ while Black et al. reported increased usage.⁶² These divergent findings indicate that community-specific assessment of tobacco and marijuana usage may be needed to help determine the urgency with which this factor needs to be addressed in individual communities. In our study, race-related differences in tobacco and marijuana usage were observed indicating that assessment by race may also be important: White participants reported increased usage of tobacco and marijuana products, while Black and African American participants decreased their usage. The latter aligns with data reported by Fucito et al. and Kowitt et al. which found that Black participants reported tobacco use reductions and intention to quit during the pandemic compared to White participants.^{63,64} For the other two lifestyle factor assessed, we observed positive trends: overall, more participants reported increasing fruit and vegetable consumption and decreasing alcohol consumption as compared to the number of participants reporting decreasing consumption of fruit and vegetables and increasing consumption of alcohol. No race or ethnicity differences were observed. Our finding that fruit and vegetable consumption increased aligns with a literature review by Johnson et al.⁶⁵ as well as studies conducted in Spain, Italy, and Poland which showed healthy eating improved during Covid-19 pandemic.⁶⁶⁻⁶⁸ It is noteworthy that Sharma et al. reported that fruit and vegetable intake decreased significantly in low-income families in the US⁶⁹ and it has been reported that food insecurity increased by approximately 10% in US households during the pandemic.⁷⁰ Based on this, it is possible that our observed increase in fruit and vegetable consumption may at least in part be due to the majority of our study participants being in higher income brackets – the most frequently reported annual income brackets were \$50,000 - \$74,999, \$75,000 - \$99,999, and \$100,000 - \$199,999. Further assessment of this metric in the lower income population in our community is needed. In contrast to our study, other survey studies have reported an overall increase in alcohol consumption during the pandemic.⁷¹⁻⁷⁴ While we observed an overall decrease, significantly more White participants reported increasing alcohol consumption compared to other races and this does align with race data reported by these other studies. The combined data indicates that targeted outreach to help reduce alcohol consumption in White adults in our community and beyond is warranted. As already noted, pharmacists receive extensive patient counseling training including smoking cessation counseling and healthy lifestyle choice counseling training and studies have demonstrated success in these areas.^{27,29-35} The motivational interviewing training which pharmacists receive provides

them with another tool to help encourage healthier habits through behavior changes and further reduce cancer risk and risk of progression and/or recurrence.

The Covid-19 pandemic also had a major impact on access to healthcare. Our survey data shows that a significant number of participants either cancelled healthcare-related appointments (22.8%) or had the appointments cancelled by their providers (25.8%). This aligns with what other studies have reported: a national survey conducted by Anderson et al. found that 41% of participants reported forgoing medical care from March to July in 2020,⁷⁵ while a national survey by Wenger et al. found that 37% of participants had cancelled or postponed appointments.²¹ No race or ethnicity-related differences were observed for these metrics in our study or the studies by Anderson et al. and only weak associations were observed by Wenger et al. Missing or postponing healthcare visits is well known to contribute to worse outcomes for all disease states including cancer and as such there is clearly an urgent need to follow up with patients who missed healthcare visits to help mitigate these likely worse health outcomes.⁷⁶ Very few of our study participants reported not being able to obtain prescription or over-the-counter medications, including cancer medications, and no race or ethnicity-related differences were observed. The number of participants who reported having ever been diagnosed with cancer was 6.4%, which is similar to the national average in the US of 5.5%.⁷⁷ Very few of our study participants (2 out of 35 participants, 5.7%) reported having delayed cancer-related healthcare visits and none reported changing oral cancer medication use. Ensuring timely cancer treatment is known to be extremely important in regards to improving patient survival rate.⁷⁸

The combined data indicates that there is an urgent need to help support increased cancer screenings, especially in minority populations, and a need to help ensure that missed medical appointments have been rescheduled and any healthcare issues addressed. In addition, our data support the need for outreach efforts in our community to help promote increased physical exercise and to reduce tobacco and marijuana consumption. Our data indicate that providing targeted outreach to help reduce alcohol consumption by White adults in our community is also warranted. The related literature which we have cited strongly supports an enhanced role for pharmacists in cancer control efforts such as these. Addressing these factors is likely to help mitigate potential worse cancer outcomes in both the near and distant future. Our next step will be to use the data to help guide targeted outreach efforts within our local community. As with this survey study, these outreach efforts will be pharmacist-led and engage pharmacy students and other students with a healthcare interest. Other studies have demonstrated pharmacists can play an important role in community outreach efforts based on their training, experience, and penetrance and accessibility in all communities^{27,29,30,37-39} and should continue to do so and further expand these efforts.

4.1. Study limitations

This is a single center study and as such the data presented may not be generalizable beyond our community. The data obtained are based on self-report from a convenience sample of the population and as such sampling bias exists; the data may not provide a true representation of our community. Participants' responses for the lifestyle questions could have been confounded by recall bias: the Covid-19 pandemic started in early 2019 and this survey study was conducted in March 2021. This factor has been highlighted by Cross et al. in response to similar survey studies.⁶ Lastly, the study was not specifically powered to detect differences between races or ethnicities and as such the subgroup analyses by race should be interpreted with caution.

Funding

This study was funded by an ASHP Pharmacy Leadership Scholars Grant to ET-L. MSC is supported in part by funding from P30CA093373-18S3 (8/1/20-6/30/21), "Impact of COVID-19 on cancer-related behaviors among non-metropolitan and minorities in inland northern California: Seeking mitigation strategies".

Study approval

This study was approved by the CNU IRB committee (protocol #2103-01-80).

Footnotes

None.

CRedit authorship contribution statement

Erika Titus-Lay: Conceptualization, Methodology, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Funding acquisition, Resources, Supervision. **Jeffrey Nehira:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Jennifer Courtney:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Jacquelyn Jee:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Marissa Kumar:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Jenny Tiet:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Vivi Le:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Blythe Durbin-Johnson:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Moon S. Chen:** Methodology, Formal analysis, Investigation, Writing – review & editing. **Ruth Vinall:** Conceptualization, Methodology, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Funding acquisition, Resources, Supervision.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors would like to thank all of the California Northstate University College of Pharmacy faculty members and students who contributed to translation of the study survey into different languages and/or helped with survey administration.

The data underlying this article will be shared on reasonable request to the corresponding author.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.rcsop.2023.100311>.

References

- GBD 2019 Cancer Risk Factors Collaborators. The global burden of cancer attributable to risk factors, 2010-19: a systematic analysis for the global burden of disease study 2019. *Lancet*. 2022;400(10352):563–591. [https://doi.org/10.1016/S0140-6736\(22\)01438-6](https://doi.org/10.1016/S0140-6736(22)01438-6).
- Islami F, et al. Proportion and number of cancer cases and deaths attributable to potentially modifiable risk factors in the United States. *CA Cancer J Clin*. 2018;68(1):31–54. <https://doi.org/10.3322/caac.21440>.
- Ugai T, et al. Is early-onset cancer an emerging global epidemic? Current evidence and future implications. *Nat Rev Clin Oncol*. 2022;19(10):656–673. <https://doi.org/10.1038/s41571-022-00672-8>.
- Loomans-Kropp HA, Umar A. Cancer prevention and screening: the next step in the era of precision medicine. *NPJ Precis Oncol*. 2019;3:3. <https://doi.org/10.1038/s41698-018-0075-9>.
- Loud JT, Murphy J. Cancer screening and early detection in the 21(st) century. *Semin Oncol Nurs*. 2017;33(2):121–128. <https://doi.org/10.1016/j.soncn.2017.02.002>.
- Cross TJ, et al. Self-reported physical activity before a COVID-19 ‘lockdown’: is it just a matter of opinion? *BMJ Open Exercise Medicine*. 2021;7(2), e001088. <https://doi.org/10.1136/bmjsem-2021-001088>.
- Oswald LB, et al. Smoking is related to worse Cancer-related symptom burden. *Oncologist*. 2022;27(2):e176–e184. <https://doi.org/10.1093/oncolo/oyab029>.
- American Cancer Society. Health risks of smoking tobacco. Accessed on February 1st, 2023 <https://www.cancer.org/healthy/stay-away-from-tobacco/health-risks-of-tobacco/health-risks-of-smoking-tobacco.html>; 2023.
- Office of the Surgeon General. Tobacco reports and publications. Accessed on February 1st, 2023 <https://www.hhs.gov/surgeongeneral/reports-and-publications/tobacco/index.html>; 2023.
- Zhong L, et al. Alcohol and health outcomes: an umbrella review of Meta-analyses base on prospective cohort studies. *Front Public Health*. 2022;10, 859947. <https://doi.org/10.3389/fpubh.2022.859947>.
- Winzer BM, et al. Physical activity and cancer prevention: a systematic review of clinical trials. *Cancer Causes Control*. 2011;22(6):811–826. <https://doi.org/10.1007/s10552-011-9761-4>.
- Schmid D, Leitzmann MF. Association between physical activity and mortality among breast cancer and colorectal cancer survivors: a systematic review and meta-analysis. *Ann Oncol*. 2014;25(7):1293–1311. <https://doi.org/10.1093/annonc/mtu012>.
- Anand P, et al. Cancer is a preventable disease that requires major lifestyle changes. *Pharm Res*. 2008;25(9):2097–2116. <https://doi.org/10.1007/s11095-008-9661-9>.
- American Association of Cancer Research. Disparities in the burden of preventable cancer risk factors. Accessed on February 1st, 2023 <https://cancerprogressreport.aacr.org/disparities/cdpr22-contents/cdpr22-disparities-in-the-burden-of-preventable-cancer-risk-factors/>; 2023.
- Centers for Disease Control and Prevention. Adult physical activity prevalence maps by race/ethnicity. Accessed on February 1st, 2023 <https://www.cdc.gov/physicalactivity/data/inactivity-prevalence-maps/index.html>; 2023.
- Institute, N.C. Cancer disparities. Accessed on 7-7-2023; Available from: <https://www.cancer.gov/about-cancer/understanding/disparities>; 2022.
- Patel MI, et al. Racial and ethnic disparities in Cancer care during the COVID-19 pandemic. *JAMA Netw Open*. 2022;5(7), e2222009. <https://doi.org/10.1001/jamanetworkopen.2022.22009>.
- Oakes AH, et al. Rates of routine Cancer screening and diagnosis before vs after the COVID-19 pandemic. *JAMA Oncol*. 2023;9(1):145–146. <https://doi.org/10.1001/jamaoncol.2022.5481>.
- Mehraeen E, et al. The impact of COVID-19 pandemic on the levels of physical activity: a systematic review. *Infect Disord Drug Targets*. 2023. <https://doi.org/10.2174/1871526523666230120143118>.
- Federal Trade Commission. Cigarette report for 2020. Accessed on February 1st, 2023 <https://www.ftc.gov/system/files/documents/reports/federal-trade-commission-cigarette-report-2020-smokeless-tobacco-report-2020/p114508fy20cigarettereport.pdf>; 2021.
- Wenger NS, et al. The impact of COVID-19 on routine medical care and cancer screening. *J Gen Intern Med*. 2022;37(6):1450–1456. <https://doi.org/10.1007/s11606-021-07254-x>.
- Amram O, et al. Socioeconomic and racial inequities in breast Cancer screening during the COVID-19 pandemic in Washington state. *JAMA Netw Open*. 2021;4(5), e2110946. <https://doi.org/10.1001/jamanetworkopen.2021.10946>.
- Sprague BL, et al. Changes in mammography use by Women’s characteristics during the first 5 months of the COVID-19 pandemic. *J Natl Cancer Inst*. 2021;113(9): 1161–1167. <https://doi.org/10.1093/jnci/djab045>.
- Klemperer EM, et al. Change in tobacco and electronic cigarette use and motivation to quit in response to COVID-19. *Nicotine Tob Res*. 2020;22(9):1662–1663. <https://doi.org/10.1093/ntr/ntaa072>.
- Gonzalez M, et al. Smokers are more likely to smoke more after the COVID-19 California lockdown order. *Int J Environ Res Public Health*. 2021;18(5). <https://doi.org/10.3390/ijerph18052582>.
- Booker A, Malcarne VL, Sadler GR. Evaluating outcomes of community-based cancer education interventions: a 10-year review of studies. *J Cancer Educ*. 2014;29(2):233–240. <https://doi.org/10.1007/s13187-013-0578-6>.
- Chen Y, et al. Improving Cancer outreach effectiveness through targeting and economic assessments: insights from a randomized field experiment. *J Mark*. 2020; 84(3):1–27. <https://doi.org/10.1177/0022242920913025>.
- National Center for Chronic Disease, P., S. Health Promotion Office on, and Health. *Reports of the surgeon general, in the health consequences of smoking—50 years of progress: a report of the surgeon general*. Atlanta (GA): Centers for Disease Control and Prevention (US); 2014.

29. Al Aqeel S, et al. Interventions for improving pharmacist-led patient counselling in the community setting: a systematic review. *Syst Rev.* 2018;7(1):71. <https://doi.org/10.1186/s13643-018-0727-4>.
30. Lowrie R, et al. A cluster randomised controlled trial of a pharmacist-led collaborative intervention to improve statin prescribing and attainment of cholesterol targets in primary care. *PLoS One.* 2014;9(11), e113370. <https://doi.org/10.1371/journal.pone.0113370>.
31. Warren GW, et al. The 2014 Surgeon General's report: "The health consequences of smoking—50 years of progress": a paradigm shift in cancer care. *Cancer.* 2014; 120(13):1914–1916. <https://doi.org/10.1002/cncr.28695>.
32. Hluhanich R, et al. Comparing hepatitis C virus screening in clinics versus the emergency department. *West J Emerg Med.* 2022;23(3):312–317. <https://doi.org/10.5811/westjem.2021.11.53870>.
33. Lowrie R, et al. Pharmacist and homeless outreach engagement and non-medical independent prescribing Rx (PHOENIX): a study protocol for a pilot randomised controlled trial. *BMJ Open.* 2022;12(12), e064792. <https://doi.org/10.1136/bmjopen-2022-064792>.
34. Biola H, et al. Reaching the hard-to-reach: outcomes of the severe hypertension outreach intervention. *Am J Prev Med.* 2020;59(5):725–732. <https://doi.org/10.1016/j.amepre.2020.05.030>.
35. Wright QE, et al. The impact of a pharmacist-led naloxone education and community distribution project on local use of naloxone. *J Am Pharm Assoc (2003).* 2020;60(3s):S56–S60. <https://doi.org/10.1016/j.japh.2019.11.027>.
36. Langley J. Motivational interviewing offers a path to improved adherence. *Pharmacy Times*; 2019. Accessed February 6, 2023 <https://www.pharmacytimes.com/view/motivational-interviewing-offers-a-path-to-improved-adherence>. Accessed February 6, 2023.
37. Havlicek AJ, Mansell H. The community pharmacist's role in cancer screening and prevention. *Can Pharm J (Ott).* 2016;149(5):274–282. <https://doi.org/10.1177/1715163516660574>.
38. Egbewande OM, et al. Roles of community pharmacists in Cancer management. *Innov Pharm.* 2022;13(3). <https://doi.org/10.24926/iip.v13i3.4946>.
39. Konya J, et al. Can early cancer detection be improved in deprived areas by involving community pharmacists? *Br J Gen Pract.* 2022;72(717):153–154. <https://doi.org/10.3399/bjgp22X718865>.
40. Federal Register Surgeon general's call to action: "community health and prosperity". 2018. Accessed February 6, 2023. <https://www.federalregister.gov/documents/2018/09/06/2018-19313/surgeon-generals-call-to-action-community-health-and-prosperity>.
41. Scarinci IC, et al. Factors associated with perceived susceptibility to COVID-19 among urban and rural adults in Alabama. *J Community Health.* 2021;46(5): 932–941. <https://doi.org/10.1007/s10900-021-00976-3>.
42. R Core Team R: *A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing; 2015. Accessed June 23, 2022 <https://www.R-project.org/>.
43. Agresti A. *Categorical Data Analysis.* 3rd. Wiley; 2012:69–103. ISBN 978-0-470-46363-5.
44. United States Census Bureau Quick facts. Sacramento County California; 2022. Accessed February 6, 2023 <https://www.census.gov/quickfacts/sacramentocountycalifornia>. Accessed February 6, 2023.
45. United States Census Bureau Quick facts, Elk Grove California 2022. ; 2023. Accessed February 1 <https://www.census.gov/quickfacts/elkgrovecitycalifornia>. Accessed February 1.
46. Demeke J, et al. Strategies that promote equity in COVID-19 vaccine uptake for latinx communities: a review. *J Racial Ethn Health Disparities.* 2022:1–9. <https://doi.org/10.1007/s40615-022-01320-8>.
47. Khubchandani J, Macias Y. COVID-19 vaccination hesitancy in Hispanics and African-Americans: a review and recommendations for practice. *Brain Behav Immun Health.* 2021;15, 100277. <https://doi.org/10.1016/j.bbih.2021.100277>.
48. Ripon RK, et al. A meta-analysis of COVID-19 vaccines acceptance among black/African American. *Heliyon.* 2022;8(12), e12300. <https://doi.org/10.1016/j.heliyon.2022.e12300>.
49. Omari A, et al. Characteristics of the moveable middle: opportunities among adults open to COVID-19 vaccination. *Am J Prev Med.* 2022. <https://doi.org/10.1016/j.amepre.2022.11.003>.
50. Glass DC, et al. A telephone survey of factors affecting willingness to participate in health research surveys. *BMC Public Health.* 2015;15:1017. <https://doi.org/10.1186/s12889-015-2350-9>.
51. Dunn KM, et al. Patterns of consent in epidemiologic research: evidence from over 25,000 responders. *Am J Epidemiol.* 2004;159(11):1087–1094. <https://doi.org/10.1093/aje/kwh141>.
52. Meiklejohn J, Connor J, Kypr K. The effect of low survey response rates on estimates of alcohol consumption in a general population survey. *PLoS One.* 2012;7 (4), e35527. <https://doi.org/10.1371/journal.pone.0035527>.
53. Booker QS, Austin JD, Balasubramanian BA. Survey strategies to increase participant response rates in primary care research studies. *Fam Pract.* 2021;38(5): 699–702. <https://doi.org/10.1093/fampra/cmab070>.
54. Bonevski B, et al. Reaching the hard-to-reach: a systematic review of strategies for improving health and medical research with socially disadvantaged groups. *BMC Med Res Methodol.* 2014;14:42. <https://doi.org/10.1186/1471-2288-14-42>.
55. Cancino RS, et al. The impact of COVID-19 on cancer screening: challenges and opportunities. *JMIR Cancer.* 2020;6(2), e21697. <https://doi.org/10.2196/21697>.
56. Luu T. Reduced Cancer screening due to lockdowns of the COVID-19 pandemic: reviewing impacts and ways to counteract the impacts. *Front Oncol.* 2022;12, 955377. <https://doi.org/10.3389/fonc.2022.955377>.
57. Zavala VA, et al. Cancer health disparities in racial/ethnic minorities in the United States. *Br J Cancer.* 2021;124(2):315–332. <https://doi.org/10.1038/s41416-020-01038-6>.
58. Brennan M. Nurses again outpace other professions for honesty, ethics. Accessed on 7-7-2023. 7-7-2023; Available from: <https://news.gallup.com/poll/245597/nurses-again-outpace-professions-honesty-%20ethics.aspx>; 2018.
59. Lemyre A, Poliakova N, Belanger RE. The relationship between tobacco and Cannabis use: a review. *Subst Use Misuse.* 2019;54(1):130–145. <https://doi.org/10.1080/10826084.2018.1512623>.
60. Sarich P, et al. Tobacco smoking changes during the first pre-vaccination phases of the COVID-19 pandemic: a systematic review and meta-analysis. *EClinicalMedicine.* 2022;47, 101375. <https://doi.org/10.1016/j.eclinm.2022.101375>.
61. Salles J, et al. Changes in Cannabis consumption during the global COVID-19 lockdown: the international COVISTRESS study. *Front Psychiatry.* 2021;12, 689634. <https://doi.org/10.3389/fpsy.2021.689634>.
62. Black JC, et al. Evaluation of Cannabis use among US adults during the COVID-19 pandemic within different legal frameworks. *JAMA Netw Open.* 2022;5(11), e2240526. <https://doi.org/10.1001/jamanetworkopen.2022.40526>.
63. Fucito LM, et al. Cigarette smoking in response to COVID-19: examining co-morbid medical conditions and risk perceptions. *Int J Environ Res Public Health.* 2022; 19(14). <https://doi.org/10.3390/ijerph19148239>.
64. Kowitz SD, et al. Tobacco quit intentions and behaviors among cigar smokers in the United States in response to COVID-19. *Int J Environ Res Public Health.* 2020;17 (15). <https://doi.org/10.3390/ijerph17155368>.
65. Johnson AN, et al. Changes in adults' eating behaviors during the initial months of the COVID-19 pandemic: a narrative review. *J Acad Nutr Diet.* 2023;123(1): 144–194 e30. <https://doi.org/10.1016/j.jand.2022.08.132>.
66. Ruggiero E, et al. Changes in the consumption of foods characterising the Mediterranean dietary pattern and major correlates during the COVID-19 confinement in Italy: results from two cohort studies. *Int J Food Sci Nutr.* 2021;72(8):1105–1117. <https://doi.org/10.1080/09637486.2021.1895726>.
67. Navarro-Perez CF, et al. Effects of COVID-19 lockdown on the dietary habits and lifestyle in a population in southern Spain: a cross-sectional questionnaire. *Eur J Clin Nutr.* 2022;76(6):883–890. <https://doi.org/10.1038/s41430-021-01034-w>.
68. Gornicka M, et al. Dietary and lifestyle changes during COVID-19 and the subsequent lockdowns among Polish adults: a cross-sectional online survey PLifeCOVID-19 study. *Nutrients.* 2020;12(8). <https://doi.org/10.3390/nu12082324>.
69. Sharma SV, et al. Social determinants of health-related needs during COVID-19 among low-income households with children. *Prev Chronic Dis.* 2020;17:E119. <https://doi.org/10.5888/pcd17.200322>.

- 70 Otten JJ, Averill MM, Spiker ML. Food security and food access during the COVID-19 pandemic: impacts, adaptations, and looking ahead. *JPEN J Parenter Enteral Nutr.* 2022. <https://doi.org/10.1002/jpen.2445>.
- 71 Pollard MS, Tucker JS, Green Jr HD. Changes in adult alcohol use and consequences during the COVID-19 pandemic in the US. *JAMA Netw Open.* 2020;3(9), e2022942. <https://doi.org/10.1001/jamanetworkopen.2020.22942>.
- 72 Grossman ER, Benjamin-Neelon SE, Sonnenschein S. Alcohol consumption during the COVID-19 pandemic: a Cross-sectional survey of US adults. *Int J Environ Res Public Health.* 2020;17(24). <https://doi.org/10.3390/ijerph17249189>.
- 73 Kerr WC, et al. Longitudinal assessment of drinking changes during the pandemic: the 2021 COVID-19 follow-up study to the 2019 to 2020 National Alcohol Survey. *Alcohol Clin Exp Res.* 2022;46(6):1050–1061. <https://doi.org/10.1111/acer.14839>.
- 74 Nordeck CD, et al. Changes in drinking days among United States adults during the COVID-19 pandemic. *Addiction.* 2022;117(2):331–340. <https://doi.org/10.1111/add.15622>.
- 75 Anderson KE, et al. Reports of forgone medical care among US adults during the initial phase of the COVID-19 pandemic. *JAMA Netw Open.* 2021;4(1), e2034882. <https://doi.org/10.1001/jamanetworkopen.2020.34882>.
- 76 McQueenie R, et al. Morbidity, mortality and missed appointments in healthcare: a national retrospective data linkage study. *BMC Med.* 2019;17(1):2. <https://doi.org/10.1186/s12916-018-1234-0>.
77. Our World in Data. Cancer. Accessed on February 1st, 2023 <https://ourworldindata.org/cancer>; 2023.
78. Hanna TP, et al. Mortality due to cancer treatment delay: systematic review and meta-analysis. *BMJ.* 2020;371, m4087. <https://doi.org/10.1136/bmj.m4087>.