

UC San Diego

UC San Diego Previously Published Works

Title

Higher-order Fourier Analysis and Applications

Permalink

<https://escholarship.org/uc/item/783251ht>

Journal

Foundations and Trends® in Theoretical Computer Science, 13(4)

ISSN

1551-305X

Authors

Hatami, Hamed

Hatami, Pooya

Lovett, Shachar

Publication Date

2019

DOI

10.1561/04000000064

Peer reviewed

Higher-order Fourier Analysis and Applications

Hamed Hatami¹, Pooya Hatami² and Shachar Lovett³

¹*McGill University; hatami@cs.mcgill.ca*

²*Ohio State University; pooyahat@gmail.com*

³*University of California, San Diego; slovett@cs.ucsd.edu*

ABSTRACT

Fourier analysis has been extremely useful in many areas of mathematics. In the last several decades, it has been used extensively in theoretical computer science. Higher-order Fourier analysis is an extension of the classical Fourier analysis, where one allows to generalize the “linear phases” to higher degree polynomials. It has emerged from the seminal proof of Gowers of Szemerédi’s theorem with improved quantitative bounds, and has been developed since, chiefly by the number theory community. In parallel, it has found applications also in theoretical computer science, mostly in algebraic property testing, coding theory and complexity theory.

The purpose of this book is to lay the foundations of higher-order Fourier analysis, aimed towards applications in theoretical computer science with a focus on algebraic property testing.

1

Introduction

The purpose of this text is to provide an introduction to the field of higher-order Fourier analysis with an emphasis on its applications to theoretical computer science. Higher-order Fourier analysis is an extension of the classical Fourier analysis. It was initiated by a seminal paper of Gowers [37] on a new proof for Szemerédi’s theorem, and has been developed by several mathematicians over the past few decades in order to study problems in an area of mathematics called additive combinatorics, which is primarily concerned with linear patterns such as arithmetic progressions in subsets of integers. While most of the developments in additive combinatorics were focused on the group \mathbb{Z} , it was quickly noticed that the analogous questions and results for the group \mathbb{F}_2^n are of great importance to theoretical computer scientists as they are related to basic concepts in areas such as property testing and coding theory.

Classical Fourier analysis is a powerful tool that studies functions by expanding them in terms of the Fourier characters, which are “linear phase functions” such as $n \mapsto e^{-\frac{2\pi i}{N}n}$ for the group \mathbb{Z}_N , or $(x_1, \dots, x_n) \mapsto (-1)^{\sum a_j x_j}$ for the group \mathbb{F}_2^n . Note that n and $\sum a_j x_j$ are both linear functions. Fourier analysis has been extremely successful in the study

of certain linear patterns such as three-term arithmetic progressions. For example, if the number of three-term arithmetic progressions in a subset $A \subseteq \mathbb{Z}_N$ deviates from the expected number of them in a random subset of \mathbb{Z}_N with the same cardinality as A , then A must have significant correlation with a linear phase function. In other words, the characteristic function of A must have a large non-principal Fourier coefficient. Roth [66] used these ideas to show that every subset of integers of positive upper density contains an arithmetic progression of length 3. However, classical Fourier analysis seems to be inadequate in detecting more complex linear patterns such as four-term or longer arithmetic progressions. Indeed, one can easily construct dense sets $A \subseteq \mathbb{Z}_N$ that do not have significant correlation with any linear phase function, and nevertheless do not contain the number of four-term arithmetic progressions that one expects by considering random subsets of the same cardinality. Hence in order to generalize Roth's theorem to arithmetic progressions of arbitrary length, Szemerédi [76, 77] departed from the Fourier analytic approach and appealed to purely combinatorial ideas. However, his proof of this major result, originally conjectured by Erdős and Turán [27], provided poor quantitative bounds on the minimal density that guarantees the existence of the arithmetic progressions of the desired length. Later Furstenberg [31] developed an ergodic-theoretic framework and gave a new proof for Szemerédi's theorem, but his proof was still qualitative. His theory is further developed by - to name a few - Host, Kra, Ziegler, Bergelson, Tao (See e.g. [51], [88], and [10, 82]), and there are important parallels between this theory and higher-order Fourier analysis. Indeed some of the terms that are commonly used in higher-order Fourier analysis such as "phase functions" or "factors" are ergodic theoretic terms.

Generalizing Roth's original proof and obtaining good quantitative bounds for Szemerédi's theorem remained a challenge until finally Gowers [37] discovered that the essential idea to overcome the obstacles described above is to consider higher-order phase functions. His proof laid the foundation for the area of higher-order Fourier analysis, where one studies a function by approximating it by a linear combination of few higher-order phase functions. Although the idea of using higher-order phase functions already appears in Gowers's work [37], it was not

until more than fifteen years later that some of the major technical difficulties in achieving a satisfactory theory of higher-order Fourier analysis have been resolved. By now, due to great contributions by prominent mathematicians such as Gowers, Green, Tao, Szegedy, Host, Kra and Ziegler (See [75] and [80] and the references there), there is a deep understanding of qualitative aspects of this theory. However, despite these major breakthroughs, still very little is known from a quantitative perspective as many of the proofs are based on soft analytic techniques, and obtaining efficient bounds is one of the major challenges in this area.

This survey will emphasize the applications of the theory of higher-order Fourier analysis to theoretical computer science, and to this end, we will present the foundations of this theory through such applications, in particular to the area of property testing. In the early nineties, it was noticed by Blum *et al.* [20] and Babai *et al.* [6] that Fourier analysis can be used to design a very efficient algorithm that distinguishes linear functions $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ from functions that are far from being linear. This initiated the area of property testing, the study of algorithms that query their input a very small number of times and with high probability decide correctly whether their input satisfies a given property or is “far” from satisfying that property. It was soon noticed that generalizing the linearity test of Blum *et al.* [20] and Babai *et al.* [6] to other properties such as the property of being a quadratic polynomial requires overcoming the same obstacles that one faces in an attempt to generalize Fourier analytic study of three-term arithmetic progressions to four-term arithmetic progressions. Hence in parallel to additive combinatorics, theoretical computer scientists have also been working on developing tools in higher-order Fourier analysis to tackle such problems. In fact some of the most basic results, such as the inverse theorem for the Gowers U^3 norm for the group \mathbb{F}_2^n , were first proved by Samorodnitsky [70] in the context of property testing for quadratic polynomials.

In **Part I** we discuss the linearity test due to Blum *et al.* [20] and its generalization to higher degree polynomials. We will see how this naturally necessitates the development of a theory of higher-order Fourier analysis. In **Part II** we present the fundamental results of the theory of higher-order Fourier analysis. Since we are interested in the

applications to theoretical computer science, we will only consider the group \mathbb{F}_p^n where p is a fixed prime, and asymptotics are as n tends to infinity. Higher-order Fourier analysis for the group \mathbb{Z}_N , which is of more interest for number theoretic applications, shares the same basic ideas but differs on some technical aspects. For this group, the higher order phase functions, rather than being exponentials of polynomials, are the so called nilsequences. We refer the interested reader to Tao [80] for more details. In [Part III](#) we use the tools developed in [Part II](#) to prove some general results about property testing for algebraic properties.

Throughout most of the text, we will consider fields of constant prime order, namely $\mathbb{F} = \mathbb{F}_p$ where p is a constant, and study functions from \mathbb{F}_p^n to \mathbb{R} , \mathbb{C} , or \mathbb{F}_p when n is growing. Our choice is mainly for simplicity of exposition, as there have been recent research that extend several of the tools from higher-order Fourier analysis to large or non-prime fields. We refer the interested reader to a paper by Bhattacharyya *et al.* [12] for treatment of non-prime fields. In [Chapter 8](#) we will discuss a paper by Bhowmick and Lovett [19] considering the case \mathbb{F}_p^n when p is allowed to grow as a function of n .

Part I

Low Degree Testing

We start [Part I](#) by presenting the Fourier analytic linearity test of Blum *et al.* [20] in [Chapter 2](#). In [Chapter 3](#), we show how this test can be generalized from linear functions to polynomials of higher degree. The focus will be on the so-called “99% regime”, where the test is designed to distinguish polynomials of a given degree d from the functions that are somewhat far from them, i.e. functions that do not match with any polynomial of degree d on more than $1 - \epsilon$ fraction of the points for a small $\epsilon > 0$. In [Chapter 4](#), we turn our attention to the more challenging case of the so-called “1% regime”, where the test is supposed to distinguish functions that have some noticeable correlation with polynomials of degree d (i.e. more than a small constant $\epsilon > 0$) from functions that, similar to a typical random function, have essentially no correlation with any polynomial of degree d . [Chapter 4](#) is focused on the case of quadratic polynomials. Polynomials of higher degrees are more complex and are discussed in [Part II](#) of the survey.

2

Fourier Analytic Property Testing

The field of property testing is the study of algorithms that query their input a very small number of times and with high probability decide correctly whether their input satisfies a given property or is “far” from satisfying that property. In this chapter we survey some of the earliest yet important results in the area of property testing that use classical Fourier analysis to design efficient tests for certain algebraic properties. In particular, we analyze the linearity test of Blum *et al.* [20] which together with a paper by Babai *et al.* [6] are often considered as the earliest explicit examples of property testing. These results inspired Rubinfeld and Sudan [69] and Goldreich *et al.* [32] to formally define this field and initiate a systematic study of testable properties. A property is called *testable*, or sometimes *strongly testable* or *locally testable*, if the number of queries can be made independent of the size of the object without affecting the correctness probability. Perhaps surprisingly, it has been found that many natural properties satisfy this strong requirement. In this survey we will only discuss algebraic properties, however we refer the reader to [29, 68, 65, 74] for a general overview of this field.

Let $[R]$ denote the set $\{1, \dots, R\}$. Let \mathbb{F} denote a finite field, and \mathbb{F}^n denote an n -dimensional vector space over \mathbb{F} . Given two functions $f, g : \mathbb{F}^n \rightarrow [R]$, their *distance* is the fraction of points on which they disagree,

$$\text{dist}(f, g) := \Pr_{x \in \mathbb{F}^n} [f(x) \neq g(x)],$$

where here and throughout $\Pr_{x \in \mathbb{F}^n}[\cdot]$ means the probability of an event given a uniform choice of $x \in \mathbb{F}^n$.

A *property* is a subset of functions, $\mathcal{P} \subset \{f : \mathbb{F}^n \rightarrow [R] : n \in \mathbb{Z}_{\geq 0}\}$. The distance of $f : \mathbb{F}^n \rightarrow [R]$ is ε -far from \mathcal{P} is its minimal distance to a function in \mathcal{P} ,

$$\text{dist}(f, \mathcal{P}) := \min_{g \in \mathcal{P}} \text{dist}(f, g),$$

where the minimum is taken over all functions $g \in \mathcal{P}$ defined over the same domain as f (namely, $g : \mathbb{F}^n \rightarrow [R]$). If $\text{dist}(f, \mathcal{P}) \leq \varepsilon$ then we say that f is ε -close to \mathcal{P} , and otherwise we say that it is ε -far from \mathcal{P} .

Definition 2.1 (Testability with one-sided error). A property \mathcal{P} is said to be *testable* with one-sided error if there are functions $q : (0, 1) \rightarrow \mathbb{Z}_{>0}$, $\delta : (0, 1) \rightarrow (0, 1)$, and an algorithm T that, given as input a parameter $\varepsilon > 0$ and oracle access to a function $f : \mathbb{F}^n \rightarrow [R]$, makes at most $q(\varepsilon)$ queries to the oracle for f , always accepts if $f \in \mathcal{P}$ and rejects with probability at least $\delta(\varepsilon)$ if f is ε -far from \mathcal{P} . If, furthermore, q is a constant function, then \mathcal{P} is said to be *proximity-obliviously testable* (*PO testable*).

In the above definition we only allow one-sided error. These are algorithms that do not err when the function satisfies the property. However, it is also natural to study tests with two-sided error, and indeed in [Chapter 17](#) we will see examples of such tests.

The term proximity-oblivious testing was coined by Goldreich and Ron [\[35\]](#). Indeed as we will see in [Section 2.1](#), the famous linearity test of Blum *et al.* [\[20\]](#) is an example of a proximity oblivious test. It shows that the linearity of a function $f : \mathbb{F}^n \rightarrow \mathbb{F}$ is testable using only 3 queries. This test accepts if f is linear and rejects with probability $\Omega(\varepsilon)$ if f is ε -far from linear.

2.1 Linearity testing

Linearity testing is a cornerstone of property testing, and in particular the birthplace of algebraic property testing. There are two equivalent definitions for linear functions. A function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ is *linear* if

- (1) **Global definition.** $f(x) = \sum_{i=1}^n a_i x_i$ for some $a_i \in \mathbb{F}_2$.
- (2) **Local definition.** For all $x, y \in \mathbb{F}_2^n$, $f(x + y) = f(x) + f(y)$.

Our goal will be to determine whether a given function f is linear, or whether it is far from it. The distance of a function f to the family of linear functions is given by

$$\min_{g: \mathbb{F}_2^n \rightarrow \mathbb{F}_2 \text{ linear}} \Pr[f(x) \neq g(x)],$$

where the probability is over a uniform choice of $x \in \mathbb{F}_2^n$. The BLR tester of Blum *et al.* [20] uses the local definition of linearity: it chooses random x and y accepts the function if $f(x + y) = f(x) + f(y)$.

BLR test. With query access to $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$:

1. Choose $x, y \in \mathbb{F}_2^n$ independently and uniformly at random.
2. Query $f(x), f(y), f(x + y)$.
3. **Accept** if $f(x) + f(y) = f(x + y)$ and **Reject** otherwise.

It is clear that the tester always accepts linear functions. It turns out that the probability that the above tester rejects is directly related to the distance of f from linearity. We follow the proof of Bellare *et al.* [9], which simplifies and improves the original analysis of Blum *et al.* [20]. The proof relies on classical Fourier analysis. As there are a number of good books on the topic, for example an excellent book of O’Donnell [63], we assume basic familiarity with classical Fourier analysis (concretely: the definition of Fourier coefficients, the Fourier inversion formula and Parseval’s identity).

Theorem 2.2 (Bellare *et al.* [9]). Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ be a function whose distance to linear functions is ε . Then the BLR test rejects f with probability between ε and 5ε .

Proof. Let $F(x) = (-1)^{f(x)}$. We can express the acceptance probability of the test in terms of the Fourier coefficients of F , which to recall are given by the Fourier decomposition $F(x) = \sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}(\alpha) (-1)^{\langle \alpha, x \rangle}$. Below, all probabilities and expectations are taken over uniformly chosen $x, y \in \mathbb{F}_2^n$.

$$\begin{aligned} \Pr_{x,y}[f(x) + f(y) = f(x + y)] &= \frac{1}{2} + \frac{1}{2} \mathbb{E}_{x,y} [F(x)F(y)F(x + y)] \\ &= \frac{1}{2} + \frac{1}{2} \sum_{\alpha, \beta, \gamma \in \mathbb{F}_2^n} \widehat{F}(\alpha) \widehat{F}(\beta) \widehat{F}(\gamma) \mathbb{E}_{x,y} [(-1)^{\langle \alpha, x \rangle + \langle \beta, y \rangle + \langle \gamma, x+y \rangle}] \\ &= \frac{1}{2} + \frac{1}{2} \sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}(\alpha)^3. \end{aligned}$$

Here, we used the fact that

$$\mathbb{E}_{x,y} [(-1)^{\langle \alpha, x \rangle + \langle \beta, y \rangle + \langle \gamma, x+y \rangle}] = \mathbb{E}_x [(-1)^{\langle \alpha + \gamma, x \rangle}] \mathbb{E}_y [(-1)^{\langle \beta + \gamma, x \rangle}]$$

is equal to 1 if $\alpha = \beta = \gamma$, and equal to 0 otherwise.

To prove the upper bound on the acceptance probability, we can bound

$$\Pr_{x,y}[f(x) + f(y) = f(x + y)] \leq \frac{1}{2} \left(1 + \max_{\alpha \in \mathbb{F}_2^n} \widehat{F}(\alpha) \right),$$

where we applied Parseval's identity $\sum \widehat{F}(\alpha)^2 = \mathbb{E} [F(x)^2] = 1$. If f has distance ε from linear functions, then for any linear function $g(x) = \langle x, \alpha \rangle$ it holds that $\Pr[f(x) \neq g(x)] \geq \varepsilon$. Then for any $\alpha \in \mathbb{F}_2^n$,

$$\widehat{F}(\alpha) = \mathbb{E}_x [F(x) (-1)^{\langle x, \alpha \rangle}] = \Pr[f(x) = g(x)] - \Pr[f(x) \neq g(x)] \leq 1 - 2\varepsilon.$$

We thus obtain that

$$\Pr_{x,y}[f(x) + f(y) = f(x + y)] \leq 1 - \varepsilon.$$

For the lower bound, if the distance of f to linear functions is ε , then there exists $\alpha^* \in \mathbb{F}_2^n$ such that $\widehat{F}(\alpha^*) = 1 - 2\varepsilon$. Thus $\widehat{F}(\alpha^*)^3 \geq 1 - 6\varepsilon$

and the contribution of all other terms is bounded by $\sum_{\alpha \neq \alpha^*} |\widehat{F}(\alpha)|^3 \leq \sum_{\alpha \neq \alpha^*} |\widehat{F}(\alpha)|^2 = 1 - \widehat{F}(\alpha^*)^2 \leq 4\varepsilon$. So

$$\begin{aligned} \Pr_{x,y}[f(x) + f(y) = f(x + y)] &= \frac{1}{2} \left(1 + \widehat{F}(\alpha^*)^3 + \sum_{\alpha \neq \alpha^*} \widehat{F}(\alpha)^3 \right) \\ &\geq 1 - 5\varepsilon. \end{aligned}$$

□

2.2 Testing for affine linearity

A very related notion to linear functions is that of being an affine linear function. Again, there are two equivalent definitions. A function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ is *affine linear* if

- (1) **Global definition.** $f(x) = a_0 + \sum_{i=1}^n a_i x_i$ for some $a_i \in \mathbb{F}_2$.
- (2) **Local definition.** For all $x, y, z \in \mathbb{F}_2^n$, $f(x) + f(y) + f(z) = f(x + y + z)$.

Let \mathcal{P}_1 denote the family of affine linear functions (equivalently, functions whose degree as an \mathbb{F}_2 -polynomial is at most 1). A very similar test to the BLR test can detect if a function is affine linear or far from it.

BLR test for affine linearity. With query access to $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$:

1. Choose $x, y, z \in \mathbb{F}_2^n$ independently and uniformly at random.
2. Query $f(x), f(y), f(z), f(x + y + z)$.
3. **Accept** if $f(x) + f(y) + f(z) = f(x + y + z)$ and **Reject** otherwise.

Theorem 2.3. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ be a function whose distance to affine linear functions is ε . Then the BLR test for affine linearity rejects f with probability between ε and 6ε .

2.2. Testing for affine linearity

Proof. Let $F(x) = (-1)^{f(x)}$. We can express the acceptance probability of the test in terms of the Fourier coefficients of F as

$$\begin{aligned} \Pr_{x,y,z}[f(x) + f(y) + f(z) = f(x + y + z)] &= \frac{1}{2} + \frac{1}{2} \mathbb{E}_{x,y,z} [F(x)F(y)F(z)F(x + y + z)] \\ &= \frac{1}{2} + \frac{1}{2} \sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}(\alpha)^4. \end{aligned}$$

For the upper bound, we can apply Parseval’s identity and bound

$$\begin{aligned} \Pr_{x,y,z}[f(x) + f(y) + f(z) = f(x + y + z)] &\leq \frac{1}{2} \left(1 + \max_{\alpha \in \mathbb{F}_2^n} |\widehat{F}(\alpha)|^2 \right) \\ &\leq \frac{1}{2} \left(1 + \max_{\alpha \in \mathbb{F}_2^n} |\widehat{F}(\alpha)| \right). \end{aligned}$$

If f has distance ε from affine linear functions, then for any linear function $g(x) = \langle x, \alpha \rangle$ it holds that $\Pr[f(x) \neq g(x)] \geq \varepsilon$ and $\Pr[f(x) \neq g(x) + 1] \geq \varepsilon$. Then for any $\alpha \in \mathbb{F}_2^n$,

$$\left| \widehat{F}(\alpha) \right| = \left| \mathbb{E}_x [F(x)(-1)^{\langle x, \alpha \rangle}] \right| \leq 1 - 2\varepsilon.$$

Thus

$$\Pr_{x,y,z}[f(x) + f(y) + f(z) = f(x + y + z)] \leq 1 - \varepsilon.$$

For the lower bound, if the distance of f to affine linear functions is ε , then there exists $\alpha^* \in \mathbb{F}_2^n$ such that $|\widehat{F}(\alpha^*)| = 1 - 2\varepsilon$. By the same analysis as that of linearity testing, $\widehat{F}(\alpha^*)^4 \geq 1 - 8\varepsilon$ and $\sum_{\alpha \neq \alpha^*} \widehat{F}(\alpha)^4 \leq 4\varepsilon$. So

$$\begin{aligned} \Pr_{x,y,z}[f(x) + f(y) + f(z) = f(x + y + z)] &= \frac{1}{2} \left(1 + \widehat{F}(\alpha^*)^4 + \sum_{\alpha \neq \alpha^*} \widehat{F}(\alpha)^4 \right) \\ &\geq 1 - 6\varepsilon. \end{aligned}$$

□

2.3 Limitations of Fourier analysis

The analysis of linearity testing makes use of the fact that functions that are close to linear, must have a large Fourier coefficient; and those that are close to affine linear, must have a large Fourier coefficient in absolute value. Unfortunately, this property does not carry over if we are interested in testing for higher-degree behavior, such as quadratic polynomials. As the following example shows, there are quadratic polynomials which have negligible Fourier coefficients. This will necessitate the introduction of higher-order analogs of Fourier analysis.

Claim 2.4. Assume n is even and consider the quadratic polynomial $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ defined as

$$f(x) = x_1x_2 + x_3x_4 + \dots + x_{n-1}x_n.$$

Let $F(x) = (-1)^{f(x)}$. Then $|\widehat{F}(\alpha)| = 2^{-n/2}$ for all $\alpha \in \mathbb{F}_2^n$.

Proof. Fix $\alpha \in \mathbb{F}_2^n$. Then

$$\begin{aligned} \widehat{F}(\alpha) &= \mathbb{E}_x \left[(-1)^{f(x) + \langle x, \alpha \rangle} \right] \\ &= \prod_{i=1}^{n/2} \mathbb{E}_{x_{2i-1}, x_{2i}} \left[(-1)^{x_{2i-1}x_{2i} + \alpha_{2i-1}x_{2i-1} + \alpha_{2i}x_{2i}} \right]. \end{aligned}$$

The proof follows as for any $a, b \in \mathbb{F}_2$,

$$\begin{aligned} \mathbb{E}_{x,y \in \mathbb{F}_2} \left[(-1)^{xy + ax + by} \right] &= (-1)^{ab} \mathbb{E}_{x,y \in \mathbb{F}_2} \left[(-1)^{(x+a)(y+b)} \right] \\ &= (-1)^{ab} \mathbb{E}_{x,y \in \mathbb{F}_2} \left[(-1)^{xy} \right] = \frac{(-1)^{ab}}{2}. \end{aligned}$$

Thus,

$$\widehat{F}(\alpha) = 2^{-n/2} \cdot (-1)^{\alpha_1\alpha_2 + \dots + \alpha_{n-1}\alpha_n}.$$

□

3

Low-degree Tests, the 99% Regime

In [Section 2.1](#) we showed that linearity can be tested by a very simple test: sample a random pair $x, y \in \mathbb{F}_2^n$ and check if $f(x+y) = f(x) + f(y)$. This tempts one to suspect that similar tests would perform well even for more complex properties which similarly have a local definition. One such property is that of being a low-degree polynomial. Given query access to a function, we are interested in testing whether the function is a low-degree polynomial or far from any low-degree polynomial. The testability of low-degree polynomials was first proved by Alon *et al.* [3], where they showed that the local characterization of low-degree polynomials can be used to design a natural test for this task. The analysis was sharpened by Bhattacharyya *et al.* [18]. In this chapter, we focus on the “99% regime”, where the goal is to distinguish functions very close to low-degree polynomials, from functions which are somewhat far from low-degree polynomials.

3.1 Basic properties of low-degree polynomials

Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$, and $d \geq 1$ be an integer. A function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ is a polynomial of degree (at most) d if one of the following two equivalent conditions hold:

- (1) **Global definition.** $f(x) = \sum_{S \subseteq [n], |S| \leq d} a_S \prod_{i \in S} x_i$ for some $a_S \in \mathbb{F}_2$.
- (2) **Local definition.** For all $x, y_1, \dots, y_{d+1} \in \mathbb{F}_2^n$, it holds that $\sum_{S \subseteq [d+1]} f(x + \sum_{i \in S} y_i) = 0$.

We denote by \mathcal{P}_d the family of all polynomials of degree *at most* d . In order to prove that the two definitions for degree d polynomials agree, we need to first prove a few basic facts about polynomials.

A very useful notion is that of directional derivatives. For $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ and $y \in \mathbb{F}_2^n$, the directional derivative of f in direction y , denoted $D_y f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$, is defined as

$$D_y f(x) := f(x + y) + f(x).$$

Note that D_y is a linear operator, that is $D_y(f + g) = D_y f + D_y g$. Iterated derivatives are defined as $D_{y_1, \dots, y_k} f := D_{y_1} \dots D_{y_k} f$. It is straightforward to verify that

$$D_{y_1, \dots, y_k} f(x) = \sum_{S \subseteq [k]} f\left(x + \sum_{i \in S} y_i\right).$$

Then, the local definition of f being a degree d polynomial is equivalent to $D_{y_1, \dots, y_{d+1}} f = 0$ for all $y_1, \dots, y_{d+1} \in \mathbb{F}_2^n$. We next prove some facts about derivatives.

Claim 3.1. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ be a polynomial of degree d (according to the global definition). Then $D_y f$ is a polynomial of degree at most $d - 1$ (according to the global definition).

Proof. It suffices to prove the claim for monomials of degree d . Let $m(x) = \prod_{i=1}^d x_i$. Then

$$\begin{aligned} D_y m(x) &= m(x + y) + m(x) \\ &= \prod_{i=1}^d (x_i + y_i) + \prod_{i=1}^d x_i \\ &= \sum_{S \subsetneq [d]} \prod_{i \in S} x_i \prod_{i \notin S} y_i. \end{aligned}$$

All the x -monomials in $D_y m$ have degree at most $d - 1$, and hence $D_y m$ is a polynomial of degree at most $d - 1$. □

Claim 3.2. Let $m(x) = \prod_{i=1}^d x_i$. Then for any $y_1, \dots, y_{d+1} \in \mathbb{F}_2^n$,

1. $D_{y_1, \dots, y_d} m(x) = \sum_{\pi} \prod_{i=1}^d y_{i, \pi(i)}$, where π ranges over all permutations of $\{1, \dots, d\}$. In particular, it is independent of x .
2. $D_{y_1, \dots, y_{d+1}} m = 0$.

Proof. The second claim follows immediately from the first, as any derivative of a constant function is zero. We prove the first claim by induction on d . We have that

$$\begin{aligned} D_{y_1, \dots, y_d} m(x) &= D_{y_1, \dots, y_{d-1}} D_{y_d} m(x) \\ &= D_{y_1, \dots, y_{d-1}} \left(\sum_{S \subseteq [d]} \prod_{i \in S} x_i \prod_{i \notin S} y_{d,i} \right). \end{aligned}$$

All the monomials in $D_{y_d} m(x)$ of degree less than $d-1$ will be annihilated by taking $d-1$ derivatives. Hence

$$D_{y_1, \dots, y_d} m(x) = D_{y_1, \dots, y_{d-1}} \left(\sum_{j \in [d]} y_{d,j} \prod_{i \in [d] \setminus \{j\}} x_i \right).$$

By induction, we have for any $j \in [d]$ that

$$D_{y_1, \dots, y_{d-1}} \left(\prod_{i \in [d] \setminus \{j\}} x_i \right) = \sum_{\sigma} \prod_{i \in [d] \setminus \{j\}} y_{i, \sigma(i)},$$

where σ enumerates all one-to-one functions from $[d-1]$ to $[d] \setminus \{j\}$. The claim follows by linearity. □

We now prove that the global and local definitions for being a polynomial of degree at most d are equivalent.

Lemma 3.3. The global and local definitions of being a degree d polynomial are equivalent.

Proof. Let $f(x)$ be a polynomial which has degree exactly d according to the global definition. We need to show that its degree according to the local definition is also exactly d . That is, we need to show that $D_{y_1, \dots, y_{d+1}} f = 0$ for all $y_1, \dots, y_{d+1} \in \mathbb{F}_2^n$, but that $D_{y_1, \dots, y_d} f \neq 0$ for some $y_1, \dots, y_d \in \mathbb{F}_2^n$.

The first claim follows immediately from [Claim 3.1](#). For the latter claim, let $f(x) = \sum_{S \subseteq [n], |S| \leq d} a_S \prod_{i \in S} x_i$ where $a_S \neq 0$ for some S with $|S| = d$. By [Claim 3.2](#),

$$D_{y_1, \dots, y_d} f(x) = \sum_{S \subseteq [n], |S|=d} a_S \sum_{\pi} \prod_{i=1}^d y_{i, \pi(i)}.$$

This is a nonzero polynomial (according to the global definition) in the variables $\{y_{i,j} : i \in [d], j \in [n]\}$, as all the monomials $\prod_{i=1}^d y_{i, \pi(i)}$ are distinct and some have nonzero coefficient. Hence, there must exist a value for y_1, \dots, y_d which makes it nonzero. For example, if $S = \{j_1, \dots, j_d\}$ satisfies $a_S \neq 0$, then we can choose $y_{i, j_i} = 1$ and $y_{i, j} = 0$ for all $j \neq j_i$. \square

We will also need the basic fact that the minimal distance of \mathcal{P}_d is 2^{-d} . For a proof see e.g. [\[62\]](#).

Claim 3.4. Let $f, g \in \mathcal{P}_d$ be distinct polynomials of degree d . Then $\Pr[f(x) \neq g(x)] \geq 2^{-d}$.

3.2 Low-degree testing

The question of testing low-degree polynomials was first studied by Alon *et al.* [\[3\]](#), where they showed that an appropriate extension of the linearity test also works for low-degree polynomials. The original test defined in [\[3\]](#) is specific to polynomials of degree at most d with $f(0) = 0$, but with minor changes it extends to the more natural family of all polynomials of degree at most d , which is the test presented below.

AKKLR(d) test. With query access to $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$:

1. Choose $x, y_1, \dots, y_{d+1} \in \mathbb{F}_2^n$ independently and uniformly at random.
2. Query $f(x + \sum_{i \in S} y_i)$ for all $S \subseteq [d + 1]$.
3. **Accept** if $\sum_{S \subseteq [d+1]} f(x + \sum_{i \in S} y_i) = 0$ and **Reject** otherwise.

The reader can verify that this is indeed an extension of the BLR test for affine linear functions, which correspond to $d = 1$. There, we tested if $f(x) + f(y) + f(z) = f(x + y + z)$ for uniformly chosen $x, y, z \in \mathbb{F}_2^n$. Here, we test that $f(x) + f(x + y_1) + f(x + y_2) + f(x + y_1 + y_2) = 0$ for uniformly chosen $x, y_1, y_2 \in \mathbb{F}_2^n$. The tests are identical as can be seen by setting $y_1 = x + y, y_2 = x + z$. We denote by $\delta_d(f)$ the distance of a function f from degree d polynomials,

$$\delta_d(f) := \text{dist}(f, \mathcal{P}_d).$$

Theorem 3.5 (Alon et al. [3]). The AKKLR(d) test rejects every function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ with probability at least $\Omega\left(\frac{\delta_d(f)}{d \cdot 2^d}\right)$.

If we want to reject functions which are ε -far from \mathcal{P}_d with constant probability, then we can simply repeat the basic AKKLR(d) test $O\left(\frac{d}{\varepsilon} \cdot 2^d\right)$ times. This gives a tester with query complexity $O\left(\frac{d}{\varepsilon} \cdot 4^d\right)$. It was shown in [3] that any tester for \mathcal{P}_d must make at least $\Omega(2^d + \frac{1}{\varepsilon})$ queries, leaving a quadratic gap between their upper and lower-bounds. More recently, Bhattacharyya et al. [18] gave a tighter analysis of this tester and closed this gap.

Theorem 3.6 (Bhattacharyya et al. [18]). Fix $1 \leq d \leq n$. For all functions $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$,

$$\Pr[\text{AKKLR}(d) \text{ test rejects } f] \geq c \cdot \min\{2^d \delta_d(f), 1\},$$

where $c > 0$ is an absolute constant.

Given the improved analysis, if one wishes to reject functions which are ε -far from \mathcal{P}_d with constant probability, a tester can repeat the AKKLR(d) test only $O\left(1 + \frac{1}{\varepsilon^{2d}}\right)$ times, obtaining an asymptotically optimal query complexity of $O\left(2^d + \frac{1}{\varepsilon}\right)$. We present the analysis of Bhattacharyya *et al.* [18] in the next section.

3.3 Analysis of the AKKLR test

In this section we present a proof of [Theorem 3.6](#). First, we describe a variant of the test which is easier to analyze, and is essentially equivalent to the AKKLR(d) test.

$(d + 1)$ -flat test. With query access to $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$:

1. Pick A , a random affine $(d + 1)$ -dimensional subspace of \mathbb{F}_2^n .
2. Query $f|_A$, the restriction of f to A .
3. Reject if $f|_A$ is not a degree d polynomial, or equivalently if $\sum_{x \in A} f(x) \neq 0$.

Theorem 3.6 follows from the following theorem analyzing the $(d + 1)$ -flat test.

Theorem 3.7. Fix $1 \leq d \leq n$. For all functions $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$,

$$\Pr[(d + 1)\text{-flat test rejects } f] \geq c \min\{2^d \delta_d(f), 1\},$$

where $c > 0$ is an absolute constant.

First we show how [Theorem 3.6](#) follows from [Theorem 3.7](#).

Proof of [Theorem 3.6](#) given [Theorem 3.7](#). We may assume that $n > d + 1$. Let $y_1, \dots, y_{d+1} \in \mathbb{F}_2^n$ be uniformly chosen. The probability that y_1, \dots, y_{d+1} are linearly independent is at least $1 - 2^{d+1-n}$. Conditioned on this event, they span a uniform $(d + 1)$ -flat. Hence,

$$\begin{aligned} \Pr[\text{AKKLR}(d) \text{ test rejects } f] &\geq \Pr[(d + 1)\text{-flat test rejects } f](1 - 2^{d+1-n}) \\ &\geq (c/2) \min(2^d \delta_d(f), 1). \end{aligned}$$

□

The proof of [Theorem 3.7](#) is split into two parts, based on proximity of f to degree d polynomials. We first analyze the case where $\delta_d(f)$ is small.

Lemma 3.8 (Small proximity to degree d polynomials). For any function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$,

$$\Pr[(d + 1)\text{-flat test rejects } f] \geq 2^{d+1} \delta_d(f)(1 - 2^{d+1} \delta_d(f)).$$

In particular, if $\delta_d(f) \leq 2^{-(d+2)}$, then

$$\Pr[(d + 1)\text{-flat test rejects } f] \geq 2^d \delta_d(f).$$

Proof. The main idea is to lower-bound the probability that in the $(d+1)$ -flat A , there is exactly one point where f and the closest polynomial of degree d differ. Note that in this case, the tester rejects f . Let $g \in \mathcal{P}_d$ be a polynomial of degree d such that $\Pr[f(x) = g(x)] = \delta_d(f)$. Consider a random $(d + 1)$ -dimensional affine subspace A , generated by picking a random $x \in \mathbb{F}_2^n$ and a full-rank matrix $M \in \mathbb{F}_2^{n \times (d+1)}$ and letting $A = x + M\mathbb{F}_2^{d+1} = \{x + My \mid y \in \mathbb{F}_2^{d+1}\}$.

For $a \in \mathbb{F}_2^{d+1}$, let E_a be the event that $f(x + Ma) \neq g(x + Ma)$, and let F_a be the event that $f(x + Ma) \neq g(x + Ma)$ and for every $b \neq a$, $f(x + Mb) = g(x + Mb)$. Note that if F_a occurs then the tester rejects f , and that the events F_a are pairwise disjoint. Hence

$$\Pr[(d + 1)\text{-flat test rejects } f] \geq \Pr[\cup F_a] = \sum_{a \in \mathbb{F}_2^{d+1}} \Pr[F_a].$$

Next, for $a \in \mathbb{F}_2^{d+1}$ we have that

$$\Pr[F_a] \geq \Pr[E_a] - \sum_{b \neq a} \Pr[E_a \cap E_b].$$

Moreover $\Pr[E_a] = \delta_d(f)$ and $\Pr[E_a \cap E_b] = \Pr[E_a] \Pr[E_b] = \delta_d(f)^2$, as the events E_a, E_b are pairwise independent (since M has full rank). So

$$\Pr[(d + 1)\text{-flat test rejects } f] \geq 2^{d+1} \delta_d(f) (1 - 2^{d+1} \delta_d(f)).$$

□

So, from now on we consider the case where f is at least $2^{-(d+2)}$ -far from \mathcal{P}_d , and show that in such a case, the $(d + 1)$ -flat test rejects f with constant probability (independent of d, n). It will be useful to introduce yet another variant of the tester, called the k -flat test, for $k \geq d + 1$, where the tester first samples a k -flat and then tests if f is a degree d polynomial when restricted to A .

k -flat test. With query access to $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$:

1. Pick A , a random affine k -dimensional subspace of \mathbb{F}_2^n .
2. Query $f|_A$, the restriction of f to A .
3. Reject if $f|_A$ is not a degree d polynomial.

The following lemma allows us to analyze the rejection probability of the k -flat test for $k = d + c$ where $c = O(1)$.

Lemma 3.9. For every n and $k \geq k' \geq d + 1$, and every $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$, we have

$$\Pr [k'\text{-flat test rejects } f] \geq \Pr [k\text{-flat test rejects } f] \cdot 2^{-(k-k')}.$$

Proof. It suffices to prove the lemma for $k' = k - 1$, as the general case follows by an iterative application. Let A be a random k -flat and let A' be a random $k - 1$ flat inside A . We will show that if the k -flat tester rejects f , namely if $f|_A$ has degree $> d$, then with probability at least $1/2$ it holds that $f|_{A'}$ also has degree $> d$, and hence reject by the $(k - 1)$ -flat tester.

So, fix A and assume towards contradiction that $f|_A$ has degree $> d$, but that for a strict majority of $(k - 1)$ -flats (namely, hyperplanes) $H \subset A$ it holds that $f|_H$ has degree $\leq d$. First, this implies that there must exist two disjoint hyperplanes H', H'' such that $f|_{H'}, f|_{H''}$ have degree d . By interpolation, this implies that $f|_A$ has degree at most $d + 1$. Decompose $f|_A(x) = f_{d+1}(x) + f_{\leq d}(x)$ where $f_{d+1}(x)$ is the homogeneous part of f of degree $d + 1$. On any hyperplane H where $f|_H$ has degree $\leq d$ it then holds that $f_{d+1}|_H$ has degree $\leq d$. The number of such hyperplanes is by assumption more than half of all hyperplanes, namely $> \frac{1}{2}2(2^k - 1)$, which is at least 2^k . Hence, there must exist k linearly independent hyperplanes H_1, \dots, H_k on which $f|_{H_i}$ has degree $\leq d$. We will show that this implies that f_{d+1} has degree $\leq d$, which implies that $f|_A$ has degree $\leq d$, contradicting our assumption. To see that, apply an affine linear transformation mapping H_i to $\{x : x_i = 0\}$. Then f_{d+1} has degree $\leq d$ whenever we set some variable $x_i = 0$. As $k \geq d + 2$, this can only happen if f_{d+1} has degree $\leq d$. \square

3.3. Analysis of the AKCLR test

So, it suffices to prove that if $\delta_d(f) \geq 2^{-(d+2)}$ then the k -flat tester rejects it with constant probability for some $k = d + O(1)$. This is exactly what the next lemma shows.

Lemma 3.10. For any $0 < \beta < 1/24$ there exist absolute constants $\gamma, \varepsilon, c > 0$ such that the following holds for any $n \geq k \geq d + c$. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ be a function with $\delta_d(f) \geq \beta 2^{-d}$. Then

$$\Pr [k\text{-flat test rejects } f] \geq \varepsilon + \gamma \cdot 2^{d-n}.$$

Proof. The proof will require choices of $\beta < 1/24$, $\varepsilon < 1/8$, $\gamma \geq 72$ and $2^c \geq \max\{\frac{4\gamma}{1-8\varepsilon}, \frac{\gamma}{1-\varepsilon}, \frac{2}{\beta}\}$. Fix $k \geq d + c$. We apply induction on $n \geq k$. For the base case of $n = k$, the k -flat test rejects f with probability 1. By our choice of parameters, $1 > \varepsilon + \gamma \cdot 2^{d-n}$.

Let \mathcal{H} denote the set of all hyperplanes of \mathbb{F}_2^n and let $N := |\mathcal{H}| = 2(2^n - 1)$. Also, let \mathcal{H}^* be the set of all hyperplanes $A \in \mathcal{H}$ such that $\delta_d(f|_A) < \beta 2^{-d}$, and let $K := |\mathcal{H}^*|$. We have

$$\Pr [k\text{-flat test rejects } f] = \mathbb{E}_{A \in \mathcal{H}} [\Pr [k\text{-flat test rejects } f|_A]].$$

By the induction hypothesis for every $A \in \mathcal{H} \setminus \mathcal{H}^*$, the k -flat test rejects $f|_A$ with probability at least $\varepsilon + \gamma \cdot \frac{2^d}{2^{n-1}}$, and thus

$$\begin{aligned} \Pr [k\text{-flat test rejects } f|_A] &\geq \left(1 - \frac{K}{N}\right) \left(\varepsilon + \gamma \cdot \frac{2^d}{2^{n-1}}\right) \\ &\geq \varepsilon + \gamma \cdot \frac{2^d}{2^{n-1}} - \frac{K}{N}. \end{aligned}$$

If $K \leq \gamma 2^d$ then we are done. So, assume from now on that $K > \gamma 2^d$. We show below in [Lemma 3.11](#) that as long as $\beta < 1/4$, this implies that

$$\delta_d(f) \leq \frac{3}{2}\beta 2^{-d} + \frac{9}{\gamma 2^d} \leq 2^{-(d+2)},$$

where the second inequality uses $\beta < 1/24$ and $\gamma \geq 72$. At this point [Lemma 3.8](#) asserts that the k -flat test (in fact, even the $(d + 1)$ -flat test), will reject f with probability at least

$$2^d \delta_d(f) \geq \beta \geq \varepsilon + \gamma 2^{-c} \geq \varepsilon + \gamma \cdot 2^{d-n}.$$

□

Lemma 3.11. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$. Assume that there exist distinct hyperplanes A_1, \dots, A_K such that $\delta_d(f|_{A_i}) \leq \alpha$. If $K > 2^{d+1}$ and $\alpha < 2^{-(d+2)}$, then

$$\delta_d(f) \leq \frac{3}{2}\alpha + \frac{9}{K}.$$

Proof. For each A_i , let P_i be a degree d polynomial α -close to $f|_{A_i}$.

Claim 3.12. If $4\alpha < 2^{-d}$, then for every i, j , $P_i|_{A_i \cap A_j} = P_j|_{A_i \cap A_j}$.

Proof. The claim is vacuously true if $A_i = \overline{A_j}$. Otherwise, $|A_i \cap A_j| = |A_i|/2 = |A_j|/2$ and

$$\text{dist}(f|_{A_i \cap A_j}, P_i|_{A_i \cap A_j}) \leq 2\alpha$$

and similarly for P_j , and thus $\text{dist}(P_i|_{A_i \cap A_j}, P_j|_{A_i \cap A_j}) \leq 4\alpha < 2^{-d}$. The claim now follows because $P_i|_{A_i \cap A_j}, P_j|_{A_i \cap A_j}$ are degree $\leq d$ polynomials, and by [Claim 3.4](#) the minimum distance of distinct degree d polynomials is 2^{-d} . \square

Observe that there are at least $\ell = \lfloor \log_2(K + 1) \rfloor > d$ linearly independent hyperplanes among A_1, \dots, A_K . Without loss of generality assume that A_1, \dots, A_ℓ are linearly independent, and that $A_i = \{x \in \mathbb{F}_2^n \mid x_i = 0\}$, by applying an appropriate affine transformation on \mathbb{F}_2^n . This way, for $i \in [\ell]$, P_i naturally corresponds to a polynomial over \mathbb{F}_2^n which does not depend on x_i . The idea of the proof now is to glue P_1, \dots, P_ℓ together to get a polynomial P close to f .

To this end, decompose $x = (x_1, \dots, x_\ell, y)$ where $x_i \in \mathbb{F}_2$ and $y \in \mathbb{F}_2^{n-\ell}$. For every i , P_i can be decomposed as

$$P_i(x_1, \dots, x_\ell, y) = \sum_{S \subseteq [\ell]} P_{i,S}(y) \prod_{j \in S} x_j,$$

where $P_{i,S}$ is a polynomial of degree at most $d - |S|$ which depends only on y . In particular $P_{i,S} = 0$ if $|S| > d$. Note that, if P_i does not depend on x_j and $j \in S$, then $P_{i,S} = 0$. The following claim follows immediately from [Claim 3.12](#).

Claim 3.13. For every $S \subseteq [\ell]$ and $i, j \in [\ell] \setminus S$, $P_{i,S}(y) = P_{j,S}(y)$.

3.3. Analysis of the AKCLR test

We can now formally glue together the polynomials P_i . For $S \subsetneq [\ell]$, define P_S to be $P_{i,S}$ for some $i \in [\ell] \setminus S$. The above claim asserts that P_S is unique. Now define the degree d polynomial P over \mathbb{F}_2^n as follows

$$P(x_1, \dots, x_\ell, y) = \sum_{S \subsetneq [\ell]} P_S(y) \prod_{i \in S} x_i.$$

Claim 3.14. For every $i \in [K]$, $P|_{A_i} = P_i|_{A_i}$.

Proof. The claim is easy to see for $i \in [\ell]$, since the coefficients of the two polynomials P and P_i are identically the same when $x_i = 0$. Now assume that $i \in K \setminus [\ell]$. First note that for any $j \in [\ell]$ and $x \in A_i \cap A_j$, it follows from Claim 3.12 that $P_i(x) = P_j(x) = P(x)$. Thus $P_i|_{(\cup_{j \in [\ell]} A_j) \cap A_i} = P|_{(\cup_{j \in [\ell]} A_j) \cap A_i}$. Now, note that since $\ell > d$ then

$$\frac{|A_i \cap (\cup_{j \in [\ell]} A_j)|}{|A_i|} \geq 1 - 2^{-\ell} > 1 - 2^{-d}.$$

The claim now follows from the minimum distance between degree d polynomials (Claim 3.4). □

Next, we will show that P is close to f . Since A_1, \dots, A_K do not necessarily cover \mathbb{F}_2^n uniformly, we show that they do so approximately uniformly. Let

$$\text{BAD} := \{z \in \mathbb{F}_2^n \mid z \text{ is contained in less than } K/3 \text{ of the hyperplanes } A_1, \dots, A_K\}.$$

Claim 3.15. $|\text{BAD}| \leq 2^n \frac{9}{K}$.

Proof. Let $z \in \mathbb{F}_2^n$ be uniformly chosen. The random variables $1_{z \in A_i}$ each have probability $1/2$, and they are pairwise independent. Let $S = \sum_{i=1}^k 1_{z \in A_i}$. Then $\mathbb{E}[S] = K/2$ and $\text{Var}[S] = K/4$. By Chebyshev's inequality,

$$\begin{aligned} \Pr[z \in \text{BAD}] &= \Pr[S \leq K/3] \leq \Pr[|S - \mathbb{E}[S]| \leq K/6] \\ &\leq \frac{K/4}{(K/6)^2} = \frac{9}{K}. \end{aligned}$$

□

The following claim now finishes the proof of the lemma.

Claim 3.16. $\text{dist}(f, P) \leq \frac{3}{2}\alpha + \frac{|\text{BAD}|}{2^n} \leq \frac{3}{2} + \frac{9}{K}$.

Proof. Pick $z \in \mathbb{F}_2^n$ and $i \in [K]$ uniformly at random. Then

$$\Pr_{i,z} [z \in A_i \wedge f(z) \neq P_i(z)] = \frac{1}{2} \mathbb{E}_i \left[\Pr_{z \in A_i} [f(z) \neq P_i(z)] \right] \leq \frac{\alpha}{2}.$$

On the other hand, since $P|_{A_i} = P_i|_{A_i}$, we have that

$$\begin{aligned} \Pr_{i,z} [z \in A_i \wedge f(z) \neq P_i(z)] &= \Pr_{i,z} [z \in A_i \wedge f(z) \neq P(z)] \\ &\geq \Pr_{i,z} [z \in A_i \wedge f(z) \neq P(z) \wedge z \notin \text{BAD}] \\ &= \Pr_z [f(z) \neq P(z) \wedge z \notin \text{BAD}] \\ &\quad \cdot \Pr_{i,z} [z \in A_i | f(z) \neq P(z) \wedge z \notin \text{BAD}] \\ &\geq \Pr_z [f(z) \neq P(z) \wedge z \notin \text{BAD}] \cdot \frac{1}{3} \\ &\geq \frac{1}{3} \left(\text{dist}(f, P) - \frac{|\text{BAD}|}{2^n} \right). \end{aligned}$$

□

□

[Theorem 3.6](#) now directly follows from [Lemma 3.8](#), [Lemma 3.9](#), and [Lemma 3.10](#).

3.4 Implications for the Gowers norms

We will introduce the Gowers uniformity norm in later chapters. For now, it suffices to know that for $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$, its $(d + 1)$ -th Gowers norm is defined as

$$\begin{aligned} \|(-1)^f\|_{U^{d+1}} \\ = \left(\Pr [\text{AKKLR}(d) \text{ accepts } f] - \Pr [\text{AKKLR}(d) \text{ rejects } f] \right)^{1/2^{d+1}}. \end{aligned}$$

It is easy to see that this expression is bounded between 0 and 1, and is equal to 1 if and only if f is a polynomial of degree d . [Theorem 3.6](#) shows that if the Gowers norm is close to 1, then f is close to a degree d polynomial.

Theorem 3.17. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$. Then for any $1 \leq d \leq n$, if $\|f\|_{U^{d+1}} \geq 1 - \varepsilon$ then $\delta_d(f) \leq c\varepsilon$, where $c > 0$ is an absolute constant.

Proof. If $\|f\|_{U^{d+1}} \geq 1 - \varepsilon$ then by definition,

$$2 \Pr[\text{AKKLR}(d) \text{ rejects } f] \leq 1 - (1 - \varepsilon)^{2^{d+1}} \leq 2^{d+1}\varepsilon.$$

Hence by [Theorem 3.6](#), we have $\delta_d(f) \leq c\varepsilon$. □

[Theorem 3.17](#) becomes trivial when $\varepsilon > 1/c$. In particular it says nothing about the case where $\|(-1)^f\|_{U^{d+1}}$ is bounded away from zero, e.g. $\|(-1)^f\|_{U^{d+1}} > 1/3$. One expects such functions to have some structure as a typical random function will satisfy $\|(-1)^f\|_{U^{d+1}} = o(1)$ with high probability. In the next section we will see a result due to Samorodnitsky [\[70\]](#) and independently Green and Tao [\[42\]](#) relating the Gowers U^3 norm to proximity to quadratic polynomials even in the regime where the distance is close to $1/2$. More precisely, they prove that if $\|(-1)^f\|_{U^3}$ is bounded away from 0, then f must have significant correlation with a quadratic polynomial. This immediately gives a tester for proximity to quadratics in the 1% regime.

4

Low-degree Tests, the 1% Regime

In [Chapter 3](#) we presented the AKKLR test for proximity to degree- d polynomials. The analysis in [Theorem 3.6](#) shows that the test is effective in the 99% regime: it can distinguish functions which are very close to degree- d polynomials from those which are somewhat far. Concretely, its rejection probability is $c \cdot \min(\delta_d(f)2^d, 1)$ for some absolute constant $c > 0$. In this chapter, we will focus on the 1% regime, where we have the more ambitious goal of detecting any non-trivial agreement with degree- d polynomials. Concretely, we would like to distinguish between functions with $\delta_d(f) \leq \frac{1}{2} - \varepsilon$ from functions with $\delta_d(f) = \frac{1}{2} - o(1)$.

Compare to the 99% regime, the 1% regime is much more complex. In the 99% regime, we are interested in functions which are close to polynomials of degree d . Hence so much of the polynomial structure is remained in these functions that by taking only few samples, we can obtain a good understanding of the global structure of the function. Indeed, as we saw in [Chapter 3](#), the proof of the AKKLR test in the 99% regime was based on such ideas. However in the 1% regime, we are concerned with functions with very little structure. For example, consider a degree d polynomial (think of it as a completely structured function), and obtain a new function from it by retaining the values of

1% of the points and assigning random values to the rest. Thus, this function is likely to agree with the polynomial on about 50.5% fraction of the inputs, only slightly more than a random function, which is likely to agree with the polynomial on about 50% of the inputs. While in this blurry picture much of the structure of the original polynomial is lost, some of it is still maintained since a typical random function cannot agree with any polynomial of degree d on 50.5% of the points. However, detecting this structure seems to be much harder as now if we sample the function, we will typically receive random values that are not related to the original polynomial.

It was conjectured independently by Samorodnitsky [70] and Green and Tao [42] that the AKKLR test can be used to detect such weak structures, and thus distinguish between functions with $\delta_d(f) \leq \frac{1}{2} - \varepsilon$ from functions with $\delta_d(f) = \frac{1}{2} - o(1)$.

Conjecture 4.1 (AKKLR test, 1% regime). Fix $d \geq 1$. For $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ let $p_d(f) = \Pr[\text{AKKLR}(d) \text{ accepts } f]$. Then

- **(Completeness)** If $\delta_d(f) \leq \frac{1}{2} - \varepsilon$ then $p_d(f) \geq \frac{1}{2} + \alpha(\varepsilon)$, where $\alpha(\varepsilon) > 0$.
- **(Soundness)** If $p_d(f) \geq \frac{1}{2} + \varepsilon$ then $\delta_d(f) \leq \frac{1}{2} - \beta(\varepsilon)$, where $\beta(\varepsilon) > 0$.

We will see that completeness holds for any $d \geq 1$ with $\alpha(\varepsilon) = \Omega(\varepsilon^{2^{d+1}})$. However, soundness turns out to be more intricate. For $d = 1$ it follows by a relatively simple extension of linearity testing. For $d = 2$ it also holds, but the proof is much more involved. It was accomplished independently by Samorodnitsky [70] and Green and Tao [42], where the analysis relies on tools from additive combinatorics. For $d = 3$ it turns out to be false! The counter-example was discovered independently by Lovett *et al.* [60] and Green and Tao [43], and will be presented in Chapter 5.

4.1 Completeness

We prove that any function at a distance noticeably smaller than $\frac{1}{2}$ from degree d polynomials, is accepted by the AKKLR test with probability noticeably larger than $\frac{1}{2}$.

Theorem 4.2. Fix $d \geq 1$. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ with $\delta_d(f) = \frac{1}{2} - \varepsilon$. Then

$$\Pr[\text{AKKLR}(d) \text{ accepts } f] \geq \frac{1}{2} \left(1 + (2\varepsilon)^{2^d} \right).$$

Proof. Let $P(x)$ be a polynomial of degree d so that $\text{dist}(f, P) = \frac{1}{2} - \varepsilon$ and hence $\mathbb{E}[(-1)^{(f+P)(x)}] = 2\varepsilon$. To recall, the AKKLR test samples $x, y_1, \dots, y_{d+1} \in \mathbb{F}_2^n$ independently and accepts f if

$$\sum_{S \subseteq [d+1]} f \left(x + \sum_{i \in S} y_i \right) = 0.$$

The test always accepts degree- d polynomials, as it holds that

$$\sum_{S \subseteq [d+1]} P \left(x + \sum_{i \in S} y_i \right) \equiv 0.$$

Thus, we can reformulate the acceptance probability of the test as

$$\begin{aligned} & \Pr[\text{AKKLR}(d) \text{ accepts } f] \\ &= \Pr_{x, y_1, \dots, y_{d+1} \in \mathbb{F}_2^n} \left[\sum_{S \subseteq [d+1]} f \left(x + \sum_{i \in S} y_i \right) = 0 \right] \\ &= \Pr_{x, y_1, \dots, y_{d+1} \in \mathbb{F}_2^n} \left[\sum_{S \subseteq [d+1]} (f + P) \left(x + \sum_{i \in S} y_i \right) = 0 \right] \\ &= \frac{1}{2} \left(1 + \mathbb{E}_{x, y_1, \dots, y_{d+1} \in \mathbb{F}_2^n} \left[(-1)^{D_{y_1, \dots, y_{d+1}}(f+P)(x)} \right] \right). \end{aligned}$$

The lemma then follows from the following claim: for any function $g : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ and any $k \geq 1$, it holds that

$$\mathbb{E}_{x, y_1, \dots, y_k \in \mathbb{F}_2^n} \left[(-1)^{D_{y_1, \dots, y_k} g(x)} \right] \geq \left(\mathbb{E}_{x \in \mathbb{F}_2^n} \left[(-1)^{g(x)} \right] \right)^{2^k}.$$

The claim holds for $k = 1$ as

$$\begin{aligned} \mathbb{E}_{x,y \in \mathbb{F}_2^n} [(-1)^{D_y g(x)}] &= \mathbb{E}_{x,y \in \mathbb{F}_2^n} [(-1)^{g(x)+g(x+y)}] \\ &= \mathbb{E}_{x,y} [(-1)^{g(x)+g(y)}] = \left(\mathbb{E}_{x \in \mathbb{F}_2^n} [(-1)^{g(x)}] \right)^2, \end{aligned}$$

and for $k > 1$ by induction:

$$\begin{aligned} \mathbb{E}_{x,y_1,\dots,y_k} [(-1)^{D_{y_1,\dots,y_k} g(x)}] &= \mathbb{E}_{y_1,\dots,y_{k-1}} \left[\mathbb{E}_{x,y_k} [(-1)^{D_{y_k} D_{y_1,\dots,y_{k-1}} g(x)}] \right] \\ &= \mathbb{E}_{y_1,\dots,y_{k-1}} \left(\mathbb{E}_x [(-1)^{D_{y_1,\dots,y_{k-1}} g(x)}] \right)^2 \\ &\geq \left(\mathbb{E}_{x,y_1,\dots,y_{k-1}} [(-1)^{D_{y_1,\dots,y_{k-1}} g(x)}] \right)^2 \\ &\geq \left(\mathbb{E}_x [(-1)^{g(x)}] \right)^{2^k}, \end{aligned}$$

where the first inequality follows by the Cauchy-Schwartz inequality, and the second by induction. \square

4.2 Soundness for $d = 1$

We show that if the AKKLR(1) test accepts a function with probability noticeably larger than $\frac{1}{2}$, then the function has a non-trivial correlation with some linear polynomial. Equivalently, it has a noticeable Fourier coefficient.

Theorem 4.3. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$. Assume that $\Pr[\text{AKKLR}(1) \text{ accepts } f] \geq \frac{1}{2} + \varepsilon$. Then $\delta_1(f) \leq \frac{1}{2} - \sqrt{\varepsilon/2}$.

Proof. As in the proof of [Theorem 2.3](#), let $F(x) = (-1)^{f(x)}$, where we expand the acceptance probability of the test in terms of the Fourier coefficients of F ,

$$\begin{aligned} \Pr[\text{AKKLR}(1) \text{ accepts } f] &= \Pr_{x,y,z \in \mathbb{F}_2^n} [f(x) + f(y) + f(z) = f(x + y + z)] \\ &= \frac{1}{2} + \frac{1}{2} \sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}(\alpha)^4. \end{aligned}$$

Then

$$2\varepsilon \leq \sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}(\alpha)^4 \leq \|\widehat{F}\|_\infty^2 \sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}(\alpha)^2 = \|\widehat{F}\|_\infty^2.$$

Thus $\|\widehat{F}\|_\infty \geq \sqrt{2\varepsilon}$. To conclude observe that $\delta_1(f) = \frac{1}{2}(1 - \|\widehat{F}\|_\infty)$. \square

Corollary 4.4. For any $\varepsilon > 0$ there is a test which makes $O(1/\varepsilon^8)$ queries to a function f and distinguishes, with high probability, between functions with $\delta_1(f) \leq \frac{1}{2} + \varepsilon$ and functions with $\delta_1(f) \geq \frac{1}{2} - o(1)$.

Proof. Consider the AKKLR(1) test. If $\delta_1(f) \leq \frac{1}{2} + \varepsilon$, then by [Theorem 4.2](#) it accepts f with probability at least $\frac{1}{2}(1 + (2\varepsilon)^4)$. If, on the other hand, $\delta_1(f) \geq \frac{1}{2} - o(1)$, then by [Theorem 4.3](#) it accepts f with probability at most $\frac{1}{2}(1 + o(1))$. Thus, if we repeat the test $\Omega(1/\varepsilon^8)$ times, then with high probability we can distinguish the two cases. \square

4.3 Soundness for $d = 2$

We show that if the AKKLR(2) test accepts a function with probability noticeably larger than $\frac{1}{2}$, then the function has a non-trivial correlation with some quadratic polynomial. We follow [\[70\]](#) below.

Theorem 4.5. Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$. Assume that AKKLR(2) accepts f with probability at least $\frac{1}{2} + \varepsilon$. Then $\delta_2(f) \leq \frac{1}{2} - \beta(\varepsilon)$, where $\beta(\varepsilon) \geq \exp(-c \cdot \log(1/\varepsilon)^4)$ for an absolute constant $c > 0$.

Fix a function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ for the remainder of this section, which we assume is accepted by the AKKLR(2) test with probability at least $\frac{1}{2} + \varepsilon$. We may assume that $n \geq c \cdot \log(1/\varepsilon)^4$, as otherwise the conclusion of [Theorem 4.5](#) is trivial. Let $F(x) = (-1)^{f(x)}$ and $F_y(x) = (-1)^{D_y f(x)} = F(x)F(x + y)$.

In order to build intuition, consider first the case that $f(x) = x^t D x$ is a quadratic polynomial. Then $F_y(x) = (-1)^{(x, (D + D^t)y)}$ and hence F_y has only one nonzero Fourier coefficient, which is at $(D + D^t)y$. Note that $D + D^t$ is a symmetric matrix with a zero diagonal. We will show that obtaining this, even approximately, implies that f is correlated with a quadratic polynomial.

Lemma 4.6. Assume there exists an $n \times n$ symmetric matrix A with a zero diagonal such that

$$\mathbb{E}_y \left[\widehat{F}_y(Ay)^2 \right] \geq \gamma.$$

Then there exists a quadratic polynomial $q(x)$ such that

$$\text{dist}(f, q) \leq \frac{1}{2} - \frac{1}{2}\sqrt{\gamma}.$$

In order to prove **Lemma 4.6** we need a few technical claims. For functions $G, H : \mathbb{F}_2^n \rightarrow \mathbb{R}$ define $GH(x) = G(x)H(x)$ and $(G * H)(x) = \mathbb{E}_{y \in \mathbb{F}_2^n} [G(y)H(x + y)]$. It is a standard fact in classical Fourier analysis that $\widehat{G * H}(\alpha) = \widehat{G}(\alpha)\widehat{H}(\alpha)$.

Claim 4.7. Let $G : \mathbb{F}_2^n \rightarrow \mathbb{R}$. Let $G_y(x) = G(x)G(x + y)$. Then $(G_y * G_y)(x) = (G_x * G_x)(y)$.

Proof. We have

$$\begin{aligned} (G_y * G_y)(x) &= \mathbb{E}_{s \in \mathbb{F}_2^n} [G_y(s)G_y(x + s)] \\ &= \mathbb{E}_{s \in \mathbb{F}_2^n} [G(s)G(s + y)G(x + s)G(x + s + y)]. \end{aligned}$$

The claim follows as the RHS is symmetric to swapping x and y . \square

Claim 4.8. Let $G, H : \mathbb{F}_2^n \rightarrow \mathbb{R}$. Let $G_y(x) = G(x)G(x + y)$ and $H_y(x) = H(x)H(x + y)$. Then

$$\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\left(\mathbb{E}_{x \in \mathbb{F}_2^n} [G_y(x)H_y(x)] \right)^2 \right] = \sum_{\alpha} \widehat{GH}(\alpha)^4.$$

Proof.

$$\begin{aligned}
 & \mathbb{E}_y \left[\left(\mathbb{E}_x [G_y(x)H_y(x)] \right)^2 \right] \\
 &= \mathbb{E}_{y,x,x'} [G_y(x)H_y(x)G_y(x')H_y(x')] \\
 &= \mathbb{E}_{y,x,x'} [G(x)G(x+y)H(x)H(x+y)G(x')G(x'+y)H(x')H(x'+y)] \\
 &= \mathbb{E}_{y,x,x'} [GH(x)GH(x+y)GH(x')GH(x'+y)] \\
 &= \mathbb{E}_y \left[\left(\mathbb{E}_x [GH(x)GH(x+y)] \right)^2 \right] \\
 &= \mathbb{E}_y [(GH * GH)(y)^2] \\
 &= \sum_{\alpha} \widehat{GH * GH}(\alpha)^2 = \sum_{\alpha} \widehat{GH}(\alpha)^4.
 \end{aligned}$$

□

Proof of Lemma 4.6. Decompose $A = D + D^t$ where D is upper triangular and let $G(x) = (-1)^{x^t D x}$. Note that $G_y(x) = G(x)G(x+y) = (-1)^{\langle x, Ay \rangle}$. We will show that FG has a noticeable Fourier coefficient. By Claim 4.8 and our assumption,

$$\begin{aligned}
 \sum_{\alpha} \widehat{FG}(\alpha)^4 &= \mathbb{E}_y \left[\left(\mathbb{E}_x [F_y(x)G_y(x)] \right)^2 \right] \\
 &= \mathbb{E}_y \left[\left(\mathbb{E}_x [F_y(x)(-1)^{\langle x, Ay \rangle}] \right)^2 \right] = \mathbb{E}_y [\widehat{F_y}(Ay)^2] \geq \gamma.
 \end{aligned}$$

This implies that FG has a large Fourier coefficient, as

$$\sum_{\alpha} \widehat{FG}(\alpha)^4 \leq \|\widehat{FG}\|_{\infty}^2 \sum_{\alpha} \widehat{FG}(\alpha)^2 = \|\widehat{FG}\|_{\infty}^2.$$

So, there exists $\alpha \in \mathbb{F}_2^n$ and $c \in \mathbb{F}_2$ such that

$$\mathbb{E}_x [F(x)G(x)(-1)^{\langle x, \alpha \rangle + c}] = |\widehat{FG}(\alpha)| \geq \sqrt{\gamma}.$$

So, the quadratic polynomial $q(x) = x^t D x + \langle x, \alpha \rangle + c$ satisfies $\text{dist}(f, q) \leq \frac{1}{2} - \frac{1}{2}\sqrt{\gamma}$. □

So, our goal from now on is to show the existence of such a matrix A for which $\widehat{F}_y(Ay)$ is noticeable for a typical y . As a first step, we show that for a typical y , F_y has some noticeable Fourier coefficients. Formally, we show that the 4th moments of \widehat{F}_y are noticeable.

Claim 4.9. $\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}_y(\alpha)^4 \right] \geq 2\epsilon.$

Proof. Let $f_y = D_y f$. We can express

$$\begin{aligned} \Pr[\text{AKKLR}(2) \text{ accepts } f] &= \Pr_{x, y_1, y_2, y_3 \in \mathbb{F}_2^n} \left[\sum_{S \subseteq [3]} f \left(x + \sum_{i \in S} y_i \right) = 0 \right] \\ &= \mathbb{E}_{y \in \mathbb{F}_2^n} \Pr_{\substack{x, y_1, y_2 \\ \in \mathbb{F}_2^n}} [f_y(x) + f_y(x + y_1) + f_y(x + y_2) + f_y(x + y_1 + y_2) = 0]. \end{aligned}$$

As in the proof of [Theorem 4.3](#), we can express the inner probability as

$$\begin{aligned} \Pr_{x, y_1, y_2 \in \mathbb{F}_2^n} [f_y(x) + f_y(x + y_1) + f_y(x + y_2) + f_y(x + y_1 + y_2) = 0] \\ = \frac{1}{2} + \frac{1}{2} \sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}_y(\alpha)^4. \end{aligned}$$

Thus

$$\Pr[\text{AKKLR}(2) \text{ accepts } f] = \frac{1}{2} \left(1 + \mathbb{E}_{y \in \mathbb{F}_2^n} \left[\sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}_y(\alpha)^4 \right] \right).$$

The claim follows from our assumption that $\Pr[\text{AKKLR}(2) \text{ accepts } f] \geq \frac{1}{2} + \epsilon.$ \square

The next step is to show that the noticeable Fourier coefficients of \widehat{F}_y have an approximate linear structure. The following lemma plays an important role in this.

Lemma 4.10.

$$\mathbb{E}_{y, z \in \mathbb{F}_2^n} \left[\sum_{\alpha, \beta \in \mathbb{F}_2^n} \widehat{F}_y(\alpha)^2 \widehat{F}_z(\beta)^2 \widehat{F}_{y+z}(\alpha + \beta)^2 \right] = \mathbb{E}_{y \in \mathbb{F}_2^n} \left[\sum_{\alpha \in \mathbb{F}_2^n} \widehat{F}_y(\alpha)^6 \right].$$

Proof. Note that $\widehat{F}_y(\alpha)^2 = \mathbb{E}_{u,u' \in \mathbb{F}_2^n} F_y(u)F_y(u')(-1)^{\langle u+u', \alpha \rangle}$. Using this, we can expand the LHS in the lemma statement as

$$\begin{aligned} \text{LHS} &= \mathbb{E}_{y,z} \left[\sum_{\alpha,\beta} \widehat{F}_y(\alpha)^2 \widehat{F}_z(\beta)^2 \widehat{F}_{y+z}(\alpha + \beta)^2 \right] \\ &= \mathbb{E}_{y,z} \sum_{\alpha,\beta} \left[\mathbb{E}_{u,u'} F_y(u)F_y(u')(-1)^{\langle u+u', \alpha \rangle} \cdot \mathbb{E}_{v,v'} F_z(v)F_z(v')(-1)^{\langle v+v', \beta \rangle} \right. \\ &\quad \left. \mathbb{E}_{w,w'} F_{y+z}(w)F_{y+z}(w') \cdot (-1)^{\langle w+w', \alpha+\beta \rangle} \right]. \end{aligned}$$

Next, note that

$$\begin{aligned} &\sum_{\alpha,\beta} (-1)^{\langle u+u', \alpha \rangle} (-1)^{\langle v+v', \beta \rangle} (-1)^{\langle w+w', \alpha+\beta \rangle} \\ &= \sum_{\alpha} (-1)^{\langle u+u'+w+w', \alpha \rangle} \sum_{\beta} (-1)^{\langle v+v'+w+w', \beta \rangle} \\ &= 2^{2n} \cdot \mathbb{1}_{[u+u'=v+v'=w+w']}. \end{aligned}$$

We can thus restrict our attention to the case that $u + u' = v + v' = w + w' = s$ for some $s \in \mathbb{F}_2^n$, and obtain a simplified expression for the LHS as

$$\begin{aligned} \text{LHS} &= \mathbb{E}_{y,z,s} \left[\mathbb{E}_u F_y(u)F_y(u+s) \cdot \mathbb{E}_v F_z(v)F_z(v+s) \right. \\ &\quad \left. \cdot \mathbb{E}_w F_{y+z}(w)F_{y+z}(w+s) \right] \\ &= \mathbb{E}_{y,z,s} [(F_y * F_y)(s)(F_z * F_z)(s)(F_{y+z} * F_{y+z})(s)] \\ &= \mathbb{E}_{y,z,s} [(F_s * F_s)(y)(F_s * F_s)(z)(F_s * F_s)(y+z)] \\ &= \mathbb{E}_s \left[\sum_{\alpha} \widehat{F_s * F_s}(\alpha)^3 \right] \\ &= \mathbb{E}_s \left[\sum_{\alpha} \widehat{F_s}(\alpha)^6 \right]. \end{aligned}$$

□

As a corollary, we obtain that the noticeable Fourier coefficients of \widehat{F}_y have an approximate linear structure, at least locally.

Corollary 4.11. $\mathbb{E}_{y,z \in \mathbb{F}_2^n} \left[\sum_{\alpha, \beta \in \mathbb{F}_2^n} \widehat{F}_y(\alpha)^2 \widehat{F}_z(\beta)^2 \widehat{F}_{y+z}(\alpha + \beta)^2 \right] \geq 4\varepsilon^2.$

Proof. By Claim 4.9 we have $\mathbb{E}_y \sum_{\alpha} \widehat{F}_y(\alpha)^4 \geq 2\varepsilon.$ By the Cauchy-Schwartz inequality

$$\begin{aligned} \sum_{\alpha} \widehat{F}_y(\alpha)^4 &= \sum_{\alpha} \widehat{F}_y(\alpha) \cdot \widehat{F}_y(\alpha)^3 \\ &\leq \sqrt{\sum_{\alpha} \widehat{F}_y(\alpha)^2 \cdot \sum_{\alpha} \widehat{F}_y(\alpha)^6} = \sqrt{\sum_{\alpha} \widehat{F}_y(\alpha)^6}. \end{aligned}$$

Thus we have

$$\mathbb{E}_y \sum_{\alpha} \widehat{F}_y(\alpha)^6 \geq \mathbb{E}_y \left(\sum_{\alpha} \widehat{F}_y(\alpha)^4 \right)^2 \geq \left(\mathbb{E}_y \sum_{\alpha} \widehat{F}_y(\alpha)^4 \right)^2 \geq 4\varepsilon^2.$$

The corollary now follows by Lemma 4.10. □

The next step is to show that a typical \widehat{F}_x has a noticeable Fourier coefficient at $\phi(x)$, where ϕ is approximately linear. This will be the starting point for finding an actual linear map for which this holds.

Lemma 4.12. There exists $X \subset \mathbb{F}_2^n$ and a map $\phi : X \rightarrow \mathbb{F}_2^n$ such that

- (i) $\widehat{F}_x(\phi(x))^2 \geq \varepsilon^2$ for all $x \in X.$
- (ii) $\Pr_{x,y \in \mathbb{F}_2^n} [x, y, x + y \in X \wedge \phi(x) + \phi(y) = \phi(x + y)] \geq \varepsilon^2/2.$

Proof. Define a random function $\phi : \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ by picking, independently for each $y \in \mathbb{F}_2^n,$ $\phi(y) = \alpha$ with probability $\widehat{F}_y(\alpha)^2.$ Note that this is indeed a probability distribution as by Parseval's identity $\sum_{\alpha} \widehat{F}_y(\alpha)^2 = 1$ for each $y.$ For $\delta = \varepsilon^2$ define

$$\begin{aligned} L(\phi) := \Pr_{x,y \in \mathbb{F}_2^n} \left[\phi(x + y) = \phi(x) + \phi(y), \widehat{F}_x(\phi(x))^2 \geq \delta, \right. \\ \left. \widehat{F}_y(\phi(y))^2 \geq \delta, \widehat{F}_{x+y}(\phi(x + y))^2 \geq \delta \right]. \end{aligned}$$

It suffices to show that $\mathbb{E}_{\phi} L(\phi) \geq \varepsilon^2/2.$ Indeed for a choice of ϕ which attains the bound, by taking

$$X = \left\{ x \in \mathbb{F}_2^n : \widehat{F}_x(\phi(x))^2 \geq \delta \right\},$$

the desired items (i),(ii) will follow by definition. In order to compute $\mathbb{E}_\phi L(\phi)$, it will be useful to define

$$\Lambda := \left\{ (\alpha, \beta) \in \mathbb{F}_2^{2n} : \widehat{F}_x(\alpha)^2 \geq \delta, \widehat{F}_y(\beta)^2 \geq \delta, \widehat{F}_{x+y}(\alpha + \beta)^2 \geq \delta \right\}.$$

Then,

$$\mathbb{E}_\phi [L(\phi)] = \mathbb{E}_{x,y} \left[\sum_{(\alpha,\beta) \in \Lambda} \Pr_\phi [\phi(x) = \alpha, \phi(y) = \beta, \phi(x+y) = \alpha + \beta] \right]$$

Next, by definition of ϕ , if $x, y, x + y$ are all distinct then

$$\begin{aligned} \Pr_\phi [\phi(x) = \alpha, \phi(y) = \beta, \phi(x+y) = \alpha + \beta] \\ &= \Pr_\phi [\phi(x) = \alpha] \Pr_\phi [\phi(y) = \beta] \Pr_\phi [\phi(x+y) = \alpha + \beta] \\ &= \widehat{F}_x(\alpha)^2 \widehat{F}_y(\beta)^2 \widehat{F}_{x+y}(\alpha + \beta)^2. \end{aligned}$$

The probability that $x, y, x + y$ are not all distinct is $O(2^{-n})$. Thus,

$$\mathbb{E}_\phi [L(\phi)] \geq \mathbb{E}_{x,y} \left[\sum_{(\alpha,\beta) \in \Lambda} \widehat{F}_x(\alpha)^2 \widehat{F}_y(\beta)^2 \widehat{F}_{x+y}(\alpha + \beta)^2 \right] - O(2^{-n}).$$

We would like to extend the sum to include all choices of α, β , so that we can then apply [Corollary 4.11](#). The sum over α for which $\widehat{F}_x(\alpha)^2 < \delta$ can be bounded by

$$\begin{aligned} \mathbb{E}_{x,y} \left[\sum_{\alpha,\beta: \widehat{F}_x(\alpha)^2 < \delta} \widehat{F}_x(\alpha)^2 \widehat{F}_y(\beta)^2 \widehat{F}_{x+y}(\alpha + \beta)^2 \right] \\ &\leq \delta \cdot \mathbb{E}_{x,y} \left[\sum_{\alpha,\beta} \widehat{F}_y(\beta)^2 \widehat{F}_{x+y}(\alpha + \beta)^2 \right] \\ &= \delta \cdot \mathbb{E}_{x,y} \left[\sum_{\alpha,\beta} \widehat{F}_y(\beta)^2 \widehat{F}_{x+y}(\alpha)^2 \right] \\ &= \delta, \end{aligned}$$

where we used Parseval's identity. We can bound the sum over α, β for which $\widehat{F}_y(\beta)^2 < \delta$ or $\widehat{F}_{x+y}(\alpha + \beta)^2 < \delta$ in a similar manner. We thus have

$$\begin{aligned} \mathbb{E}_{\phi} [L(\phi)] &\geq \mathbb{E}_{x,y} \left[\sum_{\alpha, \beta \in \mathbb{F}_2^n} \widehat{F}_x(\alpha)^2 \widehat{F}_y(\beta)^2 \widehat{F}_{x+y}(\alpha + \beta)^2 \right] - 3\delta - O(2^{-n}) \\ &\geq 4\epsilon^2 - 3\delta - O(2^{-n}) \geq \epsilon^2 - O(2^{-n}) \geq \epsilon^2/2, \end{aligned}$$

where in the last inequality we used our assumption that n is large enough. □

Next, we show that ϕ can be approximated by a linear map. To that end, we will need several results in additive combinatorics. For a set $S \subseteq \mathbb{F}_2^n$ its sumset is $S + S = \{s + s' : s, s' \in S\}$. The first result is the Balog-Szemerédi-Gowers theorem (abbreviated BSG theorem) [8, 37]. It states that if many pairs in S have their sum in a small set, then there is a large subset of S with a small sumset. The proof can be found in the original paper of [37]. See also [81]. A simplified proof for the case of $G = \mathbb{F}_2^n$ can be found in an exposition of Viola [84].

Theorem 4.13 (BSG theorem [8, 37]). Let G be an Abelian group and let $S \subseteq G$. If

$$\Pr_{s, s' \in S} [s + s' \in A] \geq \epsilon,$$

then there exists $S' \subset S$ of size $|S'| \geq c\epsilon^2|S|$ such that $|S' + S'| \leq c^{-1}\epsilon^{-5}|A|$. Here, $c > 0$ is an absolute constant.

The other ingredient required is the structure of sets S for which $S + S$ is not much larger than S . Here, the best result to date is by Sanders [71]. See also the survey by Sanders [72] for more details, and the exposition by Lovett [59] giving a simplified proof for $G = \mathbb{F}_2^n$. Below, we present the result specialized to the case of $G = \mathbb{F}_2^n$.

Theorem 4.14 (Sanders [71]). Let $S \subset \mathbb{F}_2^n$ be a set such that $|S + S| \leq K|S|$. Then there exists an affine linear subspace $V \subset \mathbb{F}_2^n$ of size $|V| \leq |S|$ such that

$$|S \cap V| \geq \exp(-c \log^4 K)|S|,$$

where $c > 0$ is an absolute constant.

With these theorems at our disposal, let $S = \{(x, \phi(x)) : x \in X\} \subset \mathbb{F}_2^{2n}$ be the graph of ϕ . By Lemma 4.12 we have that $\Pr_{s_1, s_2 \in S}[s_1 + s_2 \in S] \geq \varepsilon^2/2$. By Theorem 4.13, there exists a subset $S' \subset S$ of size $|S'| \geq c'\varepsilon^4$ such that $|S' + S'| \leq c''\varepsilon^{-10}|S'|$. By Theorem 4.14, there exists an affine linear subspace $V \subset \mathbb{F}_2^{2n}$ such that $S' \cap V$ is large and $|V| \leq |S'|$. We will only use the fact that $|V| \leq 2^n$ and that $|S \cap V| \geq \eta 2^n$ where $\eta = \exp(-c'''\log^4(1/\varepsilon))$ (here, $c', c'', c''' > 0$ are unspecified absolute constants). Next, we show that this implies the existence of a linear map which approximates ϕ .

Lemma 4.15. There exist an $n \times n$ matrix A and a vector $b \in \mathbb{F}_2^n$ such that

$$\Pr_{x \in S}[\phi(x) = Ax + b] \geq \eta^2.$$

Proof. Let $U = \{x : (x, y) \in V\}$ be the projection of V to the first n coordinates. As $|S \cap V| \geq \eta 2^n$ and S contains at most one element (x, y) for each $x \in \mathbb{F}_2^n$, it must be that $|U| \geq \eta 2^n$. Thus, we can decompose V as the disjoint union of subspaces of the form $V_i = \{(x, \ell_i(x)) : x \in U\}$, where $\ell_i : U \rightarrow \mathbb{F}_2^n$ are affine linear maps, and $1 \leq i \leq |V/U|$. Choose $\ell = \ell_i$ which maximizes $|S \cap V_i|$. For this choice, we get $|S \cap V_i| \geq \eta|S \cap V| \geq \eta^2 2^n$. If $|U| < \mathbb{F}_2^n$ then complete ℓ arbitrarily to a linear map from \mathbb{F}_2^n . The lemma follows as the fact that ℓ is an affine linear map implies $\ell(x) = Ax + b$. \square

Corollary 4.16. $\nu := \mathbb{E}_{y \in \mathbb{F}_2^n} [\widehat{F}_y(Ay + b)^2] \geq \eta^3 \varepsilon^2$.

Proof. For any $y \in S$ we have $\widehat{F}_y(\phi(y))^2 \geq \varepsilon^2$. There are $\eta^2|S|$ elements $y \in S$ for which $\phi(y) = Ay + b$. Finally, $|S| \geq |S \cap V| \geq \eta 2^n$. \square

To conclude, we would want to apply Lemma 4.6. To do so, we need to show that we can take (i) A to be symmetric with zero diagonal; and (ii) $b = 0$. The following claim will be useful for both. For $M \in \{A, A^t\}$, define $R_M : \mathbb{F}_2^n \rightarrow \mathbb{R}$ as

$$R_M(z) := \sum_{y \in \mathbb{F}_2^n} \widehat{F}_y(My + z)^2.$$

First, we prove a general claim on the Fourier coefficients of R_M .

Claim 4.17. $\widehat{R}_M(\alpha) = \widehat{F}_\alpha(M^t\alpha)^2$.

Proof.

$$\begin{aligned} \widehat{R}_M(\alpha) &= \mathbb{E}_{z \in \mathbb{F}_2^n} \left[\sum_y \widehat{F}_y(My + z)^2 (-1)^{\langle \alpha, z \rangle} \right] \\ &= \mathbb{E}_{z \in \mathbb{F}_2^n} \left[\sum_y \mathbb{E}_{u, u' \in \mathbb{F}_2^n} \left[F_y(u) F_y(u') (-1)^{\langle u+u', My+z \rangle + \langle \alpha, z \rangle} \right] \right] \\ &= \sum_y \mathbb{E}_{u, u' \in \mathbb{F}_2^n} \left[F_y(u) F_y(u') (-1)^{\langle u+u', My \rangle} \cdot \mathbb{E}_{z \in \mathbb{F}_2^n} \left[(-1)^{\langle u+u'+\alpha, z \rangle} \right] \right] \end{aligned}$$

The last term is 2^{-n} if $u' = u + \alpha$, and is zero otherwise. Thus, we may restrict to the case that $u' = u + \alpha$ and get

$$\begin{aligned} \widehat{R}_M(\alpha) &= 2^{-n} \sum_y \mathbb{E}_{u \in \mathbb{F}_2^n} \left[F_y(u) F_y(u + \alpha) (-1)^{\langle \alpha, My \rangle} \right] \\ &= \mathbb{E}_{y \in \mathbb{F}_2^n} \left[\mathbb{E}_{u \in \mathbb{F}_2^n} \left[F_y(u) F_y(u + \alpha) (-1)^{\langle \alpha, My \rangle} \right] \right] \\ &= \mathbb{E}_{y \in \mathbb{F}_2^n} \left[(F_y * F_y)(\alpha) (-1)^{\langle \alpha, My \rangle} \right] \\ &= \mathbb{E}_{y \in \mathbb{F}_2^n} \left[(F_\alpha * F_\alpha)(y) (-1)^{\langle \alpha, My \rangle} \right] \\ &= \widehat{F}_\alpha * \widehat{F}_\alpha(M^t\alpha) = \left(\widehat{F}_\alpha(M^t\alpha) \right)^2. \end{aligned}$$

□

Next, we show that we may assume $b = 0$.

Claim 4.18. $\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(Ay)^2 \right] \geq \nu$.

Proof. Let $R = R_A$. Since the Fourier coefficients of R are all non-negative, we have that $R(0) \geq R(z)$ for all $z \in \mathbb{F}_2^n$, as

$$R(0) = \sum_\alpha \widehat{R}(\alpha) \geq \sum_\alpha \widehat{R}(\alpha) (-1)^{\langle \alpha, z \rangle} = R(z).$$

In particular,

$$\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(Ay)^2 \right] = R(0) \geq R(b) = \mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(Ay + b)^2 \right] = \nu.$$

□

We continue with showing that A may be taken to be symmetric with zero diagonal. The following claim will be useful.

Claim 4.19. Let $y, \alpha \in \mathbb{F}_2^n$ with $\langle \alpha, y \rangle = 1$. Then $\widehat{F}_y(\alpha) = 0$.

Proof.

$$\begin{aligned} \widehat{F}_y(\alpha) &= \mathbb{E}_{x \in \mathbb{F}_2^n} \left[F(x)F(x+y)(-1)^{\langle \alpha, x \rangle} \right] \\ &= - \mathbb{E}_{x \in \mathbb{F}_2^n} \left[F(x)F(x+y)(-1)^{\langle \alpha, x+y \rangle} \right] = -\widehat{F}_y(\alpha). \end{aligned}$$

□

We next show that many large Fourier coefficients are supported on y for which $Ay = A^t y$.

Claim 4.20. $\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(Ay)^2 \cdot 1_{Ay=A^t y} \right] \geq \nu^2$.

Proof. Let $G(x) = (-1)^{x^t A x}$ and $R = R_{A^t}$. By [Claim 4.17](#), $\widehat{R}(y) = \widehat{F}_y(Ay)^2$. By [Claim 4.19](#), if $\widehat{F}_y(Ay) \neq 0$ then necessarily $\langle Ay, y \rangle = 0$, which is equivalent to $G(y) = 1$. Thus

$$G(y)\widehat{R}(y) = \widehat{R}(y).$$

This implies that

$$\nu = \mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{R}(y) \right] = \mathbb{E}_{y \in \mathbb{F}_2^n} G(y)\widehat{R}(y) = \mathbb{E}_{z \in \mathbb{F}_2^n} \left[\widehat{G}(z)R(z) \right].$$

Recall that by definition $R(z) \geq 0$. Moreover, by Parseval's identity

$$\mathbb{E}_{z \in \mathbb{F}_2^n} [R(z)] = 2^{-n} \sum_{y, z \in \mathbb{F}_2^n} \widehat{F}_y(My + z)^2 = 1.$$

Thus by Jensen inequality,

$$\mathbb{E}_{z \in \mathbb{F}_2^n} \left[\widehat{G}(z)^2 R(z) \right] \geq \left(\mathbb{E}_z \left[\widehat{G}(z)R(z) \right] \right)^2 = \nu^2.$$

On the other hand,

$$\mathbb{E}_{z \in \mathbb{F}_2^n} \left[\widehat{G}(z)^2 R(z) \right] = \mathbb{E}_{y \in \mathbb{F}_2^n} \left[(G * G)(y)\widehat{R}(y) \right].$$

Now,

$$\begin{aligned} (G * G)(y) &= \mathbb{E}_{z \in \mathbb{F}_2^n} \left[(-1)^{z^t A z + (z+y)^t A (z+y)} \right] \\ &= \mathbb{E}_{z \in \mathbb{F}_2^n} \left[(-1)^{\langle z, (A+A^t)y \rangle + y^t A y} \right] \\ &= 1_{Ay=A^t y} G(y). \end{aligned}$$

We thus obtained

$$\begin{aligned} \mathbb{E}_{y \in \mathbb{F}_2^n} \left[1_{Ay=A^t y} \widehat{F}_y(Ay)^2 \right] &= \mathbb{E}_{y \in \mathbb{F}_2^n} \left[1_{Ay=A^t y} \widehat{R}(y) \right] \\ &= \mathbb{E}_{y \in \mathbb{F}_2^n} \left[1_{Ay=A^t y} G(y) \widehat{R}(y) \right] \\ &= \mathbb{E}_{y \in \mathbb{F}_2^n} \left[(G * G)(y) \widehat{R}(y) \geq \nu^2 \right]. \end{aligned}$$

□

Let $W := \{x : Ax = A^t x\}$ be a linear subspace of \mathbb{F}_2^n . Restricted to W , A is symmetric. Thus, we can find a symmetric matrix A' such that $A'x = Ax$ for all $x \in W$. We have that

$$\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(A'y)^2 \right] \geq \mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(Ay)^2 \cdot 1_{y \in W} \right] \geq \nu^2.$$

It remains to deal with the diagonal of A' . Let $v \in \mathbb{F}_2^n$ be the diagonal of A' , and define

$$A'' = A' + vv^t.$$

Clearly, A'' is symmetric with zero diagonal. The follow claim is the last step in the proof.

Claim 4.21. $\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(A''y)^2 \right] \geq \mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(A'y)^2 \right]$.

Proof. We have $A''y = A'y + \langle y, v \rangle vv^t$. If $\langle y, v \rangle = 0$ then $A''y = A'y$. If $\langle y, v \rangle = 1$ then $y^t A'y = \langle y, v \rangle = 1$ and hence by [Claim 4.19](#), $\widehat{F}_y(A'y) = 0$. □

So, we have constructed a symmetric matrix with zero diagonal A'' for which

$$\mathbb{E}_{y \in \mathbb{F}_2^n} \left[\widehat{F}_y(A''y)^2 \right] \geq \nu^2.$$

By [Lemma 4.6](#) this implies the existence of a quadratic polynomial q whose distance from f is noticeably less than $1/2$. Concretely,

$$\text{dist}(f, q) \leq \frac{1}{2} - \frac{\nu}{2}.$$

5

Gowers Norms, the Inverse Gowers Conjecture and its Failure

To recall, the probability that the $\text{AKKLR}(d)$ test accepts a function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ is given by

$$\Pr[\text{AKKLR}(d) \text{ accepts } f] = \Pr_{x, y_1, \dots, y_{d+1} \in \mathbb{F}_2^n} \left[\sum_{S \subseteq [d+1]} f \left(x + \sum_{i \in S} y_i \right) = 0 \right].$$

The AKKLR test is intimately related to Gowers norms, which we now define.

5.1 Gowers norms

In its most general form, the Gowers norms are defined for functions $F : G \rightarrow \mathbb{C}$, where G is a finite Abelian group (as a motivating example, consider the case of $G = \mathbb{F}_2^n$ and $F(x) = (-1)^{f(x)}$). The (multiplicative) derivative of F in direction $y \in G$ is given by $\Delta_y F(x) = F(x+y)\overline{F(x)}$. Note that if $F(x) = (-1)^{f(x)}$ then $\Delta_y F = (-1)^{f(x+y)+f(x)} = (-1)^{D_y f}$. Iterative derivatives are defined as $\Delta_{y_1, \dots, y_d} F = \Delta_{y_1} \dots \Delta_{y_d} F$. The *Gowers norm* of order d for F is defined as the expected d -th multiplicative derivative of F in d random directions at a random point.

Definition 5.1 (Gowers norm). Let G be a finite Abelian group, $d \geq 1$. Given a function $F : G \rightarrow \mathbb{C}$, the *Gowers norm of order d* for F is given by

$$\begin{aligned} \|F\|_{U^d} &= \left| \mathbb{E}_{y_1, \dots, y_d, x \in G} [(\Delta_{y_1} \Delta_{y_2} \cdots \Delta_{y_d} F)(x)] \right|^{1/2^d} \\ &= \left| \mathbb{E}_{y_1, \dots, y_d, x \in G} \left[\prod_{S \subseteq [d]} \mathcal{C}^{d-|S|} F(x + \sum_{i \in S} y_i) \right] \right|^{1/2^d}, \end{aligned}$$

where \mathcal{C} is the conjugation operator $\mathcal{C}(z) = \bar{z}$.

Note that as $\|F\|_{U^1} = |\mathbb{E}[F]|$ the Gowers norm of order 1 is only a semi-norm. However, for $d > 1$ it turns out that $\|\cdot\|_{U^d}$ is indeed a norm [37]. It satisfies the following inequality, known as the Gowers Cauchy-Schwarz inequality.

Lemma 5.2 (Gowers Cauchy-Schwarz [37]). Consider a family of functions $F_S : G \rightarrow \mathbb{C}$, where $S \subseteq [d]$. Then

$$\left| \mathbb{E}_{x, y_1, \dots, y_d \in \mathbb{F}^n} \left[\prod_{S \subseteq [d]} F_S(x + \sum_{i \in S} y_i) \right] \right| \leq \prod_{S \subseteq [d]} \|F_S\|_{U^d}.$$

In this survey, we will only consider the case of $G = \mathbb{F}^n$ where \mathbb{F} is a finite field. Gowers norms are directly related to the acceptance probability of the AKKLR test. If $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ then

$$\|(-1)^f\|_{U^d}^{2^d} = \Pr[\text{AKKLR}(d) \text{ accepts } f] - \Pr[\text{AKKLR}(d) \text{ rejects } f]$$

In particular, if $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ is a polynomial of degree $\leq d$, then $\|(-1)^f\|_{U^{d+1}} = 1$. We can rephrase Theorem 4.2, Theorem 4.3 and Theorem 4.5 as a direct theorem and inverse theorems for the Gowers norm for functions $F : \mathbb{F}_2^n \rightarrow \{-1, 1\}$. The results and proofs generalize to bounded functions $F : \mathbb{F}_p^n \rightarrow \mathbb{C}$, where $p \geq 2$ is a fixed prime. For a polynomial $P : \mathbb{F}_p \rightarrow \mathbb{F}_p$ let $e(P) = \exp(2\pi i/p \cdot P(x))$. For two functions $F, G : \mathbb{F}_p^n \rightarrow \mathbb{C}$ define their inner product as $\langle F, G \rangle = \mathbb{E}_x F(x) \overline{G(x)}$.

Theorem 5.3. Let $p \geq 2$ be a fixed prime. Let $F : \mathbb{F}_p^n \rightarrow \mathbb{C}$ with $\|F\|_\infty \leq 1$. Then

- **(Direct theorem)** For any polynomial $P : \mathbb{F}_p^n \rightarrow \mathbb{F}_p$ of degree $\leq d - 1$, it holds that $|\langle F, e(P) \rangle| \leq \|F\|_{U^d}$.
- **(Inverse theorem, $d = 1$)** If $\|F\|_{U^2} \geq \varepsilon$, then there exists a polynomial $P : \mathbb{F}_p^n \rightarrow \mathbb{F}_p$ of degree $d = 1$ such that $|\langle F, e(P) \rangle| \geq \varepsilon^2$.
- **(Inverse theorem, $d = 2$)** If $\|F\|_{U^3} \geq \varepsilon$, then there exists a polynomial $P : \mathbb{F}_p^n \rightarrow \mathbb{F}_p$ of degree $d = 2$ such that $|\langle F, e(P) \rangle| \geq \delta$ where $\delta = \delta(p, \varepsilon) > 0$.

The (original) inverse Gowers conjecture, independently due to Samorodnitsky [70] and Green and Tao [42], states that the inverse theorem should hold for any $d \geq 1$.

Conjecture 5.4. Let $p \geq 2$ be a fixed prime, and let $d \geq 1$. Let $F : \mathbb{F}_p^n \rightarrow \mathbb{C}$ with $\|F\|_\infty \leq 1$. If $\|F\|_{U^{d+1}} \geq \varepsilon$ then there exists a polynomial $P : \mathbb{F}_p^n \rightarrow \mathbb{F}_p$ of degree $\leq d$ such that $|\langle F, e(P) \rangle| \geq \delta$ where $\delta = \delta(p, d, \varepsilon)$.

As we will shortly see, Conjecture 5.4 is false, even for $p = 2, d = 3$. To salvage the conjecture, we will need to revise our notion of polynomials. These are the so-called nonclassical polynomials, which will be introduced in Chapter 6.

5.2 The counter-example

We give a counter example to Conjecture 5.4 for $p = 2, d = 3$. It combines the bounds obtained in Lovett *et al.* [60] and Green and Tao [43]. The example is the degree 4 symmetric polynomial S_4 . In this section, all of the functions are defined on \mathbb{F}_2^n .

Definition 5.5 (Symmetric polynomials). Let $k \geq 1$ be an integer. The elementary symmetric polynomial of degree k over n variables is denoted by S_k and is defined as

$$S_k(x_1, \dots, x_n) = \sum_{S \subseteq [n], |S|=k} \prod_{i \in S} x_i.$$

Theorem 5.6 (Lovett *et al.* [60], Green and Tao [43]). Let $n \geq 1$ be sufficiently large. Then

$$\|(-1)^{S_4}\|_{U^4}^{16} = \frac{1}{8} + O(2^{-n/2}),$$

but for any polynomial $Q : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ of degree ≤ 3 it holds that

$$\left| \Pr_x[S_4(x) = Q(x)] - \frac{1}{2} \right| \leq \exp(-cn),$$

for some absolute constant $c > 0$.

5.2.1 U^4 -norm of S_4

Our goal is to prove

$$\|(-1)^{S_4}\|_{U^4}^{16} = \frac{1}{8} + O(2^{-n/2}).$$

Define the symmetric bilinear form $B : \mathbb{F}_2^n \times \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ by

$$B(a, b) = \sum_{i, j \in [n]: i \neq j} a_i b_j,$$

for $a = (a_1, \dots, a_n), b = (b_1, \dots, b_n)$ in \mathbb{F}_2^n . It is easy to verify the following identity regarding the fourth additive derivatives of S_4 .

$$D_a D_b D_c D_d S_4(x) = B(a, b)B(c, d) + B(a, c)B(b, d) + B(a, d)B(b, c).$$

Consequently

$$\|(-1)^{S_4}\|_{U^4}^{16} = \mathbb{E}_{a, b, c, d \in \mathbb{F}_2^n} (-1)^{B(a, b)B(c, d) + B(a, c)B(b, d) + B(a, d)B(b, c)}. \quad (5.1)$$

In order to understand the above quantity, we need to understand the distribution of

$$B_6(a, b, c, d) := (B(a, b), B(a, c), B(a, d), B(b, c), B(b, d), B(c, d))$$

over \mathbb{F}_2^6 , when $a, b, c, d \in \mathbb{F}_2^n$ are chosen uniformly and independently at random. The next lemma shows that for large n , they are essentially independently and uniformly chosen.

Lemma 5.7. For every $\tau \in \mathbb{F}_2^6$,

$$\Pr_{a, b, c, d \in \mathbb{F}_2^n} [B_6(a, b, c, d) = \tau] = \frac{1}{2^6} \pm O(2^{-n/2}).$$

5.2. *The counter-example*

Proof. By standard Fourier analysis, it is sufficient to show that for every collection of $\lambda_{ab}, \lambda_{ac}, \lambda_{ad}, \lambda_{bc}, \lambda_{bd}, \lambda_{cd} \in \mathbb{F}_2$, not all zero,

$$\mathbb{E}_{a,b,c,d \in \mathbb{F}_2^n} \left[(-1)^{\lambda_{ab}B(a,b) + \lambda_{ac}B(a,c) + \lambda_{ad}B(a,d) + \lambda_{bc}B(b,c) + \lambda_{bd}B(b,d) + \lambda_{cd}B(c,d)} \right] = O(2^{-n/2}).$$

We may assume $\lambda_{ab} = 1$ by symmetry. It suffices to show that for every $c, d \in \mathbb{F}_2^n$,

$$\mathbb{E}_{a,b \in \mathbb{F}_2^n} \left[(-1)^{B(a,b) + \lambda_{ac}B(a,c) + \lambda_{ad}B(a,d) + \lambda_{bc}B(b,c) + \lambda_{bd}B(b,d) + \lambda_{cd}B(c,d)} \right] = O(2^{-n/2}).$$

Since B is a bilinear form, $B(\cdot, c)$ and $B(\cdot, d)$ are linear forms and we can rewrite the above quantity as

$$\mathbb{E}_{a,b \in \mathbb{F}_2^n} \left[(-1)^{B(a,b) + L_1(a) + L_2(b)} \right],$$

where $L_1 := \lambda_{ac}B(\cdot, c) + \lambda_{ad}B(\cdot, d)$ and $L_2 := \lambda_{bc}B(\cdot, c) + \lambda_{bd}B(\cdot, d)$ are two linear forms. The term $L_2(b)$ can be removed by an application of the Cauchy-Schwarz inequality on a , and we obtain

$$\left| \mathbb{E}_{a,b \in \mathbb{F}_2^n} \left[(-1)^{B(a,b) + L_1(a) + L_2(b)} \right] \right|^2 \leq \mathbb{E}_{a,a',b \in \mathbb{F}_2^n} \left[(-1)^{B(a+a',b) + L_1(a+a')} \right].$$

Now observe that by the definition of B , $\mathbb{E}_b \left[(-1)^{B(a+a',b) + L_1(a+a')} \right] = 0$ whenever $a \neq a'$. Giving

$$\mathbb{E}_{a,a',b \in \mathbb{F}_2^n} \left[(-1)^{B(a+a',b) + L_1(a+a')} \right] = \Pr_{a,a' \in \mathbb{F}_2^n} [a = a'] = 2^{-n},$$

as was desired. □

Lemma 5.7 implies that the joint distribution of $B_6(a, b, c, d)$ is $O(2^{-n/2})$ -close in statistical distance to uniform over \mathbb{F}_2^6 . Thus, for a uniformly random $\tau = (\tau_{ab}, \tau_{ac}, \tau_{ad}, \tau_{bc}, \tau_{bd}, \tau_{cd}) \in \mathbb{F}_2^6$, we have

$$\begin{aligned} \|(-1)^{S_4}\|_{U^4}^{16} &= \mathbb{E}_{a,b,c,d \in \mathbb{F}_2^n} \left[(-1)^{B(a,b)B(c,d)+B(a,c)B(b,d)+B(a,d)B(b,c)} \right] \\ &= \mathbb{E}_{\tau \in \mathbb{F}_2^6} \left[(-1)^{\tau_{ab}\tau_{cd}+\tau_{ac}\tau_{bd}+\tau_{ad}\tau_{bc}} + O(2^{-n/2}) \right] \\ &= \left(\mathbb{E}_{u,v \in \mathbb{F}_2} [(-1)^{uv}] \right)^3 + O(2^{-n/2}) = \frac{1}{8} + O(2^{-n/2}). \end{aligned}$$

5.2.2 Bounds on correlation with cubic polynomials

The exponential bound on the correlation with cubic polynomials was obtained by Lovett *et al.* [60], the proof of which is involved. We instead present the proof of a weaker bound due to Green and Tao [43] which is still sufficient to refute **Conjecture 5.4**. Green and Tao [43] obtain the following bound using a modification of a clever Ramsey-theoretic argument by Alon and Beigel [1].

Theorem 5.8 (Green and Tao [43]). Let $n \geq 1$ be large enough. Then for any polynomial $Q : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ of degree at most 3,

$$\left| \Pr_x[S_4(x) = Q(x)] - \frac{1}{2} \right| \leq \exp(-c \log \log \log n),$$

for some absolute constant $c > 0$.

We first show that S_4 has small correlation with symmetric cubic polynomials.

Lemma 5.9. For any $c_0, c_1, c_2, c_3 \in \mathbb{F}_2$,

$$\mathbb{E}_{x \in \mathbb{F}_2^n} \left[(-1)^{S_4(x)+c_3S_3(x)+c_2S_2(x)+c_1S_1(x)+c_0} \right] \leq \exp(-cn)$$

for some absolute constant $c > 0$.

Proof. Let $|x|$ denote the hamming weight of x , that is the number of nonzero coordinates in x . Let $|x| = \sum_i b_i(x)2^i$ be its binary representation with $b_i(x) \in \{0, 1\}$. By Lucas' theorem on binomial coefficients [61],

$S_1(x) = b_0(x)$, $S_2(x) = b_1(x)$, $S_3(x) = b_0(x)b_1(x)$, and $S_4(x) = b_2(x)$. Thus,

$$\begin{aligned} \mathbb{E}_{x \in \mathbb{F}_2^n} \left[(-1)^{S_4(x) + c_3 S_3(x) + c_2 S_2(x) + c_1 S_1(x) + c_0} \right] \\ = \mathbb{E}_{x \in \mathbb{F}_2^n} \left[(-1)^{b_2(x) + c_3 b_0 b_1(x) + c_2 b_1(x) + c_1 b_0(x) + c_0} \right]. \end{aligned}$$

Note that if $b_0(x), b_1(x), b_2(x)$ were uniformly distributed in $\{0, 1\}^3$, then the average would be zero. To conclude, we need to show that they are close to uniformly distributed. Equivalently, we will show that $|x| \bmod 8 = \sum_{i=0}^2 b_i(x) 2^i$ is close to uniformly distributed in \mathbb{Z}_8 . For any $a \in \mathbb{Z}_8$ we have

$$\begin{aligned} \Pr_{x \in \mathbb{F}_2^n} [|x| \bmod 8 = a] &= 2^{-n} \sum_{k=0}^n \binom{n}{k} 1_{k \bmod 8 = a} \\ &= 2^{-n} \sum_{k=0}^n \binom{n}{k} \frac{\sum_{r=0}^7 e^{2\pi i r(k-a)/8}}{8} \\ &= \frac{1}{8} \sum_{r=0}^7 e^{-2\pi i r a/8} \left(\frac{1 + e^{-2\pi i r/8}}{2} \right)^n. \end{aligned}$$

The term corresponding for $r = 0$ is equal to $1/8$. For any $r \neq 0$, we can bound its corresponding term contribution by $|(1 + e^{-2\pi i r/8})/2|^n \leq \exp(-cn)$, for some $c > 0$. Thus,

$$\left| \Pr_{x \in \mathbb{F}_2^n} [|x| \bmod 8 = a] - \frac{1}{8} \right| \leq \exp(-cn).$$

□

The proof of [Theorem 5.8](#) follows from a Ramsey-type argument, which allows to reduce any polynomial to a symmetric polynomial with fewer variables.

Lemma 5.10. Let $Q : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ be a polynomial of degree ≤ 3 . Then there exists a symmetric polynomial $Q' : \mathbb{F}_2^{n'} \rightarrow \mathbb{F}_2$ of degree ≤ 3 , with $n' = \Omega(\log \log n)$, such that

$$\Pr_{x \in \mathbb{F}_2^{n'}} [Q'(x) = S_4(x)] \geq \Pr_{x \in \mathbb{F}_2^n} [Q(x) = S_4(x)].$$

Proof. Let Q be a degree 3 polynomial. Decompose

$$Q(x) = \sum q_{i,j,k} x_i x_j x_k + Q_2(x),$$

where Q_2 is a quadratic polynomial. Let H be a 3-uniform hypergraph with $V(H) = [n]$ and $E(H) = \{(i, j, k) : q_{i,j,k} = 1\}$. By the hypergraph Ramsey theorem [25, 24] there exists a set $A \subset [n]$ of size $|A| \geq \Omega(\log \log n)$ such that A is either a clique or an independent set for H . For $x \in \mathbb{F}_2^A$, $a \in \mathbb{F}_2^{[n] \setminus A}$ let $S_4(x, a)$ and $Q(x, a)$ be the polynomials with the appropriate inputs. By an averaging argument, there exists an assignment $a \in \mathbb{F}_2^{[n] \setminus A}$ such that

$$\Pr_{x \in \mathbb{F}_2^A} [S_4(x, a) = Q(x, a)] \geq \Pr_{x \in \mathbb{F}_2^n} [S_4(x) = Q(x)].$$

Note that $S_4(x, a)$ is a symmetric polynomial in x of degree 4, whose homogenous degree 4 part is $S_4(x)$; and $Q(x, a)$ is a polynomial of degree ≤ 3 , whose homogeneous degree 3 part is symmetric, equal either to 0 or to $S_3(x)$. Next, we make the quadratic part symmetric. Let

$$Q_2(x, a) = \sum_{i,j \in A} q'_{i,j} x_i x_j + Q_1(x),$$

where $Q_1(x)$ is a linear polynomial. Applying a similar argument, let G be a graph with $V(G) = A$ and $E(G) = \{(i, j) \in A : q'_{i,j} = 1\}$. By Ramsey's theorem for graphs [26], there exists a subset $B \subset A$ of size $|B| \geq \Omega(\log |A|)$ which is either a clique or an independent set. Thus, there exists an assignment $b \in \mathbb{F}_2^{A \setminus B}$ for which

$$\begin{aligned} \Pr_{x \in \mathbb{F}_2^B} [S_4(x, a, b) = Q(x, a, b)] &\geq \Pr_{x \in \mathbb{F}_2^A} [S_4(x, a) = Q(x, a)] \\ &\geq \Pr_{x \in \mathbb{F}_2^n} [S_4(x) = Q(x)]. \end{aligned}$$

Note that $Q(x, a, b)$ has homogeneous parts of degrees 3 and 2 both being symmetric. To conclude, let

$$Q_1(x, a, b) = \sum_{i \in B} q''_i x_i + q'''.$$

Let $C \subset B$ be a set for which q''_i for $i \in B$ are all equal, where $|C| \geq |B|/2$. There exists an assignment $c \in \mathbb{F}_2^{B \setminus C}$ such that

$$\Pr_{x \in \mathbb{F}_2^C} [S_4(x, a, b, c) = Q(x, a, b)] \geq \Pr_{x \in \mathbb{F}_2^n} [S_4(x) = Q(x)].$$

The lemma follows as $S_4(x, a, b, c) + Q(x, a, b)$ is a symmetric polynomial of degree 4 in $|C| = \Omega(\log \log \log n)$ variables, whose homogeneous part of degree 4 is equal to $S_4(x)$. \square

Proof of Theorem 5.8. Let $Q(x)$ be a cubic polynomial which maximizes $|\Pr[Q(x) = S_4(x)] - 1/2|$. By possibly replacing Q with $Q + 1$, we may assume that $\Pr[Q(x) = S_4(x)] \geq 1/2$. By Lemma 5.10, there exists a symmetric cubic polynomial Q' on $n' = \Omega(\log \log \log n)$ variables such that $\Pr[Q(x) = S_4(x)] \leq \Pr[Q'(x) = S_4(x)]$. By Lemma 5.9, $\Pr[Q'(x) = S_4(x)] \leq 1/2 + \exp(-cn')$. \square

Part II

Higher Order Fourier Analysis

In [Part II](#) of the survey we will introduce the foundations of higher-order Fourier analysis in detail. In classical Fourier analysis characters are the exponentials of linear functions. Since a typical random function $f : \mathbb{F}_p^n \rightarrow [-1, 1]$ has very small correlation with all the Fourier characters, it is natural to consider the characters as highly structured functions. This leads to a “structure” versus “pseudorandomness” dichotomy in which functions that have small correlation with all the Fourier characters are regarded as pseudorandom functions, and the ones that have noticeable correlation with a Fourier character are considered to be somewhat structured.

Consequently, the large Fourier coefficients of $f : \mathbb{F}_p^n \rightarrow [-1, 1]$ constitute the structured part of f , while the small Fourier coefficients correspond to its pseudorandom part. This can be formulated as an “approximate structure theorem” that says that, we can decompose any function f as $f = f_1 + f_2$, where f_1 is structured as it is a linear combination of few Fourier characters, and f_2 is pseudorandom in that all of the Fourier coefficients of f_2 are small. Such approximate structure theorems are very useful as for many problems such as estimating the probability of the success of linearity test, or estimating the density of 3-term arithmetic progressions in a set. The reason is that it is possible to treat f_2 as a small random noise, and extract the desired information about f from the highly structured part, namely f_1 .

In order to be able to discard f_2 safely, we need to deduce from the fact that f_2 does not have a noticeable correlation with any Fourier character that it does not contribute significantly to the estimated parameter. In other words, we need an inverse theorem which would state that significant contribution implies significant correlation with a character. Although, at first glance, it might appear that one needs to establish a separate inverse theorem for every problem that one wishes to study via this theory, fortunately, this is not the case. It suffices to prove an inverse theorem for the Gowers U^2 norm. It turns out that for many problems, a few applications of the classical Cauchy-Schwarz inequality show that f_2 can be discarded if $\|f_2\|_{U^2}$ is small.

The structure versus pseudorandomness dichotomy discussed above is not sufficiently strong to be applicable to more complex linear structures. In other words discarding f_2 , even when $\|f_2\|_{U^2}$ is very small, can

have a significant effect on certain parameters such as the probability of success of AKKLR(d) test for $d > 1$, or the density of four-term arithmetic progressions in a set. To resolve this issue, Gowers [37] defined higher uniformity norms U^d , and found a new proof for Szemerédi's theorem by partially extending the above program for these norms. Indeed, as we shall see in Chapter 11, for every linear pattern, there exists a d such that if $\|f_2\|_{U^d}$ is sufficiently small, then for the purpose of estimating the density of that pattern, the pseudorandom f_2 can be discarded without having a significant effect on the estimate. However, this fact by itself is not very useful without an inverse theorem for U^d norm as one needs an inverse theorem to extract a structured part from every function with noticeable U^d norm.

In Theorem 5.3 we established an inverse theorem for the U^3 norm. This theorem lays the foundation of quadratic Fourier analysis as it implies that every function $f : \mathbb{F}_p^n \rightarrow [-1, 1]$ can be decomposed into a structured part that is a linear combination of the exponentials of a few quadratic characters and a pseudorandom part that has small U^3 norm. Unfortunately, as we saw in Theorem 5.6 its most natural generalization Conjecture 5.4 to higher uniformity norms is false. In other words, classical polynomials are not the right structure for higher uniformity norms. Tao and Ziegler [83] proved that Conjecture 5.4 can be fixed by replacing classical polynomials with a generalization of classical polynomials, which we will refer to as non-classical polynomials.

We introduce non-classical polynomials and state Tao and Ziegler's inverse theorem in Chapter 6. We then proceed to develop the theory of higher order Fourier analysis in the remainder of this part. Concretely, in Chapter 7 we introduce the notions of rank, regularity and uniformity for polynomials, which are the basis for the structure vs pseudorandomness dichotomy discussed above. In Chapter 9 we develop the decomposition theorems. In Chapter 10 we develop a notion of homogeneous non-classical polynomials, which is useful in certain applications. In Chapter 11 we apply these to the study of linear patterns in sets. A few technical proofs from these chapters are deferred to Chapter 12.

6

Nonclassical Polynomials, and the Inverse Gowers Theorem

In [Theorem 5.6](#) we saw that the most natural generalization of the U^2 and U^3 inverse theorem to higher uniformity norms is false. That is for $d \geq 4$, there exist bounded functions that have large U^d norm but do not have significant correlation with the exponential of any polynomial of degree $d - 1$. In this chapter, we address this issue by introducing an extension of classical polynomials, called *nonclassical polynomials*. We will show that an inverse theorem for Gowers uniformity norms of any order holds for nonclassical polynomials. Below we will give some intuition on how we can arrive to the definition of nonclassical polynomials.

Let $d \geq 0$ be an integer. Similar to the situation over finite fields, there are both a global and a local definition for a low degree real-valued polynomial. A real-valued function $P : \mathbb{R}^n \rightarrow \mathbb{R}$ being a polynomial of (total) degree $\leq d$ can be defined in two equivalent ways:

- (1) **Global definition.** P is a polynomial of degree $\leq d$ if it can be written as

$$P(x_1, \dots, x_n) = \sum_{\substack{i_1, \dots, i_n \geq 0 \\ i_1 + \dots + i_n \leq d}} c_{i_1, \dots, i_n} x_1^{i_1} \cdots x_n^{i_n},$$

with coefficients $c_{i_1, \dots, i_n} \in \mathbb{R}$.

- (2) **Local definition.** P is a polynomial of degree $\leq d$ if it is $d + 1$ times differentiable and its $(d + 1)$ -th derivative vanishes everywhere.

It is easy to see by linearity of the derivative operator that (i) implies (ii). For the other direction, one can use the Taylor series expansion to go from the local to the global condition, and show that the two definitions above are equivalent. Both these definitions can be extended to the finite characteristic setting, i.e. when $P : \mathbb{F}^n \rightarrow G$ for a finite field \mathbb{F} and an Abelian group G . The global definition extends in a straightforward manner, and the local definition uses the notion of additive directional derivatives.

Definition 6.1 (Polynomials over finite fields (local definition)). For an integer $d \geq 0$, a function $P : \mathbb{F}^n \rightarrow G$ is said to be a *polynomial of degree $\leq d$* if for all $y_1, \dots, y_{d+1}, x \in \mathbb{F}^n$, it holds that

$$(D_{y_1} \cdots D_{y_{d+1}} P)(x) = 0,$$

where $D_y P(x) = P(x + y) - P(x)$ is the additive derivative of P with direction y evaluated at x . The *degree* of P is the smallest d for which the above holds.

It follows simply from the definition that for any direction $y \in \mathbb{F}^n$, $\deg(D_y P) < \deg(P)$. In the “classical” case of polynomials $P : \mathbb{F}^n \rightarrow \mathbb{F}$, it is a well-known fact that the global and local definitions coincide. However, the situation is different in more general groups. One can already suspect this from the fact that division by $d!$ may not be possible in G , and hence one cannot make use of the Taylor expansion to go from the local definition to the global one.

When the range of P is the torus \mathbb{R}/\mathbb{Z} , it turns out that the global definition must be refined to the “nonclassical polynomials”, which may have monomials that are different from the classical case. This phenomenon was first noted by Tao and Ziegler [83] in the study of Gowers norms, where it was also proved that [Conjecture 5.4](#) can be modified to hold by replacing classical polynomials with nonclassical polynomials.

6.1 Nonclassical polynomials

Let $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ denote the torus (namely, the group of addition modulo 1). Let $e : \mathbb{T} \rightarrow \mathbb{C}$ be given by $e(x) = \exp(2\pi ix)$. Fix a prime finite field $\mathbb{F} = \mathbb{F}_p$ for the remainder of this chapter. Nonclassical polynomials arise when studying functions $P : \mathbb{F}^n \rightarrow \mathbb{T}$ and their phase functions $f = e(P) : \mathbb{F}^n \rightarrow \mathbb{C}$.

Definition 6.2 (Nonclassical polynomials). For an integer $d \geq 0$, a function $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is said to be a *nonclassical polynomial of degree $\leq d$* (or simply a *polynomial of degree $\leq d$*) if for all $y_1, \dots, y_{d+1}, x \in \mathbb{F}^n$, it holds that

$$(D_{y_1} \cdots D_{y_{d+1}} P)(x) = 0. \quad (6.1)$$

The *degree* of P is the smallest d for which the above holds. A function $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is said to be a *classical polynomial of degree $\leq d$* if it is a nonclassical polynomial of degree $\leq d$ whose image is contained in $\frac{1}{p}\mathbb{Z}/\mathbb{Z}$.

Denote by $\text{Poly}(\mathbb{F}^n \rightarrow \mathbb{T})$, $\text{Poly}_d(\mathbb{F}^n \rightarrow \mathbb{T})$ and $\text{Poly}_{\leq d}(\mathbb{F}^n \rightarrow \mathbb{T})$, the set of all nonclassical polynomials over \mathbb{F}^n , all nonclassical polynomials of degree d and all nonclassical polynomials of degree $\leq d$, respectively.

The following lemma of Tao and Ziegler [83] shows that a *classical polynomial* P of degree d must always be of the form $x \mapsto \frac{|Q(x)|}{p}$, where $Q : \mathbb{F}^n \rightarrow \mathbb{F}$ is a polynomial (in the usual sense) of degree d , and $|\cdot|$ is the canonical map from $\mathbb{F} = \mathbb{F}_p$ to $\{0, 1, \dots, p-1\}$. Moreover, it provides a global characterization of the structure of nonclassical polynomials.

Lemma 6.3 (Tao and Ziegler [83], Lemma 1.7). A function $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is a polynomial of degree $\leq d$ if and only if P can be represented as

$$P(x_1, \dots, x_n) = \alpha + \sum_{\substack{0 \leq d_1, \dots, d_n < p; k \geq 0: \\ 0 < \sum_{i=1}^n d_i \leq d - k(p-1)}} \frac{c_{d_1, \dots, d_n, k} |x_1|^{d_1} \cdots |x_n|^{d_n}}{p^{k+1}} \pmod{1},$$

for a unique choice of $c_{d_1, \dots, d_n, k} \in \{0, 1, \dots, p-1\}$ and $\alpha \in \mathbb{T}$. The element α is called the *shift* of P , and the largest integer k such that there exist d_1, \dots, d_n for which $c_{d_1, \dots, d_n, k} \neq 0$ is called the *depth* of P . A depth- k polynomial P takes values in a coset of the subgroup $\mathbb{U}_{k+1} := \frac{1}{p^{k+1}}\mathbb{Z}/\mathbb{Z}$. Classical polynomials correspond to polynomials with 0 shift and 0 depth.

In many cases, for the sake of brevity, we will omit writing “mod 1” in the description of the defined nonclassical polynomials. We do not include the proof of Lemma 6.3 in this text. For a proof we refer the reader to either the paper of Tao and Ziegler [83] or to a blog post of Tao [79] describing the proof.

Example 6.4. Consider the univariate function $f : \mathbb{F}_2 \rightarrow \mathbb{T}$ given by $f(x) = \frac{|x|}{4} \pmod{1}$. It is a nonclassical polynomial of degree 2. To see this, we compute its derivatives $D_y f$ for $y \in \mathbb{F}_2$. Clearly $D_0 f = 0$ for any univariate function, so it suffices to compute $D_1 f$. One can verify that

$$\begin{aligned} D_1 f(x) &= f(x \oplus 1) - f(x) = \begin{cases} 1/4 & x = 0 \\ -1/4 & x = 1 \end{cases} \\ &= \frac{1}{4} + \frac{|x|}{2} \pmod{1}. \end{aligned}$$

Moreover,

$$D_1 D_1 f(x) = \frac{1}{2}, \quad D_1 D_1 D_1 f(x) = 0.$$

Thus, taking 3 derivatives annihilates f , but not 2 derivatives. So by definition f is a degree 2 polynomial. Similarly, one can show that $\frac{|x|}{2^k}$ is a nonclassical polynomial of degree k .

Remark 6.5. An equivalent definition of nonclassical polynomials is via functions which map \mathbb{F}_p^n to the ring \mathbb{Z}_p^{k+1} . Concretely, if $f : \mathbb{F}_p^n \rightarrow \mathbb{T}$ is

a nonclassical polynomial of degree d and depth k , then it takes values in $\frac{1}{p^{k+1}}\mathbb{Z}/\mathbb{Z}$. Thus, we can write f as

$$f(x) = \frac{F(x)}{p^{k+1}} \pmod{1},$$

where $F : \mathbb{F}_p^n \rightarrow \mathbb{Z}_{p^{k+1}}$. One can verify that F is a polynomial. However, note that the classical definition of degree for F (based on monomials) does not coincide with our definition of degree for f (based on derivatives). For example, in the univariate case $F(x) = x$ corresponds to $f(x) = \frac{x}{p^{k+1}}$ which has degree $1 + (p-1)k$, while $F(x) = px$ corresponds to $f(x) = \frac{x}{p^k}$ which has degree $1 + (p-1)(k-1)$.

Note that [Lemma 6.3](#) immediately implies the following important corollary. Below we use the following standard shorthand, which follows as \mathbb{T} is an Abelian group. For $n \in \mathbb{Z}$ and $x \in \mathbb{T}$, nx stands for $x + \dots + x$ if $n \geq 0$ and $-x - \dots - x$ otherwise, where there are $|n|$ terms in both expressions.

Corollary 6.6. Let $Q : \mathbb{F}_p^n \rightarrow \mathbb{T}$ be a polynomial of degree d and depth k . Then

- (1) If $\lambda \in \mathbb{Z}$ is co-prime to p then λQ also has degree d and depth k .
- (2) pQ has degree $\max(d - p + 1, 0)$ and depth $k - 1$. In other words, if Q is classical, then pQ vanishes, and otherwise, its degree decreases by $p - 1$ and its depth by 1.
- (3) $p^{k+1}Q = 0$. This implies that if $\lambda \equiv \lambda' \pmod{p^{k+1}}$ then $\lambda Q = \lambda' Q$.

For convenience of exposition, henceforth we will assume that the shifts of all polynomials are zero. This can be done without affecting any of the results presented in this text. Under this assumption, all polynomials of depth k take values in \mathbb{U}_{k+1} .

6.2 The inverse theorem for Gowers norms

Gowers norms, which were introduced by Gowers [37], play an important role in additive combinatorics, more specifically in the study of

polynomials of bounded degree. We have seen in the direct theorem for Gowers norms (Theorem 4.2; see also Theorem 5.3, direct theorem) that correlation with polynomials implies large Gowers norm. The proof generalizes directly to include nonclassical polynomials.

Theorem 6.7 (Direct theorem for Gowers norm). Let $f : \mathbb{F}^n \rightarrow \mathbb{C}$ be a function and $d \geq 1$ an integer. Then for every degree- d nonclassical polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$,

$$|\langle f, e(P) \rangle| \leq \|f\|_{U^{d+1}}.$$

The theorem follows from the monotonicity of the Gowers norms.

Claim 6.8. Let $f : \mathbb{F}^n \rightarrow \mathbb{C}$ and $d \geq 1$. Then $\|f\|_{U^{d+1}} \geq \|f\|_{U^d}$.

Proof. Recall that for a function $g : \mathbb{F}^n \rightarrow \mathbb{C}$ and $y \in \mathbb{F}^n$, $\Delta_y g(x) = g(x + y)\overline{g(x)}$ denotes the multiplicative derivative of g at direction y . We have

$$\begin{aligned} \|f\|_{U^{d+1}}^{2^{d+1}} &= \mathbb{E}_{x, y_1, \dots, y_{d+1}} [\Delta_{y_1, \dots, y_{d+1}} f(x)] \\ &= \mathbb{E}_{y_1, \dots, y_d} \left[\mathbb{E}_{x, y_{d+1}} [\Delta_{y_1, \dots, y_d} f(x + y_{d+1}) \cdot \overline{\Delta_{y_1, \dots, y_d} f(x)}] \right] \\ &= \mathbb{E}_{y_1, \dots, y_d} \left[\left| \mathbb{E}_x [\Delta_{y_1, \dots, y_d} f(x)] \right|^2 \right] \\ &\geq \left| \mathbb{E}_{x, y_1, \dots, y_d} [\Delta_{y_1, \dots, y_d} f(x)] \right|^2 \\ &= \|f\|_{U^d}^{2^{d+1}}. \end{aligned}$$

□

Proof of Theorem 6.7. Let $g(x) = f(x)e(P(x))$. Then

$$|\langle f, e(P) \rangle| = \|g\|_{U^1} \leq \|g\|_{U^{d+1}}.$$

Now, for any $y_1, \dots, y_{d+1} \in \mathbb{F}^n$ it holds that

$$\Delta_{y_1, \dots, y_{d+1}} g(x) = \Delta_{y_1, \dots, y_{d+1}} f(x) \cdot e(D_{y_1, \dots, y_{d+1}} P(x)) = \Delta_{y_1, \dots, y_{d+1}} f(x),$$

and hence,

$$\begin{aligned} \|g\|_{U^{d+1}}^{2^{d+1}} &= \mathbb{E}_{y_1, \dots, y_{d+1}} [\Delta_{y_1, \dots, y_{d+1}} g(x)] \\ &= \mathbb{E}_{y_1, \dots, y_{d+1}} [\Delta_{y_1, \dots, y_{d+1}} f(x)] = \|f\|_{U^{d+1}}^{2^{d+1}}. \end{aligned}$$

□

As we showed in [Section 5.2](#), the inverse direction is false, when we restrict ourselves to classical polynomials. However, Tao and Ziegler [\[83\]](#) proved that it is true if we include nonclassical polynomials. The inverse theorem only applies to bounded functions. Let $\mathbb{D} := \{z \in \mathbb{C} : |z| \leq 1\}$ be the unit disk in the complex plane.

Theorem 6.9 (Inverse theorem for Gowers norm ([\[83\]](#), Theorem 1.11)). Fix $d \geq 1$ an integer and $\varepsilon > 0$. There exists an $\delta = \delta_{6.9}(\mathbb{F}, d, \varepsilon)$ such that the following holds. For every function $f : \mathbb{F}^n \rightarrow \mathbb{D}$ with $\|f\|_{U^{d+1}} \geq \varepsilon$, there exists a polynomial $P \in \text{Poly}_{\leq d}(\mathbb{F}^n \rightarrow \mathbb{T})$ that is δ -correlated with f , that is

$$|\langle f, e(P) \rangle| \geq \delta.$$

A comment about notation. We use the following notation throughout the survey: unspecified constants (such as $\delta_{6.9}(\mathbb{F}, d, \varepsilon)$ in [Section 6.2](#)) are labeled by the number of the lemma / theorem where they are referenced.

It is easy to see that for every degree d nonclassical polynomial P , $\|e(P)\|_{U^{d+1}} = 1$. [Theorem 6.7](#) and [Theorem 6.9](#) provide a robust version of this statement, showing that the Gowers norm of a function f is large if and only if it contains some low degree structure, namely if f correlates with a low degree polynomial. We also note that [Theorem 6.9](#) only shows existence of a constant $\delta > 0$ for every ε and finding reasonable quantitative bounds or limitations to such bounds is a fascinating problem which to this day remains unsolved. In the case of quadratics $d = 1$, we have $\|f\|_{U^2} = \|\hat{f}\|_4$ which gives $\delta = \Omega(\varepsilon^2)$. The case of $d = 2$ is already nontrivial and the best known lower-bound for δ is quasi-polynomial in ε , which follows from a result of Sanders on the Bogolyubov-Ruzsa conjecture (Sanders [\[71\]](#), see also Lovett [\[58, 59\]](#)).

7

Rank, Regularity, and Other Notions of Uniformity

Consider a function $f : \mathbb{F}_p^n \rightarrow [-1, 1]$ and a positive integer d . As we shall see in [Chapter 9](#) the inverse theorems ([Theorem 5.3](#) and [Theorem 6.9](#)) allow us to approximate f in the U^{d+1} norm by a linear combination of a few higher-order *phase functions*, which are exponentials of nonclassical polynomials of degree d . We will think of this as an order- d Fourier expansion of f , and regard it as the structured part of f . Unlike classical Fourier expansion, higher-order Fourier expansions are not unique by any means. This is simply because there are too many polynomials of degree d , (there are asymptotically $2^{O(n^d)}$ classical degree $\leq d$ polynomials), and thus they cannot form a linear basis for the space of functions whose dimension is p^n . However, this might be disappointing as one of the most important and useful properties of the classical Fourier characters is that they form an orthonormal basis. While it is not possible to achieve this orthogonality in an exact way, we can still hope to obtain an approximate version of it by approximating f with a linear combination of a few higher-order phase functions which are pairwise almost orthogonal. We will see how to accomplish this through the notions of *rank* and *regularity*, which are the topic of this chapter. We will present these topics for the general case of an arbitrary fixed value

of d , and unfortunately this leads to many technicalities that might be overwhelming for the readers who see these materials for the first time. Some of these technicalities are unnecessary for the case of the quadratic Fourier analysis, i.e. $d = 2$, and many definitions and statements are more intuitive and familiar in this case, and for example the definition of rank coincides with the familiar notion of rank from linear algebra. Thus we recommend those readers who find this chapter too technical and unintuitive to first consult the excellent lecture notes of Ben Green on quadratic Fourier analysis [41].

A property of Fourier characters is that they behave like a random function, for example their value is uniformly distributed when the input is drawn uniformly at random. Having this in mind, we can demand several such random-like behaviors from higher-degree polynomials. We refer to such properties as *regularity*. One property of random functions is that they cannot be expressed as a function of few low-degree polynomials. We will capture this property by defining the notion of *rank* of a polynomial, intended to capture its complexity according to lower degree polynomials.

Definition 7.1 (Rank of a polynomial). Given a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ and an integer $d \geq 1$, the d -rank of P , denoted $\text{rank}_d(P)$, is defined to be the smallest integer r such that there exist polynomials $Q_1, \dots, Q_r : \mathbb{F}^n \rightarrow \mathbb{T}$ of degree $\leq d - 1$ and a function $\Gamma : \mathbb{T}^r \rightarrow \mathbb{T}$ satisfying $P(x) = \Gamma(Q_1(x), \dots, Q_r(x))$. If $d = 1$, then 1-rank is defined to be ∞ if P is non-constant, and 0 otherwise.

The *rank* of a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is its $\deg(P)$ -rank. We say that P is r -regular if $\text{rank}(P) \geq r$.

Note that for an integer $1 \leq \lambda \leq p - 1$, $\text{rank}(P) = \text{rank}(\lambda P)$. A high-rank polynomial of degree d is, intuitively, a “generic” degree- d polynomial; there are no unexpected ways to decompose it into lower degree polynomials. For future use, we record here a simple lemma stating that restrictions of high rank polynomials to hyperplanes generally preserve degree and high rank.

Lemma 7.2. Suppose $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is a polynomial of degree d and rank $\geq r$, where $r > p + 1$. Let A be a hyperplane in \mathbb{F}^n , and denote by P' the restriction of P to A . Then, P' is a polynomial of degree d and rank $\geq r - p$, unless $d = 1$ and P is constant on A .

Proof. For the case $d = 1$, we can check directly that either P' is constant or else, P' is a non-constant degree-1 polynomial and so has rank infinity.

So, assume $d > 1$. By applying an affine transformation, we can assume without loss of generality that A is the hyperplane $\{x : x_1 = 0\}$. Let $\pi : \mathbb{F}^n \rightarrow \mathbb{F}^{n-1}$ be the projection to A given by $\pi(x_1, x_2, \dots, x_n) = (0, x_2, \dots, x_n)$. Let $P'' = P - P' \circ \pi$. Clearly, P'' is zero on A . For $a \in \mathbb{F} \setminus \{0\}$, let $h_a = (a, 0, \dots, 0) \in \mathbb{F}^n$. Note that $D_{h_a}P''$ is of degree $\leq d - 1$ and that $(D_{h_a}P'')(y) = P''(y + h_a)$ for all $y \in A$. Hence, for every $a \in \mathbb{F} \setminus \{0\}$, P'' on $A + h_a$ agrees with a polynomial Q_a of degree $\leq d - 1$. So, for a function $\Gamma : \mathbb{T}^{p+1} \rightarrow \mathbb{T}$, we can write $P(x) = \Gamma(|x_1|/p, P'(x), Q_1(x), Q_2(x), \dots, Q_{p-1}(x))$, where $|x_1|/p, Q_1, \dots, Q_{p-1}$ are of degree $\leq d - 1$.

Now, if P' itself is of degree $d - 1$, then P is of rank $\leq p + 1 < r$, a contradiction. If P' is of rank $< r - p$, then again P is of rank $< r - p + p = r$, a contradiction. □

7.1 Polynomial factors

Next, we will formalize the notion of a generic collection of polynomials. Intuitively, it should mean that there are no unexpected algebraic dependencies among the polynomials. First, we need to set up some notation.

Definition 7.3 (Factors). If X is a finite set, then by a *factor* \mathcal{B} we simply mean a partition of X into finitely many parts called *atoms*.

A finite collection of functions ϕ_1, \dots, ϕ_C from X to some other space Y naturally define a factor $\mathcal{B} = \mathcal{B}_{\phi_1, \dots, \phi_C}$ whose atoms are sets of the form $\{x \in X : (\phi_1(x), \dots, \phi_C(x)) = (y_1, \dots, y_C)\}$ for some $(y_1, \dots, y_C) \in Y^C$. By an abuse of notation we also use \mathcal{B} to denote the map $x \mapsto (\phi_1(x), \dots, \phi_C(x))$, thus also identifying the atom containing x with $(\phi_1(x), \dots, \phi_C(x))$.

Definition 7.4 (Polynomial factors). If $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$ is a sequence of polynomials, then the factor $\mathcal{B}_{P_1, \dots, P_C}$ is called a *polynomial factor*.

The *complexity* of \mathcal{B} , denoted $|\mathcal{B}| := C$, is the number of defining polynomials. The *degree* of \mathcal{B} is the maximum degree among its defining polynomials P_1, \dots, P_C . If P_1, \dots, P_C are of depths k_1, \dots, k_C , respectively, then the number of atoms of \mathcal{B} is at most $\|\mathcal{B}\| := \prod_{i=1}^C p^{k_i+1}$.

Note that it makes a difference whether we define atoms according to the ordered set of evaluations $(P_1(x), \dots, P_C(x))$ rather than the multiset $\{P_1(x), \dots, P_C(x)\}$, and we choose the former in our definition.

We say that a function $f : \mathbb{F}^n \rightarrow \mathbb{T}$ is *measurable* in a polynomial factor $\mathcal{B}_{P_1, \dots, P_C}$ (or *\mathcal{B} -measurable* in short) if there exists a $\Gamma : \mathbb{T}^C \rightarrow \mathbb{T}$ such that

$$f = \Gamma(P_1, \dots, P_C),$$

or in other words the value of $f(x)$ can be determined by knowing only the values of $P_1(x), \dots, P_C(x)$. Note that, here Γ is an arbitrary map with no restriction on its degree or complexity. Equivalently, we say that f is *\mathcal{B} -measurable* if f is constant over each atom of \mathcal{B} .

Example 7.5. The function $f(x) := \frac{x_1 x_2 + x_1 x_3}{2} + \frac{x_3}{4}$ is measurable in the factor defined by $P_1 = \frac{x_1}{2}$, $P_2 = \frac{x_2 + x_3}{2}$ and $P_3 = \frac{x_3}{4}$.

Next we define conditional expectation over a factor, which results in a function that is constant on each atom of the factor.

Definition 7.6 (Conditional expectation over factors). Let \mathcal{B} be a polynomial factor defined by $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$. For $f : \mathbb{F}^n \rightarrow \mathbb{C}$, the conditional expectation of f with respect to \mathcal{B} , denoted $\mathbb{E}[f|\mathcal{B}] : \mathbb{F}^n \rightarrow \mathbb{C}$, is

$$\mathbb{E}[f|\mathcal{B}](x) := \mathbb{E}_{y \in \mathbb{F}^n} [f(y) | P_1(y) = P_1(x), \dots, P_C(y) = P_C(x)].$$

Namely, $\mathbb{E}[f|\mathcal{B}]$ is constant on every atom of \mathcal{B} , and takes the average value of f over this atom.

Note that $\mathbb{E}[f|\mathcal{B}]$ is \mathcal{B} -measurable. The following is a simple observation stating that $E[f|\mathcal{B}]$ has the same correlation as f with any \mathcal{B} -measurable function.

Remark 7.7. Let $f : \mathbb{F}^n \rightarrow \mathbb{C}$. Let \mathcal{B} be a polynomial factor defined by polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$, and let $g : \mathbb{F}^n \rightarrow \mathbb{C}$ be a \mathcal{B} -measurable function. Then

$$\langle f, g \rangle = \langle \mathbb{E}(f|\mathcal{B}), g \rangle.$$

Finally, we define the rank of a polynomial factor. We require that every nonzero linear combination of the polynomials which define the factor has high rank. Recall that by [Corollary 6.6](#), if $P : \mathbb{F}_p^n \rightarrow \mathbb{T}$ is a polynomial of depth k then for $\lambda \in \mathbb{Z}$, the depth of λP depends only on $\lambda \pmod{p^{k+1}}$.

Definition 7.8 (Rank of a factor). Let \mathcal{B} be a polynomial factor defined by a sequence of polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$ with respective depths k_1, \dots, k_C . The rank of \mathcal{B} is the least integer r , for which there exists $(\lambda_1, \dots, \lambda_C) \in \mathbb{Z}^C$, with $(\lambda_1 \pmod{p^{k_1+1}}, \dots, \lambda_C \pmod{p^{k_C+1}}) \neq 0^C$, such that $\text{rank}_d(\sum_{i=1}^C \lambda_i P_i) \leq r$, where $d = \max_i \deg(\lambda_i P_i)$.

Given a polynomial factor \mathcal{B} and a function $r : \mathbb{N} \rightarrow \mathbb{N}$, we say that \mathcal{B} is r -regular if the rank of \mathcal{B} is larger than $r(|\mathcal{B}|)$.

Notice that by the definition of rank, for a degree- d polynomial P of depth k we have

$$\text{rank}(\{P\}) = \min \left\{ \text{rank}_d(P), \text{rank}_{d-(p-1)}(pP), \dots, \text{rank}_{d-k(p-1)}(p^k P) \right\},$$

where $\{P\}$ is a polynomial factor consisting of one polynomial P .

Regular factors indeed do behave like a generic collection of polynomials, and we will establish this in a precise sense in [Section 7.4](#). Thus, given any factor \mathcal{B} that is not regular, it is often useful to *regularize* \mathcal{B} , that is to find a refinement \mathcal{B}' of \mathcal{B} that is regular up to our desires. Various regularity lemmas for polynomials will be discussed in [Section 7.5](#)

7.2 Analytic measures of uniformity

Regularity defined by the notion of rank is an algebraic/combinatorial notion of pseudorandomness. There are several cases where an analytic notion would be much more useful. In many cases, what is needed from the notion of regularity is that the polynomials defining the

factor, when evaluated jointly on a uniform input, would behave as independent random variables. This can be equivalently expressed as the condition that any nonzero linear combination of the polynomials is almost uniformly distributed. We will accomplish this through the analytic notion of bias.

Definition 7.9 (Bias). The bias of a function $f : \mathbb{F}^n \rightarrow \mathbb{T}$ is defined to be

$$\text{bias}(f) := \mathbb{E}_{x \in \mathbb{F}^n} [e(f(x))].$$

Note that a function f that takes every value from \mathbb{U}_k equally likely, will satisfy $\text{bias}(f) = 0$, and that a random polynomial will have bias very close to 0, and thus having small bias can be potentially used as an analytic notion of regularity.

It turns out that the bias and the rank of a polynomial are closely related. The following theorem was first proven for the case of $d < |\mathbb{F}|$ by Green and Tao [43], and then extended to the general case by Kaufman and Lovett [54].

Theorem 7.10 (Green and Tao [43] and Kaufman and Lovett [54]). Fix a prime finite field \mathbb{F} , an integer $d \geq 1$ and a real $\varepsilon > 0$. There exists $r = r_{7.10}(\mathbb{F}, d, \varepsilon)$ such that the following is true. If $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is a degree- d polynomial with rank greater than r , then $|\mathbb{E}_x[e(P(x))]| < \varepsilon$.

Kaufman and Lovett originally proved [Theorem 7.10](#) for classical polynomials. However, their proof extends to nonclassical ones without modification. Note that $r_{7.10}(\mathbb{F}, d, \varepsilon)$ does not depend on the dimension n of \mathbb{F}^n , but depends in an unspecified way to $\mathbb{F}, d, \varepsilon$. The original proof had terrible dependency on all three parameters (Ackerman-type), and hence only applies for constant-size prime finite fields, constant degree and constant bias.

Remark 7.11. [Theorem 7.10](#) was extended in two ways by subsequent papers. Bhattacharyya *et al.* [12] extended it to constant-size non-prime finite fields. Bhowmick and Lovett [19] refined the dependency on the field size and bias to be polynomial. This allows the theorem to be applicable for large finite fields, of size possibly growing with n . We present this in [Chapter 8](#). In this section, we restrict our attention to constant-size prime finite fields.

Next, motivated by [Theorem 7.10](#) we define unbiasedness for polynomial factors.

Definition 7.12 (Unbiased factor). Let $\varepsilon : \mathbb{N} \rightarrow \mathbb{R}^+$ be a decreasing function. A polynomial factor \mathcal{B} defined by a sequence of polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$ with respective depths k_1, \dots, k_C is said to be ε -unbiased if for every collection $(\lambda_1, \dots, \lambda_C) \in \mathbb{Z}^C$, with $(\lambda_1 \bmod p^{k_1+1}, \dots, \lambda_C \bmod p^{k_C+1}) \neq 0^C$, it holds that

$$\left| \mathbb{E}_x \left[e \left(\sum_i \lambda_i P_i(x) \right) \right] \right| < \varepsilon(|\mathcal{B}|).$$

Using Gowers norms, one can define the following analytic notion of uniformity for polynomials which is stronger than unbiasedness.

Definition 7.13 (Uniformity). Let $\varepsilon > 0$ be a real. A degree- d polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is said to be ε -uniform if

$$\|e(P)\|_{U^d} < \varepsilon.$$

Tao and Ziegler [[83](#)] used [Theorem 7.10](#) to show that high rank polynomials have small Gowers norm.

Theorem 7.14 (Tao and Ziegler [[83](#)], Theorem 1.20). Fix a prime finite field \mathbb{F} , an integer $d \geq 1$ and $\varepsilon > 0$. There exists $r = r_{7.14}(\mathbb{F}, d, \varepsilon)$ such that the following is true. For every nonclassical polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ of degree $\leq d$, if $\|e(P)\|_{U^d} \geq \varepsilon$, then $\text{rank}_d(P) \leq r$.

This immediately implies that a high-rank polynomial is also uniform in the sense of [Definition 7.13](#).

Corollary 7.15. Let $\mathbb{F}, d, \varepsilon$ and $r(\mathbb{F}, d, \varepsilon)$ be as in [Theorem 7.14](#). Every r -regular polynomial P of degree d is also ε -uniform.

The next claim, which is a standard application of Fourier analysis, shows that the converse of this is true at least qualitatively.

Claim 7.16. Fix a prime finite field \mathbb{F} , and integers $d, r \geq 1$. There exists $\varepsilon = \varepsilon_{7.16}(\mathbb{F}, d, r)$ such that the following is true. For every $P : \mathbb{F}^n \rightarrow \mathbb{T}$, if $\text{rank}_d(P) \leq r$ then $\|e(P)\|_{U^d} \geq \varepsilon$.

Proof. We will show that low rank functions cannot be very uniform. Let $\mathbb{F} = \mathbb{F}_p$. Assume that $P(x) = \Gamma(Q_1(x), \dots, Q_r(x))$, where each $Q_i : \mathbb{F}_p^n \rightarrow \mathbb{T}$ is a polynomial of degree $\leq d-1$ and depth k_i , and where $\Gamma : \prod_{i=1}^r \mathbb{U}_{k_i+1} \rightarrow \mathbb{T}$. Shorthand $G := \prod_{i=1}^r \mathbb{U}_{k_i+1}$ and let $G' := \prod_{i=1}^r \mathbb{Z}_p^{k_i+1}$ be the dual group. The Fourier decomposition of $e(\Gamma) : G \rightarrow \mathbb{C}$ is given by

$$e(\Gamma(z)) = \sum_{\alpha \in G'} \widehat{\Gamma}(\alpha) e(\langle \alpha, z \rangle),$$

where $|\widehat{\Gamma}(\alpha)| \leq 1$ for all α . Let $Q_\alpha := \sum \alpha_i Q_i(x)$ for $\alpha \in G'$, which is a polynomial of degree $\leq d-1$. Then

$$e(P(x)) = \sum_{\alpha \in G'} \widehat{\Gamma}(\alpha) e(Q_\alpha(x)). \tag{7.1}$$

Thus

$$1 = |\langle e(P(x)), e(P(x)) \rangle| \leq \sum_{\alpha \in G'} |\langle e(P(x)), e(Q_\alpha(x)) \rangle|,$$

which shows that for some $\alpha^* \in G'$,

$$|\langle e(P), e(Q_{\alpha^*}) \rangle| \geq \frac{1}{|G|}.$$

Next, observe that by the direct theorem for the Gowers norm ([Theorem 6.7](#)), as Q_{α^*} has degree $\leq d-1$ we have

$$\|e(P)\|_{U^d} \geq |\langle e(P), e(Q_{\alpha^*}) \rangle| \geq |G|^{-1}.$$

To conclude, a simple calculation gives that

$$|G| = p^{(k_1+1)+\dots+(k_r+1)} \leq p^{(1+\lceil (d-1)/(p-1) \rceil)r}$$

as $k_i(p-1) \leq d-1$. Thus we conclude that

$$\|e(P)\|_{U^d} \geq \varepsilon(\mathbb{F}_p, d, r) := p^{-(1+\lceil (d-1)/(p-1) \rceil)r}.$$

□

Remark 7.17. The expansion in [Equation \(7.1\)](#) is referred to as the *higher-order Fourier expansion* of $e(P)$, and will be discussed further in [Section 9.2](#). Throughout this text we will repeatedly use the higher-order Fourier expansion of a function g to argue that g must be correlated with some low-degree polynomial.

It is straightforward to extend the notion of uniformity from a single polynomial (Definition 7.13) to a polynomial factor.

Definition 7.18 (Uniform factor). Let $\varepsilon : \mathbb{N} \rightarrow \mathbb{R}^+$ be a decreasing function. A polynomial factor \mathcal{B} defined by a sequence of polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$ with respective depths k_1, \dots, k_C is said to be ε -uniform if for every collection $(\lambda_1, \dots, \lambda_C) \in \mathbb{Z}^C$, with $(\lambda_1 \bmod p^{k_1+1}, \dots, \lambda_C \bmod p^{k_C+1}) \neq (0, \dots, 0)$, it holds that

$$\left\| e \left(\sum_i \lambda_i P_i \right) \right\|_{U^d} < \varepsilon(|\mathcal{B}|),$$

where $d = \max_i \deg(\lambda_i P_i)$.

Remark 7.19 (Equivalence between regularity and uniformity). Similar to Corollary 7.15 it also follows from Theorem 7.14 that an r -regular degree- d factor \mathcal{B} is also ε -uniform when $r = r_{7.14}(\mathbb{F}, d, \varepsilon)$ is as in Theorem 7.14. By Claim 7.16 the converse of this also holds and we have an approximate equivalence between regularity and uniformity.

7.3 The derivative polynomial

Let $P : \mathbb{F}^n \rightarrow \mathbb{T}$ be a nonclassical polynomial of degree d . The order- d Gowers norm of $e(P)$, which we have seen to control its correlation with lower degree polynomials, can be expressed as the bias of the *derivative polynomial* of P .

Definition 7.20 (Derivative polynomial). Let $P : \mathbb{F}^n \rightarrow \mathbb{T}$ be a degree- d polynomial. The derivative polynomial $\partial P : (\mathbb{F}^n)^d \rightarrow \mathbb{T}$ of P is defined as

$$\partial P(h_1, \dots, h_d) := D_{h_1} \cdots D_{h_d} P(0),$$

where $h_1, \dots, h_d \in \mathbb{F}^n$.

Remark 7.21. Note that we could equivalently define $\partial P(h_1, \dots, h_d) := D_{h_1} \cdots D_{h_d} P(y)$ for an arbitrary $y \in \mathbb{F}^n$. This is due to the fact that $D_{h_1} \cdots D_{h_d} P$ is a degree 0 function, which is a constant function.

A closed form for ∂P , applying the derivatives iteratively, is

$$\partial P(h_1, \dots, h_d) = \sum_{S \subseteq [d]} (-1)^{d-|S|} P \left(\sum_{i \in S} h_i \right).$$

The following lemma shows some useful properties of the derivative polynomial.

Lemma 7.22. Let $P : \mathbb{F}^n \rightarrow \mathbb{T}$ be a degree- d polynomial. Then

- (i) $\partial P(h_1, \dots, h_d)$ is a classical homogeneous polynomial of degree d .
- (ii) $\partial P(h_1, \dots, h_d)$ is invariant under permutations of h_1, \dots, h_d .
- (iii) $\partial P(h_1, \dots, h_d)$ is linear in each of h_1, \dots, h_d .
- (iv) For any $x \in \mathbb{F}^n$, $D_{h_1} \cdots D_{h_d} P(x) = \partial P(h_1, \dots, h_d)$.

Proof. The proof follows by the properties of the additive derivative D_h . By definition, $\partial P(h_1, \dots, h_d)$ is a (possibly nonclassical) polynomial of degree d . Item (i) follows since ∂P is annihilated by multiplication by p , since

$$p\partial P(h_1, h_2, \dots, h_d) = pD_{h_1} \cdots D_{h_d}(P)(0) = D_{h_1} \cdots D_{h_d}(pP)(0) = 0,$$

as we have $\deg(pP) = \max(d - (p - 1), 0) < d$ (see [Corollary 6.6](#)). Item (ii) holds since $D_h D_{h'} Q = D_{h'} D_h Q$ for every function $Q : \mathbb{F}^n \rightarrow \mathbb{T}$ and every $h, h' \in \mathbb{F}^n$. Item (iii) holds as every monomial of ∂P must depend on each of the variable sets h_i , as for example if we set $h_1 = 0$ then $\partial P(0, h_2, \dots, h_d) = 0$ since $D_0 Q = 0$ for any function Q . As $\deg(\partial P) \leq d$, each monomial must contain exactly one variable from each h_i . Hence ∂P is linear in each h_i and has the form

$$\partial P(h_1, \dots, h_d) = \sum_{i_1, \dots, i_d=1}^n c_{i_1, \dots, i_d} \prod_{j=1}^d (h_j)_{i_j},$$

where c_{i_1, \dots, i_d} depends only on the multiset $\{i_1, \dots, i_d\}$. Item (iv) follows as

$$D_{h_1} \cdots D_{h_d} P(x + h) - D_{h_1} \cdots D_{h_d} P(x) = D_{h_1} \cdots D_{h_d} D_h P(x) = 0.$$

□

Corollary 7.23. Let $P : \mathbb{F}^n \rightarrow \mathbb{T}$ be a polynomial of degree $\leq d$. Then P has low rank if and only if its derivative polynomial is biased. Formally, there exists functions $\varepsilon_1(\mathbb{F}, d, r_1)$ and $r_2(\mathbb{F}, d, \varepsilon_2)$ such that $\text{rank}_d(P) \leq r_1$ implies $\text{bias}(\partial P) \geq \varepsilon_1$ and $\text{bias}(\partial P) > \varepsilon_2$ implies $\text{rank}_d(P) \leq r_2$.

Proof. The corollary follows from [Theorem 7.14](#) and [Claim 7.16](#), since

$$\begin{aligned} \|e(P)\|_{U^d}^{2^d} &= \mathbb{E}_{h_1, \dots, h_d, x} [e(D_{h_1} \cdots D_{h_d} P(x))] \\ &= \mathbb{E}_{h_1, \dots, h_d} [e(D_{h_1} \cdots D_{h_d} P(0))] \\ &= \text{bias}(\partial P). \end{aligned}$$

□

7.4 Equidistribution of regular factors

In this section, we make precise the intuition that a high-rank collection of polynomials, evaluated on a joint input, behaves close to a collection of independent random variables. The key technical tool is the connection between the combinatorial notion of rank and the analytic notion of bias ([Theorem 7.10](#)).

Using a standard observation that relates the bias of a function to its distribution on its range, [Theorem 7.10](#) implies the following.

Lemma 7.24 (Size of atoms). Given $\varepsilon > 0$, let \mathcal{B} be a polynomial factor of degree $d \geq 1$, complexity C , and rank $r_{7.10}(\mathbb{F}, d, \varepsilon)$, defined by a tuple of polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$ having respective depths k_1, \dots, k_C . Suppose $b = (b_1, \dots, b_C) \in \mathbb{U}_{k_1+1} \times \cdots \times \mathbb{U}_{k_C+1}$. Then

$$\Pr_{x \in \mathbb{F}^n} [\mathcal{B}(x) = b] = \frac{1}{\|\mathcal{B}\|} \pm \varepsilon.$$

In particular, for $\varepsilon < \frac{1}{\|\mathcal{B}\|}$, $\mathcal{B}(x)$ attains every possible value in its range and thus has $\|\mathcal{B}\|$ atoms.

Proof. We can express the fraction of inputs in an atom b as

$$\begin{aligned} & \Pr_{x \in \mathbb{F}^n} [\mathcal{B}(x) = b] \\ &= \mathbb{E}_x \left[\prod_{i=1}^C \mathbf{1}_{[P_i(x)=b_i]} \right] \\ &= \mathbb{E}_x \left[\prod_{i=1}^C \frac{1}{p^{k_i+1}} \sum_{\lambda_i=0}^{p^{k_i+1}-1} e(\lambda_i(P_i(x) - b_i)) \right] \\ &= \prod_i p^{-(k_i+1)} \cdot \sum_{(\lambda_1, \dots, \lambda_C) \in \prod_{i=1}^C [0, p^{k_i+1}-1]} \mathbb{E}_x \left[e \left(\sum_{i=1}^C \lambda_i(P_i(x) - b_i) \right) \right]. \end{aligned}$$

The second equality uses the fact that $P_i(x) - b_i$ is in \mathbb{U}_{k_i+1} and that for every nonzero $x \in \mathbb{U}_{k_i+1}$, $\sum_{\lambda=0}^{p^{k_i+1}-1} e(\lambda x) = 0$. By our assumption that \mathcal{B} is r -regular, for every $(\lambda_1, \dots, \lambda_C) \neq \vec{0}$ we have that $\sum_{i=1}^C \lambda_i P_i(x)$ has rank at least r , which by [Theorem 7.10](#) implies that its bias is at most ε . Thus

$$\Pr_{x \in \mathbb{F}^n} [\mathcal{B}(x) = b] = \prod_{i=1}^C p^{-(k_i+1)} \cdot \left(1 \pm \varepsilon \prod_{i=1}^C p^{k_i+1} \right) = \frac{1}{\|\mathcal{B}\|} \pm \varepsilon.$$

□

An almost identical proof implies a similar statement for unbiased factors instead of regular factors.

Lemma 7.25 (Equidistribution for unbiased factors). Suppose that $\varepsilon : \mathbb{N} \rightarrow \mathbb{R}^+$ is a decreasing function. Let \mathcal{B} be an ε -unbiased factor of degree $d \geq 1$, defined by a tuple of polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$ having respective depths k_1, \dots, k_C . Suppose $b = (b_1, \dots, b_C) \in \mathbb{U}_{k_1+1} \times \dots \times \mathbb{U}_{k_C+1}$. Then

$$\Pr_{x \in \mathbb{F}^n} [\mathcal{B}(x) = b] = \frac{1}{\|\mathcal{B}\|} \pm \varepsilon(|\mathcal{B}|).$$

7.5 Regularization of factors

Due to the generic properties of regular factors, it is often useful to *refine* a given polynomial factor to a regular one. We will first formally define what we mean by refining a polynomial factor.

Definition 7.26 (Refinement). A factor \mathcal{B}' is called a *refinement* of \mathcal{B} , and denoted $\mathcal{B}' \succeq \mathcal{B}$, if the induced partition by \mathcal{B}' is a combinatorial refinement of the partition induced by \mathcal{B} . In other words, if for every $x, y \in \mathbb{F}^n$, $\mathcal{B}'(x) = \mathcal{B}'(y)$ implies $\mathcal{B}(x) = \mathcal{B}(y)$.

One needs to be careful about distinguishing between two types of refinements.

Definition 7.27 (Semantic and syntactic refinements). \mathcal{B}' is called a *syntactic refinement* of \mathcal{B} , and denoted $\mathcal{B}' \succeq_{syn} \mathcal{B}$, if the sequence of polynomials defining \mathcal{B}' extends that of \mathcal{B} . It is called a *semantic refinement*, and denoted $\mathcal{B}' \succeq_{sem} \mathcal{B}$ if the induced partition is a combinatorial refinement of the partition induced by \mathcal{B} . In other words, if for every $x, y \in \mathbb{F}^n$, $\mathcal{B}'(x) = \mathcal{B}'(y)$ implies $\mathcal{B}(x) = \mathcal{B}(y)$.

Remark 7.28. Clearly, being a syntactic refinement is stronger than being a semantic refinement. But observe that if \mathcal{B}' is a semantic refinement of \mathcal{B} , then there exists a syntactic refinement \mathcal{B}'' of \mathcal{B} that induces the same partition of \mathbb{F}^n as \mathcal{B}' , and for which $|\mathcal{B}''| \leq |\mathcal{B}'| + |\mathcal{B}|$. To construct \mathcal{B}'' , simply add the defining polynomials of \mathcal{B} to those of \mathcal{B}' .

The following lemma by Green and Tao [43] shows that every classical polynomial factor can be refined to a regular factor. The basic idea is simple: if some polynomial has low rank, decompose it to a few lower degree polynomials, and repeat. Formally, it follows by transfinite induction on the number of polynomials of each degree that define the polynomial factor.

Lemma 7.29 (Regularity lemma for classical polynomials [43]). Fix a prime finite field \mathbb{F} , an integer $d \geq 1$ and a non-decreasing function $r : \mathbb{N} \rightarrow \mathbb{N}$. There exists a function $C_{7.29}^{\mathbb{F}, d, r} : \mathbb{N} \rightarrow \mathbb{N}$ for which the following holds.

Suppose \mathcal{B} is a polynomial factor of degree $\leq d$ and complexity C , defined by classical polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{F}$. Then there exists an r -regular factor \mathcal{B}' that semantically refines \mathcal{B} , where \mathcal{B}' has degree $\leq d$ and complexity $C' \leq C_{7.29}^{(\mathbb{F}, d, r)}(C)$.

Furthermore, if \mathcal{B} is itself a syntactic refinement of some \mathcal{B}_0 that has rank $> r(C')$, then \mathcal{B}' can be taken to be a syntactic refinement of \mathcal{B}_0 .

Proof. Let \mathcal{B} be a polynomial factor defined by classical polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree at most d . Define $M(\mathcal{B}) := (M_d, \dots, M_1) \in \mathbb{N}^d$ where M_i is the number of polynomials of degree i among P_1, \dots, P_C . Note that $M_1 + \dots + M_d = C = |\mathcal{B}|$. We define the lexicographic order on \mathbb{N}^d , where $M > M'$ if $M_i > M'_i$ for some $1 \leq i \leq d$ and $M_j = M'_j$ for all $j > i$. The proof will be by transfinite induction on M under the lexicographic order. Namely, we will apply the fact that \mathbb{N}^d under the lexicographic order is Noetherian, that is, it does not contain an infinite decreasing sequence.

Let \mathcal{B} be a polynomial factor, and assume that \mathcal{B} is not r -regular. Then by definition, some linear combination of the polynomials that define \mathcal{B} has rank less than $r(C)$. Let $P(x) = \sum \lambda_i P_i(x)$ with $\lambda_i \in \mathbb{F}$, not all zero, such that $\text{rank}(P) \leq r(C)$. By definition of rank, we can decompose P as a function of $r(C)$ lower degree polynomials. That is,

$$P(x) = \Gamma(Q_1(x), \dots, Q_{r(C)}(x))$$

where $\deg(Q_i) \leq \deg(P) - 1$ and where $\Gamma : \mathbb{F}^{r(C)} \rightarrow \mathbb{F}$ is some function. Let $i^* \in [C]$ be chosen so that $\lambda_{i^*} \neq 0$ and $e = \deg(P_{i^*})$ is maximal. In particular, $\deg(P_{i^*}) \geq \deg(P)$. We can express P_{i^*} as a linear combination of $\{P_i : i \neq i^*\}$ and $\Gamma(Q_1, \dots, Q_{r(C)})$. Define $\mathcal{B}_1 = \mathcal{B} \setminus \{P_{i^*}\} \cup \{Q_1, \dots, Q_{r(C)}\}$. We claim that:

- (i) \mathcal{B}_1 is a semantic refinement of \mathcal{B} .
- (ii) $M(\mathcal{B}_1) < M(\mathcal{B})$.

The first item follows by definition, and the second since in order to construct \mathcal{B}_1 , we removed one polynomial of degree e from \mathcal{B} and added $r(|\mathcal{B}|)$ many lower degree polynomials. If \mathcal{B}_1 is still not r -regular, we can repeat this process and obtain \mathcal{B}_2 which is a semantic refinement of \mathcal{B}_1 , and so on. By transfinite induction, this process must halt after a finite number of steps. More formally, the number of steps depends only on $M(\mathcal{B})$ as in the argument above, we can provide an upper bound

M' which depends only on $M(\mathcal{B})$ such that $M(\mathcal{B}_1) \leq M' < M(\mathcal{B})$, by taking $M' = (M_d, \dots, M_{e+1}, M_e - 1, M_{e-1} + r(|\mathcal{B}|), \dots, M_1 + r(|\mathcal{B}|))$.

For the final part, assume that \mathcal{B} is a syntactic refinement of a polynomial factor \mathcal{B}_0 , defined without loss of generality by the $C_0 = |\mathcal{B}_0|$ polynomials $P_1, \dots, P_{C_0} : \mathbb{F}^n \rightarrow \mathbb{F}$. In the regularization process, we will attempt to choose $P_{i^*} \notin \{P_1, \dots, P_{C_0}\}$ if possible. If we can achieve this at every step, then at the end we retain all the original polynomials that define \mathcal{B}_0 , and hence we obtain a syntactic refinement of \mathcal{B}_0 . If we fail, then at some stage we obtained a polynomial factor \mathcal{B}'' for which some linear combination of polynomials from P_1, \dots, P_{C_0} has rank at most $r(|\mathcal{B}''|) \leq r(C')$. This contradicts our assumption that \mathcal{B}_0 has high rank. \square

The bounds obtained on $C_{7.29}^{\mathbb{F},d,r}$ have Ackermann-type dependence on the degree d , even when $r(\cdot)$ is a “reasonable” function. As such, it gives nontrivial results only for constant degrees. The extension of [Lemma 7.29](#) to nonclassical polynomials is more involved, and was proved by Tao and Ziegler [[83](#)] as part of their proof of the inverse Gowers theorem ([Theorem 6.9](#)).

Theorem 7.30 (Regularity lemma for nonclassical polynomials [[83](#)]). Fix a prime finite field \mathbb{F} , an integer $d \geq 1$ and a non-decreasing function $r : \mathbb{N} \rightarrow \mathbb{N}$. There exists a function $C_{7.30}^{\mathbb{F},d,r} : \mathbb{N} \rightarrow \mathbb{N}$ for which the following holds.

Suppose \mathcal{B} is a polynomial factor of degree $\leq d$ and complexity C , defined by nonclassical polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{T}$. Then there exists an r -regular factor \mathcal{B}' that semantically refines \mathcal{B} , where \mathcal{B}' has degree $\leq d$ and complexity $C' \leq C_{7.29}^{(\mathbb{F},d,r)}(C)$.

Furthermore, if \mathcal{B} is itself a syntactic refinement of some \mathcal{B}_0 that has rank $> r(C')$, then \mathcal{B}' can be taken to be a syntactic refinement of \mathcal{B}_0 .

We sketch the proof of [Theorem 7.30](#). The basic approach is the same as that of [Lemma 7.29](#). Let \mathcal{B} be a polynomial factor defined by nonclassical polynomials of degree $\leq d$. It is said to be *extended* if whenever $P \in \mathcal{B}$, then either $pP = 0$ or $pP \in \mathcal{B}$. Clearly, every polynomial factor can be made extended by adding the appropriate polynomials to

it. If \mathcal{B} is extended and defined by polynomials P_1, \dots, P_C , then for it to have rank r , it suffices if $\text{rank}(\sum \lambda_i P_i) \geq r$ for all $\lambda_i \in \{0, \dots, p-1\}$, not all zero. The main idea in the proof of [Theorem 7.30](#) is to apply the same inductive argument as in the proof of [Lemma 7.29](#), while maintaining an extended factor throughout the proof. However, this raises the following challenge: assume that P_i has low rank, and we wish to replace it with a few lower degree polynomials. However, assume that also $P_i = pP_j$, for some other polynomial P_j defining the factor. Then, we must remove P_j from the factor as well. If P_j also had low rank, that would suffice. However, this need not be true. What is true is that P_j can be decomposed as a function of a few polynomials, each of which either has lower degree or the same degree and lower depth as that of P_j . We summarize this in a lemma below, for which we omit the proof. With that lemma in hand, the proof goes through as before, except that now we need to keep track of the number of polynomials of a given depth and a given degree.

Lemma 7.31 (Tao and Ziegler [[83](#)]). Let $P : \mathbb{F}^n \rightarrow \mathbb{T}$ be a polynomial of degree d and rank r . Assume that $P = pQ$, where Q is a polynomial of degree $d + (p-1)$. Then $Q(x) = \Gamma(Q_1(x), \dots, Q_c(x))$, where for each i , either $\deg(Q_i) < \deg(Q)$ or $\deg(Q_i) = \deg(Q)$ and $\text{depth}(Q_i) < \text{depth}(Q)$. Furthermore, $c \leq C_{7.31}(\mathbb{F}, d, r)$.

7.6 Strong equidistribution of regular factors

One of the important and useful properties of the classical Fourier characters is that they form an orthonormal basis. [Theorem 7.10](#), [Lemma 7.24](#) and [Lemma 7.25](#) provide an approximate version of this phenomenon that is useful for several applications. However, for certain applications we need a stronger notion of orthogonality, one where each polynomial of the factor is evaluated not just on a single input, but on a collection of inputs given by linear forms.

A linear form in k variables is a vector $L = (\lambda_1, \dots, \lambda_k) \in \mathbb{F}^k$. It defines a map $L : (\mathbb{F}_n)^k \rightarrow \mathbb{F}_n$ by $L(x_1, \dots, x_k) = \sum \lambda_i x_i$. Often, we would need to analyze averages of f evaluated on several linear forms with joint variables. Consider for example the BLR linearity

test, explored in [Chapter 2](#). Given a function $P : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$, let $f(x) = (-1)^{P(x)}$. The acceptance probability of the BLR linearity test on P is equal to

$$\frac{1}{2} + \frac{1}{2} \mathbb{E}_{x,y \in \mathbb{F}^n} [f(x)f(y)f(x+y)].$$

Thus, in order to analyze it, we needed to evaluate f on three related inputs: $x, y, x + y$.

The general form is as follows. Let $L_1, \dots, L_m \in \mathbb{F}^k$ be linear forms, where to recall $L_i : (\mathbb{F}^n)^k \rightarrow \mathbb{F}^n$. The goal is to analyze, for $f : \mathbb{F}^n \rightarrow \mathbb{C}$, averages of the form

$$\mathbb{E}_{X \in (\mathbb{F}^n)^k} [f(L_1(X)) \cdots f(L_m(X))].$$

The first step in such an analysis will be to find a suitable polynomial factor \mathcal{B} , such that we can replace f with $\mathbb{E}[f|\mathcal{B}]$ without affecting the average by much. Let us assume that we have already done so. Then we can specialize our treatment to functions of the form $f(x) = \Gamma(P_1(x), \dots, P_C(x))$, where $P_i : \mathbb{F}^n \rightarrow \mathbb{T}$ are bounded degree polynomials. This in turn would require to analyze the joint distribution of P_1, \dots, P_C on the linear forms $L_1(X), \dots, L_m(X)$. That is, we would need to understand the distribution of the random variable

$$A_{\mathcal{B}, \mathcal{L}}(X) := \begin{pmatrix} P_1(L_1(X)) & P_2(L_1(X)) & \dots & P_C(L_1(X)) \\ P_1(L_2(X)) & P_2(L_2(X)) & \dots & P_C(L_2(X)) \\ \vdots & & & \vdots \\ P_1(L_m(X)) & P_2(L_m(X)) & \dots & P_C(L_m(X)) \end{pmatrix},$$

where $X \in (\mathbb{F}^n)^k$ is uniformly distributed, $\mathcal{B} = \{P_1, \dots, P_C\}$ is a family of nonclassical polynomials and $\mathcal{L} = \{L_1, \dots, L_m\} \subset \mathbb{F}^k$ is a system of linear forms.

Our analysis so far (see [Lemma 7.24](#)) have shown that if we restrict our attention to a single linear form, namely a single row of $A_{\mathcal{B}, \mathcal{L}}(X)$, then if we assume that \mathcal{B} is regular enough, then its values are distributed close to uniform in their range $\prod_{j=1}^C \mathbb{U}_{k_j+1}$. Thus, a first guess might be that if \mathcal{B} is regular enough, then $A_{\mathcal{B}, \mathcal{L}}$ should be uniformly distributed over all the $m \times C$ matrices $A \in \prod_{i=1}^m \prod_{j=1}^C \mathbb{U}_{k_j+1}$.

However, this is false. The reason is that nonclassical polynomials of a given degree satisfy various linear identities, governed by the fact that

they are annihilated by taking enough derivatives. For example, if P_1 is a linear polynomial then $P_1(x_1 + x_2 + x_3) - P_1(x_1 + x_2) - P_1(x_1 + x_3) + P_1(x_1) = 0$ holds, and hence the four random variables $(P_1(x_1 + x_2 + x_3), P_1(x_1 + x_2), P_1(x_1 + x_3), P_1(x_1))$ are far from independent. Thus, the rows of $A_{\mathcal{B}, \mathcal{L}}(X)$ satisfy certain linear constraints. As we will shortly see, strong equidistribution implies that $A_{\mathcal{B}, \mathcal{L}}(X)$ is close to uniform on all values that satisfy the necessary linear constraints. In particular, the columns of $A_{\mathcal{B}, \mathcal{L}}(X)$ are nearly independent, with each column being uniform modulo the required linear dependencies.

The development of the strong equidistribution theorem occurred in several steps. Hatami and Lovett [48] established a strong near-equidistribution for factors of classical polynomials, when the characteristic of the field \mathbb{F} is greater than the degree of the polynomial factor. Bhattacharyya *et al.* [14] later extended the result to the general characteristic case and nonclassical polynomials, but under an extra assumption that the system of linear forms is affine, i.e. there is a variable that appears with coefficient 1 in all the linear forms. Bhattacharyya *et al.* [17] used the same proof technique to replace the condition that each L_i is affine, with the condition that all the coefficients of L_i are in $\{0, 1\}$. Finally, in the work of Hatami *et al.* [46] the joint distribution of the matrix $A_{\mathcal{B}, \mathcal{L}}$ was fully characterized for nonclassical polynomials without any extra assumptions on the linear forms. Instead, it requires that the polynomials are *homogeneous*.

We formally define and analyze homogeneous nonclassical polynomials in Chapter 10. For the time being, it is sufficient to note that each nonclassical polynomial $P(x)$ of degree d can be decomposed as a sum of homogeneous nonclassical polynomials of degrees $1, \dots, d$; and that in the regularization process (as in Theorem 7.30) one can require that the resulting polynomials are all homogeneous. Thus, one can always assume that a polynomial factor is defined by homogeneous nonclassical polynomials, and these can be made high rank by regularization.

Theorem 7.32 (Near orthogonality over linear forms). Let \mathbb{F} be a prime field, $d \geq 1$, $\varepsilon > 0$. Let $\{L_1, \dots, L_m\}$ be a system of linear forms. Let $\mathcal{B} = \{P_1, \dots, P_C\}$ be a polynomial factor of degree at most d and $\text{rank}(\mathcal{B}) > r_{7.32}(\mathbb{F}, d, \varepsilon)$. Assume furthermore that each P_i is a

nonclassical homogeneous polynomial (defined in [Chapter 10](#)). For every set of coefficients $\Lambda = \{\lambda_{i,j} \in \mathbb{Z} : i \in [C], j \in [m]\}$, let

$$P_\Lambda(x) := \sum_{i=1}^C \sum_{j=1}^m \lambda_{i,j} P_i(L_j(x)).$$

Then one of the following two cases holds:

- $P_\Lambda \equiv 0$. In this case, for every $i \in [C]$ and any nonclassical homogeneous polynomials $Q_i : \mathbb{F}^n \rightarrow \mathbb{T}$ with $\deg(Q_i) = \deg(P_i)$ and $\text{depth}(Q_i) \leq \text{depth}(P_i)$ it holds that $\sum_{j \in [m]} \lambda_{i,j} Q_i(L_j(\cdot)) \equiv 0$.
- $P_\Lambda \not\equiv 0$. In this case, $|\mathbb{E}[e(P_\Lambda)]| < \varepsilon$.

We defer the proof of [Theorem 7.32](#) to [Section 12.1](#), after we cover some more necessary background material.

Next, we state a special version of [Theorem 7.32](#) useful in some applications. A system of linear forms L_1, \dots, L_m is said to be an *affine constraint* if $L_{i,1} = 1$ for all $i \in [m]$. Such systems arise naturally in the study of affine invariant properties, which we discuss in [Chapter 14](#). Here, we state a special case of [Theorem 7.32](#) for affine constraints, where the polynomials are not required to be homogeneous. For the proof see [\[14\]](#).

Theorem 7.33 (Near orthogonality over affine constraints). Let \mathbb{F} be a prime field, $d \geq 1$, $\varepsilon > 0$. Let $\{L_1, \dots, L_m\}$ be an affine constraint. Let $\mathcal{B} = \{P_1, \dots, P_c\}$ be a polynomial factor of degree at most d and $\text{rank}(\mathcal{B}) > r_{7.33}(\mathbb{F}, d, \varepsilon)$. For every set of coefficients $\Lambda = \{\lambda_{i,j} \in \mathbb{Z} : i \in [c], j \in [m]\}$, let

$$P_\Lambda(x) := \sum_{i=1}^c \sum_{j=1}^m \lambda_{i,j} P_i(L_j(x)).$$

Then one of the following two cases holds:

- $P_\Lambda \equiv 0$. In this case, for every $i \in [C]$ and any nonclassical polynomials $Q_i : \mathbb{F}^n \rightarrow \mathbb{T}$ with $\deg(Q_i) \leq \deg(P_i)$ and $\text{depth}(Q_i) \leq \text{depth}(P_i)$, it holds that $\sum_{j \in [m]} \lambda_{i,j} Q_i(L_j(\cdot)) \equiv 0$.
- $P_\Lambda \not\equiv 0$. In this case, $|\mathbb{E}[e(P_\Lambda)]| < \varepsilon$.

In the next section, as a corollary of [Theorem 7.32](#), we determine the distribution of $A_{\mathcal{B}, \mathcal{L}}$ when \mathcal{B} is of sufficiently high rank.

7.7 Joint distribution of high-rank polynomials over linear forms

In order to understand the joint distribution of polynomials over linear forms, we first need to understand the necessary linear constraints. The following definition formalizes this.

Definition 7.34. Let $\mathbb{F} = \mathbb{F}_p$ be a prime field. Given a system of linear forms L_1, \dots, L_m over \mathbb{F} in ℓ variables. and integers $d, k > 0$, the (d, k) -dependency set of L_1, \dots, L_m is the set of tuples $(\lambda_1, \dots, \lambda_m) \in \mathbb{Z}_{p^{k+1}}^m$ such that $\sum_{i=1}^m \lambda_i P(L_i(x_1, \dots, x_\ell)) \equiv 0$ for every homogeneous polynomial $P : \mathbb{F}^\ell \rightarrow \mathbb{T}$ of degree d and depth k .

Observe that the (d, k) -dependency set of L_1, \dots, L_m is closed under addition, and hence is a subgroup of $\mathbb{Z}_{p^{k+1}}^m$. We do not currently have an explicit description of it, except for the implicit definition given in Definition 7.34.

Theorem 7.32 characterizes exactly when for a regular factor \mathcal{B} it holds that $P_\Lambda \equiv 0$. We record this in the language of dependency sets.

Corollary 7.35. Let L_1, \dots, L_m be a system of linear forms over $\mathbb{F} = \mathbb{F}_p$ in ℓ variables. Fix an integer $c > 0$, tuples $(d_1, \dots, d_c) \in \mathbb{Z}_{>0}^c$ and $(k_1, \dots, k_c) \in \mathbb{Z}_{\geq 0}^c$. For $i \in [c]$, let Λ_i be the (d_i, k_i) -dependency set of L_1, \dots, L_m .

Then, for every polynomial factor \mathcal{B} defined by homogeneous polynomials $P_1, \dots, P_c : \mathbb{F}^\ell \rightarrow \mathbb{T}$, where P_i has degree d_i and depth k_i , and \mathcal{B} has rank $> r_{7.14} \left(\mathbb{F}, \max_i d_i, \frac{1}{2} \right)$, it is the case that a tuple $(\lambda_{i,j})_{i \in [c], j \in [m]}$ satisfies

$$\sum_{i=1}^c \sum_{j=1}^m \lambda_{i,j} P_i(L_j(x_1, \dots, x_\ell)) \equiv 0$$

if and only if for every $i \in [c]$,

$$(\lambda_{i,1} \bmod p^{k_i+1}, \dots, \lambda_{i,m} \bmod p^{k_i+1}) \in \Lambda_i.$$

Proof. The “if” direction is obvious. For the “only if” direction, we use Theorem 7.32 to conclude that if $\sum_{i,j} \lambda_{i,j} P_i(L_j(\cdot)) \equiv 0$, it must be that for every $i \in [c]$, $\sum_j \lambda_{i,j} Q_i(L_j(\cdot)) \equiv 0$ for every homogeneous polynomial Q_i with degree d_i and depth k_i . This is equivalent to saying $(\lambda_{i,1} \bmod p^{k_i+1}, \dots, \lambda_{i,m} \bmod p^{k_i+1}) \in \Lambda_i$. \square

Remark 7.36. For large characteristic fields, Hatami and Lovett [48] showed that the analog of Corollary 7.35 is true even without the rank condition.

The joint distribution of $(P_i(L_j(x_1, \dots, x_\ell)) : i \in [c], j \in [m])$ is only going to be supported on atoms which respect the constraints imposed by dependency sets. This is obvious: if P is a polynomial of degree d and depth k , such that $P(L_j(x_1, \dots, x_\ell)) = b_j$, then for every $(\lambda_1, \dots, \lambda_m)$ in the (d, k) -dependency set of (L_1, \dots, L_m) it must be the case that $\sum_j \lambda_j b_j = 0$. We call atoms which respect this constraint for all P_i in a factor, *consistent*.

Definition 7.37 (Consistency). Let \mathbb{F} be a prime field. Let L_1, \dots, L_m be a system of linear forms over \mathbb{F} in ℓ variables. A sequence of elements $b_1, \dots, b_m \in \mathbb{T}$ are said to be (d, k) -consistent with L_1, \dots, L_m if $b_1, \dots, b_m \in \mathbb{U}_{k+1}$ and for every tuple $(\lambda_1, \dots, \lambda_m)$ in the (d, k) -dependency set of L_1, \dots, L_m , it holds that $\sum_{i=1}^m \lambda_i b_i = 0$.

Given vectors $\mathbf{d} = (d_1, \dots, d_c) \in \mathbb{Z}_{>0}^c$ and $\mathbf{k} = (k_1, \dots, k_c) \in \mathbb{Z}_{\geq 0}^c$, a sequence of vectors $b_1, \dots, b_m \in \mathbb{T}^c$ are said to be (\mathbf{d}, \mathbf{k}) -consistent with L_1, \dots, L_m if for every $i \in [c]$, the elements $b_{1,i}, \dots, b_{m,i}$ are (d_i, k_i) -consistent with L_1, \dots, L_m .

If \mathcal{B} is a polynomial factor, the term \mathcal{B} -consistent with L_1, \dots, L_m is a synonym for (\mathbf{d}, \mathbf{k}) -consistent with L_1, \dots, L_m where $\mathbf{d} = (d_1, \dots, d_c)$ and $\mathbf{k} = (k_1, \dots, k_c)$ are respectively the degree and depth sequences of polynomials defining \mathcal{B} .

Theorem 7.32 implies that for a polynomial factor of large enough rank, the joint distribution $(P_i(L_j(x_1, \dots, x_\ell)))$ is equi-distributed on all consistent values.

Theorem 7.38. Given $\varepsilon > 0$, let \mathcal{B} be a polynomial factor of degree $d > 0$ and rank at least $r_{7.10}(\mathbb{F}, d, \varepsilon)$. Assume that \mathcal{B} is defined by a sequence of homogeneous polynomials $P_1, \dots, P_c : \mathbb{F}^n \rightarrow \mathbb{T}$ having respective degrees d_1, \dots, d_c and respective depths k_1, \dots, k_c . Let L_1, \dots, L_m be a system of linear forms over \mathbb{F} in ℓ variables, and let Λ_i be the (d_i, k_i) -dependency set of L_1, \dots, L_m .

7.7. Joint distribution of high-rank polynomials over linear forms 331

Suppose $b_1, \dots, b_m \in \mathbb{T}^c$ are atoms of \mathcal{B} that are \mathcal{B} -consistent with L_1, \dots, L_m . We denote $b_i = (b_{i,1}, \dots, b_{i,c})$. Then

$$\Pr_{x_1, \dots, x_\ell \in \mathbb{F}^n} [P_i(L_j(x_1, \dots, x_\ell)) = b_{j,i} \ \forall i \in [c], j \in [m]] = \frac{\prod_{i=1}^c |\Lambda_i|}{\|\mathcal{B}\|^m} \pm \varepsilon.$$

Proof. The proof is similar to that of [Lemma 7.24](#). Let $x = (x_1, \dots, x_\ell)$.

$$\begin{aligned} & \Pr_x [P_i(L_j(x)) = b_{j,i} \ \forall i \in [c], \forall j \in [m]] \\ &= \mathbb{E}_x \left[\prod_{i \in [c], j \in [m]} \frac{1}{p^{k_i+1}} \sum_{\lambda_{i,j}=0}^{p^{k_i+1}-1} e(\lambda_{i,j}(P_i(L_j(x)) - b_{j,i})) \right] \\ &= p^{-m \sum_{i=1}^c (k_i+1)} \cdot \sum_{(\lambda_{i,j}) \in \prod_{i,j} [0, p^{k_i+1}-1]} e\left(-\sum_{i,j} \lambda_{i,j} b_{j,i}\right) \mathbb{E} \left[e\left(\sum_{i,j} \lambda_{i,j} P_i(L_j(x))\right) \right] \\ &= p^{-m \sum_{i=1}^c (k_i+1)} \cdot \left(\prod_{i=1}^c |\Lambda_i| \pm \varepsilon p^{m \sum_{i=1}^c (k_i+1)} \right). \end{aligned}$$

To see the last equality, observe that by [Corollary 7.35](#), $\sum_{i,j} \lambda_{i,j} P_i(L_j(\cdot))$ is identically zero for $\prod_i |\Lambda_i|$ many tuples $(\lambda_{i,j})$ and, in that case, $\sum_{i,j} \lambda_{i,j} b_{j,i} = 0$ because of the consistency requirement. For any other tuple $(\lambda_{i,j})$, the expectation in the third line is bounded by ε in absolute value. □

8

Bias vs Low Rank in Large Fields

Theorem 7.10 exhibited the “bias implies low rank” phenomena: every biased polynomial has low rank. Concretely, if $P : \mathbb{F}^n \rightarrow \mathbb{F}$ is a degree d polynomial, and $|\mathbb{E}_x[e(P(x))]| \geq \varepsilon$, then $\text{rank}(P) \leq r_{7.10}(\mathbb{F}, d, \varepsilon)$. In this section, we describe an improved theorem due to Bhowmick and Lovett [19] which allows for large fields (of size growing with n) as well as for errors which are polynomially small in the field order.

Theorem 8.1. Let $d, s \in \mathbb{N}$. Let \mathbb{F} be a finite field of characteristic $\text{char}(\mathbb{F}) > d$. Let $P : \mathbb{F}^n \rightarrow \mathbb{F}$ be a polynomial of degree d . Suppose that $|\mathbb{E}_{x \in \mathbb{F}^n}[e(P(x))]| \geq |\mathbb{F}|^{-s}$. Then, $\text{rank}(P) \leq r_{8.1}(d, s)$.

Observe that here $r_{8.1}(d, s)$ does not depend on the field. Also, the error is allowed to be polynomially small in the field size. For concreteness, we focus on the proof where \mathbb{F} is a prime field, where we assume throughout that $|\mathbb{F}| > d$. For the proof for non-prime fields we refer the interested reader to the original paper by Bhowmick and Lovett [19].

8.1 Bias implies low rank approximation

We start by showing that biased polynomials can be approximated by a few lower degree polynomials.

Lemma 8.2. Let $d, s, t \in \mathbb{N}$. Let $P : \mathbb{F}^n \rightarrow \mathbb{F}$ be a polynomial of degree d . Suppose that $|\mathbb{E}_{x \in \mathbb{F}^n}[e(P(x))]| \geq |\mathbb{F}|^{-s}$. Then, there exist polynomials $Q_1, \dots, Q_c : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree at most $d - 1$, where $c = c(d, s, t) = \binom{d+t+2s+3}{d}$, and a function $\Gamma : \mathbb{F}^c \rightarrow \mathbb{F}$, such that

$$\Pr_{x \in \mathbb{F}^n} [P(x) \neq \Gamma(Q_1(x), \dots, Q_c(x))] \leq |\mathbb{F}|^{-t}.$$

We prove Lemma 8.2 in this section. Fix a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree d , and let $\mu := \mathbb{E}_{x \in \mathbb{F}^n}[e(P(x))]$ be its bias, where we assume $|\mu| \geq |\mathbb{F}|^{-s}$. We begin with the following claim.

Claim 8.3. For all $x \in \mathbb{F}^n$,

$$\mu \cdot e(-P(x)) = \mathbb{E}_{y \in \mathbb{F}^n} [e(D_y P(x))].$$

Proof. We have

$$\begin{aligned} \mathbb{E}_{y \in \mathbb{F}^n} [e(D_y P(x))] &= \mathbb{E}_{y \in \mathbb{F}^n} [e(P(x+y))e(-P(x))] \\ &= \mathbb{E}_{y \in \mathbb{F}^n} [e(P(y))] \cdot e(-P(x)) \\ &= \mu \cdot e(-P(x)). \end{aligned}$$

□

Fix $x \in \mathbb{F}^n$. Pick $z = (z_1, \dots, z_k) \in (\mathbb{F}^n)^k$ uniformly for some k to be specified later. For $a \in \mathbb{F}^k, z \in (\mathbb{F}^n)^k$, we shorthand $a \cdot z = \sum_{i=1}^k a_i z_i \in \mathbb{F}^n$. For $a \in \mathbb{F}^k \setminus \{0\}$, let $W_{x,a}(z)$ be the random variable (over the choice of z) defined as

$$W_{x,a}(z) := e(D_{a \cdot z} P(x)).$$

For $a \neq 0^k$, we have

$$\mathbb{E}_{z \in (\mathbb{F}^n)^k} [W_{x,a}(z)] = \mathbb{E}_{y \in \mathbb{F}^n} [e(D_y P(x))].$$

Also, observe that for distinct $\alpha, \beta \in \mathbb{F}$,

$$|e(\alpha) - e(\beta)| \geq |\mathbb{F}|^{-1}.$$

We have the following.

Claim 8.4. Fix $x \in \mathbb{F}^n$. If for $z \in (\mathbb{F}^n)^k$ it holds that

$$\left| \frac{1}{|\mathbb{F}|^k - 1} \sum_{a \in \mathbb{F}^k \setminus \{0^k\}} W_{x,a}(z) - \mathbb{E}_y [e(D_y P(x))] \right| < \frac{1}{2|\mathbb{F}|^{s+1}},$$

then

$$P(x) = \Gamma_0 \left(D_{a,z} P(x) : a \in \mathbb{F}^k \setminus \{0\} \right)$$

where $\Gamma_0 : \mathbb{F}^{|\mathbb{F}|^k - 1} \rightarrow \mathbb{F}$ is defined as

$$\Gamma_0(v_1, \dots, v_{|\mathbb{F}|^k - 1}) = \arg \min_{\alpha \in \mathbb{F}} \left| \frac{1}{|\mathbb{F}|^k - 1} \sum_{i=1}^{|\mathbb{F}|^k - 1} e(v_i) - e(-\alpha) \mu \right|.$$

Proof. By Claim 8.3, we have $\mu \cdot e(-P(x)) = \mathbb{E}_y [e(D_y P(x))]$. Observe that for any $\beta \in \mathbb{F} \setminus \{-P(x)\}$,

$$|\mu \cdot e(-P(x)) - \mu \cdot e(\beta)| \geq \mu \cdot |\mathbb{F}|^{-1} \geq |\mathbb{F}|^{-(s+1)}.$$

Thus by the assumption of the claim, the value of $-P(x)$ (and hence $P(x)$) is determined by finding α for which, $\mu \cdot e(\alpha)$ is closest to $\frac{1}{|\mathbb{F}|^k - 1} \sum_{a \in \mathbb{F}^k \setminus \{0^k\}} W_{x,a}(z)$. It is easy to check that this is captured by Γ_0 , namely

$$\Gamma_0 \left(D_{a,z} P(x) : a \in \mathbb{F}^k \setminus \{0\} \right) = P(x).$$

□

Observe that the random variables $\{W_{x,a}(z) : a \in \mathbb{F}^k \setminus \{0\}\}$ are pairwise independent. That is, for any distinct a, a' the random variable $(W_{x,a}(z), W_{x,a'}(z))$ is uniformly distributed in \mathbb{F}^2 , when $z \in \mathbb{F}^k$ is uniformly chosen. Thus, we can apply Chebychev's inequality and obtain that, for $k = t + 2s + 3$, it holds that

$$\begin{aligned} \Pr_{z \in (\mathbb{F}^n)^k} \left[\left| \frac{1}{|\mathbb{F}|^k - 1} \sum_{a \neq 0} W_{x,a}(z) - \mathbb{E}_y [e(D_y P(x))] \right| \geq \frac{1}{2|\mathbb{F}|^{s+1}} \right] \\ \leq \frac{4|\mathbb{F}|^{2s+2}}{|\mathbb{F}|^k - 1} \leq \frac{1}{|\mathbb{F}|^t}. \end{aligned} \quad (8.1)$$

Thus, for all $x \in \mathbb{F}^n$,

$$\Pr_{z \in (\mathbb{F}^n)^k} [\Gamma_0 \left(D_{a,z} P(x) : a \in \mathbb{F}^k \setminus \{0\} \right) = P(x)] \geq 1 - |\mathbb{F}|^{-t}.$$

Therefore, by an averaging argument, there exists a choice of $z \in (\mathbb{F}^n)^k$ for which

$$\Pr_{x \in \mathbb{F}^n} [\Gamma_0 (D_{a \cdot z} P(x) : a \in \mathbb{F}^k \setminus \{0\}) = P(x)] \geq 1 - |\mathbb{F}|^{-t}. \tag{8.2}$$

We now prove that we only need a constant number of derivatives in order to approximate P , instead of a number which is polynomial in $|\mathbb{F}|$. Let $|\cdot| : \mathbb{F} \rightarrow \mathbb{N}$ be the canonical map $|x| = x$ for $x = \{0, 1, \dots, |\mathbb{F}| - 1\}$.

Claim 8.5. Let $B := \{b \in \mathbb{F}^k : \sum_{j=1}^k |b_j| \leq d\}$. Then for any $a \in \mathbb{F}^k$,

$$D_{a \cdot z} P(x) = \sum_{b \in B} \lambda_{a,b} D_{b \cdot z} P(x)$$

for some $\lambda_{a,b} \in \mathbb{F}$.

Proof. Let $|a| = \sum_{i=1}^k |a_i|$. We prove the claim by induction on $|a|$. If $|a| \leq d$, the claim is straightforward, so assume $|a| > d$. As P is a degree d polynomial, we have for any $m > d$ and $y_1, \dots, y_m \in \mathbb{F}^n$ that

$$D_{y_1} \dots D_{y_m} P \equiv 0.$$

This translates to

$$\sum_{c \in \{0,1\}^m} (-1)^{\sum c_i} P(x + \sum c_i y_i) = 0.$$

As the sum of the coefficients is zero, this implies

$$\begin{aligned} \sum_{c \in \{0,1\}^m} (-1)^{\sum c_i} \left(P(x + \sum c_i y_i) - P(x) \right) \\ = \sum_{c \in \{0,1\}^m} (-1)^{\sum c_i} D_{c \cdot y} P(x) = 0, \end{aligned}$$

for $y = (y_1, \dots, y_m)$. Apply this for $m = |a|$ and y_1, \dots, y_m set to z_1 repeated a_1 times, z_2 repeated a_2 times, up to z_k repeated a_k times. Then we obtain that

$$\sum_{a' \leq a} \nu_{a'} D_{a' \cdot z} P(x) = 0,$$

where the sum is over all $a' \in \mathbb{F}^k$ such that $|a'_i| \leq |a_i|$ for all $1 \leq i \leq k$, and $\nu_{a'} = (-1)^{|a'|} \prod_{i=1}^k \binom{a_i}{a'_i}$. In particular, $\nu_a \in \{-1, +1\}$. This implies that $D_{a \cdot z} P(x)$ is a linear combination of $D_{a' \cdot z} P(x)$ for $a' \in \mathbb{F}^k$ with $|a'| < |a|$. The claim now follows by applying the induction hypothesis. \square

This concludes the proof of [Lemma 8.2](#). By [Claim 8.5](#) we may rewrite

$$\Gamma_0 \left(D_{a,z}P(x) : a \in \mathbb{F}^k \setminus \{0\} \right) = \Gamma \left(D_{a,z}P(x) : a \in B \right),$$

where $c = |B|$ and $\Gamma : \mathbb{F}^c \rightarrow \mathbb{F}$ is defined as $\Gamma(v_b : b \in B) = \Gamma_0 \left((\sum_{b \in B} \lambda_{a,b} v_b) : a \in \mathbb{F}^k \setminus \{0\} \right)$. Let Q_1, \dots, Q_c in the statement of [Lemma 8.2](#) be $\{D_{a,z}P(x) : a \in B\}$. Clearly, these are polynomials of degree $\leq d - 1$. To conclude, we need to bound $c = |B|$. We have

$$|B| = \sum_{i=1}^d \binom{k}{d} \leq \binom{d+k}{d}.$$

[Lemma 8.2](#) follows as we set $k = t + 2s + 3$.

8.2 Bias implies low rank exact computation

We prove [Theorem 8.1](#) in this section. The proof is by induction on the degree d and follows along the lines of [Theorem 1.7](#) of [\[43\]](#). We sketch the proof below.

We first show that [Theorem 8.1](#) implies a similar theorem, where instead of assuming that a polynomial is biased, we assume that it has a noticeable Gowers uniformity norm.

Lemma 8.6 (Large Gowers norm implies low rank). Suppose [Theorem 8.1](#) is true up to degree d . Let $P : \mathbb{F}^n \rightarrow \mathbb{F}$ be a polynomial of degree d . Suppose that $\|e(P)\|_{U^d} \geq |\mathbb{F}|^{-s}$. Then $\text{rank}(P) \leq r_{(8.6)}(d, s)$.

Proof. We have

$$\left| \mathbb{E}_{x, y_1, \dots, y_d \in \mathbb{F}^n} [e(D_{y_1, \dots, y_d}P(x))] \right| = \|e(P)\|_{U^d}^{2^d} \geq |\mathbb{F}|^{-2^d s}.$$

Define $Q : \mathbb{F}^{n(d+1)} \rightarrow \mathbb{F}$ as

$$Q(x, y_1, \dots, y_d) := D_{y_1, \dots, y_d}P(x).$$

By [Theorem 8.1](#),

$$\text{rank}(Q) \leq r_{(8.1)}(d, 2^d s).$$

Applying the Taylor approximation theorem to P , as we assume $d < |\mathbb{F}|$, we have

$$P(x) = \frac{1}{d!} D_{x, \dots, x} P(0) + R(x) = \frac{1}{d!} Q(0, x, \dots, x) + R(x),$$

where R is a polynomial of degree $\leq d - 1$. We conclude that $\text{rank}(P) \leq \text{rank}(Q) + 1$. This concludes the lemma by setting

$$r_{(8.6)}(d, s) = r_{(8.1)}(d, 2^d s) + 1. \quad \square$$

The next lemma shows that a regular factor has atoms of roughly equal size. We shorthand $\alpha \pm \delta$ to denote any element in the interval $[\alpha - \delta, \alpha + \delta]$.

Lemma 8.7 (Size of atoms). Suppose [Theorem 8.1](#) is true up to degree d . Let $\mathcal{B} = \{Q_1, \dots, Q_c\}$ be a polynomial factor of degree at most d . Given $s \in \mathbb{N}$, assume that \mathcal{B} has rank at least $r_{(8.1)}(d, s)$. Then for every $b \in \mathbb{F}^c$,

$$\Pr_{x \in \mathbb{F}^n} [\mathcal{B}(x) = b] = \frac{1}{|\mathbb{F}|^c} \pm \frac{1}{|\mathbb{F}|^s}.$$

Proof. For any $b \in \mathbb{F}^c$,

$$\begin{aligned} \Pr[\mathcal{B}(x) = b] &= \frac{1}{|\mathbb{F}|^c} \sum_{a \in \mathbb{F}^c} \mathbb{E}_x \left[e \left(\sum_i a_i (Q_i(x) - b_i) \right) \right] \\ &= \frac{1}{|\mathbb{F}|^c} \pm \frac{1}{|\mathbb{F}|^c} \sum_{0 \neq a \in \mathbb{F}^c} \left| \mathbb{E}_x \left[e \left(\sum_i a_i Q_i(x) \right) \right] \right| \\ &= \frac{1}{|\mathbb{F}|^c} \pm \frac{1}{|\mathbb{F}|^s} \end{aligned}$$

The last line follows because of the following argument. Suppose for some $a \neq 0$, $|\mathbb{E}_x [e(\sum_i a_i Q_i(x))]| > \frac{1}{|\mathbb{F}|^s}$. Then by [Theorem 8.1](#), $\text{rank}(\sum_i a_i Q_i) \leq r_{(8.1)}(d, s)$. This contradicts the assumption on the rank of \mathcal{B} . \square

Recall that a linear form $L = (\ell_1, \dots, \ell_k) \in \mathbb{F}^k$ is said to be *affine* if $\ell_1 = 1$. Next, we argue that a high rank factor evaluated on a system of affine linear forms is near orthogonal.

Lemma 8.8 (Near orthogonality of affine linear forms). Suppose [Theorem 8.1](#) is true up to degree d . Let $c, d, k, s, m \in \mathbb{N}$. Let $\mathcal{B} = \{Q_1, \dots, Q_c\}$ be a polynomial factor of degree at most d . Assume \mathcal{B} has rank at least $r_{(8.8)}(d, k, s)$. Let (L_1, \dots, L_m) be a system of affine linear forms on k variables. Let $\Lambda = (\lambda_{ij})_{i \in [c], j \in [m]}$ be a tuple of integers. Define

$$Q_\Lambda(x_1, \dots, x_k) = \sum_{i \in [c], j \in [m]} \lambda_{ij} Q_i(L_j(x_1, \dots, x_k)).$$

Then, one of the following is true.

1. $Q_\Lambda \equiv 0$. Moreover, for every $i \in [c]$, and Q_i of degree at most d , it holds that $\sum_{j=1}^m \lambda_{ij} Q_i(L_j(\cdot)) \equiv 0$.
2. $Q_\Lambda \not\equiv 0$. Moreover, $|\mathbb{E}[e(Q_\Lambda(x_1, \dots, x_k)))]| \leq |\mathbb{F}|^{-s}$.

The proof is identical to the proof of [Theorem 7.33](#), specialized to only allow classical polynomials, and taking care of the improved dependence on $|\mathbb{F}|$ guaranteed by [Lemma 8.6](#). As a corollary, we state the above result for the case of parallelepipeds, which will be needed for the inductive proof of [Theorem 8.1](#). We first set up some notations, following Section 4 of [\[43\]](#).

Let $\mathcal{B} = \{Q_1, \dots, Q_c\}$ be a polynomial factor of degree at most d . We assume \mathcal{B} has rank at least $r_{(8.6)}(d, s)$. For $i \in [d]$, let M_i denote the number of polynomials in \mathcal{B} of degree exactly equal to i .

Definition 8.9 (Faces and lower faces). Let $k \in \mathbb{N}$ and $0 \leq k' \leq k$. A set $F \subseteq \{0, 1\}^k$ is called a *face* of dimension k' if

$$F = \{b \in \{0, 1\}^k : b_i = \delta_i, i \in I\},$$

where $I \subseteq [k]$, $|I| = k - k'$ and $\delta_i \in \{0, 1\}$. If $\delta_i = 0$ for all $i \in I$, then F is called a *lower face*.

Let $\Sigma := \otimes_{i \in [d]} \mathbb{F}^{M_i}$, so that $\Sigma^{\{0,1\}^k} \cong |\mathbb{F}|^{2^k \sum_{i \in [d]} |M_i|}$. We will consider vectors $v \in \Sigma^{\{0,1\}^k}$ indexed as $v(i, j, w)$ for $i \in [d], j \in [M_i], w \in \{0, 1\}^k$.

Definition 8.10 (Face vectors and parallelepiped constraints). Let $i_0 \in [d]$, $j_0 \in [M_{i_0}]$ and $F \subseteq \{0, 1\}^k$. Let $v[i_0, j_0, F] \in \Sigma^{\{0,1\}^k}$ be given by

8.2. Bias implies low rank exact computation

$v(i, j, \omega) = (-1)^{|\omega|}$ if $i = i_0, j = j_0$ and $\omega \in F$ and zero otherwise. This is called a face vector. If F is a lower face, then it corresponds to a lower face vector. If $\dim(F) \geq i_0 + 1$, then it is a relevant (lower) face vector. A vector $t \in \Sigma^{\{0,1\}^k}$ satisfies the parallelepiped constraints if it is orthogonal to all the relevant lower face vectors.

Let $\Sigma_0 \subseteq \Sigma^{\{0,1\}^k}$ be the subspace of vectors satisfying the parallelepiped constraints. Below we use the following shorthand: $\binom{k}{\leq i} := \sum_{j=0}^i \binom{k}{j}$.

Claim 8.11 (Dimension of Σ_0 (Green and Tao [43], Lemma 4.4)). Let $d < k$. Then,

$$\dim(\Sigma_0) = \sum_{i=1}^d M_i \binom{k}{\leq i}.$$

Lemma 8.12 (Equidistribution of parallelepipeds). Suppose Theorem 8.1 is true up to degree d . Given $s, d < k \in \mathbb{N}$, let \mathcal{B} be a polynomial factor of rank at least $r_{(8.12)}(k, s)$ defined by polynomials $Q_1, \dots, Q_c : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree at most d . Then for every $t \in \Sigma_0$ and every $x \in \mathbb{F}^n$ such that $\mathcal{B}(x) = t(0)$, it holds that

$$\Pr_{y \in (\mathbb{F}^n)^k} [\mathcal{B}(x + \omega \cdot y) = t(\omega) \ \forall \omega \in \{0, 1\}^k] = \frac{1}{|\mathbb{F}| \sum_{i=1}^d M_i \binom{k}{\leq i}} \pm \frac{1}{|\mathbb{F}|^s}.$$

Proof. This immediately follows from the dimension of Σ_0 (Claim 8.11) and Lemma 8.8 applied to the parallelepiped. \square

We are now ready to prove Theorem 8.1, but before doing so, let us explain the high-level approach. Given a biased degree- d polynomial P , we use Lemma 8.2 to approximate P by a function that is measurable in a degree- $(d - 1)$ polynomial factor \mathcal{B} . We then regularize this factor to a sufficiently regular polynomial factor \mathcal{B}' of degree $d - 1$. The final step of the proof is to show that P itself is measurable in \mathcal{B}' . To show this, we divide atoms of \mathcal{B}' into two categories: we call an atom A of \mathcal{B}' “good” if P is almost a constant over A , and otherwise we call it “bad”. We first argue that P is in fact constant in each good atom. Let A be a good atom and let c_A be P ’s most popular value over A . We argue that for every $x \in A$, there are directions y_1, \dots, y_{d+1} such that for any

nonempty $S \subseteq [d + 1]$, $P(x + \sum_{i \in S} y_i) = c_A$. As a result, observing that $D_{y_1, \dots, y_{d+1}} P(x) \equiv 0$, and all terms but $P(x)$ in $D_{y_1, \dots, y_{d+1}} P(x)$ are equal to c_A , we get that $P(x) = c_A$. If A is a bad atom, we first find a collection of good atoms A_w for $w \in \{0, 1\}^{d+1} \setminus \{0\}^{d+1}$, such that the evaluation vector of \mathcal{B}' over this collection of atoms satisfies the parallelepiped constraints. Let c_w be the constant value that P takes on the good atom A_w . Having found such a collection of atoms, given an arbitrary $x \in A$, we may invoke [Lemma 8.12](#) to show that there exists y_1, \dots, y_{d+1} such that for all nonempty $S \subseteq [d + 1]$, $x + \sum_{i \in S} y_i \in A_w$. Finally, similar to the good atom case, we use the $d + 1$ -th derivative of P to argue that $P(x)$ a fixed value determined by c_w for $w \in \{0, 1\}^{d+1} \setminus \{0\}^{d+1}$

Proof of Theorem 8.1. The base case of $d = 1$ is trivial. Indeed, if a linear polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ satisfies $|\mathbb{E}[e(P(x))]| \geq |\mathbb{F}|^{-s}$, then by orthogonality of linear polynomials, we have $P(x)$ is a constant and hence has rank 0.

So, suppose that the hypothesis is true for degrees up to $d - 1$, and we will prove it for d . Let $t \in \mathbb{N}$ depending on d be specified later. Recall that we assume that $|\mathbb{E}[e(P(x))]| \geq |\mathbb{F}|^{-s}$. By [Lemma 8.2](#), there exists a polynomial factor $\mathcal{B} = \{Q_1, \dots, Q_c\}$ of degree $d - 1$ where $c = c(d, s, t)$, and $\Gamma : \mathbb{F}^c \rightarrow \mathbb{F}$, such that

$$\Pr_{x \in \mathbb{F}^n} [P(x) \neq \Gamma(Q_1(x), \dots, Q_c(x))] \leq |\mathbb{F}|^{-t}.$$

Let $r : \mathbb{N} \rightarrow \mathbb{N}$ be a growth function that depends on d and will be specified later. Regularize \mathcal{B} to an r -regular polynomial factor $\mathcal{B}' = \{Q'_1, \dots, Q'_{c'}\}$, $c' \leq C_{r,d}^{(7.29)}(c)$. Thus, we have for an appropriate $\Gamma' : \mathbb{F}^{c'} \rightarrow \mathbb{F}$ that

$$\Pr_{x \in \mathbb{F}^n} [P(x) \neq \Gamma'(Q'_1(x), \dots, Q'_{c'}(x))] \leq |\mathbb{F}|^{-t}.$$

We will now prove that P is \mathcal{B}' -measurable, namely, P is constant on each atom of \mathcal{B}' . This will conclude the proof of [Theorem 8.1](#).

We will assume that $r(j) \geq r_{(8.1)}(d, 2t + j)$ for all $j \in \mathbb{N}$. Say that an atom A of \mathcal{B}' is *good* if

$$\Pr_{x \in A} [P(x) \neq \Gamma'(Q'_1(x), \dots, Q'_{c'}(x))] \leq |\mathbb{F}|^{-t/4}.$$

By Markov's inequality and Lemma 8.7, at least $1 - |\mathbb{F}|^{-t/4}$ fraction of the atoms A of \mathcal{B}' are good. The first step is to prove that on such atoms, P is constant. Fix a good atom A and let $A' \subseteq A$ be the set where $P(x) = \Gamma'(Q'_1(x), \dots, Q'_{c'}(x))$. Observe that $\Gamma'(Q'_1(x), \dots, Q'_{c'}(x))$ is constant on A , and hence $P(x) = c_A$ for some constant c_A for all $x \in A'$.

Lemma 8.13. Let t be large enough depending on d . Let $x \in A$ be arbitrary. Then there exists $h \in (\mathbb{F}^n)^{d+1}$ such that $x + \omega \cdot h \in A'$ for all $\omega \in \{0, 1\}^{d+1} \setminus 0^{d+1}$.

We omit the proof of Lemma 8.13, and note for the interested reader that it is identical to the proof of Lemma 5.2 of [43]. Continuing, since P is a degree d polynomial, we have

$$\sum_{\omega \in \{0,1\}^{d+1}} (-1)^{|\omega|} P(x + \omega \cdot h) = 0.$$

Now, Lemma 8.13 has guaranteed that $P(x + \omega \cdot h) = c_A$ for all $\omega \neq 0$. Plugging in c_A for all the terms except $P(x)$ in the above equation implies $P(x) = c_A$. Since x was an arbitrary point in A , we have proved that P is constant on the entire atom A . This finishes the first step.

We have shown for $1 - |\mathbb{F}|^{-t/4}$ fraction of the atoms A of \mathcal{B}' , i.e. the good atoms, that $P(x) = c_A$ for all $x \in A$. The final step shows that for any arbitrary atom A , there exist good atoms A_ω , for $\omega \in \{0, 1\}^{d+1} \setminus \{0\}^{d+1}$, such that the vector $t = \mathcal{B}'(A_\omega) \in \Sigma^{\{0,1\}^{d+1}}$ satisfies the parallelepiped constraints. To achieve this, it is enough to find one parallelepiped for which $x + \omega \cdot h$ lie in good atoms for all $\omega \neq 0$. Indeed, let $x \in A$ be arbitrary. Pick $h_1, \dots, h_{d+1} \in \mathbb{F}^n$ randomly. The probability that for a fixed $\omega \neq 0$, $x + \omega \cdot h$ lies in a good atom is at least $1 - |\mathbb{F}|^{-t/4} > 1 - 2^{-2d}$ for t large enough. The result now follows by a union bound over $\omega \in \{0, 1\}^{d+1}$. Let $t \in \Sigma^{\{0,1\}^{d+1}}$ be defined by this parallelepiped. By Lemma 8.12, there is a choice of r such that, for every $x \in A$,

$$\Pr_{y_1, \dots, y_k} \left[x + w \cdot y \in A_w \ \forall w \in \{0, 1\}^k \right] > 0.$$

Now as all A_w but A are chosen to be good atoms, the value of $P(x)$ for $x \in A$ is a fixed value determined by c_w for $w \in \{0, 1\}^{d+1} \setminus \{0\}^{d+1}$, concretely $P(x) = \sum_{w \in \{0,1\}^{d+1} \setminus \{0\}^{d+1}} (-1)^{|w|+1} c_w$. \square

9

Decomposition Theorems

“Decomposition theorems” [38, 78, 43] are important consequences of inverse theorems. They allow one to decompose an arbitrary function into a “structured” part and a “pseudorandom” part. The structured part is simple enough to be analyzed directly (often when assuming regularity and equidistribution as a result of regularization lemmas), and the pseudorandomness requirements are usually set up so that the pseudorandom part has little effect on the analysis and can often be ignored as small noise. We refer the interested reader to the papers by Gowers [38] and Reingold *et al.* [64] for a detailed discussion of decomposition theorems of this type and how the finite-dimensional Hahn-Banach theorem can be used to give short and transparent proofs of many results of these kinds.

9.1 Basic decomposition theorem

The following decomposition theorem is a direct consequence of the inverse theorem for Gowers norms, [Theorem 6.9](#). Recall that $\mathbb{D} = \{z \in \mathbb{C} : |z| \leq 1\}$ denotes the unit disk in the complex plane.

Theorem 9.1 (Basic decomposition theorem). Let \mathbb{F} be a prime finite field, $d \geq 1, \varepsilon > 0$ and let $r : \mathbb{N} \rightarrow \mathbb{N}$ be an arbitrary function. Every function $f : \mathbb{F}^n \rightarrow \mathbb{D}$ can be decomposed as

$$f = g + h,$$

where

- (i) $g = \mathbb{E}[f|\mathcal{B}]$, where \mathcal{B} is an r -regular polynomial factor of degree at most d and complexity $C \leq C_{9.1}(\mathbb{F}, d, r, \varepsilon)$.
- (ii) $\|h\|_{U^{d+1}} < \varepsilon$.

Furthermore, if we are given an initial polynomial factor \mathcal{B}_1 of degree at most d , then we may assume that \mathcal{B} is a refinement of \mathcal{B}_1 , and in which case we have that $C = |\mathcal{B}| \leq C_{9.1}(\mathbb{F}, d, r, \varepsilon, |\mathcal{B}_1|)$.

Proof. We define a sequence $\mathcal{B}_1, \mathcal{B}_2, \dots$ of r -regular polynomial factors of degree at most d , where if \mathcal{B}_1 is not given then we take $\mathcal{B}_1 = \emptyset$. Let $g_i = \mathbb{E}[f|\mathcal{B}_i]$ and $h_i = f - g_i$. We will show that for a bounded value of i we have $\|h_i\|_{U^{d+1}} < \varepsilon$ in which case we are done, as we can take $g = g_i, h = h_i$.

So, consider $i \geq 1$ with $\|h_i\|_{U^{d+1}} \geq \varepsilon$. By [Theorem 6.9](#), there exists a polynomial $P_i : \mathbb{F}^n \rightarrow \mathbb{T}$ of degree $\leq d$ such that $|\langle h_i, e(P_i) \rangle| \geq \delta$, where $\delta = \delta_{6.9}(\mathbb{F}, d, \varepsilon)$. In this case, let $\mathcal{B}'_i = \mathcal{B}_i \cup \{P_i\}$ and let \mathcal{B}_{i+1} be an r -regular factor refining \mathcal{B}'_i , as given by [Theorem 7.30](#). Note that $|\mathcal{B}_{i+1}|$ is bounded by a function of $|\mathcal{B}_i|, d, r$. Concretely, $|\mathcal{B}_{i+1}| \leq C_{7.30}^{\mathbb{F}, d, r} (|\mathcal{B}_i| + 1)$. In order to show that this process terminates after a bounded number of steps, we will show that $\|g_{i+1}\|_2^2 \geq \|g_i\|_2^2 + \delta^2$, and hence the process must terminate after at most $1/\delta^2$ steps.

To see that, first note that $\langle g_i, h_i \rangle = 0$, as the average of $h_i = f - \mathbb{E}[f|\mathcal{B}_i]$ in any atom of \mathcal{B}_i is zero, while g_i is \mathcal{B}_i -measurable, and hence is constant on any atom of \mathcal{B}_i . In particular, this implies that $\|g_i\|_2^2 + \|h_i\|_2^2 = \|f\|_2^2 \leq 1$. Next, let $g'_i = \mathbb{E}[f|\mathcal{B}'_i]$ and $h'_i = f - g'_i$. As \mathcal{B}'_i is a refinement of \mathcal{B}_i we also have that $\langle g_i, g'_i - g_i \rangle = 0$. Thus

$$\|g'_i\|_2^2 = \|g_i\|_2^2 + \|g'_i - g_i\|_2^2 = \|g_i\|_2^2 + \|h'_i - h_i\|_2^2.$$

Next, note that $|\langle h_i, e(P_i) \rangle| \geq \delta$ but $\langle h'_i, e(P_i) \rangle = 0$, since in any atom of \mathcal{B}'_i we have that P_i is constant and the average of h'_i is zero. Thus

$$\begin{aligned} \|h'_i - h_i\|_2^2 &= \mathbb{E}_x |h'_i(x) - h_i(x)|^2 \\ &= \mathbb{E}_x |(h'_i(x) - h_i(x))e(P_i(x))|^2 \\ &\geq |\mathbb{E}_x [(h'_i(x) - h_i(x))e(P_i(x))]|^2 \\ &\geq \delta^2. \end{aligned}$$

To conclude the proof, note that as \mathcal{B}_{i+1} is a refinement of \mathcal{B}'_i we have that $\|g_{i+1}\|_2^2 = \|g'_i\|_2^2 + \|g_{i+1} - g'_i\|_2^2 \geq \|g'_i\|_2^2$. \square

In Section 12.2 we will use a multifunction version of Theorem 9.1, namely decomposing a collection of functions using a single regular polynomial factor. This can be achieved by a proof almost identical to the above.

9.2 Higher-order Fourier expansion

Let $f : \mathbb{F}^n \rightarrow \mathbb{D}$ be a function, \mathcal{B} a polynomial factor and let $g = \mathbb{E}[f|\mathcal{B}]$. It is possible to express g as a function of the polynomials which define \mathcal{B} . Assume that \mathcal{B} is defined by polynomials $\{P_1, \dots, P_C\}$. Then as g is constant on the atoms of \mathcal{B} , we can express

$$g(x) = \Gamma(P_1(x), \dots, P_C(x)),$$

for some function $\Gamma : \mathbb{T}^c \rightarrow \mathbb{T}$. More concretely, assume that P_i has depth k_i , and hence takes values in \mathbb{U}_{k_i+1} . Let $G = \prod_{i=1}^C \mathbb{U}_{k_i+1}$ and let $G' = \prod_{i=1}^C \mathbb{Z}_p^{k_i+1}$ be the dual group. Applying standard Fourier decomposition to $\Gamma : G \rightarrow \mathbb{C}$ gives

$$\Gamma(z) = \sum_{\alpha \in G'} \hat{\Gamma}(\alpha) e(\langle \alpha, z \rangle).$$

As $\|g\|_\infty \leq \|f\|_\infty \leq 1$ we have $\|\Gamma\|_\infty \leq 1$. Hence by Parseval's identity, $\sum |\hat{\Gamma}(\alpha)|^2 \leq 1$. We can apply the same decomposition to g . Define polynomials $P_\alpha(x) = \sum \alpha_i P_i(x)$ for $\alpha \in G'$. Then the higher-order Fourier expansion of g is given by

$$g(x) = \sum_{\alpha \in G'} \hat{\Gamma}(\alpha) e(P_\alpha(x)).$$

Note that the polynomials P_α play the role of characters in standard Fourier analysis. In standard Fourier analysis, characters are orthogonal, which plays an important role. If we assume that the polynomial factor \mathcal{B} has a high enough rank, then the same approximately holds for higher-order Fourier expansions.

Claim 9.2 (claim). Assume that $\text{rank}(\mathcal{B}) \geq r_{9.2}(\mathbb{F}, d, \varepsilon, |\mathcal{B}|)$. Then

$$\left| \langle g, e(P_\alpha) \rangle - \hat{\Gamma}(\alpha) \right| \leq \varepsilon.$$

Proof. Assume that \mathcal{B} is r -regular for $r = r_{7.10}(\mathbb{F}, d, \varepsilon/|G|)$, where note that $|G| \leq p^{\lceil (d-1)/(p-1) \rceil |\mathcal{B}|}$. By the regularity assumption, $\text{bias}(P_{\alpha'}) \leq \varepsilon/|G|$ for all $\alpha' \neq 0$. Thus

$$\begin{aligned} \langle g, e(P_\alpha) \rangle &= \mathbb{E}_x [g(x)e(-P_\alpha(x))] \\ &= \sum_{\alpha'} \hat{\Gamma}(\alpha') \mathbb{E}_x [e(P_{\alpha'}(x) - P_\alpha(x))] \\ &= \sum_{\alpha'} \hat{\Gamma}(\alpha') \text{bias}(P_{\alpha'-\alpha}) \end{aligned}$$

and

$$\left| \langle g, e(P_\alpha) \rangle - \hat{\Gamma}(\alpha) \right| \leq \sum_{\alpha' \neq \alpha} |\hat{\Gamma}(\alpha')| |\text{bias}(P_{\alpha'-\alpha})| \leq \varepsilon.$$

□

Via a similar analysis one can obtain an approximate version of Plancherel’s theorem when the expansion has high rank.

9.3 Strong decomposition theorems

In many applications, once we decompose $f = g + h$ where $g = \mathbb{E}[f|\mathcal{B}]$ and $\|h\|_{U^{d+1}} < \varepsilon$, then it is necessary to make ε small in terms of the complexity of \mathcal{B} . It turns out that this is possible, if we allow another L_2 error term.

Theorem 9.3 (Strong decomposition theorem). Let \mathbb{F} be a prime finite field, $d \geq 1, \delta > 0$, and let $\varepsilon : \mathbb{N} \rightarrow \mathbb{R}^+$ and $r : \mathbb{N} \rightarrow \mathbb{N}$ be arbitrary functions. Any function $f : \mathbb{F}^n \rightarrow \mathbb{D}$ can be decomposed as

$$f = f_1 + f_2 + f_3,$$

where

(i) $f_1 = \mathbb{E}[f|\mathcal{B}]$, where \mathcal{B} is an r -regular polynomial factor of degree at most d and complexity $C \leq C_{9.1}(\mathbb{F}, d, r, \varepsilon, \delta)$.

(ii) $\|f_2\|_{U^{d+1}} < \varepsilon(|\mathcal{B}|)$ and $\|f_2\|_\infty \leq 2$.

(iii) $\|f_3\|_2 \leq \delta$.

Furthermore, if we are given an initial polynomial factor \mathcal{B}_1 of degree at most d , then we may assume that \mathcal{B} is a refinement of \mathcal{B}_1 , in which case we have that $C = |\mathcal{B}| \leq C_{9.1}(\mathbb{F}, d, r, \varepsilon, \delta, |\mathcal{B}_1|)$.

Proof. Define a sequence $\mathcal{B}_1, \mathcal{B}_2, \dots$ of polynomial factors as follows. The first factor \mathcal{B}_1 is given as input, and otherwise let $\mathcal{B}_1 = \emptyset$. To obtain \mathcal{B}_{i+1} , apply [Theorem 9.1](#) with initial factor \mathcal{B}_i and error $\varepsilon_i = \varepsilon(|\mathcal{B}_i|)$. Define as before $g_i = \mathbb{E}[f|\mathcal{B}_i]$ and $h_i = f - g_i$. Note that as \mathcal{B}_{i+1} is a refinement of \mathcal{B}_i we have that $\langle g_i, g_{i+1} - g_i \rangle = 0$ and hence $\|g_{i+1}\|_2^2 = \|g_i\|_2^2 + \|g_{i+1} - g_i\|_2^2$. Let $i \leq 1/\delta^2$ be minimal such that $\|g_{i+1}\|_2^2 \leq \|g_i\|_2^2 + \delta^2$. We then take

$$\mathcal{B} = \mathcal{B}_i, \quad f_1 = g_i, \quad f_2 = h_{i+1}, \quad f_3 = h_i - h_{i+1}.$$

Note that by definition, $f_1 = \mathbb{E}[f|\mathcal{B}]$ and $\|f_2\|_{U^{d+1}} \leq \varepsilon(|\mathcal{B}|)$. It is also simple to verify that $\|f_2\|_\infty \leq \|f\|_\infty + \|g_{i+1}\|_\infty \leq 2$. Finally by our construction we have that $\|f_3\|_2^2 = \|h_i - h_{i+1}\|_2^2 = \|g_i - g_{i+1}\|_2^2 = \|g_{i+1}\|_2^2 - \|g_i\|_2^2 \leq \delta^2$. \square

If we specialize [Theorem 9.4](#) to Boolean functions, we can have more control on the ranges of f_1, f_2, f_3 .

Theorem 9.4 (Strong decomposition theorem, Boolean functions). Let $f : \mathbb{F}^n \rightarrow \{0, 1\}$. Under the same conditions as that of [Theorem 9.3](#), we obtain the same decomposition $f = f_1 + f_2 + f_3$, which furthermore satisfies that f_1 and $f_1 + f_3$ take values in $[0, 1]$; and f_2 and f_3 take values in $[-1, 1]$.

Proof. We have $f_1 = g_i = \mathbb{E}[f|\mathcal{B}_i]$ and $f_1 + f_3 = g_{i+1} = \mathbb{E}[f|\mathcal{B}_{i+1}]$, which guarantees that they take values in $[0, 1]$. We have $f_2 = f - \mathbb{E}[f|\mathcal{B}_i]$ and $f_3 = \mathbb{E}[f|\mathcal{B}_i] - \mathbb{E}[f|\mathcal{B}_{i+1}]$, which guarantees that they take values in $[-1, 1]$. \square

Theorem 9.3 can also be extended to decompose multiple functions with a single polynomial factor, which is useful in certain applications. The proof is identical and is omitted.

Theorem 9.5 (Strong decomposition theorem, multiple functions). Let \mathbb{F} be a prime finite field, $d \geq 1, m \geq 1, \delta > 0$ and let $\varepsilon : \mathbb{N} \rightarrow \mathbb{R}^+$ and $r : \mathbb{N} \rightarrow \mathbb{N}$ be arbitrary functions. Any m functions $f^{(1)}, \dots, f^{(m)} : \mathbb{F}^n \rightarrow \mathbb{D}$ can be decomposed as

$$f^{(i)} = f_1^{(i)} + f_2^{(i)} + f_3^{(i)}$$

where

- $f_1^{(i)} = \mathbb{E}[f^{(i)}|\mathcal{B}]$, where \mathcal{B} is an r -regular polynomial factor of degree at most d and complexity $C \leq C_{9.5}(\mathbb{F}, d, m, r, \varepsilon, \delta)$.
- $\|f_2^{(i)}\|_{U^{d+1}} < \varepsilon(|\mathcal{B}|)$ and $\|f_2^{(i)}\|_\infty \leq 2$.
- $\|f_3^{(i)}\|_2 \leq \delta$.

Furthermore, if we are given an initial polynomial factor \mathcal{B}_1 of degree at most d , then we may assume that \mathcal{B} is a refinement of \mathcal{B}_1 , and in which case we have that $C = |\mathcal{B}| \leq C_{9.5}(\mathbb{F}, d, m, r, \varepsilon, \delta, |\mathcal{B}_1|)$.

9.4 Sub-atom selection

It turns out that the strong decomposition theorems are not sufficiently strong for certain applications in algebraic property testing. These require to control all the error terms, including the L_2 error term, by a function of the complexity of the polynomial factor. This is impossible if we have just one polynomial factor, but becomes possible when working in parallel with two factors: a polynomial factor and a refinement of it.

First, we need to define the notion of a polynomial factor representing another polynomial factor for a function. As all the applications of these tools apply to Boolean functions, we specialize the treatment below for Boolean functions, and note that it can be easily extended to bounded complex functions.

Definition 9.6 (Polynomial factor represents another factor). Let $f : \mathbb{F}^n \rightarrow \{0, 1\}$, \mathcal{B} a polynomial factor and \mathcal{B}' a refinement of it. For $0 < \zeta < 1$, we say that \mathcal{B}' ζ -represents \mathcal{B} with respect to f , if for at most ζ fraction of atoms c of \mathcal{B} , more than ζ fraction of the atoms c' of \mathcal{B}' lying inside c satisfy $|\mathbb{E}[f|c] - \mathbb{E}[f|c']| > \zeta$.

We now state the following “Two-level decomposition theorem” proved by Bhattacharyya *et al.* [15] (it is referred to as “super decomposition theorem” in [15]).

Theorem 9.7 (Two-level decomposition theorem (Bhattacharyya *et al.* [15], Theorem 4.9)). Let $d \geq 1, \zeta > 0$ and let $\varepsilon, \delta : \mathbb{N} \rightarrow \mathbb{R}^+$ and $r : \mathbb{N} \rightarrow \mathbb{N}$ be arbitrary functions. Given any function $f : \mathbb{F}^n \rightarrow \{0, 1\}$ there exists a polynomial factor \mathcal{B} of degree d , and a refinement of it \mathcal{B}' of degree d , both r -regular and of complexity at most $C_{9.7}(\mathbb{F}, d, r, \varepsilon, \delta, \zeta)$, such that the following holds. We can decompose

$$f = f_1 + f_2 + f_3$$

where

- (i) $f_1 = \mathbb{E}[f|\mathcal{B}']$.
- (ii) $\|f_2\|_{U^{d+1}} < \varepsilon(|\mathcal{B}'|)$.
- (iii) $\|f_3\|_2 \leq \delta(|\mathcal{B}|)$.
- (iv) f_1 and $f_1 + f_3$ take values in $[0, 1]$; and f_2, f_3 take values in $[-1, 1]$.
- (v) \mathcal{B}' ζ -represents \mathcal{B} with respect to f .

Proof. Define a sequence $\mathcal{B}_1, \mathcal{B}_2, \dots$ of polynomial factors as follows, where initially \mathcal{B}_1 is empty, and \mathcal{B}_{i+1} is obtained by applying [Theorem 9.4](#) with the parameters d, r, ε and $\delta_i = \delta(|\mathcal{B}_i|)$. If \mathcal{B}_{i+1} ζ -represents \mathcal{B}_i with respect to f , then the theorem follows with $\mathcal{B} = \mathcal{B}_i, \mathcal{B}' = \mathcal{B}_{i+1}$, and f_1, f_2, f_3 as given in the decomposition with \mathcal{B}_{i+1} . If not, then one can verify that

$$\mathbb{E}[f|\mathcal{B}_{i+1}]^2 \geq \mathbb{E}[f|\mathcal{B}_i]^2 + \zeta^3.$$

Hence, for some $i \leq 1/\zeta^3$, the condition will hold. □

Although the above two-level decomposition theorem may be useful by itself for some applications, certain algebraic property testing applications require the ability to choose a sub-atom in \mathcal{B}' inside each atom of \mathcal{B} , such that the error of f_3 on all sub-atoms is bounded, and most sub-atoms represent their atoms. Moreover, we would need this choice to be algebraically consistent.

To define this latter condition formally, assume that \mathcal{B}' is a syntactic refinement of \mathcal{B} , and thus is defined by adding new polynomials $Q_1, \dots, Q_{|\mathcal{B}'|-|\mathcal{B}|}$ to the polynomials defining \mathcal{B} . Thus, we can describe atoms of \mathcal{B}' as (c, s) , where $c \in \mathbb{T}^{|\mathcal{B}|}$ describes an atom of \mathcal{B} and $s \in \mathbb{T}^{|\mathcal{B}'|-|\mathcal{B}|}$. The choice of s allows to choose a sub-atom (c, s) of the finer factor within an atom c of the coarser partition.

We require that there exists a *fixed* $s \in \mathbb{T}^{|\mathcal{B}'|-|\mathcal{B}|}$, such that

- For all atoms c in \mathcal{B} , the L^2 error term f_3 within the corresponding sub-atom is small.
- For most atoms c in \mathcal{B} , the sub-atom (c, s) represents the atom c , in the sense that $\mathbb{E}[f|c] \approx \mathbb{E}[f|(c, s)]$.

The following theorem formalizes this. It is presented for one function, but can be easily extended to allow multiple functions, in the same way that [Theorem 9.5](#) extends [Theorem 9.3](#) to multiple functions.

Theorem 9.8 (Subatom selection (Bhattacharyya *et al.* [15], Theorem 4.12)). Let $f : \mathbb{F}^n \rightarrow \{0, 1\}$. Under the same conditions as in [Theorem 9.7](#), there exists a polynomial factor \mathcal{B} and a syntactic refinement of it \mathcal{B}' , whose atoms are indexed by (c, s) with $c \in \mathbb{T}^{|\mathcal{B}|}$, $s \in \mathbb{T}^{|\mathcal{B}'|-|\mathcal{B}|}$, such that in addition to the guarantees of [Theorem 9.7](#), there exists a choice of $s \in \mathbb{T}^{|\mathcal{B}'|-|\mathcal{B}|}$ for which the following is true:

- (vi) For every atom c of \mathcal{B} , the sub-atom (c, s) of \mathcal{B}' satisfies that

$$\mathbb{E} \left[|f_3(x)|^2 \mid \mathcal{B}'(x) = (c, s) \right] \leq \delta(|\mathcal{B}|)^2.$$

- (vii) For at most a ζ fraction of atoms c in \mathcal{B} it holds that

$$|\mathbb{E}[f|c] - \mathbb{E}[f|(c, s)]| > \zeta.$$

Proof sketch. The proof is very similar to the proof of [Theorem 9.7](#). The fact that \mathcal{B}' is a syntactic refinement of \mathcal{B} can be guaranteed by making sure that \mathcal{B} has high enough rank. Choose s uniformly among the possible set of values. Condition (vi) holds with high probability by a union bound on all the $|\mathcal{B}|$ possible atoms, by choosing $\delta(|\mathcal{B}|) \ll 1/|\mathcal{B}|$ small enough. Condition (vii) follows by condition (v) and Markov inequality. \square

10

Homogeneous Nonclassical Polynomials

The main difficulty in dealing with fields of low characteristic is that in the higher-order Fourier expansions, instead of the exponentials of classical polynomials, one has to work with exponentials of nonclassical polynomials. These do not share many of the convenient properties of classical polynomials. To overcome these difficulties, we develop a theory of nonclassical homogeneous polynomials which will enable us to confine to this simpler class of nonclassical polynomials. Recall that the need for homogeneous nonclassical polynomials arises in [Theorem 7.32](#), which characterizes the joint distribution of multiple high-rank polynomials evaluated in parallel on multiple linear forms.

A classical degree d polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ is called homogeneous if all of its monomials are of degree d . A simple consequence of homogeneity is that every homogeneous classical polynomial $P(x)$ satisfies $P(cx) = c^d P(x)$ for every $c \in \mathbb{F}$. It turns out that this consequence is the “right” way to generalize homogeneity for nonclassical polynomials.

Definition 10.1 (Homogeneity). A (nonclassical) polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ is called homogeneous if, for every $c \in \mathbb{F}$, there exists $\sigma_c \in \mathbb{Z}$ such that $P(cx) = \sigma_c P(x)$ for all $x \in \mathbb{F}^n$.

We start with a simple yet useful constraint on σ_c .

Claim 10.2. Let $P : \mathbb{F}^n \rightarrow \mathbb{T}$ be a degree d polynomial. Assume that $P(cx) = \sigma_c P(x)$ for some $\sigma_c \in \mathbb{Z}$. Then

$$\sigma_c = |c|^{\deg(P)} \pmod{p}.$$

Proof. Recall that ∂P is the derivative polynomial of P (see [Definition 7.20](#)). Define $Q(x) = P(cx) - \sigma_c P(x)$, where by assumption $Q = 0$. Then

$$0 = \partial Q(h_1, \dots, h_d) = \partial P(ch_1, \dots, ch_d) - \sigma_c \partial P(h_1, \dots, h_d).$$

By [Lemma 7.22](#) ∂P is a classical homogeneous polynomial of degree d which is linear in each of h_1, \dots, h_d . Thus $\partial P(ch_1, \dots, ch_d) = |c|^d \partial P(h_1, \dots, h_d)$. So we obtained that

$$0 = (|c|^d - \sigma_c) \partial P(h_1, \dots, h_d).$$

As ∂P is a nonzero classical polynomial this implies that $|c|^d - \sigma_c = 0 \pmod{p}$. □

Notice that for a polynomial P to be homogeneous, it suffices that for some generator ζ of F^* (the multiplicative group of the field \mathbb{F}) it holds that $P(\zeta x) = \sigma P(x)$. This is since any nonzero $c \in F$ can be represented as $c = \zeta^m$ for some $0 \leq m \leq p - 1$, and hence $P(cx) = \sigma^m P(x)$.

If P has depth k , then we can assume that $\sigma \in \{0, \dots, p^{k+1} - 1\}$ since $p^{k+1} P \equiv 0$. Obviously, the value of σ depends on the choice of ζ . However, the following lemma shows that for a fixed ζ , the value of σ is uniquely determined for all homogeneous polynomials of degree d and depth k . Henceforth, we will denote this unique value by $\sigma(d, k)$.

Let \mathbf{Z}_p denote the p -adic integers. We will show that there exists a choice of $\sigma(d) \in \mathbf{Z}_p$ such that $\sigma(d, k) = \sigma(d) \pmod{p^{k+1}}$ for all $k \geq 0$. This is related to the so-called Teichmüller characters (see e.g. Section 4.3 of [\[23\]](#)). However, in order to keep the presentation elementary, we avoid exploiting this connection.

Lemma 10.3. Fix a prime finite field $\mathbb{F} = \mathbb{F}_p$ and a generator $\zeta \in \mathbb{F}^*$. For every $d \geq 1, k \geq 0$ there exists a unique $\sigma = \sigma(d, k) \in \{0, \dots, p^{k+1} - 1\}$, such that for every homogeneous polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ of degree d and depth k , it holds that $P(\zeta x) = \sigma P(x)$. Furthermore, there exists $\sigma(d) \in \mathbf{Z}_p$ such that $\sigma(d, k) = \sigma(d) \pmod{p^{k+1}}$.

Proof. Let P be a homogeneous polynomial of degree d and depth k , and let $\sigma \in \{0, \dots, p^{k+1} - 1\}$ be such that $P(\zeta x) = \sigma P(x)$. By Claim 10.2 we know that $\sigma = |\zeta|^d \pmod p$. Observe that as $\zeta^{p-1} = 1$ it holds that $P(x) = P(\zeta^{p-1}x) = \sigma^{p-1}P(x)$, from which and the assumption that P has depth k it follows that $\sigma^{p-1} \equiv 1 \pmod{p^{k+1}}$.

We claim that $\sigma \in \{0, \dots, p^{k+1} - 1\}$ is uniquely determined by these two properties:

- (i) $\sigma = |\zeta|^d \pmod p$;
- (ii) $\sigma^{p-1} = 1 \pmod{p^{k+1}}$.

Suppose to the contrary that there exist two nonzero values $\sigma_1, \sigma_2 \in \mathbb{Z}_{p^{k+1}}$ that satisfy the above two properties, and choose $t \in \mathbb{Z}_{p^{k+1}}$ such that $\sigma_1 = t\sigma_2$. It follows from (i) that $t = 1 \pmod p$ and from (ii) that $t^{p-1} = 1$. We will show that $t = 1$ is the only possible such value in $\mathbb{Z}_{p^{k+1}}$.

Let $a_1, \dots, a_{p^k} \in \mathbb{Z}_{p^{k+1}}$ be all the possible solutions to $x = 1 \pmod p$ in $\mathbb{Z}_{p^{k+1}}$. Note that ta_1, \dots, ta_{p^k} is a permutation of the first sequence and thus

$$t^{p^k} \prod a_i = \prod a_i.$$

Consequently $t^{p^k} = 1 \pmod{p^{k+1}}$, which combined with $t^p = t$ implies $t = 1 \pmod{p^{k+1}}$.

For the last assertion, note that $\sigma(d, k)^{p-1} = 1 \pmod{p^{k+1}}$ implies $\sigma(d, k)^{p-1} = 1 \pmod{p^{\ell+1}}$ for every $\ell < k$. By the uniqueness of $\sigma(d, \ell)$, this implies that $\sigma(d, k) = \sigma(d, \ell) \pmod{p^{\ell+1}}$. We can thus take $\sigma(d) \in \mathbf{Z}_p$ given by the equations $\sigma(d) \pmod{p^{k+1}} = \sigma(d, k)$. □

Lemma 6.3 allows us to express every nonclassical polynomial as a linear span of monomials of the form $\frac{|x_1|^{d_1} \dots |x_n|^{d_n}}{p^{k+1}}$. Unfortunately, unlike in the classical case, these monomials are not necessarily homogeneous, and for some applications it is important to express a polynomial as a linear span of homogeneous polynomials. We show that this is possible as homogeneous nonclassical polynomials form a basis for the space of nonclassical polynomials.

10.1 A homogeneous basis for nonclassical polynomials

In this section we will prove that homogeneous polynomials span the space of all nonclassical polynomials.

Theorem 10.4. There is a basis for $\text{Poly}(\mathbb{F}^n \rightarrow \mathbb{T})$ consisting only of homogeneous multivariate polynomials.

This theorem allows us to make the extra assumption in decomposition theorems, that the resulting polynomial factor \mathcal{B} consists only of homogeneous polynomials. In order to achieve that, we decompose each polynomial of degree d and depth k as a sum of homogeneous polynomials of degree d and depths $1, \dots, k$. If one of these polynomials has low rank, we decompose it to a few lower degree polynomials, and repeat the process.

We start by proving the following simple observation.

Claim 10.5. Let $P : \mathbb{F} \rightarrow \mathbb{T}$ be a univariate polynomial of degree d . Then for every $c \in \mathbb{F} \setminus \{0\}$,

$$\deg(P(cx) - |c|^d P(x)) < d.$$

Proof. By [Lemma 6.3](#) it suffices to prove the claim for a monomial $m(x) = \frac{|x|^s}{p^{k+1}}$ with $k(p-1) + s = d$. Note that $m(cx) - |c|^d m(x)$ takes values in $\frac{1}{p^k} \mathbb{Z} / \mathbb{Z}$ as by Fermat's little theorem

$$|cx|^s - |c|^d |x|^s \equiv |x|^s |c|^s (1 - |c|^{k(p-1)}) \equiv 0 \pmod{p}.$$

It follows then from [Lemma 6.3](#) that $m(cx) - |c|^d m(x)$ is of depth at most $k-1$, and hence

$$\deg(m(cx) - |c|^d m(x)) \leq (p-1)(k-1) + (p-1) < d. \quad (10.1)$$

□

It is not difficult to show that the above claim holds also for any multivariate polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$. We provide a proof of this fact, although the univariate case suffices for our purposes in this section.

Claim 10.6. Let $P : \mathbb{F}^n \rightarrow \mathbb{T}$ be a multivariate polynomial of degree d . Then for every $c \in \mathbb{F} \setminus \{0\}$,

$$\deg \left(P(cx) - |c|^d P(x) \right) < d.$$

Proof. Notice that the claim is trivial for classical polynomials, since in this case, if R denotes the homogeneous degree- d part of P , then $R(cx) - |c|^d R(x) = 0$. We prove the statement for nonclassical polynomials. Let $Q(x) := P(cx)$, and note $\deg(Q) = d$. We will inspect the derivative polynomial of Q . Recall from [Definition 7.20](#) and [Lemma 7.22](#) that the derivative polynomial of Q ,

$$\partial Q(y_1, \dots, y_d) = D_{y_1} \cdots D_{y_d} Q(0),$$

is a degree- d classical homogeneous multi-linear polynomial which is invariant under permutations of (y_1, \dots, y_d) . In particular

$$\begin{aligned} |c|^{-d} \partial Q(y_1, \dots, y_d) &= \partial Q(c^{-1}y_1, \dots, c^{-1}y_d) \\ &= D_{c^{-1}y_1} D_{c^{-1}y_2} \cdots D_{c^{-1}y_d} Q(0) \\ &= \sum_{S \subseteq [d]} (-1)^{d-|S|} Q \left(c^{-1} \sum_{i \in S} y_i \right) \\ &= \sum_{S \subseteq [d]} (-1)^{d-|S|} P \left(\sum_{i \in S} y_i \right) \\ &= \partial P(y_1, \dots, y_d). \end{aligned}$$

This implies that $\partial(Q - |c|^d P) \equiv 0$ and thus $\deg(Q - |c|^d P) < d$. \square

10.1.1 A homogeneous basis, the univariate case

First, we prove [Theorem 10.4](#) for univariate polynomials.

Lemma 10.7. There is a basis of homogeneous univariate polynomials for $\text{Poly}(\mathbb{F} \rightarrow \mathbb{T})$.

Proof. We will prove by induction on d that there is a basis $\{h_1, \dots, h_d\}$ of homogeneous univariate polynomials for $\text{Poly}_{\leq d}(\mathbb{F} \rightarrow \mathbb{T})$ for every d . Let ζ be a fixed generator of \mathbb{F}^* . For any degree $d > 0$, we will build a degree- d homogeneous polynomial $h_d(x)$ such that $h_d(\zeta x) =$

$\sigma_d h_d(x)$ for some integer σ_d . The base case of $d \leq p - 1$ is trivial as $\text{Poly}_{\leq p-1}(\mathbb{F} \rightarrow \mathbb{T})$ consists of only classical polynomials, and those are spanned by $h_0(x) := \frac{1}{p}, h_1(x) := \frac{|x|}{p}, \dots, h_{p-1}(x) := \frac{|x|^{p-1}}{p}$. Now suppose that $d = s + (p - 1)(k - 1)$ with $0 < s \leq p - 1$, and $k > 1$. It suffices to show that the degree- d monomial $\frac{|x|^s}{p^k}$ can be expressed as a linear combination of homogeneous polynomials. Consider the function

$$f(x) := \frac{|\zeta x|^s}{p^k} - \frac{|\zeta|^d |x|^s}{p^k}.$$

Claim 10.5 implies that $\deg(f) < d$. Using the induction hypothesis, we can express $f(x)$ as a linear combination of $\frac{|x|^s}{p^\ell}$ for $\ell = 0, \dots, k - 1$, and h_e for $e < d$ with $e \not\equiv s \pmod{p - 1}$:

$$f(x) = \sum_{\ell=1}^{k-1} a_\ell \frac{|x|^s}{p^\ell} + \sum_{\substack{e < d, \\ e \not\equiv d \pmod{p-1}}} b_e h_e(x).$$

Set $A := |\zeta|^d + \sum_{\ell=1}^{k-1} a_\ell p^{k-\ell}$, so that

$$\frac{|\zeta x|^s}{p^k} - A \frac{|x|^s}{p^k} = \sum_{\substack{e < d, \\ e \not\equiv d \pmod{p-1}}} b_e h_e(x). \tag{10.2}$$

By the induction hypothesis, for $e < d$, $h_e(\zeta x) = \sigma_e h_e(x)$ where $\sigma_e = |\zeta|^e \pmod{p}$, and thus as $A = |\zeta|^d \pmod{p}$, we have $\sigma_e \not\equiv A \pmod{p}$ when $e \not\equiv d \pmod{p - 1}$. Consequently,

$$\begin{aligned} \sum_{\substack{e < d, \\ e \not\equiv d \pmod{p-1}}} b_e h_e(x) &= \sum_{\substack{e < d, \\ e \not\equiv d \pmod{p-1}}} \frac{b_e}{\sigma_e - A} (\sigma_e - A) h_e(x) \\ &= \sum_{\substack{e < d, \\ e \not\equiv d \pmod{p-1}}} \frac{b_e}{\sigma_e - A} (h_e(\zeta x) - A h_e(x)). \end{aligned}$$

Combining this with (10.2) we conclude that

$$h_d(x) := \frac{|x|^s}{p^k} - \sum_{\substack{e < d, \\ e \not\equiv d \pmod{p-1}}} \frac{b_e}{\sigma_e - A} h_e(x),$$

satisfies

$$h_d(\zeta x) = A h_d(x).$$

□

10.1.2 A homogeneous basis, the multivariate case

We are now ready to prove the main result of this section.

Theorem 10.4 (restated). *There is a basis for $\text{Poly}(\mathbb{F}^n \rightarrow \mathbb{T})$ consisting only of homogeneous multivariate polynomials.*

Proof. We will show by induction on the degree d and k , that every degree d monomial of depth k can be written as a linear combination of homogeneous polynomials. The base case of $d < p, k = 0$ is trivial as such monomials are classical and thus homogeneous themselves. Consider a nonclassical monomial $m(x_1, \dots, x_n) = \frac{|x_1|^{s_1} \dots |x_n|^{s_n}}{p^{k+1}}$ of degree $d = s_1 + \dots + s_n + (p-1)k$. For every $i \in [n]$ let $g_i(x_i) := h_{s_i+(p-1)k}(x_i)$ where $h_{s_i+(p-1)k}$ is the homogeneous univariate polynomial from Lemma 10.7.

Every g_i takes values in $\frac{1}{p^{k+1}}\mathbb{Z}/\mathbb{Z}$, and thus corresponds to a polynomial $G_i : \mathbb{F} \rightarrow \mathbb{Z}_{p^{k+1}}$. That is,

$$g_i(x) = \frac{G_i(x)}{p^{k+1}} \pmod 1.$$

Define $H : \mathbb{F}^n \rightarrow \mathbb{Z}_{p^{k+1}}$ as

$$H(x_1, \dots, x_n) := G_1(x_1) \cdots G_n(x_n),$$

and $h : \mathbb{F}^n \rightarrow \mathbb{T}$ as

$$h(x_1, \dots, x_n) := \frac{H(x_1, \dots, x_n)}{p^{k+1}}.$$

We claim that

- (i) h is a homogeneous polynomial.
- (ii) $\text{deg}(h) = d$.
- (iii) $\text{deg}(h - m) \leq d - 1$.

Thus by induction m can be written as a sum of homogeneous polynomials. To conclude the proof we verify (i)-(iii).

To verify (i) note that as g_i is a homogeneous univariate polynomial it holds that $g_i(\zeta x_i) = \sigma_i g_i(x_i) \pmod 1$ for some $\sigma_i \in \mathbb{Z}$. Thus $G_i(\zeta x_i) = \sigma_i G_i(x_i) \pmod{p^{k+1}}$. This implies that $H(\zeta x) = (\prod \sigma_i) H(x) \pmod{p^{k+1}}$,

and hence $h(\zeta x) = (\prod \sigma_i)h(x) \pmod{1}$. To verify (ii) and (iii) note that by construction, $G_i(x_i) = x_i^{s_i} + P_i(x_i) + pQ_i(x_i)$ where $\deg(P_i) < s_i$. One can then verify that $H(x)$ contains the monomial $\prod x_i^{s_i}$, and all other monomials are of the form $p^e \prod x_i^{t_i}$, where $\sum t_i \leq d - 1 + (p - 1)e$. Thus $h(x)$ contains a single monomial of degree d , which is $m(x)$, and all the remaining monomials have degree $\leq d - 1$. Thus $\deg(h - m) \leq d - 1$. \square

11

Complexity of Systems of Linear Forms

Gowers norms control the density of linear patterns in subsets of an Abelian group. For example, given a linear pattern, for a sufficiently large d , if the characteristic function of two sets are close in the U^d norm, then they contain almost the same number of copies of this pattern. For example, if $A, B \subset \mathbb{F}^n$ are two subsets whose indicator functions $1_A, 1_B : \mathbb{F}^n \rightarrow \{0, 1\}$ satisfy that $\|1_A - 1_B\|_{U^d}$ is negligible, then the number of $(d+1)$ -term arithmetic progressions $x, x+y, x+2y, \dots, x+dy$ that fall in A is approximately the same as the number that falls in B (see [Corollary 11.3](#) below).

More generally, in order to study the density of linear patterns in a set, one can use a decomposition theorem to first decompose the function into a structured part and a pseudorandom part, and then use this property of Gowers norms to discard the pseudorandom part and restrict the analysis to the simpler structured part. Since for smaller values of d , decomposition theorems provide a simpler structured part, a natural question arises:

Given a linear pattern, what is the smallest value of d for which the above statements hold?

Investigating such questions lead to various notions of complexity associated with a collection of linear forms. In this chapter we study such notions.

11.1 Cauchy-Schwarz complexity

Let A be a subset of \mathbb{F}^n with the indicator function $\mathbf{1}_A : \mathbb{F}^n \rightarrow \{0, 1\}$. Let L_1, \dots, L_m be linear forms over \mathbb{F} in ℓ variables. Let $X \in (\mathbb{F}^n)^\ell$ be chosen uniformly at random. The probability that $L_1(X), \dots, L_m(X)$ all fall in A , can be expressed as

$$\mathbb{E}_X [\mathbf{1}_A(L_1(X)) \cdots \mathbf{1}_A(L_m(X))].$$

Roughly speaking, we say $A \subseteq \mathbb{F}^n$ is *pseudorandom* with regards to $\mathcal{L} = (L_1, \dots, L_m)$ if

$$\mathbb{E}_X \left[\prod_{i=1}^m \mathbf{1}_A(L_i(X)) \right] \approx \left(\frac{|A|}{p^n} \right)^m;$$

That is if the probability that all $L_1(X), \dots, L_m(X)$ fall in A is close to what we would expect if A was a random subset of \mathbb{F}^n of cardinality $|A|$. Let $\alpha := |A|/|\mathbb{F}^n|$ be the density of A , and define $f := \mathbf{1}_A - \alpha$. We have

$$\begin{aligned} \mathbb{E}_X \left[\prod_{i=1}^m \mathbf{1}_A(L_i(X)) \right] &= \mathbb{E}_X \left[\prod_{i=1}^m (\alpha + f(L_i(X))) \right] \\ &= \alpha^m + \sum_{S \subseteq [m], S \neq \emptyset} \alpha^{m-|S|} \mathbb{E}_X \left[\prod_{i \in S} f(L_i(X)) \right]. \end{aligned}$$

Therefore, a sufficient condition for A to be pseudorandom with regards to \mathcal{L} is that $\mathbb{E}_X [\prod_{i \in S} f(L_i(X))]$ is negligible for all nonempty subsets $S \subseteq [m]$. Green and Tao [45] showed that a sufficient condition for this to occur is that $\|f\|_{U^{s+1}}$ is small enough, where s is the *Cauchy-Schwarz complexity* of the system of linear forms, defined below.

Definition 11.1 (Cauchy-Schwarz complexity [45]). Let $\mathcal{L} = \{L_1, \dots, L_m\}$ be a system of linear forms over a field \mathbb{F} in ℓ variables. The *Cauchy-Schwarz complexity* of \mathcal{L} is the minimal s such that the following holds.

For every $1 \leq i \leq m$, we can partition $\{L_j\}_{j \in [m] \setminus \{i\}}$ into $s + 1$ subsets, such that L_i does not belong to the linear span of any of the subsets.

The reason for the term *Cauchy-Schwarz complexity* is the following lemma due to Green and Tao [45] whose proof is based on carefully chosen iterative applications of the Cauchy-Schwarz inequality.

Lemma 11.2 (Green and Tao [45], see also Gowers and Wolf [39], Theorem 2.3). Let $f_1, \dots, f_m : \mathbb{F}^n \rightarrow \mathbb{D}$. Let $\mathcal{L} = \{L_1, \dots, L_m\}$ be a system of linear forms in ℓ variables of Cauchy-Schwarz complexity s . Then

$$\left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[\prod_{i=1}^m f_i(L_i(X)) \right] \right| \leq \min_{1 \leq i \leq m} \|f_i\|_{U^{s+1}}.$$

Note that the Cauchy-Schwarz complexity of any system of m linear forms in which every two linear forms are linearly independent (i.e. one is not a multiple of the other) is at most $m - 2$, since we can always partition $\{L_j\}_{j \in [m] \setminus \{i\}}$ into the $m - 1$ singleton subsets.

As an example, we prove the statement made in the beginning of the chapter regarding arithmetic progressions.

Corollary 11.3. Fix a prime finite field \mathbb{F} , $d \geq 2, \varepsilon > 0$. Let $A \subset \mathbb{F}^n$ of density $\alpha = |A|/|\mathbb{F}|^n$. Assume that $\|1_A - \alpha\|_{U^d} \leq \varepsilon$. Then the number of $(d + 1)$ -term arithmetic progressions in A is

$$\left| \Pr_{x,y \in \mathbb{F}^n} [x, x + y, \dots, x + dy \in A] - \alpha^{d+1} \right| \leq \varepsilon(1 + \alpha)^{d+1}.$$

Proof. Let $f = 1_A - \alpha$. Consider the system of $m = d + 1$ linear forms $L_i(x, y) = x + iy$ for $i = 0, \dots, d$. Its Cauchy-Schwarz complexity is $s = d - 1$. Lemma 11.2 gives that for every nonempty $S \subseteq [d + 1]$,

$$\mathbb{E}_{x,y \in \mathbb{F}^n} \left[\prod_{i \in S} f(x + iy) \right] \leq \|f\|_{U^{s+1}} \leq \varepsilon.$$

Thus,

$$\begin{aligned} & \Pr_{x,y \in \mathbb{F}^n} [x, x + y, \dots, x + dy \in A] \\ &= \mathbb{E}_{x,y \in \mathbb{F}^n} \left[\prod_{i=0}^d 1_A(L_i(x, y)) \right] \\ &= \alpha^{d+1} + \sum_{S \subseteq [d+1], S \neq \emptyset} \alpha^{d+1-|S|} \mathbb{E}_{x,y \in \mathbb{F}^n} \left[\prod_{i \in S} f(x + iy) \right]. \end{aligned}$$

And,

$$\begin{aligned} & \left| \Pr_{x,y \in \mathbb{F}^n} [x, x + y, \dots, x + dy \in A] - \alpha^{d+1} \right| \\ & \leq \varepsilon \sum_{S \subseteq [d+1], S \neq \emptyset} \alpha^{d+1-|S|} \leq \varepsilon(1 + \alpha)^{d+1}. \end{aligned}$$

□

11.2 The True Complexity

The Cauchy-Schwarz complexity of \mathcal{L} gives an upper bound on s , such that if $\|f\|_{U^{s+1}}$ is small enough for some function $f : \mathbb{F}^n \rightarrow \mathbb{D}$, then f is pseudorandom with regards to \mathcal{L} . Gowers and Wolf [39] defined the *true complexity* of a system of linear forms as the minimal s such that the above condition holds for all $f : \mathbb{F}^n \rightarrow \mathbb{D}$.

Definition 11.4 (True complexity [39]). Let $\mathcal{L} = \{L_1, \dots, L_m\}$ be a system of linear forms over \mathbb{F} in ℓ variables. The true complexity of \mathcal{L} is the smallest $d \in \mathbb{N}$ with the following property. For every $\varepsilon > 0$ and $\alpha \geq 0$, there exists $\delta > 0$ such that if $f : \mathbb{F}^n \rightarrow \mathbb{D}$ is any function with $\|f - \alpha\|_{U^{d+1}} \leq \delta$, then

$$\left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[\prod_{i=1}^m f(L_i(X)) \right] - \alpha^m \right| \leq \varepsilon.$$

An obvious upper bound on the true complexity is the Cauchy-Schwarz complexity of the system. However, there are cases where this is not tight. Gowers and Wolf conjectured that the true complexity of a system of linear forms can be characterized by a simple linear

algebraic condition: the smallest $d \geq 1$ such that $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent, where the d -th tensor power of a linear form $L = (\lambda_1, \dots, \lambda_\ell)$ is defined as

$$L^d = \left(\prod_{j=1}^d \lambda_{i_j} : i_1, \dots, i_d \in [\ell] \right) \in \mathbb{F}^{\ell^d}.$$

Remark 11.5. Note that if $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent, then for every $e \geq d$, the vectors $L_1^{e+1}, \dots, L_m^{e+1}$ are also linearly independent. Thus, for a system of linear forms L_1, \dots, L_m , it is natural to consider the smallest d for which $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent.

Conjecture 11.6 (True complexity characterization [39]. Resolved below in [Theorem 11.8](#)). The true complexity of a system of linear forms $\mathcal{L} = \{L_1, \dots, L_m\}$ is the smallest d such that $\{L_1^{d+1}, \dots, L_m^{d+1}\}$ are linearly independent.

Example 11.7. Consider the collection of linear forms $L_1 = (1, 0, 0)$, $L_2 = (1, 1, 0)$, $L_3 = (1, 0, 1)$, $L_4 = (1, 1, 1)$, $L_5 = (1, 1, -1)$, $L_6 = (1, -1, 1)$. It is easy to check that here the Cauchy-Schwarz complexity is 2, while as observed by Gowers and Wolf, the true complexity of this system of linear forms is 1.

Gowers and Wolf [40] verified their conjecture in the case where $|\mathbb{F}|$ is sufficiently large; more precisely when $|\mathbb{F}|$ is at least the Cauchy-Schwarz complexity of the system of linear form. [Conjecture 11.6](#) in its full generality was settled by Hatami *et al.* [47] through the following theorem.

Theorem 11.8 (Hatami *et al.* [47]). Let $\mathcal{L} = \{L_1, \dots, L_m\}$ be a system of linear forms over \mathbb{F} in ℓ variables. Assume that $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent. Then for every $\varepsilon > 0$, there exists $\delta > 0$ such that for any collection of functions $f_1, \dots, f_m : \mathbb{F}^n \rightarrow \mathbb{D}$ with $\min_{i \in [m]} \|f_i\|_{U^{d+1}} \leq \delta$, we have

$$\left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[\prod_{i=1}^m f_i(L_i(X)) \right] \right| \leq \varepsilon.$$

We defer the proof of [Theorem 11.8](#) to [Section 12.2](#). [Remark 11.5](#) and [Theorem 11.8](#) imply that the smallest d for which $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent equals the true complexity of the system.

To see the usefulness of [Theorem 11.8](#), let $\mathcal{L} = \{L_1, \dots, L_m\}$ be a system of linear forms of true complexity d . Assume that we are given bounded functions f_1, \dots, f_m . A version of [Theorem 9.1](#) for multiple functions (a simpler version of [Theorem 9.5](#)), allows us to find a high rank polynomial factor \mathcal{B} of degree d , such that we can decompose $f_i = g_i + h_i$ with $g_i = \mathbb{E}[f|\mathcal{B}]$ and $\|h_i\|_{U^{d+1}} \leq \delta$. The following corollary to [Theorem 11.8](#) shows that we may simply replace f_i with g_i when counting linear patterns. In particular, choosing $f_1 = \dots = f_m = f$ and $g_1 = \dots = g_m = \alpha$ proves the Gowers-Wolf conjecture ([Conjecture 11.6](#)) in its full generality.

Corollary 11.9. Let $\mathcal{L} = \{L_1, \dots, L_m\}$ be a system of linear forms over \mathbb{F} in ℓ variables. Assume that $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent. Then for every $\varepsilon > 0$, there exists $\delta > 0$ such that for any functions $f_1, \dots, f_m, g_1, \dots, g_m : \mathbb{F}^n \rightarrow \mathbb{D}$ with $\|f_i - g_i\|_{U^{d+1}} \leq \delta$, we have

$$\left| \mathbb{E}_X \left[\prod_{i=1}^m f_i(L_i(X)) \right] - \mathbb{E}_X \left[\prod_{i=1}^m g_i(L_i(X)) \right] \right| \leq \varepsilon.$$

Proof. Choosing $\delta = \delta(\varepsilon')$ as in [Theorem 11.8](#) for $\varepsilon' := \varepsilon/m$, we have

$$\begin{aligned} & \left| \mathbb{E}_X \left[\prod_{i=1}^m f_i(L_i(X)) \right] - \mathbb{E}_X \left[\prod_{i=1}^m g_i(L_i(X)) \right] \right| \\ &= \left| \sum_{i=1}^m \mathbb{E}_X \left[(f_i - g_i)(L_i(X)) \cdot \prod_{j=1}^{i-1} g_j(L_j(X)) \cdot \prod_{j=i+1}^m f_j(L_j(X)) \right] \right| \\ &\leq \sum_{i=1}^m \left| \mathbb{E}_X \left[(f_i - g_i)(L_i(X)) \cdot \prod_{j=1}^{i-1} g_j(L_j(X)) \cdot \prod_{j=i+1}^m f_j(L_j(X)) \right] \right| \\ &\leq m \cdot \varepsilon' \leq \varepsilon, \end{aligned}$$

where the second inequality follows from [Theorem 11.8](#) since $\|f_i - g_i\|_{U^{d+1}} \leq \delta$. □

12

Deferred Technical Proofs

In this chapter we prove [Theorem 7.32](#) and [Theorem 11.8](#). The proofs are somewhat technical, but they demonstrate the power of the techniques that we have developed so far.

12.1 Near-orthogonality: Proof of [Theorem 7.32](#)

Theorem 7.32 (restated). *Let \mathbb{F} be a prime field, $d \geq 1$, $\varepsilon > 0$. Let $\{L_1, \dots, L_m\}$ be a system of linear forms, and $\mathcal{B} = \{P_1, \dots, P_C\}$ be a polynomial factor of degree at most d and $\text{rank}(\mathcal{B}) > r_{7.32}(\mathbb{F}, d, \varepsilon)$. Assume that each P_i is a nonclassical homogeneous polynomial. For every set of coefficients $\Lambda = \{\lambda_{i,j} \in \mathbb{Z} : i \in [C], j \in [m]\}$, define*

$$P_\Lambda(x) := \sum_{i=1}^C \sum_{j=1}^m \lambda_{i,j} P_i(L_j(x)).$$

Then, one of the following two cases holds:

- $P_\Lambda \equiv 0$. In this case, for every $i \in [C]$, and any nonclassical homogeneous polynomials $Q_i : \mathbb{F}^n \rightarrow \mathbb{T}$ with $\deg(Q_i) = \deg(P_i)$ and $\text{depth}(Q_i) \leq \text{depth}(P_i)$, we have $\sum_{j \in [m]} \lambda_{i,j} Q_i(L_j(\cdot)) \equiv 0$.
- $P_\Lambda \not\equiv 0$. In this case, $|\mathbb{E}[e(P_\Lambda)]| < \varepsilon$.

We present the proof given by Hatami *et al.* [47]. By a linear form we will always mean $L \in \mathbb{F}^\ell$, i.e. a linear form over a prime field $\mathbb{F} = \mathbb{F}_p$ in ℓ variables. For a linear form $L = (\lambda_1, \dots, \lambda_\ell)$ define $|L| = \sum_{i=1}^\ell |\lambda_i|$.

Claim 12.1. Let $d > 0$ be an integer, and $L = (\lambda_1, \dots, \lambda_\ell) \in \mathbb{F}^\ell$ be a linear form. Then there exist linear forms $L_i = (\lambda_{i,1}, \dots, \lambda_{i,\ell}) \in \mathbb{F}^\ell$ for $i = 1, \dots, m$, and integer coefficients $a_1, \dots, a_m \in \mathbb{Z}$ with $m \leq |\mathbb{F}|^\ell$ such that

- $P(L(X)) = \sum_{i=1}^m a_i P(L_i(X))$ for every degree- d polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$;
- $|L_i| \leq d$ for every $i \in [m]$;
- $|\lambda_{i,j}| \leq |\lambda_j|$ for every $i \in [m]$ and $j \in [\ell]$.

Proof. The proof proceeds by simplifying $P(L(X))$ using identities that are valid for every polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ of degree d .

We prove the statement by induction on $|L|$. For the base case $|L| \leq d$ there is nothing to prove. Consider $|L| > d$. Since P is of degree d then deriving it $d + 1$ times yields the zero polynomial. As $|L| > d$, the same holds if we derive it $|L|$ times. So, for every choice of $y_1, \dots, y_{|L|} \in \mathbb{F}^n$, we have

$$\sum_{S \subseteq [|L|]} (-1)^{|L|-|S|} P\left(\sum_{i \in S} y_i\right) = (D_{y_1} \dots D_{y_{|L|}} P)(0) \equiv 0. \tag{12.1}$$

Let $X = (x_1, \dots, x_\ell) \in (\mathbb{F}^n)^\ell$. Setting $|\lambda_i|$ of the vectors $y_1, \dots, y_{|L|}$ to be equal to x_i for every $i \in [\ell]$, Equation (12.1) gives linear forms $M_i = (\tau_{i,1}, \dots, \tau_{i,\ell})$ for which

$$P(L(X)) = \sum_i \alpha_i P(M_i(X)), \tag{12.2}$$

12.1. Near-orthogonality: Proof of Theorem 7.32

for all i , $|M_i| \leq |L| - 1$, and for every $j \in [\ell]$, $|\tau_{i,j}| \leq |\lambda_j|$. Since $|M_i| \leq |L| - 1$, we can apply the induction hypothesis to obtain the identities

$$P(M_i(X)) = \sum_{j=1}^{m_j} a_{ij} P(M_{ij}(X))$$

where the linear forms $M_{ij} = (\rho_{ij1}, \dots, \rho_{ij\ell})$ satisfy $|M_{ij}| \leq d$, and $|\rho_{ijk}| \leq |\tau_{ij}| \leq |\lambda_j|$ for every i, j, k . Substituting these identities in Equation (12.2) yields the desired result. \square

The next claim shows that we can further simplify the expression given in Claim 12.1. Let $\mathcal{L}_d \subseteq \mathbb{F}^\ell$ denote the set of nonzero linear forms L with $|L| \leq d$ and with the first (left-most) nonzero coefficient equal to 1, e.g. $(0, 1, 0, 2) \in \mathcal{L}_3$ but $(2, 1, 0, 0) \notin \mathcal{L}_3$.

Claim 12.2. For any linear form $L \in \mathbb{F}^\ell$ and integer $d > 0$, there exists a collection of integer coefficients $\{a_{M,c} \in \mathbb{Z}\}_{M \in \mathcal{L}_d, c \in \mathbb{F}^*}$ such that for every degree- d polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$,

$$P(L(X)) = \sum_{M \in \mathcal{L}_d, c \in \mathbb{F}^*} a_{M,c} P(cM(X)). \tag{12.3}$$

Proof. Similar to the proof of Claim 12.1 we simplify $P(L(X))$ using identities that are valid for every polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ of degree d .

We use induction on the number of nonzero entries of L . The case when L has only one nonzero entry is trivial. For the induction step, choose $c \in \mathbb{F}$ so that the leading nonzero coefficient of $L' = c \cdot L$ is equal to 1. Assume that $L' = (\lambda_1, \dots, \lambda_\ell)$. If $|L'| \leq d$ we are done. Assume otherwise that $|L'| > d$. Applying Claim 12.1 for the degree- d polynomial $R(x) := P(c^{-1}x)$ and the linear form L' we can write

$$\begin{aligned} P(L(X)) &= P(c^{-1}L'(X)) = R(L'(X)) \\ &= \sum_i \beta_i R(M_i(X)) = \sum_i \beta_i P(c^{-1}M_i(X)), \end{aligned} \tag{12.4}$$

where for every i , $\beta_i \in \mathbb{Z}$ and $M_i = (\lambda_{i,1}, \dots, \lambda_{i,\ell})$ satisfies $|M_i| \leq d$; and $|\lambda_{i,j}| \leq |\lambda_j|$ for every $j \in [\ell]$. Let $\mathcal{I} = \{i : M_i \in \mathcal{L}_d\}$. Then

$$P(L(x)) = \sum_{i \in \mathcal{I}} \alpha_i P(c^{-1}M_i(X)) + \sum_{i \notin \mathcal{I}} \alpha_i P(c^{-1}M_i(X)). \tag{12.5}$$

In order to conclude the proof, we need to handle the sum over $i \notin \mathcal{I}$. Observe that if $i \notin \mathcal{I}$, then as the leading coefficient of L' is 1, then [Claim 12.1](#) implies in particular that M_i has smaller support than L . Thus, we may apply our induction hypothesis for $c^{-1}M_i$ for all $i \notin \mathcal{I}$, which concludes the proof. \square

[Claim 12.2](#) applies to all polynomials of degree d . If we also specify the depth, then we can obtain a stronger statement. However, in this statement we shall need to assume that the polynomials are homogeneous.

Claim 12.3. Let $d \geq 1, k \geq 0, L_1, \dots, L_m \in \mathbb{F}^\ell$ be linear forms, and let $\{\lambda_i \in \mathbb{Z}\}_{i \in [m]}$ be integer coefficients. Then there exist integer coefficients $\{a_M \in \mathbb{Z}\}_{M \in \mathcal{L}_d}$ such that the following is true. For every homogeneous polynomial $P : \mathbb{F}^n \rightarrow \mathbb{T}$ of degree d and depth $\leq k$,

- $\sum_{i=1}^m \lambda_i P(L_i(X)) \equiv \sum_{M \in \mathcal{L}_d} a_M P(M(X));$
- For every M with $a_M \neq 0$, we have $|M| \leq \deg(a_M P)$.

Proof. The proof is similar to that of [Claim 12.2](#), except that now we repeatedly apply [Claim 12.2](#) to every term of the form $\lambda P(L(X))$ to express it as an integer linear combination of $P(cM(X))$ for $M \in \mathcal{L}_{\deg(\lambda P)}$ and $c \in \mathbb{F}^*$. We use the assumption that P is homogeneous of degree d to replace $P(cM(X))$ with $\sigma_c P(M(X))$, where if $c = \zeta^i$ for the fixed generator $\zeta \in \mathbb{F}^*$ then $\sigma_c = \sigma(d, k)^i$. Observe that the condition $a_M P \neq 0$ depends only on the depth of P (that is, if it is zero for some polynomial P of depth k , then it is also zero for all polynomials of depth $\leq k$). Thus, the above process depends only on the assumption that P is homogeneous, its degree d , and a bound on its depth k . By repeating this procedure we arrive at the desired expansion. \square

We are now ready for the proof of [Theorem 7.32](#). For a linear form $L = (\lambda_1, \dots, \lambda_\ell)$, let $\text{lc}(L)$ denote the index of its first nonzero entry, namely $\text{lc}(L) := \min_{i: \lambda_i \neq 0} i$.

Proof of Theorem 7.32. Let d be the degree of the factor. For every $i \in [C]$, by Claim 12.3 we have

$$\sum_{j=1}^m \lambda_{i,j} P_i(L_j(X)) = \sum_{M \in \mathcal{L}_d} \lambda'_{i,M} P_i(M(X)) \tag{12.6}$$

for some integers $\lambda'_{i,M}$ such that if $\lambda'_{i,M} \neq 0$ then $|M| \leq \deg(\lambda'_{i,M} P_i)$. The simplifications of Claim 12.3 depend only on the degrees and depths of the polynomials. Hence, if $\lambda'_{i,M} P_i \equiv 0$ for all $M \in \mathcal{L}_d$, then $(\lambda_{i,1}, \dots, \lambda_{i,m})$ is in the (d_i, k'_i) -dependency set for all $k'_i \leq k_i$ (see Definition 7.34 for the definition). This implies that $\sum_{j=1}^m \lambda_{i,j} Q_i(L_j(X))$ for all homogeneous polynomials Q_i of degree d_i and depth $\leq k_i$.

So to prove the theorem, it suffices to show that P_Λ has small bias if $\lambda'_{i,M} P_i \not\equiv 0$ for some $i \in [C]$ and $M \in \mathcal{L}_d$. Suppose this is true, and thus there exists a nonempty set $\mathcal{M} \subseteq \mathcal{L}_d$ such that

$$P_\Lambda(X) = \sum_{i \in [C], M \in \mathcal{M}} \lambda'_{i,M} P_i(M(X)),$$

and for every $M \in \mathcal{M}$, there is at least one index $i \in [C]$ for which $\lambda'_{i,M} P_i \not\equiv 0$. Choose $i^* \in [C]$ and $M^* \in \mathcal{M}$ in the following manner.

- First, let $M^* \in \mathcal{M}$ be such that $\text{lc}(M^*) = \min_{M \in \mathcal{M}} \text{lc}(M)$, and among these, $|M^*|$ is maximal.
- Then, let $i^* \in [C]$ be such that $\deg(\lambda'_{i^*,M^*} P_{i^*})$ is maximized.

Without loss of generality assume that $i^* = 1$, $\text{lc}(M^*) = 1$, and let $d^* := \deg(\lambda'_{1,M^*} P_1) \leq d$. We claim that if $\sum_{j \in [m]} \lambda_{1,j} P_1(L_j(X))$ is not the zero polynomial, then $\deg(P_\Lambda) \geq d^*$, and moreover, P_Λ has small bias. We prove this by deriving P_Λ in specific directions in a manner that all the terms but $\lambda'_{1,M^*} P_1(M^*(X))$ vanish.

The following definition will be useful to that affect.

Definition 12.4 (Derivative according to pair). Let $P : (\mathbb{F}^n)^\ell \rightarrow \mathbb{T}$. For a vector $\alpha \in \mathbb{F}^\ell$ and an element $y \in \mathbb{F}^n$, the derivative of P according to the pair (α, y) is defined as

$$D_{(\alpha,y)} P(x_1, \dots, x_\ell) := P(x_1 + \alpha_1 y, \dots, x_\ell + \alpha_\ell y) - P(x_1, \dots, x_\ell). \tag{12.7}$$

For $P_i : \mathbb{F}^n \rightarrow \mathbb{T}$ and a linear form $M \in \mathbb{F}^\ell$, define $P_i \circ M : (\mathbb{F}^n)^\ell \rightarrow \mathbb{T}$ as

$$(P_i \circ M)(x_1, \dots, x_\ell) := P_i(M(x_1, \dots, x_\ell)).$$

Note that for every $M \in \mathcal{M}$,

$$\begin{aligned} D_{(\alpha,y)}(P_i \circ M)(x_1, \dots, x_\ell) &= P_i(M(x_1, \dots, x_\ell) + M(\alpha_1, \dots, \alpha_\ell)y) - P_i(M(x_1, \dots, x_\ell)) \\ &= (D_{\langle M, \alpha \rangle y} P_i)(M(x_1, \dots, x_\ell)). \end{aligned}$$

Thus if α is chosen such that $\langle M, \alpha \rangle = 0$ then $D_{(\alpha,y)}(P_i \circ M) \equiv 0$. Thus, our goal is to carefully choose a set of directions that will annihilate all linear forms but M^* , which will greatly simplify the analysis.

Assume that $M^* = (w_1^*, \dots, w_\ell^*)$, where $w_1^* = 1$. Let $t := |M^*|$. By [Claim 12.3](#) we have $t \leq d^*$, since if $\lambda'_{i,M^*} \neq 0$ then necessarily $|M^*| \leq \deg(\lambda'_{i,M^*} P_i) = d^*$. As a first step, take $\alpha_1 := e_1 = (1, 0, 0, \dots, 0) \in \mathbb{F}^\ell$. Choose additional $t - 1$ vectors as follows: for each $j = 2, \dots, \ell$, and each $w = 1, \dots, w_j^* - 1$, pick $\alpha_{j,w} := -we_1 + e_j = (-w, 0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{F}^\ell$, where the 1 is in the j -th coordinate. Observe that there are indeed $|M^*| - 1 = t - 1$ such vectors, and for convenience number them as $\alpha_2, \dots, \alpha_t \in \mathbb{F}^\ell$.

The following claim shows that deriving P_Λ iteratively according to the pairs $(\alpha_1, y_1), \dots, (\alpha_t, y_t)$ annihilates all linear forms except M^* .

Claim 12.5.

$$\begin{aligned} &D_{(\alpha_1,y_1)} \cdots D_{(\alpha_t,y_t)} P_\Lambda(X) \\ &= \left(D_{\langle M^*, \alpha_1 \rangle y_1} \cdots D_{\langle M^*, \alpha_t \rangle y_t} \left(\sum_{i=1}^C \lambda'_{i,M^*} P_i \right) \right) (M^*(X)). \end{aligned} \tag{12.8}$$

Proof. We need to show that for every $M = \mathcal{M} \setminus \{M^*\}$, there exists $i \in [t]$ such that $\langle M, \alpha_i \rangle = 0$. Assume that $M = (w_1, \dots, w_\ell)$. If $w_1 = 0$ then $\langle M, \alpha_1 \rangle = 0$. Otherwise by our assumption $w_1 = 1$. As $|M^*|$ was chosen to be maximal among all M with $\text{lc}(M) = \text{lc}(M^*)$, there must exist some $j = 2, \dots, \ell$ for which $w_j < w_j^*$. But then the appropriate $\alpha_i = \alpha_{j,w_j}$ satisfies $\langle M, \alpha_i \rangle = 0$. □

Next, we choose additional $d^* - t$ vectors $\alpha_{t+1}, \dots, \alpha_{d^*} \in \mathbb{F}^\ell$ to only keep polynomials P_i in the sum for which $\lambda'_{i,M^*} P_i$ has maximal

degree, namely d^* . To do so, simply choose $\alpha_{t+1} = \dots = \alpha_{d^*} := e_1 \in \mathbb{F}^\ell$. The next claim shows that deriving P_Λ according to the pairs $(\alpha_1, y_1), \dots, (\alpha_{d^*}, y_{d^*})$ annihilates all linear forms but M^* , and furthermore keeps only polynomials P_i where $\lambda'_{i, M^*} P_i$ has degree d .

Claim 12.6.

$$\begin{aligned}
 & D_{(\alpha_1, y_1)} \cdots D_{(\alpha_{d^*}, y_{d^*})} P_\Lambda(X) \\
 &= \left(D_{\langle M^*, \alpha_1 \rangle y_1} \cdots D_{\langle M^*, \alpha_{d^*} \rangle y_{d^*}} \sum_{\substack{i \in [C]: \\ \deg(\lambda'_{i, M^*} P_i) = d^*}} \lambda'_{i, M^*} P_i \right) (M^*(X)).
 \end{aligned}
 \tag{12.9}$$

Proof. We already know by Claim 12.5 that

$$\begin{aligned}
 & D_{(\alpha_1, y_1)} \cdots D_{(\alpha_t, y_t)} P_\Lambda(X) \\
 &= \left(D_{\langle M^*, \alpha_1 \rangle y_1} \cdots D_{\langle M^*, \alpha_t \rangle y_t} \left(\sum_{i=1}^C \lambda'_{i, M^*} P_i \right) \right) (M^*(X)).
 \end{aligned}$$

Thus, we just need to additionally derive this expression according to the pairs $(\alpha_{t+1}, y_{t+1}), \dots, (\alpha_{d^*}, y_{d^*})$. We have

$$\begin{aligned}
 & D_{(\alpha_1, y_1)} \cdots D_{(\alpha_{d^*}, y_{d^*})} P_\Lambda(X) \\
 &= \left(D_{\langle M^*, \alpha_1 \rangle y_1} \cdots D_{\langle M^*, \alpha_{d^*} \rangle y_{d^*}} \left(\sum_{i=1}^C \lambda'_{i, M^*} P_i \right) \right) (M^*(X)).
 \end{aligned}$$

The claim follows as by the choice of d^* we have $\deg(\lambda'_{i, M^*} P_i) \leq d^*$ for all i . If $\deg(\lambda'_{i, M^*} P_i) < d^*$ then the d^* derivatives $D_{\langle M^*, \alpha_1 \rangle y_1} \cdots D_{\langle M^*, \alpha_{d^*} \rangle y_{d^*}}$ annihilate $\lambda'_{i, M^*} P_i$. Thus, we only retain the polynomials P_i for which $\lambda'_{i, M^*} P_i$ has maximal degree, namely d^* . \square

Let $Q(x) := \sum_{i \in [C]: \deg(\lambda'_{i, M^*} P_i) = d^*} \lambda'_{i, M^*} P_i(x)$. Claim 12.6 implies that

$$\begin{aligned}
 & \mathbb{E}_{y_1, \dots, y_{d^*} \in \mathbb{F}^n, X \in (\mathbb{F}^n)^\ell} \left[e \left(D_{(\alpha_1, y_1)} \cdots D_{(\alpha_{d^*}, y_{d^*})} P_\Lambda \right) (X) \right] \\
 &= \mathbb{E}_{y_1, \dots, y_{d^*} \in \mathbb{F}^n, X \in (\mathbb{F}^n)^\ell} \left[e \left(\left(D_{\langle M^*, \alpha_1 \rangle y_1} \cdots D_{\langle M^*, \alpha_{d^*} \rangle y_{d^*}} Q \right) (M^*(X)) \right) \right].
 \end{aligned}$$

Next, define $x = M^*(X) \in \mathbb{F}^n$ and $z_j = \langle M^*, \alpha_j \rangle y_j \in \mathbb{F}^n$. It is simple to verify from our construction that $\langle M^*, \alpha_j \rangle \neq 0$ for all j . Thus, the joint distribution of $x, z_1, \dots, z_{d^*} \in \mathbb{F}^n$ is uniform and independent, and hence

$$\begin{aligned} \mathbb{E}_{\substack{y_1, \dots, y_{d^*} \in \mathbb{F}^n, \\ X \in (\mathbb{F}^n)^\ell}} \left[e \left(D_{(\alpha_1, y_1)} \cdots D_{(\alpha_{d^*}, y_{d^*})} P_\Lambda \right) (X) \right] \\ &= \mathbb{E}_{x, z_1, \dots, z_{d^*} \in \mathbb{F}^n} \left[e \left((D_{z_1} \cdots D_{z_{d^*}} Q) (x) \right) \right] \\ &= \|Q\|_{U^{d^*}}^{2^{d^*}}. \end{aligned}$$

Recall that \mathcal{B} is assumed to be of high rank. Concretely, we require \mathcal{B} to have rank at least $r_{7.32}(\mathbb{F}, d, \varepsilon) := r_{7.14}(\mathbb{F}, d, \varepsilon)$. By definition, any linear combination of the polynomials defining \mathcal{B} has such rank. In particular, this holds for Q , which by [Theorem 7.14](#) implies that $\|Q\|_{U^{d^*}} \leq \varepsilon$. Thus

$$\mathbb{E}_{y_1, \dots, y_{d^*}, X} \left[e \left(D_{(\alpha_1, y_1)} \cdots D_{(\alpha_{d^*}, y_{d^*})} P_\Lambda \right) (x_1, \dots, x_\ell) \right] \leq \varepsilon^{2^d},$$

In order to conclude the proof, we need to relate the LHS of the above expression to the bias of P_Λ . This can be achieved via a repeated application of the Cauchy-Schwarz inequality. The following claim appeared first in [\[14\]](#).

Claim 12.7 ([Bhattacharyya et al. \[14\]](#), Claim 3.4). For any nonzero $\alpha_1, \dots, \alpha_{d^*} \in \mathbb{F}^\ell$,

$$\begin{aligned} \mathbb{E}_{y_1, \dots, y_{d^*} \in \mathbb{F}^n, X \in (\mathbb{F}^n)^\ell} \left[e \left((D_{(\alpha_1, y_1)} \cdots D_{(\alpha_{d^*}, y_{d^*})} P_\Lambda) (X) \right) \right] \\ &\geq \left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} [e(P_\Lambda(X))] \right|^{2^d}. \end{aligned}$$

Proof. It suffices to show that for any function $P : (\mathbb{F}^n)^\ell \rightarrow \mathbb{T}$ and nonzero $\alpha = (a_1, \dots, a_\ell) \in \mathbb{F}^\ell$,

$$\left| \mathbb{E}_{y \in \mathbb{F}^n, X \in (\mathbb{F}^n)^\ell} \left[e \left((D_{\langle \alpha, y \rangle} P) (X) \right) \right] \right| \geq \left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} [e(P(X))] \right|^2.$$

12.1. Near-orthogonality: Proof of Theorem 7.32

Recall that $(D_{(\alpha,y)}P)(x_1, \dots, x_\ell) = P(x_1 + a_1y, \dots, x_\ell + a_\ell y) - P(x_1, \dots, x_\ell)$. Without loss of generality, suppose $a_1 \neq 0$. We make a change of coordinates so that α can be assumed to be $(1, 0, \dots, 0)$. More precisely, define $P' : (\mathbb{F}^n)^\ell \rightarrow \mathbb{T}$ as

$$P'(x_1, \dots, x_\ell) := P\left(x_1, \frac{x_2 + a_2x_1}{a_1}, \frac{x_3 + a_3x_1}{a_1}, \dots, \frac{x_\ell + a_\ell x_1}{a_1}\right).$$

Observe that with this definition,

$$P(x_1, \dots, x_\ell) = P'(x_1, a_1x_2 - a_2x_1, a_1x_3 - a_3x_1, \dots, a_1x_\ell - a_\ell x_1)$$

and

$$\begin{aligned} (D_{(\alpha,y)}P)(x_1, \dots, x_\ell) &= P'(x_1 + a_1y, a_1x_2 - a_2x_1, \dots, a_1x_\ell - a_\ell x_1) \\ &\quad - P'(x_1, a_1x_2 - a_2x_1, \dots, a_1x_\ell - a_\ell x_1). \end{aligned}$$

Therefore

$$\begin{aligned} &\mathbb{E}_{y,x_1,\dots,x_\ell \in \mathbb{F}} \left[e \left((D_{(\alpha,y)}P)(x_1, \dots, x_\ell) \right) \right] \\ &= \mathbb{E}_{y,x_1,\dots,x_\ell \in \mathbb{F}} \left[e \left(P'(x_1 + a_1y, a_1x_2 - a_2x_1, \dots, a_1x_\ell - a_\ell x_1) \right. \right. \\ &\quad \left. \left. - P'(x_1, a_1x_2 - a_2x_1, \dots, a_1x_\ell - a_\ell x_1) \right) \right] \\ &= \mathbb{E}_{y,x_1,\dots,x_\ell \in \mathbb{F}} \left[e \left(P'(x_1 + a_1y, x_2, \dots, x_\ell) - P'(x_1, x_2, \dots, x_\ell) \right) \right] \\ &= \mathbb{E}_{x_2,\dots,x_\ell \in \mathbb{F}} \left| \mathbb{E}_{x_1 \in \mathbb{F}} \left[e \left(P'(x_1, x_2, \dots, x_\ell) \right) \right] \right|^2 \end{aligned}$$

We can thus conclude the proof as, by the Cauchy-Schwartz inequality, it holds that

$$\begin{aligned} &\left| \mathbb{E}_{y,x_1,\dots,x_\ell \in \mathbb{F}} \left[e \left((D_{(\alpha,y)}P)(x_1, \dots, x_\ell) \right) \right] \right| \\ &\leq \left| \mathbb{E}_{x_1,x_2,\dots,x_\ell \in \mathbb{F}} \left[e \left(P'(x_1, x_2, \dots, x_\ell) \right) \right] \right|^2 \\ &= \left| \mathbb{E}_{x_1,x_2,\dots,x_\ell \in \mathbb{F}} \left[e \left(P(x_1, x_2, \dots, x_\ell) \right) \right] \right|^2. \end{aligned}$$

□

□

Remark 12.8. The proof of [Theorem 7.32](#) also shows the following. Suppose, under the same conditions of [Theorem 7.32](#), that for every polynomial P_i for $i \in [C]$ and any linear form L_j for $j \in [m]$, one the following two conditions holds:

- $|L_j| \leq \deg(\lambda_{i,j}P_i)$; or
- $\lambda_{i,j}P_i = 0$.

Now, if $\lambda_{i,j} \equiv 0 \pmod{p^{\text{depth}(P_i)+1}}$ for all $i \in [C], j \in [m]$, then clearly $\lambda_{i,j}P_i = 0$ for all i, j and hence $P_\Lambda = 0$. In all other cases, we obtain that $P_\lambda \neq 0$ and hence $\text{bias}(P_\Lambda) < \varepsilon$. The proof is identical to the proof of [Theorem 7.32](#), except that there is no need to transform $\lambda_{i,j}$ to $\lambda'_{i,j}$ in the beginning of the proof.

12.2 Proof of [Theorem 11.8](#)

Theorem 11.8 (restated). Let $\mathcal{L} = \{L_1, \dots, L_m\}$ be a system of linear forms in ℓ variables, such that $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent. For every $\varepsilon > 0$, there exists $\delta > 0$ such that for any collection of functions $f_1, \dots, f_m : \mathbb{F}^n \rightarrow \mathbb{D}$ with $\min_{i \in [m]} \|f_i\|_{U^{d+1}} \leq \delta$, we have

$$\left| \mathbb{E}_{X \in (\mathbb{F}^n)^k} \left[\prod_{i=1}^m f_i(L_i(X)) \right] \right| \leq \varepsilon. \tag{12.10}$$

We present the proof due to Hatami *et al.* [\[47\]](#). First, note that since $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent, it must be the case that L_1, \dots, L_m are pairwise linearly independent. That is, it is not the case that $L_i = cL_j$ for some distinct $i, j \in [m]$ and $c \in \mathbb{F}$. Consequently, $\mathcal{L} = \{L_1, \dots, L_m\}$ is of finite Cauchy-Schwarz complexity s for some $s \leq m - 2 < \infty$, because as pointed out earlier, for every $i \in [m]$, the partition of $\{L_j\}_{j \in [m] \setminus \{i\}}$ into $m - 1$ singletons satisfies the requirement of [Definition 11.1](#). The case when $s \leq d$ follows from [Lemma 11.2](#). Thus we are left with the case when $s > d$. We will prove that $\|f_1\|_{U^{d+1}} \leq \delta$

where δ is sufficiently small, implies Equation (12.10). The theorem then follows by symmetry in the choice of f_1 .

As a first step, we decompose the functions as $f_i = g_i + h_i$, where $g_i = \mathbb{E}[f_i|\mathcal{B}]$ for a suitably chosen polynomial factor \mathcal{B} of sufficiently high rank, and where h_i has negligible $(s + 1)$ -th Gowers norm. In order to do so, let $r : \mathbb{N} \rightarrow \mathbb{N}$ be a large enough rank bound to be chosen later. Apply a multifunction version of Theorem 9.1 along with Theorem 10.4 and Theorem 7.30, to obtain a simultaneous decomposition $f_i = g_i + h_i$ for all $i \in [m]$ where

1. $g_i = \mathbb{E}[f_i|\mathcal{B}]$, where \mathcal{B} is an r -regular polynomial factor, defined by homogeneous polynomials, of degree at most s and complexity $C \leq C_{\max}(|\mathbb{F}|, s, m, \varepsilon, r(\cdot))$.
2. $\|h_i\|_{U^{s+1}} \leq \frac{\varepsilon}{2m}$.

We will assume that the rank $r(C)$ is chosen large enough, so that \mathcal{B} is $\nu(C)$ -uniform for $\nu(C) > 0$ to be chosen later (See Remark 7.19). We first show that we may replace f_i 's in Equation (12.10) with g_i 's, using the same proof as Corollary 11.9.

Claim 12.9. $\left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[\prod_{i=1}^m f_i(L_i(X)) \right] - \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[\prod_{i=1}^m g_i(L_i(X)) \right] \right| \leq \frac{\varepsilon}{2}$.

Proof. We have

$$\begin{aligned} & \left| \mathbb{E}_X \left[\prod_{i=1}^m f_i(L_i(X)) \right] - \mathbb{E}_X \left[\prod_{i=1}^m g_i(L_i(X)) \right] \right| \\ &= \left| \sum_{i=1}^m \mathbb{E}_X \left[h_i(L_i(X)) \cdot \prod_{j=1}^{i-1} g_j(L_j(X)) \cdot \prod_{j=i+1}^m f_j(L_j(X)) \right] \right| \\ &\leq \sum_{i=1}^m \left| \mathbb{E}_X \left[h_i(L_i(X)) \cdot \prod_{j=1}^{i-1} g_j(L_j(X)) \cdot \prod_{j=i+1}^m f_j(L_j(X)) \right] \right| \\ &\leq \sum_{i=1}^m \|h_i\|_{U^{s+1}} \leq \frac{\varepsilon}{2}, \end{aligned}$$

where the second inequality follows from Lemma 11.2 as the Cauchy-Schwarz complexity of \mathcal{L} is s . □

Thus it is sufficient to bound $\left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[\prod_{i=1}^m g_i(L_i(X)) \right] \right| \leq \varepsilon/2$. For each i , $g_i = \mathbb{E}[f_i | \mathcal{B}]$ and thus

$$g_i(x) = \Gamma_i(P_1(x), \dots, P_C(x)),$$

where P_1, \dots, P_C are the nonclassical homogeneous polynomials of degree $\leq s$ defining \mathcal{B} and $\Gamma_i : \mathbb{T}^C \rightarrow \mathbb{D}$ is a function. Let $k_i = \text{depth}(P_i)$ so that by [Lemma 6.3](#), each P_i takes values in $\mathbb{U}_{k_i+1} = \frac{1}{p^{k_i+1}} \mathbb{Z}/\mathbb{Z}$. Let $\Sigma := \mathbb{Z}_{p^{k_1+1}} \times \dots \times \mathbb{Z}_{p^{k_C+1}}$.

Using the Fourier transform on $\mathbb{U}_{k_1+1} \times \dots \times \mathbb{U}_{k_C+1} \cong \Sigma$, for every $\tau = (\tau_1, \dots, \tau_C) \in \Sigma$ we have

$$\Gamma_i(\tau) = \sum_{\Lambda=(\lambda_1, \dots, \lambda_C) \in \Sigma} \widehat{\Gamma}_i(\Lambda) \cdot e \left(\sum_{j=1}^C \lambda_j \tau_j \right), \tag{12.11}$$

where

$$\widehat{\Gamma}_i(\Lambda) := \mathbb{E}_{\tau} \left[\Gamma_i(\tau) e \left(\sum_{j=1}^C \lambda_j \tau_j \right) \right]$$

is the Fourier coefficient of Γ_i corresponding to Λ . Observe that $|\widehat{\Gamma}_i(\Lambda)| \leq 1$. Consequently,

$$g_i(x) = \sum_{\Lambda=(\lambda_1, \dots, \lambda_C) \in \Sigma} \widehat{\Gamma}_i(\Lambda) \cdot e \left(\sum_{j=1}^C \lambda_j P_j(x) \right). \tag{12.12}$$

Let $P_{\Lambda} := \sum_{j=1}^C \lambda_j P_j(x)$ for the sake of brevity so that we may write

$$\mathbb{E}_X \left[\prod_{i=1}^m g_i(L_i(X)) \right] = \sum_{\Lambda_1, \dots, \Lambda_m \in \Sigma} \left(\prod_{i=1}^m \widehat{\Gamma}_i(\Lambda_i) \right) \cdot \mathbb{E}_X \left[e \left(\sum_{i=1}^m P_{\Lambda_i}(L_i(X)) \right) \right]. \tag{12.13}$$

We will show that each term in [Equation \(12.13\)](#) can be bounded by $\frac{\varepsilon}{2^{|\Sigma|^m}}$, thus concluding the proof by the triangle inequality.

We first show that the terms for which $\text{deg}(P_{\Lambda_1}) \leq d$ are small. Recall that by our assumption $\|f_1\|_{U^{d+1}} \leq \delta$, where δ is small enough to be determined later.

Claim 12.10. Let $\Lambda_1 \in \Sigma$ be such that $\deg(P_{\Lambda_1}) \leq d$. Then

$$\left| \widehat{\Gamma}_1(\Lambda_1) \right| \leq \delta + |\Sigma| \nu(C).$$

Proof. It follows from Equation (12.12) that

$$\widehat{\Gamma}_1(\Lambda_1) = \mathbb{E}_{x \in \mathbb{F}^n} [g_1(x)e(-P_{\Lambda_1}(x))] - \sum_{\Lambda \neq \Lambda_1} \widehat{\Gamma}_1(\Lambda) \cdot \mathbb{E}_{x \in \mathbb{F}^n} [e(P_{\Lambda}(x) - P_{\Lambda_1}(x))].$$

We bound the first term.

$$\begin{aligned} \left| \mathbb{E}_x [g_1(x)e(-P_{\Lambda_1}(x))] \right| &= \left| \mathbb{E}_x [f_1(x)e(-P_{\Lambda_1}(x))] \right| \\ &\leq \|f_1 e(-P_{\Lambda_1})\|_{U^{d+1}} \\ &= \|f_1\|_{U^{d+1}} \leq \delta. \end{aligned}$$

The first equality follows as $g_1 = \mathbb{E}[f_1|\mathcal{B}]$ and P_{Λ_1} is \mathcal{B} -measurable.

Next, we bound each summand in the second term. Fix $\Lambda \neq \Lambda_1$. Recall that the polynomial factor \mathcal{B} is $\nu(C)$ -uniform by our construction. In particular the bias of $P_{\Lambda} - P_{\Lambda_1} = P_{\Lambda - \Lambda_1}$ is at most $\nu(C)$. That is, $|\mathbb{E}_{x \in \mathbb{F}^n} [e(P_{\Lambda}(x) - P_{\Lambda_1}(x))]| \leq \nu(C)$. As also $|\widehat{\Gamma}_1(\Lambda)| \leq 1$, each summand in the sum in the second term is bounded by $\nu(C)$. \square

Claim 12.10 allows us to bound the contribution of the terms in Equation (12.13) corresponding to tuples $(\Lambda_1, \dots, \Lambda_m) \in \Sigma^m$ with $\deg(P_{\Lambda_1}) \leq d$.

$$\sum_{\substack{\Lambda_1, \dots, \Lambda_m \in \Sigma: \\ \deg(P_{\Lambda_1}) \leq d}} \left(\prod_{i=1}^m \widehat{\Gamma}_i(\Lambda_i) \right) \mathbb{E}_X \left[e \left(\sum_{i=1}^m P_{\Lambda_i}(L_i(X)) \right) \right] \leq |\Sigma|^m (\delta + |\Sigma| \nu(C)). \tag{12.14}$$

Next, we bound the terms for which $\deg(P_{\Lambda_1}) > d$. We will need the following lemma.

Lemma 12.11. Assume that $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent, and let $(\Lambda_1, \dots, \Lambda_m) \in \Sigma^m$ be such that $\deg(P_{\Lambda_1}) \geq d + 1$. Then

$$\sum_{i=1}^m P_{\Lambda_i}(L_i(X)) \neq 0.$$

Before proving [Lemma 12.11](#) let us first describe why it suffices to complete the proof of [Theorem 11.8](#). [Theorem 7.32](#) gives that, if we that the rank $r(\cdot)$ is chosen so that $r(C) \geq r_{7.32}(\mathbb{F}, s, \delta)$, then under the conclusion of [Lemma 12.11](#), we in fact have that

$$\left| \mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[e \left(\sum_{i=1}^m P_{\Lambda_i}(L_i(X)) \right) \right] \right| \leq \delta.$$

We may thus conclude that

$$\sum_{\substack{\Lambda_1, \dots, \Lambda_m \in \Sigma: \\ \deg(P_{\Lambda_1}) \geq d+1}} \left(\prod_{i=1}^m \widehat{\Gamma}_i(\Lambda_i) \right) \cdot \mathbb{E}_X \left[e \left(\sum_{i=1}^m P_{\Lambda_i}(L_i(X)) \right) \right] \leq \delta |\Sigma|^m. \tag{12.15}$$

Combining [\(12.14\)](#) and [\(12.15\)](#) allows us to conclude that

$$\sum_{\Lambda_1, \dots, \Lambda_m \in \Sigma} \left(\prod_{i=1}^m \widehat{\Gamma}_i(\Lambda_i) \right) \cdot \mathbb{E}_X \left[e \left(\sum_{i=1}^m P_{\Lambda_i}(L_i(X)) \right) \right] \leq (\delta + |\Sigma| \nu(C)) |\Sigma|^m.$$

To conclude we set parameters. Observe that $|\Sigma| = |\mathbb{F}|^{\sum_{i \in [C]} \text{depth}(P_i)+1} \leq |\mathbb{F}|^{\sum_{i \in [C]} \text{deg}(P_i)} \leq |\mathbb{F}|^{Cs}$. Thus, we may choose $r(\cdot)$ so that $\nu(C) |\Sigma|^{m+1} \leq \varepsilon/4$. To conclude, after choosing $r(\cdot)$ we have an upper bound on the complexity of \mathcal{B} , namely $C \leq C_{\max} = C_{\max}(|\mathbb{F}|, s, m, \varepsilon, r(\cdot))$. We set $\delta := (\varepsilon/4) |\mathbb{F}|^{C_{\max} sm}$ and conclude the proof.

Thus, we are left with proving [Lemma 12.11](#). This is the only place in the proof where we actually use the assumption that $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent.

Proof of [Lemma 12.11](#). Assume to the contrary that $\sum_{i=1}^m P_{\Lambda_i}(L_i(X))$ is identically zero. Denoting the coordinates of Λ_i by $(\lambda_{i,1}, \dots, \lambda_{i,C}) \in \Sigma^C$ we have

$$\sum_{i=1}^m P_{\Lambda_i}(L_i(X)) = \sum_{i \in [m], j \in [C]} \lambda_{i,j} P_j(L_i(X)) \equiv 0.$$

We apply [Theorem 7.32](#). This requires assuming that the polynomial factor \mathcal{B} has high enough rank, which we can achieve by requiring that $r(C) \geq r_{7.32}(\mathbb{F}, s, 1/2)$, say. Thus, for every $j \in [C]$ we must have

$$\sum_{i=1}^m \lambda_{i,j} P_j(L_i(X)) \equiv 0. \tag{12.16}$$

In fact, we know more: for any nonclassical homogeneous polynomials Q_j with $\deg(Q_j) = \deg(P_j)$ and $\text{depth}(Q_j) \leq \text{depth}(P_j)$ it holds that

$$\sum_{i=1}^m \lambda_{i,j} Q_j(L_i(X)) \equiv 0.$$

Next, as we assume that $\deg(P_{\Lambda_1}) \geq d + 1$, there must exist $j \in [C]$ such that $\deg(\lambda_{1,j} P_j) \geq d + 1$. Let $j^* \in [C]$ be such that $d^* := \deg(\lambda_{1,j^*} P_{j^*})$ is maximized. By our assumptions, $d^* \geq d + 1$. Let $t \geq 0$ be the largest integer such that $p^t = |\mathbb{F}|^t$ divides λ_{i,j^*} for all $i \in [m]$. Define $\mu_i := \lambda_{i,j^*} / p^t \in \mathbb{Z}$ and $P(x) = p^t P_{j^*}(x)$. Then

$$\sum_{i=1}^m \mu_i P(L_i(X)) = \sum_{i=1}^m \frac{\lambda_{i,j^*}}{p^t} p^t P_{j^*}(L_i(X)) \equiv 0.$$

The polynomial P belongs to \mathcal{B} . As such, $\{P\}$ has rank at least $r(C)$ as well. Thus by another application of Theorem 7.32, for any homogeneous polynomial Q of degree $\deg(Q) = \deg(P)$ and $\text{depth}(Q) \leq \text{depth}(P)$ it holds that

$$\sum_{i=1}^m \mu_i Q(L_i(X)) \equiv 0. \tag{12.17}$$

In particular, this holds for any homogeneous classical polynomial Q of degree $D := \deg(P) \geq \deg(\lambda_{1,j} P_j) = d^* \geq d + 1$. That is, if $R : \mathbb{F}^n \rightarrow \mathbb{F}$ is a homogeneous classical polynomial of degree D then, by setting $Q(x) = \frac{R(x)}{p}$ we get

$$\sum_{i=1}^m \mu_i R(L_i(X)) \equiv 0 \pmod{p}. \tag{12.18}$$

We apply it to the homogeneous classical monomial $R(z) = z_1 z_2 \dots z_D$. In order to compute $R(L_i(X))$, let $X = (x_1, \dots, x_\ell) \in (\mathbb{F}^n)^\ell$ where $x_i = (x_{i,1}, \dots, x_{i,n}) \in \mathbb{F}^n$. Let $L_i = (\lambda_{i,1}, \dots, \lambda_{i,\ell}) \in \mathbb{F}^\ell$. Then

$$R(L_i(X)) = \prod_{a=1}^D \left(\sum_{b=1}^{\ell} \lambda_{i,b} x_{b,a} \right).$$

In particular, for $b_1, \dots, b_D \in [\ell]$, the coefficient of the monomial $\prod_{a=1}^D x_{b_a,a}$ in $R(L_i(X))$ is $\prod_{a=1}^D \lambda_{i,b_a}$, which is the (b_1, \dots, b_D) coef-

ficient of L_i^D . Thus (12.18) implies that

$$\sum_{i=1}^m \mu_i L_i^D \equiv 0 \pmod{p}.$$

However, by our choice of t , there must exist $i^* \in [m]$ for which $p^t \lambda_{i^*, j^*}$ does not divide by p . That is, $\mu_{i^*} \neq 0 \pmod{p}$. But then we get that L_1^D, \dots, L_m^D are linearly dependent. This contradicts the assumption that the assumption that $L_1^{d+1}, \dots, L_m^{d+1}$ are linearly independent. \square

13

Algorithmic Regularity

This chapter is concerned with the algorithmic versions of regularity lemmas for polynomials over finite fields. The regularity lemmas proved by Green and Tao [43] and Kaufman and Lovett [54] as discussed in Section 7.5 show that one can modify a given collection of polynomials $\mathcal{B} = \{P_1, \dots, P_c\}$ into a regular collection $\mathcal{B}' = \{P_1, \dots, P_{c'}\}$ of polynomials of same or lower degree. Here, we mean regularity in the sense of Definition 7.8. These lemmas are central to higher-order Fourier analysis and have various applications. Bhattacharyya *et al.* [17] studied algorithmic versions of these theorems and showed that analytic notions of regularity such as the ones defined in Section 7.2 allow for efficient algorithms.

13.1 A lemma of Bogdanov and Viola

A key first step in proofs of inverse theorems for Gowers norms and that biased polynomials have small rank is an elegant argument due to Bogdanov and Viola [21], proving that if a polynomial of degree d is *biased*, then it can be approximated by a bounded set of polynomials of lower degree. In Chapter 8 we saw a more efficient version of this lemma due to Bhowmick and Lovett [19]. It was observed by Bhattacharyya

et al. [17] that the Bogdanov-Viola lemma can be made algorithmic due to its probabilistic proof.

Lemma 13.1 (Algorithmic Bogdanov-Viola lemma [17]). Let $d \geq 0$ be an integer, and $\delta, \sigma, \beta \in (0, 1]$ be parameters. There exists a randomized algorithm, that given a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree d with

$$\text{bias}(P) \geq \delta,$$

runs in time $O_{\delta, \beta, \sigma}(n^d)$, and with probability $1 - \beta$ returns functions $\tilde{P} : \mathbb{F}^n \rightarrow \mathbb{F}$ and $\Gamma : \mathbb{F}^C \rightarrow \mathbb{F}$, and a set of polynomials P_1, \dots, P_C , where $C \leq \frac{|\mathbb{F}|^5}{\delta^2 \sigma \beta}$ and $\deg(P_i) < d$ for all $i \in [C]$, for which

- $\Pr_x(P(x) \neq \tilde{P}(x)) \leq \sigma$, and
- $\tilde{P}(x) = \Gamma(P_1(x), \dots, P_C(x))$.

The idea is to use derivatives of P at a few randomly chosen directions in order to approximate the value of P .

Proof. The proof will be an adaptation of the proof from the paper by Green and Tao [43]. Given query access to the polynomial P , we can compute the explicit description of P in $O(n^d)$ queries. For every $a \in \mathbb{F}$ define the measure $\mu_a(t) := \Pr(P(x) = a + t)$. It is easy to see that if $\text{bias}(P(x)) \geq \delta$ then, for every $a \neq b$,

$$\|\mu_a - \mu_b\|_\infty \geq \frac{4\delta}{|\mathbb{F}|}. \tag{13.1}$$

We will try to estimate each of these distributions. Let

$$\tilde{\mu}_a(t) := \frac{1}{C} \sum_{1 \leq i \leq C} \mathbb{1}_{P(x_i)=a+t},$$

where $C > \frac{|\mathbb{F}|^5}{\delta \cdot \beta_1}$, and $x_1, x_2, \dots, x_C \in \mathbb{F}^n$ are chosen uniformly at random. Therefore by an application of Chebyshev's inequality

$$\Pr \left(|\tilde{\mu}_a(t) - \mu_a(t)| > \frac{\delta}{2|\mathbb{F}|^2} \right) < \frac{\beta_1}{|\mathbb{F}|},$$

for all $t \in \mathbb{F}$ and therefore

$$\Pr \left(\|\tilde{\mu}_a - \mu_a\|_\infty > \frac{\delta}{2|\mathbb{F}|^2} \right) < \beta_1. \tag{13.2}$$

Now we will focus on approximating $P(x)$. Remember that $D_h P(x) = P(x + h) - P(x)$ is the additive derivative of $P(x)$ in direction h . We have

$$\Pr_h(D_h P(x) = r) = \Pr_h(P(x + h) - P(x) = r) = \mu_{P(x)}(r),$$

where $h \in \mathbb{F}^n$ is chosen uniformly at random. Let $\mathbf{h} = (h_1, \dots, h_C) \in (\mathbb{F}^n)^C$ be chosen uniformly at random, where C is a sufficiently large constant to be chosen later. Define the corresponding “observed” distribution as

$$\mu_{\text{obs}}^{(x)}(t) := \frac{1}{C} \sum_{1 \leq i \leq C} \mathbb{1}_{D_{h_i} P(x) = t},$$

and let

$$\tilde{P}_h(x) := \arg \min_{r \in \mathbb{F}} \|\tilde{\mu}_r - \mu_{\text{obs}}^{(x)}\|_\infty.$$

Now choosing $C \geq \frac{|\mathbb{F}|^5}{\delta^2 \sigma \beta_2}$, another application of Chebyshev’s inequality gives

$$\Pr_h(\tilde{P}_h(x) \neq P(x)) \leq \Pr_h\left(\|\mu_{\text{obs}}^{(x)} - \mu_{P(x)}\|_\infty \geq \frac{\delta}{|\mathbb{F}|}\right) \leq \sigma \beta_2, \tag{13.3}$$

where the first inequality follows from (13.1) and (13.2). Therefore

$$\Pr_{x,h}(\tilde{P}_h(x) \neq P(x)) = \mathbb{E}_x \mathbb{E}_h \mathbb{1}_{\tilde{P}_h(x) \neq P(x)} \leq \sigma \beta_2,$$

and thus

$$\Pr_h \left[\Pr_x \left(\tilde{P}_h(x) \neq P(x) \right) \geq \sigma \right] \leq \beta_2.$$

Let $P_i := D_{h_i} P$, so that P_i is of degree $\leq d$ and \tilde{P}_h is a function of P_1, \dots, P_C . Now setting $\beta_1 := \frac{\beta}{2|\mathbb{F}|^2}$ and $\beta_2 := \frac{\beta}{2}$ finishes the proof. \square

13.2 Algorithmic regularity lemmas

Lemma 13.1 implies an algorithmic analogue of the Lemma 7.29, with the caveat that the refinement is approximate. The proof is by an induction similar to that of Lemma 7.29 with the difference that at each step one has to control the errors that are introduced through the use of Lemma 13.1 and the probability of correctness.

Definition 13.2. Let $\mathcal{F} = \{P_1, \dots, P_m\}$ be a polynomial factor of degree d . We say that a polynomial factor $\mathcal{F}' = \{Q_1, \dots, Q_M\}$ is σ -close to being a refinement of \mathcal{F} , if there exists a map Γ for which $\Pr_{x \in \mathbb{F}^n}[\mathcal{F}(x) \neq \Gamma(\mathcal{F}'(x))] \leq \sigma$.

Lemma 13.3 (Unbiased almost refinement). Let $d \geq 1$ be an integer. Suppose $\gamma : \mathbb{N} \rightarrow \mathbb{R}^+$ is a decreasing function and $\sigma, \rho \in (0, 1]$. There is a randomized algorithm that given a factor \mathcal{F} of degree d , runs in time $O_{\gamma, \rho, \sigma, \dim(\mathcal{F})}(n^d)$ and with probability $1 - \rho$ returns a γ -unbiased factor \mathcal{F}' with $\dim(\mathcal{F}') = O_{\gamma, \rho, \sigma, \dim(\mathcal{F})}(1)$, such that \mathcal{F}' is σ -close to being a refinement of \mathcal{F} .

Proof. The proof idea is similar to that of Lemma 7.29 in the sense that we use the same type of induction. The difference is that at each step we will have to control the errors that we introduce and the probability of correctness. At all steps in the proof without loss of generality we will assume that the polynomials in the factor are linearly independent, because otherwise we can always detect such a linear combination in $O_{\gamma, \rho, \sigma, \dim(\mathcal{F})}(n^d)$ time and remove a polynomial that can be written as a linear combination of the rest of the polynomials in the factor.

The base case for $d = 1$ is simple, a linearly independent set of non-constant linear polynomials is not biased at all, namely it is 0-unbiased. If \mathcal{F} is γ -biased, then there exists a set of coefficients $\{c_{i,j} \in \mathbb{F}\}_{1 \leq i \leq d, 1 \leq j \leq M_i}$ such that

$$\text{bias}\left(\sum_{i,j} c_{i,j} P_{i,j}\right) \geq \gamma(\dim(\mathcal{F})).$$

To detect this, we will use the following algorithm:

We will estimate bias of each of the $|\mathbb{F}|^{\dim(\mathcal{F})}$ linear combinations and check whether it is greater than $\frac{3\gamma(\dim(\mathcal{F}))}{4}$. To do so, for each linear combination $\sum_{i,j} c_{i,j} P_{i,j}$ independently select a set of vectors x_1, \dots, x_C uniformly at random from \mathbb{F}^n , and let $\widetilde{\text{bias}}(\sum_{i,j} c_{i,j} P_{i,j}) := \left| \frac{1}{C} \sum_{\ell \in [C]} e_{\mathbb{F}}(y_{\ell}) \right|$, where $y_{\ell} = \sum_{i,j} c_{i,j} \cdot P_{i,j}(x_{\ell})$. Choosing

$$C = O_{\dim(\mathcal{F})} \left(\frac{1}{\gamma(\dim(\mathcal{F}))^2} \log\left(\frac{1}{\rho}\right) \right),$$

we can distinguish bias $\geq \gamma$ from bias $\leq \frac{\gamma}{2}$, with probability $1 - \rho'$, where $\rho' := \frac{\rho}{4|\mathbb{F}|^{\dim(\mathcal{F})}}$. Let $\sum_{i,j} c_{i,j} P_{i,j}$ be such that the estimated bias

was above $\frac{3\gamma(\dim(\mathcal{F}))}{4}$ and k be its degree. We will stop if there is no such linear combination or if the factor is of degree 1. Since by a union bound with probability at least $1 - \frac{\rho}{4}$, $\text{bias}(\sum_{i,j} c_{i,j} P_{i,j}) \geq \frac{\gamma(\dim(\mathcal{F}))}{2}$, by [Lemma 13.1](#) we can find, with probability $1 - \frac{\rho}{4}$, a set of polynomials Q_1, \dots, Q_r of degree $k - 1$ such that

- $\sum_{i,j} c_{i,j} P_{i,j}$ is $\frac{\sigma}{2}$ -close to a function of Q_1, \dots, Q_r ,
- $r \leq \frac{16|\mathbb{F}|^5}{\gamma(\dim(\mathcal{F}))^2 \cdot \sigma \cdot \rho}$.

We replace one polynomial of highest degree that appears in $\sum_{i,j} c_{i,j} P_{i,j}$ with polynomials Q_1, \dots, Q_r .

We will prove by the induction that our algorithm satisfies the statement of the lemma. For the base case, if \mathcal{F} is of degree 1, our algorithm does not refine \mathcal{F} by design. Again, since we have removed all linear dependencies, \mathcal{F} is in fact 0-unbiased in this case.

Now given a factor \mathcal{F} , if \mathcal{F} is γ -biased, then with probability $1 - \rho'$ our algorithm will refine \mathcal{F} . With probability $1 - \frac{\rho}{4}$ the linear combination used for the refinement is $\frac{\gamma(\dim(\mathcal{F}))}{2}$ -biased. Let $\tilde{\mathcal{F}}$ be the outcome of one step of our algorithm. With probability $1 - \frac{\rho}{4}$, $\tilde{\mathcal{F}}$ is $\frac{\sigma}{2}$ -close to being a refinement of \mathcal{F} . Using the induction hypothesis with parameters $\gamma, \frac{\sigma}{2}, \frac{\rho}{4}$ we can find, with probability $1 - \frac{\rho}{4}$, a γ -unbiased factor \mathcal{F}' which is $\frac{\sigma}{2}$ -close to being a refinement of $\tilde{\mathcal{F}}$ and therefore, with probability at least $1 - (\frac{\rho}{4} + \frac{\rho}{4} + \frac{\rho}{4} + \rho') > 1 - \rho$, is σ -close to being a refinement of \mathcal{F} . □

When the field order is large, this approximate refinement can be made exact if we work with uniform factors (see [Definition 7.18](#)).

Lemma 13.4 (Uniform refinement [17]). Suppose $d < |\mathbb{F}|$ is a positive integer and $\rho \in (0, 1]$ is a parameter. There is a randomized algorithm that, takes as input a factor \mathcal{F} of degree d over \mathbb{F}^n , and a decreasing function $\gamma : \mathbb{N} \rightarrow \mathbb{R}^+$, runs in time $O_{\rho, \gamma, |\mathcal{F}|}(n^d)$, and with probability $1 - \rho$ outputs a γ -uniform factor $\tilde{\mathcal{F}}$, such that $\tilde{\mathcal{F}}$ is a refinement of \mathcal{F} of the same degree d , and $|\tilde{\mathcal{F}}| \ll_{\sigma, \gamma, |\mathcal{F}|} 1$.

We only sketch the proof of [Lemma 13.4](#), which is by induction on the dimension vector of \mathcal{F} . For the induction step, one checks whether

there is a linear combination of polynomials in \mathcal{F} that has large Gowers norm. One then uses this to replace a polynomial P from \mathcal{F} with a set of lower degree polynomials. To do this, we first approximate P with a few lower degree polynomials Q_1, \dots, Q_r , then use the induction hypothesis to refine $\{Q_1, \dots, Q_r\}$ to a uniform factor $\{\tilde{Q}_1, \dots, \tilde{Q}_{r'}\}$ and use an argument similar to that seen in Section 8.2 that approximation by a sufficiently uniform factor implies exact computation to conclude that P is measurable in $\{\tilde{Q}_1, \dots, \tilde{Q}_{r'}\}$.

A similar argument combined with Lemma 13.1 gives the following.

Lemma 13.5. Suppose that an integer d satisfies $0 \leq d < |\mathbb{F}|$. Let $\delta, \beta \in (0, 1]$. There is a randomized algorithm that given a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree d such that $\text{bias}(P) \geq \delta > 0$, runs in $O_{\delta, \beta}(n^d)$ and with probability $1 - \beta$, returns a polynomial factor $\mathcal{F} = \{P_{i,j}\}_{1 \leq i \leq d-1, 1 \leq j \leq M_i}$ of degree $d - 1$ and $|\mathcal{F}| = O_{\delta, \beta}(1)$ such that P is measurable in \mathcal{F} .

Algorithmic regularity in low characteristic. Unfortunately, Gowers uniformity for polynomial factors fails to address “bias implies low rank” phenomena in the case when \mathbb{F} has small characteristic. Kaufman and Lovett [54] introduce a stronger notion of regularity in order to handle the general case. This notion is rather technical and we omit introducing it here.

Algorithmic versions of the results of Kaufman and Lovett [54] were proved by Bhattacharyya *et al.* [17] by using a similar but stronger notion of regular factors referred to as “strong unbiased factors”. The reason for the need of this new notion is likewise that uniform factors (Definition 7.18) fail to address fields of low characteristic, for the reason that previously in order to refine a factor to a uniform factor we made use of division by $d!$ which is not possible in fields with $|\mathbb{F}| \leq d$.

The main result of Bhattacharyya *et al.* [17] in the low-characteristic setting is an algorithm for regularization of factors in low characteristic.

Lemma 13.6 (Strongly unbiased refinement (informal)). Suppose that $\gamma : \mathbb{N} \rightarrow \mathbb{R}^+$ is a regularity parameter. There is a deterministic algorithm that given a factor $\mathcal{F} = \{P_1, \dots, P_m\}$ of degree d , runs in $O_\gamma(n^{O(d)})$, returns a strongly γ -unbiased degree $\leq d$ factor $\mathcal{F}' \succeq \mathcal{F}$.

13.3 Algorithmic inverse theorem for polynomials

From [Lemma 13.5](#) one can deduce an algorithmic version of an inverse theorem for Gowers norm ([Theorem 6.9](#)) for polynomials in the high characteristic setting.

Theorem 13.7 (Algorithmic inverse theorem in high characteristic). Suppose that $|\mathbb{F}| > d \geq 2$ and that $\epsilon, \beta \in (0, 1]$. There is an $\eta_{\epsilon, \beta, d} \in (0, 1]$, and a randomized algorithm that given a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree d with $\|e_{\mathbb{F}}(P(x))\|_{U^{k+1}} \geq \epsilon$, runs in $O_{\delta, \beta}(n^d)$ and with probability $1 - \beta$, returns a polynomial Q of degree $\leq k$ such that

$$|\langle e_{\mathbb{F}}(P), e_{\mathbb{F}}(Q) \rangle| \geq \eta.$$

This theorem follows from the following proposition via a basic Fourier analytic argument, along with a theorem of Goldreich and Levin [\[34\]](#).

Proposition 13.1 (Computing polynomials with high Gowers norm). Suppose that $|\mathbb{F}| > d \geq 2$ and that $\delta, \beta \in (0, 1]$. There is a randomized algorithm that given a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree d with $\|e_{\mathbb{F}}(P(x))\|_{U^d} \geq \delta$, runs in $O_{\delta, \beta}(n^d)$ and with probability $1 - \beta$, returns a polynomial factor \mathcal{F} of degree $d - 1$ such that

- There is a function $\Gamma : \mathbb{F}^{|\mathcal{F}|} \rightarrow \mathbb{F}$ such that $P = \Gamma(\mathcal{F})$.
- $|\mathcal{F}| = O_{d, \delta, \beta}(1)$.

Proof. Write $\partial^d P(h_1, \dots, h_d) := D_{h_1} \cdots D_{h_d} P(x)$. Since P has degree d , $\partial^d P$ does not depend on x . From the definition of the U^d norm, we have

$$\text{bias}(\partial^d P) = \|e(P)\|_{U^d}^{2d} \geq \delta^{2d}.$$

Applying [Lemma 13.5](#) to $\partial^d P$, with probability $1 - \frac{\beta}{2}$, we can find a factor $\tilde{\mathcal{F}}$ of degree $d - 1$, such that $|\tilde{\mathcal{F}}| = O_{\delta, \beta, d}(1)$ and $\partial^d P$ is measurable in $\tilde{\mathcal{F}}$. It is easy to check that since $|\mathbb{F}| > d$, we have the following Taylor expansion

$$P(x) = \frac{1}{d!} \partial^d P(x, \dots, x) + Q(x),$$

where Q is a polynomial of degree $\leq d - 1$. We can find an explicit description of P , and therefore one of Q in $O(n^d)$. Thus letting $\mathcal{F}' := \tilde{\mathcal{F}} \cup \{Q\}$ finishes the proof. \square

We are now ready to see the proof of [Theorem 13.7](#).

Proof of Theorem 13.7. By [Proposition 13.1](#), with probability $1 - \frac{\beta}{3}$ we can find a polynomial factor $\tilde{\mathcal{F}}$ of degree $d - 1$ such that P is measurable in $\tilde{\mathcal{F}}$. Let $\gamma : \mathbb{N} \rightarrow \mathbb{R}^+$ be a decreasing function to be specified later. By [Lemma 13.4](#), with probability $1 - \frac{\beta}{3}$, we can refine $\tilde{\mathcal{F}}$ to a γ -uniform factor $\mathcal{F} = \{P_1, \dots, P_m\}$ of the same degree $d - 1$, with $\dim(\mathcal{F}) = O_{\gamma, \beta}(1)$. Since P is measurable in \mathcal{F} , there exists $\Gamma : \mathbb{F}^{\dim(\mathcal{F})} \rightarrow \mathbb{F}$ such that $P = \Gamma(\mathcal{F})$. Using the Fourier decomposition of $e_{\mathbb{F}}(\Gamma)$ we can write

$$f(x) := e_{\mathbb{F}}(P(x)) = \sum_{i=1}^L c_i e_{\mathbb{F}}(\langle \alpha^{(i)}, \mathcal{F} \rangle(x)), \tag{13.4}$$

where $L = |\mathbb{F}|^{\dim(\mathcal{F})} = O_{\gamma, \beta}(1)$, $\alpha^{(i)} \in \mathbb{F}^{\dim(\mathcal{F})}$, and

$$\langle \alpha^{(i)}, \mathcal{F} \rangle(x) := \sum_{j=1}^m \alpha_j^{(i)} \cdot P_j(x).$$

Notice that the terms in [\(13.4\)](#), unlike Fourier characters, are not orthogonal. But since the factor is γ -uniform, [Theorem 7.33](#) ensures approximate orthogonality. Let $Q_i := \langle \alpha^{(i)}, \mathcal{F} \rangle$. Choose $\gamma(u) \leq \frac{\sigma}{|\mathbb{F}|^{2u}}$, so that $\gamma(\dim(\mathcal{F})) \leq \frac{\sigma}{L^2}$, where $\sigma := \frac{\epsilon^{2k+1}}{4}$. It follows from the near orthogonality of the terms in [\(13.4\)](#) by [Theorem 7.33](#) that

$$|c_i - \langle f, e_{\mathbb{F}}(Q_i) \rangle| \leq \frac{\sigma}{L}, \tag{13.5}$$

and

$$\left| \|f\|_2^2 - \sum_{i=1}^L c_i^2 \right| \leq \sigma. \tag{13.6}$$

Claim 13.8. There exists $\delta'(\epsilon, |\mathbb{F}|) \in (0, 1]$ such that the following holds. Assume that f and \mathcal{F} are as above. Then there is $i \in [L]$, for which $\deg(Q_i) \leq k$ and $|\langle f, e_{\mathbb{F}}(Q_i) \rangle| \geq \delta'$.

Proof. We will induct on the degree of \mathcal{F} . Assume for the base case that \mathcal{F} is of degree k , i.e. $d = k + 1$. The following Cauchy-Schwarz inequality

$$\epsilon^{2k+1} \leq \|f\|_{\mathcal{U}^{k+1}}^{2k+1} \leq \|f\|_2^2 \|f\|_{\infty}^{2k+1-2}, \tag{13.7}$$

and (13.6) imply that there exists $i \in [L]$ such that $c_i^2 \geq \frac{\epsilon^{2^{k+1}} - \sigma}{L} = \frac{3\epsilon^{2^{k+1}}}{4L}$, which combined with (13.5) implies that $|\langle f, e_{\mathbb{F}}(Q_i) \rangle| \geq \frac{\epsilon^{2^{k+1}}}{2L}$.

Now for the induction step, assume that $d > k+1$. We will decompose (13.4) into two parts, first part consisting of the terms of degree $\leq k$ and the second part consisting of the terms of degree strictly higher than k . Namely, letting $S := \{i \in [L] : \deg(Q_i) \leq k\}$ we write $f = g + h$ where $g := \sum_{i \in S} c_i e_{\mathbb{F}}(Q_i)$ and $h := \sum_{i \in [L] \setminus S} c_i e_{\mathbb{F}}(Q_i)$. Notice that by the triangle inequality of Gowers norm, our choice of γ , and the fact that \mathcal{F} is γ -uniform

$$\|h\|_{U^{k+1}} \leq \sum_{i \in [L] \setminus S} |c_i| \cdot \|e_{\mathbb{F}}(Q_i)\|_{U^{k+1}} \leq L \cdot \frac{\epsilon^{2^{k+1}}}{4L^2} = \frac{\epsilon^{2^{k+1}}}{4L},$$

and thus

$$\|g\|_{U^{k+1}} \geq \frac{\epsilon}{2}.$$

Now the claim follows by the base case. □

Let $\delta'(\epsilon, |\mathbb{F}|)$ be as in the above claim. We will use the following theorem of [34] which gives an algorithm to find all the large Fourier coefficients of $e_{\mathbb{F}}(\Gamma)$.

Theorem 13.9 (Goldreich-Levin theorem [34]). Let $\zeta, \rho \in (0, 1]$. There is a randomized algorithm, which given oracle access to a function $\Gamma : \mathbb{F}^m \rightarrow \mathbb{F}$, runs in time $O(m^2 \log m \cdot \text{poly}(\frac{1}{\zeta}, \log(\frac{1}{\rho})))$ and outputs a decomposition

$$\Gamma = \sum_{i=1}^{\ell} b_i \cdot e_{\mathbb{F}}(\langle \eta_i, x \rangle) + \Gamma',$$

with the following guarantee:

- $\ell = O(\frac{1}{\zeta^2})$.
- $\Pr [\exists i : |b_i - \hat{\Gamma}(\eta_i)| > \zeta/2] \leq \rho$.
- $\Pr \left[\forall \alpha \text{ such that } |\hat{f}(\alpha)| \geq \zeta, \exists i \eta_i = \alpha \right] \geq 1 - \rho$.

We will use the above theorem with parameters $\zeta := \frac{\delta'}{2}$ and $\rho := \frac{\beta}{3}$. By Claim 13.8 there is $i \in [L]$ such that $\hat{\Gamma}(\alpha) = c_i \geq \frac{3\delta'}{4}$. With probability $1 - \frac{\beta}{6}$ there is j such that $\eta_j = \alpha_i$ and with probability at least $1 - \frac{\beta}{6}$, $|b_j - c_i| \leq \frac{\zeta}{2} \leq \frac{\delta'}{4}$, and therefore $c_i \geq \frac{\delta'}{2}$.

By a union bound, adding up the probabilities of the errors, with probability at least $1 - \beta$, we find Q_i such that

$$|\langle f, e_{\mathbb{F}}(Q_i) \rangle| \geq \frac{\delta'}{4}.$$

□

13.4 Derandomization via PRGs for polynomials

In this section we will discuss how all the algorithms that we have discussed above can be derandomized to efficient deterministic algorithms.

The key point here is that our only uses of randomness above have been in evaluating a collection of polynomials on set of inputs that are chosen uniformly at random from \mathbb{F}^n . We will use known efficient constructions of *pseudorandom generators* for polynomials [85, 57].

Definition 13.10. A distribution D on \mathbb{F}^n is said to ϵ -fool degree d polynomials in n variables over \mathbb{F}^n if for every degree d polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$,

$$|\mathbb{E}_D[e(P(D))] - \mathbb{E}_U[e(P(U))]| \leq \epsilon,$$

where $U \in \mathbb{F}^n$ is uniformly distributed.

The following lemma shows that a distribution that fools a single polynomial can be used to fool a collection of polynomials with a slightly worse error term.

Lemma 13.11. If D ϵ -fools degree d polynomials, then for every collection of C degree d polynomials P_1, \dots, P_C ,

$$\sum_{b_1, \dots, b_C \in \mathbb{F}^n} \left| \Pr_D[P_1(D) = b_1 \wedge \dots \wedge P_C(D) = b_C] - \Pr_U[P_1(U) = b_1 \wedge \dots \wedge P_C(U) = b_C] \right| \leq p^C \epsilon,$$

where $p = |\mathbb{F}|$.

We will use Viola’s explicit construction of pseudorandom generators for degree d polynomials.

Theorem 13.12 (Viola [85]). There is an explicit generator $g : \mathbb{F}^s \rightarrow \mathbb{F}^n$ with $s = d \log_p n + O(d \cdot 2^d \cdot \log(1/\epsilon))$, such that for uniform $Z \in \mathbb{F}^s$, $g(Z)$ ϵ -fools degree d polynomials.

Derandomized Bogdanov-Viola lemma. In the proof of [Lemma 13.1](#) we have used randomness in two steps.

First, we estimated μ_a in the statistical distance by sampling P on a constant number of randomly selected inputs. This step can be simply derandomized using [Theorem 13.12](#), by evaluating P on $g(z)$ for $z \in \mathbb{F}^s$. This requires $2^s = O(n^d)$ queries to P .

Second, we chose $h = (h_1, \dots, h_C) \in (\mathbb{F}^n)^C$ uniformly at random, and obtained a distribution μ_{obs} and \tilde{P}_h such that for every x

$$\Pr_{h_1, \dots, h_C} (\tilde{P}_h(x) \neq P(x)) \leq \Pr_h \left(\|\mu_{\text{obs}}^{(x)} - \mu_{P(x)}\| \geq \frac{\delta}{|\mathbb{F}|} \right) \leq \sigma\beta_2.$$

Suppose x is fixed, and consider polynomials $Q_i(h_1, \dots, h_C) := P(x + h_i)$. By [Lemma 13.11](#) and [Theorem 13.12](#), there is a map $g : \mathbb{F}^s \rightarrow (\mathbb{F}^n)^C$ such that

$$\Pr_{Z \in \mathbb{F}^s} (\tilde{P}_{g(Z)}(x) \neq P(x)) \leq \Pr_{Z \in \mathbb{F}^s} \left(\|\mu_{\text{obs}}^{(x)} - \mu_{P(x)}\| \geq \frac{\delta}{|\mathbb{F}|} \right) \leq \sigma\beta_2 + p^C \cdot \epsilon.$$

Therefore, choosing ϵ sufficiently small, there exists $Z \in \mathbb{F}^s$ such that

$$\Pr_x (\tilde{P}_{g(Z)}(x) \neq P(x)) \leq \sigma\beta_2 + p^C \cdot \epsilon \leq \sigma. \tag{13.8}$$

Thus we can iterate through all the $p^s = \text{Poly}(n)$ choices of Z , and check whether [Equation \(13.8\)](#) holds for that value of Z . Note that [Equation \(13.8\)](#) can itself be deterministically decided in $\text{Poly}(n)$ time using pseudorandom generators for low-degree polynomials.

Derandomized regularity lemmas. Having access to a deterministic Bogdanov-Viola lemma, consequently [Lemma 13.3](#), [Lemma 13.4](#), [Lemma 13.5](#), [Lemma 13.6](#), [Theorem 13.7](#) and [Proposition 13.1](#) can also be made deterministic. The key observation is that all these results use

their randomness only in the application of Bogdanov-Viola lemma and towards estimating bias or Gowers norm of a given polynomial. [Theorem 13.12](#) can be used to derandomize any step that requires estimating bias of a low-degree polynomial, and [Theorem 13.12](#) combined with [Lemma 13.11](#) can be used in order to estimate the Gowers norm of a low-degree polynomial.

13.5 Algorithmic Decomposition Theorems

Given a positive integer k , a vector of positive integers $\Delta = (\Delta_1, \Delta_2, \dots, \Delta_k)$ and a function $\Gamma : \mathbb{F}^k \rightarrow \mathbb{F}$, we say that a function $P : \mathbb{F}^n \rightarrow \mathbb{F}$ is (k, Δ, Γ) -structured if there exist polynomials $P_1, P_2, \dots, P_k : \mathbb{F}^n \rightarrow \mathbb{F}$ with each $\deg(P_i) \leq \Delta_i$ such that for all $x \in \mathbb{F}^n$,

$$P(x) = \Gamma(P_1(x), P_2(x), \dots, P_k(x)).$$

The polynomials P_1, \dots, P_k are said to form a (k, Δ, Γ) -decomposition (or simply, a polynomial decomposition). For instance, an n -variate polynomial over the field \mathbb{F} of total degree d factors non-trivially exactly when it is $(2, (d-1, d-1), \text{prod})$ -structured where $\text{prod}(a, b) = a \cdot b$. Later in [Chapter 16](#) we will see that the property of being (k, Δ, Γ) -structured is constant query testable with one-sided error even when P does not necessarily have bounded degree. However, the problem is harder if we ask for finding a decomposition realizing the (k, Δ, Γ) -structure.

Somewhat surprisingly, using the algorithmic regularity lemmas it can be shown that every degree-structural property can be decided in polynomial time.

Theorem 13.13 (Bhattacharyya [11]). For every finite field \mathbb{F} of prime order, positive integers $d < |\mathbb{F}|$, k , every vector of positive integers $\Delta = (\Delta_1, \Delta_2, \dots, \Delta_k)$ and every function $\Gamma : \mathbb{F}^k \rightarrow \mathbb{F}$, there is a deterministic algorithm $\mathcal{A}_{k, \Delta, \Gamma}$ that takes as input a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree d , runs in time polynomial in n , and outputs a (k, Δ, Γ) -decomposition of P if one exists while otherwise returning NO.

The algorithm is quite simple. Given a polynomial $P : \mathbb{F}^n \rightarrow \mathbb{F}$, we first use a corollary of [Lemma 13.4](#) and [Lemma 13.5](#) to write P as a function of a uniform polynomial factor $\{Q_1, \dots, Q_m\}$, i.e. $P(x) =$

$G(Q_1(x), \dots, Q_m(x))$ where $m = O(1)$ and $G : \mathbb{F}^m \rightarrow \mathbb{F}$. Now, the proof technique of Bhattacharyya [11]¹ shows that the only way P can have a (k, Δ, Γ) -decomposition is if there are functions $G_1, \dots, G_k : \mathbb{T}^m \rightarrow \mathbb{T}$ such that $G(z_1, \dots, z_m) = \Gamma(G_1(z_1, \dots, z_m), \dots, G_k(z_1, \dots, z_m))$ and also, for every $i \in [k]$, $G_i(Q_1(x), \dots, Q_m(x))$ is a polynomial of degree at most Δ_i . Since $m = O(1)$, there are only a constant number of possible G_1, \dots, G_k , and so the whole algorithm runs in polynomial time.

We will use the following corollary of Lemma 13.4 and Remark 7.19.

Theorem 13.14. Suppose $d < |\mathbb{F}|$, $\rho \in (0, 1)$ and $R : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ is a non-decreasing function. There is a function $C : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ and an efficient algorithm that takes as input a polynomial factor \mathcal{B} of degree $\leq d$ outputs an R -regular factor $\tilde{\mathcal{B}}$ where $\tilde{\mathcal{B}}$ is a refinement of \mathcal{B} , is of degree d , and $\|\tilde{\mathcal{B}}\| \leq C(|\mathcal{B}|)$. Additionally, if \mathcal{B} is defined by polynomials P_1, \dots, P_m , then we can find functions $\Gamma_1, \dots, \Gamma_m$ such that $P_i(x) = \Gamma_i(\tilde{\mathcal{B}}(x))$ for every $i \in [m]$.

Moreover, if \mathcal{B} itself is a syntactic refinement of some polynomial factor \mathcal{B}' of rank at least $R(|\mathcal{B}|) + 1$, then $\tilde{\mathcal{B}}$ will also be a syntactic refinement of \mathcal{B}' .

Proof of Theorem 13.13. Let $R : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ be chosen so that $R(m) = r(C(m+k)) + C(m+k) + |\mathbb{F}|$ for a function $r : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ to be fixed later, where C is as in Theorem 13.14. Applying Theorem 13.14 to the factor defined by the single polynomial P , we find an R -regular polynomial factor \mathcal{B} of degree d defined by polynomials $P_1, \dots, P_C : \mathbb{F}^n \rightarrow \mathbb{F}$ such that $P(x) = G(\mathcal{B}(x))$ for some $G : \mathbb{F}^C \rightarrow \mathbb{F}$. Note that here $C = O(1)$. Note that for small $n \leq Cd$, we can decide whether f is (k, Δ, Γ) -structured by brute force in $O(1)$ time, and further find such a decomposition if it exists.

Suppose $n > Cd$. From each P_i , pick a monomial m_i with degree equal to $\deg(P_i)$. Since $n > Cd$, there exists $i_0 \in [n]$ such that x_{i_0} does not appear in any of the m_i 's. Let \mathcal{B}' be the polynomial factor defined by polynomials P'_1, \dots, P'_C , which are respectively the restrictions of P_1, \dots, P_C to $x_{i_0} = 0$. Moreover, by Lemma 7.2, \mathcal{B}' is $(R - |\mathbb{F}|)$ -regular.

¹Bhattacharyya *et al.* [14] used a similar reasoning in their proof of testability of degree structural properties.

Recursively, decide (k, Δ, Γ) -structure for the polynomial $P' := P|_{x_{i_0}=0}$ on $n - 1$ variables. Note that

$$P'(x) = G(P'_1(x), \dots, P'_C(x)).$$

It is easy to see that if P' does not admit (k, Δ, Γ) -structure, then the same is true for P . So, in this case we will safely output NO. Assume otherwise that we have found

$$P'(x) = \Gamma(S_1(x), \dots, S_k(x)),$$

where $\deg(S_i) \leq \Delta_i$. We will show how in this case P is (k, Δ, Γ) -structured, and in fact we can find such a decomposition efficiently. We will use [Theorem 13.14](#) on the factor defined by $P'_1, \dots, P'_C, S_1, \dots, S_k$ to find a refinement \mathcal{B}' of rank at least $r(\mathcal{B}')$. Note that the setting of our parameters and [Theorem 13.14](#) guarantee that \mathcal{B}' is a syntactic refinement of P'_1, \dots, P'_C . That is \mathcal{B}' is defined by P'_1, \dots, P'_C and S'_1, \dots, S'_D where for each $i \in [k]$, $S_i(x) = G_i(P'_1(x), \dots, P'_C(x), S'_1(x), \dots, S'_D(x))$ for some function G_i . Thus we have for all x

$$\begin{aligned} G(P'_1(x), \dots, P'_C(x)) = \\ \Gamma(G_1(P'_1(x), \dots, P'_C(x), S'_1(x), \dots, S'_D(x)), \\ \dots, G_k(P'_1(x), \dots, P'_C(x), S'_1(x), \dots, S'_D(x))) \end{aligned}$$

Suppose r is set large enough so that by [Lemma 7.24](#) all the atoms of \mathcal{B}' are nonempty. This means that $\mathcal{B}'(x)$ acquires all possible values in its range \mathbb{F}^{C+D} . This allows us to deduce from the above equation that

$$\begin{aligned} G(a_1, \dots, a_C) \\ = \Gamma(G_1(a_1, \dots, a_C, b_1, \dots, b_D), \dots, G_k(a_1, \dots, a_C, b_1, \dots, b_D)), \end{aligned}$$

for all $a_1, \dots, a_C, b_1, \dots, b_D \in \mathbb{F}$. Define

$$Q_i(x) := G_i(P_1(x), \dots, P_C(x), 0, \dots, 0)$$

for $i \in [k]$. We will use [Corollary 16.6](#) which will be proved in [Chapter 16](#), to show that for each $i \in [k]$, $\deg(Q_i) \leq \deg(S_i) \leq \Delta_i$. Thus we have

$$P(x) = \Gamma(Q_1(x), \dots, Q_k(x)),$$

13.5. *Algorithmic Decomposition Theorems*

395

is a (k, Δ, Γ) decomposition of P .

The final algorithm is by recursively applying the above process until n is a constant, and building back the decomposition step by step. \square

Part III

Algebraic Property Testing

In [Part I](#) of this text we discussed linearity testing and more generally tests for being a polynomial of a given degree. These are all instances of “algebraic properties”, a general class that we are going to define shortly. It includes more complex properties such as being a product of two low-degree polynomials, or having sparse Fourier representation. In this part of the survey, we will use the tools developed in [Part II](#) to establish a general result showing that all algebraic properties that have local definitions are testable. We will focus on one-sided testable properties and for the treatment of the two-sided testable properties we refer the reader to Yoshida [[87](#), [86](#)], where nonstandard analysis is used to give a characterization.

14

Algebraic Properties

The study of property testing originated from the linearity test of Blum, Luby, and Rubinfeld [20] and was formally coined by Rubinfeld and Sudan [69] and Goldreich *et al.* [32]. A substantial treatment of testable combinatorial properties due to Goldreich *et al.* [32] was later followed by seminal subsequent work that lead to an in-depth understanding of testable graph properties [2, 4, 22, 5, 30]. There has also been a great deal of success in testing algebraic properties including testing Reed-Muller codes (in other words, bounded degree polynomials) [6, 7, 28, 69, 3], testing Fourier sparsity [36], testing BCH codes [53], and testing of functions satisfying functional equations [67]. Kaufman and Sudan [56] noted affine-invariance as a common characteristic of majority of the studied algebraic properties, and initiated the fruitful study of testability of affine-invariant properties [13, 73, 16, 15, 14].

14.1 Affine and linear invariance

Recall that $\mathbb{F} = \mathbb{F}_p$ for a fixed prime p . Let $[R] = \{1, \dots, R\}$ be a constant sized set. A property of functions $f : \mathbb{F}^n \rightarrow [R]$ is simply a subset of all such functions, namely $\mathcal{P} \subseteq \{\mathbb{F}^n \rightarrow [R]\}$. Typically, we assume that such a property is defined for all $n \geq 1$.

Note that a generic property can completely ignore the algebraic structure of \mathbb{F}^n and treat it as an arbitrary set of size $|\mathbb{F}|^n$. Hence, any notion of “algebraic property” must require some assumptions on \mathcal{P} that take the algebraic structure of the set \mathbb{F}^n into account. This is typically done by requiring that the property \mathcal{P} is invariant under certain algebraic transformations. Indeed this is analogous to other combinatorial settings such as graph properties, or hypergraph properties, where the properties are assumed to be invariant under permutation of the vertices.

We say that a property $\mathcal{P} \subseteq \{\mathbb{F}^n \rightarrow [R]\}$ is *linear-invariant* if it is the case that for any $f \in \mathcal{P}$ and for any linear transformation $L : \mathbb{F}^n \rightarrow \mathbb{F}^n$, it holds that $f \circ L \in \mathcal{P}$, where $(f \circ L)(x) = f(L(x))$. Similarly, an *affine-invariant* property is closed under composition with affine transformation $A : \mathbb{F}^n \rightarrow \mathbb{F}^n$ (an affine transformation A is of the form $A(x) = Lx + c$ where L is linear and c is a constant). Both linear invariance and affine invariance are very natural candidates for defining algebraic properties.

The property of a function $f : \mathbb{F}^n \rightarrow \mathbb{F}$ being affine is testable by a simple reduction to linearity testing [20], and is itself affine-invariant. Other well-studied examples of affine-invariant (and hence, linear-invariant) properties include Reed-Muller codes (in other words, bounded degree polynomials) [6, 7, 28, 69, 3] and Fourier sparsity [36]. In fact, affine invariance seems to be a common feature of most interesting properties that one would classify as algebraic. Kaufman and Sudan [56] made explicit note of this phenomenon and initiated a general study of the testability of affine-invariant properties. See also [33]. In particular, they asked for necessary and sufficient conditions for the testability of affine-invariant properties.

14.2 Locally characterized properties

Let $\mathcal{P} \subseteq \{\mathbb{F}^n \rightarrow [R]\}$. Let us first assume that \mathcal{P} is testable in the strongest possible way, and see what this entails. To recall, we say that \mathcal{P} is one-sided proximity-oblivious q -query testable if there is a local test which makes q queries to a function $f : \mathbb{F}^n \rightarrow [R]$, always accepts if $f \in \mathcal{P}$ and otherwise rejects f with probability related to the distance of f from \mathcal{P} . See Definition 2.1 for a more formal definition.

So, assume \mathcal{P} is one-sided proximity-oblivious q -query testable. Every such test has the following structure:

- Step 1: Randomly pick points $(x_1, \dots, x_q) \in (\mathbb{F}^n)^q$ according to some joint distribution μ .
- Step 2: Query the values of $f(x_1), \dots, f(x_q)$ and accordingly output “ $f \in \mathcal{P}$ ” or “ $f \notin \mathcal{P}$ ”.

Suppose that the queries result in $(f(x_1), \dots, f(x_q)) = \sigma \in [R]^q$. Since the algorithm is only allowed to make a one-sided error, if there is a function $g \in \mathcal{P}$ with $(g(x_1), \dots, g(x_q)) = \sigma$, then the algorithm must return “ $f \in \mathcal{P}$ ”. On the other hand, if there is no such function $g \in \mathcal{P}$, then it can safely return “ $f \notin \mathcal{P}$ ”. In such a case we say that $\mathcal{C} = (x_1, x_2, \dots, x_q; \sigma)$ forms a q -local constraint for \mathcal{P} (equivalently, a q -local witness for non-membership in \mathcal{P}), as if $f \in \mathcal{P}$ then necessarily $(f(x_1), \dots, f(x_q)) \neq \sigma$.

Definition 14.1 (Local constraint). A q -local constraint for a property $\mathcal{P} \subset \{\mathbb{F}^n \rightarrow [R]\}$ is $\mathcal{C} = (x_1, \dots, x_q; \sigma)$, where $x_1, \dots, x_q \in \mathbb{F}^n$, $\sigma \in [R]^q$ such that for all $g \in \mathcal{P}$ it holds that $(g(x_1), \dots, g(x_q)) \neq \sigma$.

Let $\mathcal{C} = (x_1, \dots, x_q; \sigma)$ be a local constraint for \mathcal{P} . If $f : \mathbb{F}^n \rightarrow [R]$ satisfies that $(f(x_1), \dots, f(x_q)) = \sigma$, then we say that f violates the constraint \mathcal{C} . In particular, we know that $f \notin \mathcal{P}$. Let $\mathcal{C}_1, \dots, \mathcal{C}_m$ be the set of all the q -local constraints for the above test. One can rewrite the above test as in the following:

- Step 1: Randomly pick points $(x_1, \dots, x_q) \in (\mathbb{F}^n)^q$ according to some joint distribution μ .
- Step 2: If $(x_1, \dots, x_q; f(x_1), \dots, f(x_q))$ equals some \mathcal{C}_i then output “ $f \notin \mathcal{P}$ ”. Otherwise output “ $f \in \mathcal{P}$ ”.

Note that if $f \notin \mathcal{P}$, then according to [Definition 2.1](#), the tester must reject f with positive probability. Consequently, there must exist at least one q -local constraint among $\mathcal{C}_1, \dots, \mathcal{C}_m$ that is violated by f . This provides a “local characterization” for \mathcal{P} . Namely,

$$\mathcal{P} = \{f : \mathbb{F}^n \rightarrow [R] \text{ that do not violate any of } \mathcal{C}_1, \dots, \mathcal{C}_m\}.$$

The following definition formalizes this notion.

Definition 14.2 (Locally characterized property). A property $\mathcal{P} \subset \{F^n \rightarrow [R]\}$ is q -locally characterized if there exist q -local constraints $\mathcal{C}_1, \dots, \mathcal{C}_m$ such that

$$\mathcal{P} = \{f : \mathbb{F}^n \rightarrow [R] \text{ that do not violate any of } \mathcal{C}_1, \dots, \mathcal{C}_m\}.$$

We say \mathcal{P} is *locally characterized* if it is q -locally characterized for some constant q . It follows from the above discussion that if \mathcal{P} is one-sided proximity-oblivious q -query testable then it is also q -locally characterized.

We now give some examples of locally characterized affine-invariant properties. Consider the property of $f : \mathbb{F}^n \rightarrow \mathbb{F}$ being an affine function (namely, a degree 1 polynomial). It is 4-locally characterized because a function f is affine if and only if $f(x) - f(x+y) - f(x+z) + f(x+y+z) = 0$ for every $x, y, z \in \mathbb{F}^n$. Note that this characterization naturally suggests a 4-query test: pick random $x, y, z \in \mathbb{F}^n$ and check whether the identity holds or not for that choice of x, y, z . As we saw in [Chapter 2](#), this is indeed a local test for the property of affine functions (we proved it for $\mathbb{F} = \mathbb{F}_2$ in [Theorem 2.3](#), and the proof can be extended to any finite field).

More generally, consider the property of being a polynomial of degree at most d , for some fixed integer $d > 0$. If $f : \mathbb{F}^n \rightarrow \mathbb{F}$ has degree $\leq d$, then it is annihilated by taking $d + 1$ directional derivatives (See [Section 3.1](#)). This implies that this property is also locally characterizable. Independent papers by Kaufman and Ron [55] and Jutla *et al.* [52] has obtained the optimal value of q for general fields \mathbb{F}_p : the property of being a degree $\leq d$ polynomial is q -locally characterized for $q = p^{\lceil \frac{d+1}{p-1} \rceil}$. These papers also showed that the property is testable. Again, the test is simply to pick a random constraint and check if it is violated.

Indeed, for any q -locally characterized property \mathcal{P} defined by constraints $\mathcal{C}_1, \dots, \mathcal{C}_m$, one can design the following q -query test: choose a constraint \mathcal{C}_i uniformly at random and reject only if the input function violates \mathcal{C}_i . Clearly, if the input function f is in \mathcal{P} , the test always accepts. The main challenge is in showing that if f is ε -far from \mathcal{P} , then this test rejects f with some probability $\delta = \delta(\varepsilon) > 0$. Equivalently

put, f is ε -far from \mathcal{P} then at least δm of the local constraints are violated. This was conjectured by Kaufman and Sudan [56] and proved in a sequence of papers [13, 73, 16, 15, 14].

Theorem 14.3 (Bhattacharyya *et al.* [14]). Every q -locally characterized affine-invariant property is proximity obviously testable with q queries.

The proof of [Theorem 14.3](#) is presented in [Chapter 15](#).

14.3 Locality of affine invariant properties via linear forms

In the context of linear-invariant and affine-invariant properties, we can define the notion of local characterization in a more algebraic way. As most of the research has focused on affine invariant properties we also limit our discussion to such properties. Many of the observations below naturally extend to linear invariant properties.

Consider an affine-invariant property $\mathcal{P} \subset \{\mathbb{F}^n \rightarrow [R]\}$ that is locally characterized by a collection $\mathcal{C} = \{\mathcal{C}_1, \dots, \mathcal{C}_m\}$ of q -local constraints. As \mathcal{P} is affine-invariant, one can assume that \mathcal{C} is also invariant under affine transformations. In other words, if $(a_1, a_2, \dots, a_q; \sigma) \in \mathcal{C}$, then also $(Aa_1, Aa_2, \dots, Aa_q; \sigma) \in \mathcal{C}$ for every affine transformation $A : \mathbb{F}^n \rightarrow \mathbb{F}^n$.

This can be rephrased using linear forms. Assume that $a_1, \dots, a_q \in \mathbb{F}^n$ span a linear space of dimension $r \leq q$, and let $b_1, \dots, b_r \in \mathbb{F}^n$ be a basis for this subspace. Let $\{\lambda_{i,j} \in \mathbb{F} : i \in [q], j \in [r]\}$ be such that $a_i = \sum_{j=1}^r \lambda_{i,j} b_j$. The assumption that \mathcal{P} is affine invariant means that for every linear map $L : \mathbb{F}^n \rightarrow \mathbb{F}^n$ and every $c \in \mathbb{F}^n$, we also have that $(La_1 + c, \dots, La_q + c; \sigma) \in \mathcal{C}$. Let $L_1, \dots, L_q \in \mathbb{F}^{r+1}$ be linear forms given by $L_i = (1, \lambda_{i,1}, \dots, \lambda_{i,r})$. Then the following are also q -local constraints for \mathcal{P} :

$$(L_1(x_0, x_1, \dots, x_r), \dots, L_q(x_0, \dots, x_r); \sigma) \quad \forall x_0, \dots, x_r \in \mathbb{F}^n.$$

The linear forms L_1, \dots, L_q are *affine forms*, where we recall the definition that a linear form is affine if its first coordinate is 1.

Next, we formalize this notion of affine local constraints.

Definition 14.4 (Affine local constraints). An *affine constraint of size q on k variables* is a tuple $A = (L_1, \dots, L_q; \sigma)$ of q affine forms $L_1, \dots, L_q \in \mathbb{F}^k$ and $\sigma \in [R]^q$.

An affine constraint can be used to define affine invariant properties in the following way.

Definition 14.5 (Properties defined by affine constraints). Consider an affine constraint (A, σ) of size q on k variables. A function $f : \mathbb{F}^n \rightarrow [R]$ is said to be (A, σ) -free if there exist no $x_1, \dots, x_k \in \mathbb{F}^n$ such that

$$(f(L_1(x_1, \dots, x_k)), \dots, f(L_q(x_1, \dots, x_k))) = \sigma.$$

On the other hand, if such x_1, \dots, x_k exist, we say that f induces (A, σ) at x_1, \dots, x_k .

Given a (possibly infinite) collection $\mathcal{A} = \{(A^1, \sigma^1), (A^2, \sigma^2), \dots\}$ of affine constraints, a function $f : \mathbb{F}^n \rightarrow [R]$ is said to be \mathcal{A} -free if it is (A^i, σ^i) -free for every $i \geq 1$.

The above discussion shows that an affine-invariant property is q -locally characterized if and only if it can be described using a bounded number of affine constraints of size q .

14.4 Subspace hereditary properties

Just as a necessary condition for proximity oblivious testability with one-sided error is local characterization, one can formulate a natural condition that is (almost) necessary for testability in general. In the context of affine-invariant properties, the condition can be succinctly stated as follows. We will assume throughout that \mathcal{P} is a property of functions $f : \mathbb{F}^n \rightarrow [R]$ for all $n \geq 1$.

Definition 14.6 (Subspace hereditary properties). An affine-invariant property \mathcal{P} is said to be *affine-subspace hereditary* if for any $f : \mathbb{F}^n \rightarrow [R]$ satisfying \mathcal{P} , the restriction of f to any affine subspace of \mathbb{F}^n also satisfies \mathcal{P} .

Bhattacharyya *et al.* [16] showed that every affine-invariant property testable by a “natural” tester is very “close” to a subspace hereditary property¹. Thus, if we gloss over some technicalities, subspace heredi-

¹We omit the technical definitions of “natural” and “close”, since they are unimportant here. Informally, the behavior of a “natural” tester is independent of the size of the domain and “close” means that the property deviates from an actual affine subspace hereditary property on functions over a finite domain. See [16] for details, or [4] for the analogous definitions in a graph-theoretic context.

tariness is a necessary condition for testability. In the opposite direction, Bhattacharyya *et al.* [16] conjecture the following:

Conjecture 14.7 (Bhattacharyya *et al.* [16]). Every affine-subspace hereditary property is testable with one-sided error.

Resolving [Conjecture 14.7](#) would yield a combinatorial *characterization* of the (natural) one-sided testable affine-invariant properties, similar to the characterization for testable dense graph properties [4].

Affine subspace hereditariness and affine constraints are related through the following simple observation.

Observation 14.8. An affine-invariant property \mathcal{P} is subspace hereditary if and only if it is equivalent to the property of \mathcal{A} -freeness for some fixed collection \mathcal{A} of affine constraints (possibly infinite).

Proof. Given an affine invariant subspace hereditary property \mathcal{P} , a simple (though inefficient) way to obtain the set \mathcal{A} is the following. For every $m \geq 1$, every function $g : \mathbb{F}^m \rightarrow [R]$ that is not in \mathcal{P} , and an m -dimensional subspace V of \mathbb{F}^n , include the constraint that the restriction of $f : \mathbb{F}^n \rightarrow [R]$ to V is not equal to g .

This can be expressed as an affine constraint. For every $a \in \mathbb{F}^m$ let $L_a \in \mathbb{F}^{m+1}$ be an affine form given by $L_a = (1, a_1, \dots, a_m)$. The affine constraint we add is

$$\mathcal{C}_g := ((L_a : a \in \mathbb{F}^m); (g(a) : a \in \mathbb{F}^m)).$$

In particular, g is not \mathcal{C}_g -free, as g induces \mathcal{C}_g at $x_1 = e_1, \dots, x_m = e_m$, where e_i is the i -th unit vector. Hence the property defined by \mathcal{A} is contained in \mathcal{P} . The containment in the other direction follows from \mathcal{P} being affine-invariant and hereditary.

The other direction of the observation is trivial. □

[Conjecture 14.7](#) is not yet confirmed or refuted, however Bhattacharyya *et al.* [14] prove testability under an additional assumption of “bounded complexity”. Define the *Cauchy-Schwarz complexity* (see [Definition 11.1](#)) of an affine constraint (A, σ) to be simply the Cauchy-Schwarz complexity of A . Let the Cauchy-Schwarz complexity of a collection $\mathcal{A} = \{(A^1, \sigma^1), (A^2, \sigma^2), \dots\}$ of affine constraints to be the

maximum of Cauchy-Schwarz complexity of the constraints (A^i, σ^i) . We will see a proof of the following theorem in [Chapter 15](#).

Theorem 14.9 (Bhattacharyya *et al.* [14]). Every subspace hereditary property of bounded Cauchy-Schwarz complexity is testable with one-sided error.

All natural affine-invariant properties that we know of have bounded complexity (in fact, most are locally characterized). However, testing the subspace hereditary properties not covered by [Theorem 14.9](#) is of theoretical interest.

14.5 Locality dimension

The notion of q -local characterization uses the number of queries to measure the complexity of the characterization. In the algebraic setting of affine-invariant properties it is as natural to consider the dimension of the affine subspace containing those queries.

Definition 14.10 (Locality dimension). The *locality dimension* of an affine-invariant property $\mathcal{P} \subseteq \{\mathbb{F}^n \rightarrow [R]\}$ is the smallest K such that the following holds. There exists a collection $\mathcal{F} \subset \{\mathbb{F}^K \rightarrow [R]\}$ such that for $f : \mathbb{F}^n \rightarrow [R]$, $f \in \mathcal{P}$ if and only if $f|_V \notin \mathcal{F}$ for all K -dimensional affine subspaces $V \subseteq \mathbb{F}^n$.

As discussed above, every q -locally characterized \mathcal{P} is equivalent to \mathcal{A} -freeness, where \mathcal{A} is a collection of affine constraints with each constraint on at most $q + 1$ variables. Obviously one can extend this to include all the affine linear forms on these variables, or equivalent query f on the q -dimensional affine subspace containing these $q + 1$ variables. Thus every q -locally characterized property has locality dimension of at most q . In the other direction, if \mathcal{P} is of locality dimension K , then it is equivalent to \mathcal{A} -freeness, where \mathcal{A} is a finite collection of affine constraints, with each constraint of size $|\mathbb{F}|^K$ on $K + 1$ variables.

The notion of locality dimension is even more natural for affine-subspace hereditary properties.

Observation 14.11 (Locality dimension for hereditary properties). The locality dimension of a hereditary affine-invariant property $\mathcal{P} \subseteq \{\mathbb{F}^n \rightarrow [R]\}$ is the smallest K such that for $f : \mathbb{F}^n \rightarrow [R]$, $f \in \mathcal{P}$ if and only if $f|_V \in \mathcal{P}$ for all K -dimensional affine subspaces $V \subseteq \mathbb{F}^n$.

15

One-Sided Algebraic Property Testing

In this Chapter we establish [Theorem 14.3](#) and [Theorem 14.9](#). That is, we show that every q -locally characterized affine-invariant property is proximity obviously testable with one-sided error using q queries, and more generally we show that every affine invariant subspace hereditary property of bounded Cauchy-Schwarz complexity is testable with one-sided error.

15.1 Proof overview

Before delving into the technical details let us first present an overview of the proof. For simplicity, assume for now that \mathcal{A} consists only of a single affine constraint (A, σ) where A is the tuple of affine linear forms (L_1, \dots, L_m) , each over ℓ variables, and $\sigma \in [R]^m$. Our goal is to show that, when f is ε -far from being (A, σ) -free, then f contains many instances of violation of (A, σ) -freeness. The decomposition theorems presented in [Chapter 9](#) allow us to find a polynomial factor partition of the domain \mathbb{F}^n such that, for the purpose of counting the number of violations of (A, σ) freeness, f looks uniform in most of the parts. If the notion of ε -farness would have allowed to remove points from the domain of f , then the proof would have become much simpler.

Let us first consider this simpler case. Suppose that if remove any given ε -fraction of points from \mathbb{F}^n , there are still copies of (A, σ) in f . Now we look at the polynomial factor decomposition of f and remove all the points that are not in the uniform cells. We also remove all the points that have unpopular values in the uniform cells. That is for every cell, if there is a value that appears on very few points in that cell, then we remove all the points from that cell with that value. Now since the number of non-uniform cells is small, and also the number of unpopular points is small (by their definition), we have removed only few points from the domain. Hence according the modified notion of ε -farness, there are still copies of (A, σ) remaining in f . Now every point in any such copy must belong to a uniform cell and its value is not unpopular in that cell. It follows from these two properties that there are many such copies of (A, σ) in f .

Unfortunately, the natural notion of ε -farness does not allow removing points from the domain of f , and instead one is only allowed to change the value of f on at most ε -fraction of the points. This makes proving the theorem considerably more difficult. Now one has to appeal to the more technical decomposition theorems such as [Theorem 9.8](#). This decomposition theorem, roughly speaking, shows the existence of a polynomial factor partition \mathcal{B} of the domain such that for every atom c of \mathcal{B} , there is a sub-atom c' that comes from a finer partition \mathcal{B}' such that most sub-atoms provide good statistical samples of the demography of the atom that contains them. Now the most popular values inside sub-atoms will play an important role, and instead of removing points from the domain we will change the value of f to such values. In other words, unpopular values and mis-represented values inside every atom are changed to the most popular value inside the corresponding sub-atom.

Again, the number of changes will be small (less than ε -fraction of points). So the new function will still contain at least one copy of (A, σ) . Note that the values in this copy are abundant in the corresponding sub-atoms as we changed the unpopular values to the most popular values in the sub-atoms. Then, a key property of sub-atoms (i.e. the fact that s is fixed in [Theorem 9.8](#)) and the uniformity of sub-atoms allows us to find many copies of (A, σ) inside the corresponding sub-atoms.

15.2 Big picture functions

Suppose we have a function $f : \mathbb{F}^n \rightarrow [R]$, and we want to find out whether it induces a particular affine constraint (A, σ) , where $A = (L_1, \dots, L_m)$ is a sequence of affine forms on ℓ variables and $\sigma \in [R]^m$. Now, suppose \mathbb{F}^n is partitioned by a polynomial factor \mathcal{B} defined by nonclassical polynomials P_1, \dots, P_C of degrees d_1, \dots, d_C and depths k_1, \dots, k_C , respectively. Then, observe that if $b_1, \dots, b_m \in \mathbb{T}^C$ denote the atoms of \mathcal{B} containing $L_1(x_1, \dots, x_\ell), \dots, L_m(x_1, \dots, x_\ell)$ respectively, it must be the case that b_1, \dots, b_m are \mathcal{B} -consistent with A (as defined in [Definition 7.37](#)). Thus, to locate where f might induce (A, σ) , we should restrict our search to sequences of atoms consistent with A .

It will be convenient to “blur” the given function f so as to retain only atom-level information about it. That is, for every atom c of \mathcal{B} , we will define $f_{\mathcal{B}}(c) \subseteq [R]$ to be the set of all values that f takes within c . We denote by $\mathcal{P}([R])$ the power set of $[R]$.

Definition 15.1. Given a function $f : \mathbb{F}^n \rightarrow [R]$ and a polynomial factor \mathcal{B} , the *big picture function* of f is the function $f_{\mathcal{B}} : \mathbb{T}^{|\mathcal{B}|} \rightarrow \mathcal{P}([R])$, defined by $f_{\mathcal{B}}(c) = \{f(x) : \mathcal{B}(x) = c\}$.

On the other hand, given any function $g : \mathbb{T}^C \rightarrow \mathcal{P}([R])$, and a vector of degrees $\mathbf{d} = (d_1, \dots, d_C)$ and depths $\mathbf{k} = (k_1, \dots, k_C)$ (which we think of as corresponding to the degrees and depths of some polynomial factor of complexity C), we will define what it means for such a function to “induce” a copy of a given constraint.

Definition 15.2 (Partially induce). Suppose we are given $\mathbf{d} = (d_1, \dots, d_C) \in \mathbb{Z}_{>0}^C$, $\mathbf{k} = (k_1, \dots, k_C) \in \mathbb{Z}_{\geq 0}^C$, $g : \prod_{i \in [C]} \mathbb{U}_{k_i+1} \rightarrow \mathcal{P}([R])$, and an affine constraint (A, σ) of size m . We say that g *partially* (\mathbf{d}, \mathbf{k}) -induces (A, σ) if there exist a sequence $b_1, \dots, b_m \in \mathbb{T}^C$ such that

- (i) (b_1, \dots, b_m) is (\mathbf{d}, \mathbf{k}) -consistent with A (see [Definition 7.37](#)).
- (ii) $\sigma_j \in g(b_j)$ for each $j \in [m]$.

[Definition 15.2](#) is justified by the following trivial observation.

Observation 15.3. If $f : \mathbb{F}^n \rightarrow [R]$ induces a constraint (A, σ) , then for a factor \mathcal{B} defined by polynomials of respective degrees $(d_1, \dots, d_{|\mathcal{B}|}) = \mathbf{d}$ and respective depths $(k_1, \dots, k_{|\mathcal{B}|}) = \mathbf{k}$, the big picture function $f_{\mathcal{B}}$ partially (\mathbf{d}, \mathbf{k}) -induces (A, σ) .

To handle a possibly infinite collection \mathcal{A} of affine constraints (in order to prove [Theorem 14.9](#)), we will employ a compactness argument, analogous to one used by Alon and Shapira [\[5\]](#) in the context of graph properties, to bound the size of the constraint partially induced by the big picture function. Let us make the following definition:

Definition 15.4 (The compactness function). Suppose we are given positive integers C and d , and a possibly infinite collection of affine constraints $\mathcal{A} = \{(A^1, \sigma^1), (A^2, \sigma^2), \dots\}$, where (A^i, σ^i) is of size m_i . For fixed $\mathbf{d} = (d_1, \dots, d_C) \in [d]^C$ and $\mathbf{k} = (k_1, \dots, k_C) \in \left[0, \left\lfloor \frac{d-1}{p-1} \right\rfloor\right]^C$, denote by $\mathcal{G}(\mathbf{d}, \mathbf{k})$ the set of functions $g : \prod_{i=1}^C \mathbb{U}_{k_i+1} \rightarrow \mathcal{P}([R])$ that partially (\mathbf{d}, \mathbf{k}) -induce some $(A^i, \sigma^i) \in \mathcal{A}$. The compactness function is defined as

$$\Psi_{\mathcal{A}}(C, d) = \max_{\mathbf{d}, \mathbf{k}} \max_{g \in \mathcal{G}(\mathbf{d}, \mathbf{k})} \min_{\substack{(A^i, \sigma^i) \text{ partially} \\ (\mathbf{d}, \mathbf{k})\text{-induced by } g}} m_i$$

where the outer max is over vectors $\mathbf{d} = (d_1, \dots, d_C) \in [d]^C$ and $\mathbf{k} = (k_1, \dots, k_C) \in \left[0, \left\lfloor \frac{d-1}{p-1} \right\rfloor\right]^C$. Whenever $\mathcal{G}(\mathbf{d}, \mathbf{k})$ is empty, we set the corresponding maximum to 0.

Note that $\Psi_{\mathcal{A}}(C, d)$ is indeed finite, as the number of possible degree and depth sequences are bounded by d^{2C} , and the size of $\mathcal{G}(\mathbf{d}, \mathbf{k})$ is bounded by $2^{Rp^{dC}}$.

Remark 15.5. Note that if a function $g : \mathbb{T}^C \rightarrow \mathcal{P}([R])$ partially (\mathbf{d}, \mathbf{k}) -induces some constraint from \mathcal{A} where $\mathbf{d} \in [d]^C$, then g must belong to $\mathcal{G}(\mathbf{d}, \mathbf{k})$, and consequently it will necessarily partially induce some $(A^i, \sigma^i) \in \mathcal{A}$ whose size is at most $\Psi_{\mathcal{A}}(C, d)$.

15.3 Proof of testability

In this section we establish our main result [Theorem 14.9](#) which in turn implies [Theorem 14.3](#). It suffices to prove the following.

Theorem 15.6. Let $d > 0$ be an integer. Suppose we are given a possibly infinite collection of affine constraints $\mathcal{A} = \{(A^1, \sigma^1), (A^2, \sigma^2), \dots\}$ where each (A^i, σ^i) is an affine constraint of Cauchy-Schwarz complexity $\leq d$, and of size m_i on ℓ_i variables. Then, there are functions $\ell_{\mathcal{A}} : (0, 1) \rightarrow \mathbb{Z}_{>0}$ and $\delta_{\mathcal{A}} : (0, 1) \rightarrow (0, 1)$ such that the following is true. For every $\varepsilon \in (0, 1)$, if a function $f : \mathbb{F}^n \rightarrow [R]$ is ε -far from being \mathcal{A} -free, then f induces at least $\delta_{\mathcal{A}}(\varepsilon)|\mathbb{F}|^{n\ell_i}$ many copies of some (A^i, σ^i) with $\ell_i < \ell_{\mathcal{A}}(\varepsilon)$.

Moreover, if \mathcal{A} is q -locally characterized, then $\ell_{\mathcal{A}}(\varepsilon) \leq q$ and hence is bounded by a constant independent of ε .

[Theorem 14.9](#) follow immediately. Given \mathcal{A} as in the statement of the theorems, let $f : \mathbb{F}^n \rightarrow [R]$ which is ε -far from \mathcal{A} . Let $\mathcal{A}_0 \subset \mathcal{A}$ be the subset of all $(A^i, \sigma^i) \in \mathcal{A}$ with $\ell_i \leq \ell_{\mathcal{A}}(\varepsilon)$. Observe that \mathcal{A}_0 is a finite set, as we may assume that $m_i \leq p^{\ell_i}$.

Consider the following test. Pick a uniform $(A^i, \sigma^i) \in \mathcal{A}_0$, pick uniformly $X = (x_1, \dots, x_{\ell_i}) \in \mathbb{F}^{n\ell_i}$ and test if $(f(L_1(X)), \dots, f(L_{m_i}(X))) = \sigma^i$. By [Theorem 15.6](#), if f is ε -far from \mathcal{A} -freeness, this test rejects with probability at least $|\mathcal{A}_0|^{-1}\delta_{\mathcal{A}}(\varepsilon)$. For [Theorem 14.3](#), note that if \mathcal{A} is q -locally characterized, then the test is a one-sided proximity oblivious q -local test. Below, we prove [Theorem 15.6](#).

Fix a function $f : \mathbb{F}^n \rightarrow [R]$ that is ε -far from being \mathcal{A} -free. For $i \in [R]$, define $f^{(i)} : \mathbb{F}^n \rightarrow \{0, 1\}$ so that $f^{(i)}(x)$ equals 1 when $f(x) = i$ and equals 0 otherwise. Additionally, set the following parameters, where $\Psi_{\mathcal{A}}$ is the compactness function from [Definition 15.4](#):

$$\begin{aligned} \zeta &= \frac{\varepsilon}{8R}, \\ \alpha(C) &= p^{-2dC\Psi_{\mathcal{A}}(C,d)}, & r(C) &= r_{7.14}(\mathbb{F}, d, \alpha(C)) \\ \Delta(C) &= \frac{1}{16}\zeta^{\Psi_{\mathcal{A}}(C,d)}, & \eta(C) &= \frac{1}{8p^{dC\Psi_{\mathcal{A}}(C,d)}} \left(\frac{\varepsilon}{24R}\right)^{\Psi_{\mathcal{A}}(C,d)}. \end{aligned}$$

Intuitively, think of ζ as a small enough constant (depending only on ε, R), of $r(\cdot)$ as a large enough rank, and of $\alpha(\cdot), \Delta(\cdot), \eta(\cdot)$ as taking small enough positive values.

Decomposing by regular factors. As a first step, we apply (the multi-function generalization) of [Theorem 9.8](#) to the functions $f^{(1)}, f^{(2)}, \dots, f^{(R)}$. By the theorem, there exists a polynomial factor \mathcal{B} of degree d , and a

refinement of it \mathcal{B}' of degree d , both r -regular of complexity at most $C_{9.8}(\mathbb{F}, d, r, \eta, \Delta, \zeta)$, such that the following holds. We can decompose each $f^{(i)}$ as

$$f^{(i)} = f_1^{(i)} + f_2^{(i)} + f_3^{(i)}$$

where

- (i) $f_1^{(i)} = \mathbb{E}[f|\mathcal{B}']$.
- (ii) $\|f_2^{(i)}\|_{U^{d+1}} < \eta(|\mathcal{B}'|)$.
- (iii) $\|f_3^{(i)}\|_2 \leq \Delta(|\mathcal{B}|)$.
- (iv) f_1 and $f_1 + f_3$ take values in $[0, 1]$; and f_2, f_3 take values in $[-1, 1]$.
- (v) \mathcal{B}' ζ -represents \mathcal{B} with respect to f .

Moreover, assume that the atoms of \mathcal{B}' are indexed by (c, s) with $c \in \mathbb{T}^{|\mathcal{B}|}, s \in \mathbb{T}^{|\mathcal{B}'|-|\mathcal{B}|}$. Then exists a choice of $s \in \mathbb{T}^{|\mathcal{B}'|-|\mathcal{B}|}$ for which the following is true:

- (vi) For every atom c of \mathcal{B} , the sub-atom (c, s) of \mathcal{B}' satisfies that

$$\mathbb{E} \left[|f_3^{(i)}(x)|^2 \mid \mathcal{B}'(x) = (c, s) \right] \leq \Delta(|\mathcal{B}|)^2.$$

- (vii) For at most a ζ fraction of atoms c in \mathcal{B} it holds that

$$\left| \mathbb{E}[f^{(i)}|c] - \mathbb{E}[f^{(i)}|(c, s)] \right| > \zeta.$$

We denote the sequence of polynomials defining \mathcal{B}' by $P_1, \dots, P_{|\mathcal{B}'|}$. Denote the degree of P_i by d_i and the depth of P_i by k_i . Since \mathcal{B}' is a syntactic refinement of \mathcal{B} , we may assume \mathcal{B} is generated by the polynomials $P_1, \dots, P_{|\mathcal{B}|}$. We denote $C = |\mathcal{B}|$ and $C' = |\mathcal{B}'|$.

Note that $\|\mathcal{B}\| < p^{(k_{\max}+1)C} \leq p^{dC}$, where $k_{\max} \leq \lfloor (d-1)/(p-1) \rfloor$ is the maximum depth of a polynomial in \mathcal{B} .

Cleanup. Based on \mathcal{B}' and \mathcal{B} , we construct a function $F : \mathbb{F}^n \rightarrow [R]$ that is $\frac{\zeta}{2}$ -close to f and hence, still violates \mathcal{A} -freeness. The “cleaner” structure of F will help us locate the induced constraint violated by f .

The function F is the same as f except for the following. For every atom c of \mathcal{B} , let $t_c = \arg \max_{j \in [R]} \Pr[f(x) = j \mid \mathcal{B}'(x) = (c, s)]$ be the most popular value inside the corresponding subatom (c, s) .

- **Poorly-represented atoms:** If there exists $i \in [R]$ such that $|\Pr[f(x) = i \mid \mathcal{B}(x) = c] - \Pr[f(x) = i \mid \mathcal{B}'(x) = (c, s)]| > \zeta$, then set $F(z) = t_c$ for every z in the atom c .
- **Unpopular values:** For any $i \in [R]$ such that $\Pr_x[f(x) = i \mid \mathcal{B}'(x) = (c, s)] < \zeta$, if z in the atom c satisfies $f(z) = i$, then set $F(z) = t_c$.
- For all other z in the atom c , set $F(z) = f(z)$.

A key property of the cleanup function F is that it supports a value inside an atom c of \mathcal{B} only if the original function f acquires the value on at least an ζ fraction of the subatom (c, s) . Furthermore as the following lemma shows it is $\varepsilon/2$ -close to f , and therefore as f is ε -far from \mathcal{A} -freeness, we have that F is not \mathcal{A} -free.

Lemma 15.7. The cleanup function F is $\varepsilon/2$ -close to f , and therefore, it is not \mathcal{A} -free.

Proof. The first step (fixing poorly represented atoms) applies to at most $\zeta R \|\mathcal{B}\|$ atoms, since \mathcal{B}' ζ -represents \mathcal{B} with respect to each $f^{(1)}, \dots, f^{(R)}$. By Lemma 7.24, each atom of \mathcal{B} occupies at most $\frac{1}{\|\mathcal{B}\|} + \alpha(C)$ fraction of the entire domain. So, the fraction of points whose values are set in the first step is at most $\zeta R \|\mathcal{B}\| (\frac{1}{\|\mathcal{B}\|} + \alpha(C)) < 2\zeta R = \varepsilon/4$.

In the second step (changing unpopular values), if $\Pr[f(x) = i \mid \mathcal{B}'(x) = (c, s)] < \zeta$, then $\Pr[f(x) = i \mid \mathcal{B}(x) = c] < \Pr[f(x) = i \mid \mathcal{B}'(x) = (c, s)] + \zeta < 2\zeta$. Hence, the fraction of the points z whose values are set in the second step is at most $2\zeta R = \varepsilon/4$.

Thus, the distance of F from f is bounded by $\varepsilon/2$. □

Locating a violated constraint. We now want to use F to “locate” a popular affine constraint induced in f . Setting $\mathbf{d} = (d_1, \dots, d_C)$ and $\mathbf{k} = (k_1, \dots, k_C)$, we have by Observation 15.3 that the big picture function $F_{\mathcal{B}}$ of F will partially (\mathbf{d}, \mathbf{k}) -induce some constraint from \mathcal{A} , and hence by Remark 15.5, it will partially (\mathbf{d}, \mathbf{k}) -induce some $(A, \sigma) \in \mathcal{A}$ of size $m \leq \Psi_{\mathcal{A}}(C, d)$ on ℓ variables. We will show that the original function f violates many instances of this constraint.

Denote the affine forms in A by (L_1, \dots, L_m) and the vector σ by $(\sigma_1, \dots, \sigma_m)$. Since we can assume $\ell \leq m$ (without loss of generality by making a change of variables), we can now define

$$\ell_{\mathcal{A}}(\varepsilon) = \Psi_{\mathcal{A}}(C_{9.8}(\mathbb{F}, d, r, \eta, \Delta, \zeta), d). \tag{15.1}$$

Let $b_1, \dots, b_m \in \prod_{i=1}^C \mathbb{U}_{k_i+1}$ correspond to the atoms of \mathcal{B} where (A, σ) is partially (\mathbf{d}, \mathbf{k}) -induced by $F_{\mathcal{B}}$. That is, b_1, \dots, b_m are consistent with A , and $\sigma_i \in F_{\mathcal{B}}(b_i)$ for every $i \in [m]$. Also, let $b'_1, \dots, b'_m \in \prod_{i=1}^{C'} \mathbb{U}_{k_i+1}$ index the associated subatoms in \mathcal{B}' , obtained by letting $b'_j = (b_j, s)$ for every $j \in [m]$.

Lemma 15.8. The subatoms b'_1, \dots, b'_m are consistent with A .

Proof. Since b_1, \dots, b_m are already consistent with A , we only need to show that for every $i \in [C + 1, C']$, the sequence $(b'_{1,i}, \dots, b'_{m,i}) = (s_{i-C}, s_{i-C}, \dots, s_{i-C})$ is (d_i, k_i) -consistent. That is, we need to show that there exists a homogeneous nonclassical polynomial P of degree d and depth k , and a choice of $x_1, \dots, x_{\ell} \in \mathbb{F}^n$, such that $P(L_j(x_1, \dots, x_{\ell})) = s_{i-C}$ for all $j \in [m]$. This is where we use the assumption that the linear forms are affine. Take $x_2 = \dots = x_{\ell} = 0$. Then $L_j(x_1, 0, \dots, 0) = x_1$. So we only need to show the existence of P as above and of $x_1 \in \mathbb{F}^n$ for which $P(x_1) = s_{i-C}$. But this clearly holds, say by taking $P = P_i$ and x_1 any value in the (c, s) subatom. \square

The main analysis. Let $X = (x_1, \dots, x_{\ell})$ where $x_1, \dots, x_{\ell} \in \mathbb{F}^n$ are independently and uniformly chosen. Our goal is to prove a lower bound on

$$\begin{aligned} \Pr_X [f(L_1(X)) = \sigma_1 \wedge \dots \wedge f(L_m(X)) = \sigma_m] \\ = \mathbb{E}_X \left[f^{(\sigma_1)}(L_1(X)) \dots f^{(\sigma_m)}(L_m(X)) \right]. \end{aligned} \tag{15.2}$$

Theorem 15.6 follows if the above expectation is larger than the respective $\delta_{\mathcal{A}}(\varepsilon)$. We rewrite the expectation as

$$\begin{aligned} (15.2) = \mathbb{E}_X \left[(f_1^{(\sigma_1)} + f_2^{(\sigma_1)} + f_3^{(\sigma_1)})(L_1(X)) \right. \\ \left. \dots (f_1^{(\sigma_m)} + f_2^{(\sigma_m)} + f_3^{(\sigma_m)})(L_m(X)) \right]. \end{aligned} \tag{15.3}$$

We can expand the expression inside the expectation as a sum of 3^m terms. The expectation of any term involving $f_2^{(\sigma_j)}$ for any $j \in [m]$ is bounded in magnitude by $\|f_2^{(\sigma_j)}\|_{U^{d+1}} \leq \eta(|\mathcal{B}'|)$, by Lemma 11.2 and the assumption that the Cauchy-Schwarz complexity of A is bounded by d . Hence, the expression (15.3) is at least

$$(15.3) \geq \mathbb{E}_X \left[(f_1^{(\sigma_1)} + f_3^{(\sigma_1)})(L_1(X)) \cdots (f_1^{(\sigma_m)} + f_3^{(\sigma_m)})(L_m(X)) \right] - 3^m \eta(|\mathcal{B}'|).$$

Next, because of the non-negativity of $f_1^{(\sigma_j)} + f_3^{(\sigma_j)}$ for every $j \in [m]$, we may further require that any event holds. We will require that $\mathcal{B}'(L_j(X)) = b'_j$ for all j . So

$$(15.3) \geq \mathbb{E}_X \left[(f_1^{(\sigma_1)} + f_3^{(\sigma_1)})(L_1(X)) \cdots (f_1^{(\sigma_m)} + f_3^{(\sigma_m)})(L_m(X)) \cdot \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(X))=b'_j]} \right] - 3^m \eta(|\mathcal{B}'|), \quad (15.4)$$

where $1_{[\mathcal{B}'(L_j(X))=b'_j]}$ is the indicator function of the event $\mathcal{B}'(L_j(X)) = b'_j$. In other words, now we are only counting patterns that arise from the selected subatoms b'_1, \dots, b'_m .

Next, expand the product inside the expectation into 2^m terms. We will show that the contribution from each of the $2^m - 1$ terms involving $f_3^{(\sigma_k)}$ for any $k \in [m]$ is small. Each such term is trivially bounded from above by

$$\mathbb{E}_X \left[\left| f_3^{(\sigma_k)}(L_k(X)) \right| \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(X))=b'_j]} \right]. \quad (15.5)$$

Here, we used fact that $|f_1^{(i)}(x)| \leq 1$ for all $i \in [R]$, which follows from (i) in our construction. The next lemma bounds the expression in (15.5). Let Λ_i denote the (d_i, k_i) -dependency set of L_1, \dots, L_m for $i \in [C']$.

Lemma 15.9.

$$\mathbb{E}_X \left[\left| f_3^{(\sigma_k)}(L_k(X)) \right| \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(X))=b'_j]} \right] \leq 2\Delta(C) \frac{\prod_{i \in [C']} |\Lambda_i|}{\|\mathcal{B}'\|^m}.$$

Proof. In order to simplify the proof, apply a change of basis on L_1, \dots, L_m so that $L_k = e_1$. Namely, $L_k(X) = x_1$. To simplify notation let us also denote $x = x_1$ and $\mathbf{x} = (x_2, \dots, x_\ell)$.

As a first step, we apply the Cauchy-Schwartz inequality and obtain

$$\begin{aligned} & \left(\mathbb{E}_{X \in (\mathbb{F}^n)^\ell} \left[\left| f_3^{(\sigma_k)}(x_1) \right| \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(X))=b'_j]} \right] \right)^2 \\ &= \left(\mathbb{E}_{x \in \mathbb{F}^n, \mathbf{x} \in (\mathbb{F}^n)^{\ell-1}} \left[\left| f_3^{(\sigma_k)}(x) \right| 1_{[\mathcal{B}'(x)=b'_k]} \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(x, \mathbf{x}))=b'_j]} \right] \right)^2 \\ &\leq \mathbb{E}_x \left[\left| f_3^{(\sigma_k)}(x) \right|^2 1_{[\mathcal{B}'(x)=b'_k]} \right] \mathbb{E}_x \left(\mathbb{E}_{\mathbf{x}} \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(x, \mathbf{x}))=b'_j]} \right)^2. \end{aligned} \tag{15.6}$$

We first bound the first term in the right hand side. By (vi) in our construction and [Lemma 7.24](#), we have

$$\begin{aligned} & \mathbb{E}_x \left[\left| f_3^{(\sigma_k)}(x) \right|^2 1_{[\mathcal{B}'(x)=b'_k]} \right] \\ &= \mathbb{E}_x \left[\left| f_3^{(\sigma_k)}(x) \right|^2 \mid \mathcal{B}'(x) = b'_k \right] \Pr[\mathcal{B}'(x) = b'_k] \\ &\leq \Delta^2(C) \Pr_x[\mathcal{B}'(x) = b'_k] \leq \Delta^2(C) \left(\frac{1}{\|\mathcal{B}'\|} + \alpha(C') \right) \leq \frac{2\Delta^2(C)}{\|\mathcal{B}'\|}. \end{aligned} \tag{15.7}$$

The second term in the right hand side of [\(15.6\)](#) is equal to

$$\begin{aligned} & \mathbb{E}_x \left(\mathbb{E}_{\mathbf{x}} \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(x, \mathbf{x}))=b'_j]} \right)^2 \\ &= \frac{1}{\|\mathcal{B}'\|^{2m}} \mathbb{E}_x \left[\left(\mathbb{E}_{\mathbf{x}} \prod_{\substack{i \in [C'] \\ j \in [m]}} \frac{1}{p^{k_i+1}} \sum_{\lambda_{i,j}=0}^{p^{k_i+1}-1} e \left(\lambda_{i,j} \cdot (P_i(L_j(x, \mathbf{x})) - b'_{i,j}) \right) \right)^2 \right] \\ &= \frac{1}{\|\mathcal{B}'\|^{2m}} \mathbb{E}_x \left[\left(\sum_{(\lambda_{i,j}) \in \prod_{i,j} [0, p^{k_i+1}-1]} e \left(- \sum_{\substack{i \in [C'] \\ j \in [m]}} \lambda_{i,j} b'_{i,j} \right) \right)^2 \right] \end{aligned}$$

$$\begin{aligned}
 & \cdot \mathbb{E}_{\mathbf{x}} e \left(\sum_{\substack{i \in [C'] \\ j \in [m]}} \lambda_{i,j} P_i(L_j(x, \mathbf{x})) \right) \Bigg]^2 \\
 \leq & \frac{1}{\|\mathcal{B}'\|^{2m}} \sum_{\substack{(\lambda_{i,j}), (\tau_{i,j}) \in \\ \prod_{i,j} [0, p^{k_i+1}-1]}} \left| \mathbb{E}_{x, \mathbf{x}, \mathbf{y}} \left[e \left(\sum_{\substack{i \in [C'] \\ j \in [m]}} \lambda_{i,j} P_i(L_j(x, \mathbf{x})) \right) \right. \right. \\
 & \left. \left. \cdot e \left(- \sum_{\substack{i \in [C'] \\ j \in [m]}} \tau_{i,j} P_i(L_j(x, \mathbf{y})) \right) \right] \right|. \tag{15.8}
 \end{aligned}$$

Here, $\mathbf{y} = (y_2, \dots, y_\ell)$ where $y_2, \dots, y_\ell \in \mathbb{F}^n$ are new independent uniform random variables.

We can bound the above using [Theorem 7.32](#). Let A' denote the set of $2m$ linear forms: $\{L_j(x_1, x_2, \dots, x_\ell) \mid j \in [m]\} \cup \{L_j(x_1, y_2, \dots, y_\ell) \mid j \in [m]\}$ in variables $x_1, \dots, x_\ell, y_2, \dots, y_\ell$. Let Λ'_i denote the (d_i, k_i) -dependency set of A' .

Applying [Theorem 7.32](#) (just as in the proof of [Theorem 7.38](#)), we get that

$$(15.8) \leq \frac{\prod_{i \in [C']} |\Lambda'_i|}{\|\mathcal{B}'\|^{2m}} + \alpha(C') \leq \frac{2 \prod_{i \in [C']} |\Lambda'_i|}{\|\mathcal{B}'\|^{2m}}. \tag{15.9}$$

Thus, the next step is to compute $|\Lambda'_i|$.

Claim 15.10. For each $i \in [C']$ it holds that $|\Lambda'_i| = |\Lambda_i|^2 \cdot p^{k_i+1}$.

Proof. Recall that by our initial change of basis, $L_k(x, \mathbf{x}) = L_k(x, \mathbf{y}) = x$. For any $\lambda, \tau \in \Lambda_i$ and any $\alpha \in \mathbb{Z}_p^{k_i+1}$, note that $(\lambda_1 + \alpha, \lambda_2, \dots, \lambda_m, \tau_1 - \alpha, \tau_2, \dots, \tau_m) \in \Lambda'_i$. Hence, $|\Lambda'_i| \geq |\Lambda_i|^2 \cdot p^{k_i+1}$. To show $|\Lambda'_i| \leq |\Lambda_i|^2 \cdot p^{k_i+1}$, we give a map from Λ'_i to $\Lambda_i \times \Lambda_i$ that is p^{k_i+1} -to-1.

Suppose that

$$\sum_{j=1}^m \lambda_j Q(L_j(x_1, x_2, \dots, x_\ell)) + \sum_{j=1}^m \tau_j Q(L_j(x_1, y_2, \dots, y_\ell)) \equiv 0$$

for every nonclassical homogeneous polynomial Q of degree d_i and depth k_i . Setting $x_2 = \dots = x_\ell = 0$ shows that

$$\sum_{j=1}^m \tau_j Q(L_j(x_1, y_2, \dots, y_\ell)) = - \left(\sum_{j=1}^m \lambda_j \right) Q(x_1),$$

which implies that

$$\left(- \sum_{j=2}^m \lambda_j, \lambda_2, \dots, \lambda_m \right) \in \Lambda_i.$$

Similarly, setting $y_2 = \dots = y_\ell = 0$ shows that

$$\sum_{j=1}^m \lambda_j Q(L_j(x_1, x_2, \dots, x_\ell)) = - \left(\sum_{j=1}^m \tau_j \right) Q(x_1),$$

which implies that

$$\left(- \sum_{j=2}^m \tau_j, \tau_2, \dots, \tau_m \right) \in \Lambda_i.$$

Consequently,

$$(\lambda, \tau) \mapsto \left(\left(- \sum_{j=2}^m \lambda_j, \lambda_2, \dots, \lambda_m \right), \left(- \sum_{j=2}^m \tau_j, \tau_2, \dots, \tau_m \right) \right)$$

is a map from Λ'_i to $\Lambda_i \times \Lambda_i$. To see that it is p^{k_i+1} -to-1, note that if $(\lambda_1, \dots, \lambda_m, \tau_1, \dots, \tau_m) \in \Lambda'_i$ then also

$$(\lambda_1 + \gamma, \lambda_2, \dots, \lambda_m, \tau_1 - \gamma, \tau_2, \dots, \tau_m) \in \Lambda'_i$$

for every $\gamma \in \mathbb{Z}_{p^{k_i+1}}$, and that these p^{k_i+1} elements are all mapped to the same element in $\Lambda_i \times \Lambda_i$ by our map. □

Thus

$$(15.9) \leq \frac{2 \prod_{i \in [C']} |\Lambda_i|^2 p^{k_i+1}}{\|\mathcal{B}'\|^{2m}} = \frac{2 \prod_{i \in [C']} |\Lambda_i|^2}{\|\mathcal{B}'\|^{2m-1}}.$$

Combining this with Equation (15.7) and Equation (15.6), we obtain

$$(15.6) \leq 4\Delta^2(C) \frac{\prod_{i \in [C']} |\Lambda_i|^2}{\|\mathcal{B}'\|^{2m}}. \tag{15.10}$$

This concludes the proof of the lemma. □

Finally, we turn to the main term in the expansion of Equation (15.4). We know from Lemma 15.8 that the subatoms b'_1, \dots, b'_m are consistent with A . Thus

$$\begin{aligned} & \mathbb{E}_X \left[f_1^{(\sigma_1)}(L_1(X)) \cdots f_1^{(\sigma_m)}(L_m(X)) \cdot \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(X))=b'_j]} \right] \\ &= \Pr_X[\mathcal{B}'(L_1(X)) = b'_1 \wedge \cdots \wedge \mathcal{B}'(L_m(X)) = b'_m] \\ & \quad \mathbb{E}_X \left[f_1^{(\sigma_1)}(L_1(X)) \cdots f_1^{(\sigma_m)}(L_m(X)) \mid \forall j \in [m], \mathcal{B}'(L_j(X)) = b'_j \right] \\ & \geq \left(\frac{\prod_{i=1}^{C'} |\Lambda_i|}{\|\mathcal{B}'\|^m} - \alpha(C') \right) \zeta^m. \end{aligned} \tag{15.11}$$

Let us justify the last line. The first term is due to the lower bound on the probability from Theorem 7.38. The second term in (15.11) follows since each $f_1^{(\sigma_j)}$ is constant on the atoms of \mathcal{B}' , and because by construction, the big picture function $F_{\mathcal{B}}$ of the cleanup function F , on which (A, σ) was partially induced, supports a value inside an atom b of \mathcal{B} only if the original function f acquires the value on at least an ζ fraction of the subatom (c, s) . By our choice of $\alpha(C')$, we can further deduce that

$$\mathbb{E}_X \left[f_1^{(\sigma_1)}(L_1(X)) \cdots f_1^{(\sigma_m)}(L_m(X)) \cdot \prod_{j \in [m]} 1_{[\mathcal{B}'(L_j(X))=b'_j]} \right] \geq \frac{\prod_{i=1}^{C'} |\Lambda_i|}{2\|\mathcal{B}'\|^m}. \tag{15.12}$$

Setting $\beta = \|\mathcal{B}'\|^{-m} \cdot \prod_{i=1}^{C'} |\Lambda_i|$ and combining the bounds from (15.4), Lemma 15.9 and (15.12), we conclude

$$(15.2) \geq \beta \left(\frac{1}{2} \left(\frac{\varepsilon}{8R} \right)^m - 2^{m+1} \Delta(C) \right) - 3^m \cdot \eta(C').$$

To complete the proof, we need to specify our choices of parameters. We have $\|\mathcal{B}'\| \leq p^{dC'}$, and $m \leq \Psi_{\mathcal{A}}(C, d)$. Thus, $\beta \geq \|\mathcal{B}'\|^{-\Psi_{\mathcal{A}}(C, d)}$. We choose $\Delta(C) = \frac{1}{16} \left(\frac{\varepsilon}{8R} \right)^{\Psi_{\mathcal{A}}(d, C)}$, $\eta(C') < \frac{1}{8\|\mathcal{B}'\|^{\Psi_{\mathcal{A}}(C, d)}} \left(\frac{\varepsilon}{24R} \right)^{\Psi_{\mathcal{A}}(C, d)}$, and both C and C' are upper-bounded by $C_{9.8}(\mathbb{F}, d, r, \eta, \Delta, \zeta)$. Finally, we conclude by setting

$$\delta_{\mathcal{A}}(\varepsilon) = \frac{1}{4} p^{-d\Psi_{\mathcal{A}}(C_{9.8}(\Delta, \eta, \rho, \zeta, R))} C_{9.8}(\Delta, \eta, \rho, \zeta, R) \cdot \left(\frac{\varepsilon}{8R} \right)^{\Psi_{\mathcal{A}}(C_{9.8}(\Delta, \eta, \rho, \zeta, R), d)}.$$

16

Degree Structural Properties

[Theorem 14.3](#) shows that every locally characterizable property is proximity-oblivious testable with one-sided error. The condition of being locally characterizable is quite general, and as a result, we expect that there are many such interesting algebraic properties. This, in fact, turns out to be the case. We show that a class of properties that we call *degree-structural* are all locally characterized and are, hence, proximity-obliviously testable by [Theorem 14.3](#).

Before giving the formal definition, let us first list some examples of degree-structural properties. Let d be a fixed positive integer, $\mathbb{F} = \mathbb{F}_p$ a fixed prime finite field. Each of the following definitions defines a degree-structural property.

- **Degree $\leq d$:** All polynomials $F : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree $\leq d$;
- **Product of linear forms:** Polynomials $F : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree $\leq d$ which are the product of at most d linear functions;
- **Composite polynomials:** Polynomials $F : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree $\leq d$ which factor as $F = GH$ where G, H are polynomials of degree $1 \leq \deg(G), \deg(H) \leq d - 1$;

- **Having square root:** Polynomials $F : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree $\leq d$ which can be factored as $F = G^2$ for a polynomial G of degree $\leq d/2$;
- **Sum of two products:** Polynomials $F : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree $\leq d$ which can be decomposed as $F = G_1G_2 + G_3G_4$, where $1 \leq \deg(G_i) \leq d - 1$ for all $i \in \{1, 2, 3, 4\}$;
- **Low d -rank:** For fixed integers $r \geq 1, R \geq 2$, a function $F : \mathbb{F}^n \rightarrow [R]$ has d -rank at most r if there exist classical polynomials $G_1, \dots, G_r : \mathbb{F}^n \rightarrow \mathbb{F}$ of degree $\leq d - 1$ and a function $\Gamma : \mathbb{F}^r \rightarrow [R]$ such that $F = \Gamma(G_1, \dots, G_r)$.

In fact, roughly speaking, any property that can be described as the property of being decomposable into a known structure of low-degree polynomials is degree-structural. The following definition formalizes this notion.

Definition 16.1 (Degree-structural property). Let $\mathbb{F} = \mathbb{F}_p$ be a prime finite field and let $R \geq 2$. Given an integer $c \geq 1$, a vector of non-negative integers $\mathbf{d} = (d_1, \dots, d_c) \in \mathbb{Z}_{\geq 0}^c$, and a function $\Gamma : \mathbb{F}^c \rightarrow [R]$, define the (c, \mathbf{d}, Γ) -structured property to be the collection of functions $F : \mathbb{F}^n \rightarrow [R]$ for which there exist polynomials $P_1, \dots, P_c : \mathbb{F}^n \rightarrow \mathbb{F}$ of degrees $\deg(P_i) \leq d_i$ for all $i \in [c]$, such that $F(x) = \Gamma(P_1(x), \dots, P_c(x))$ for all $x \in \mathbb{F}^n$.

We say a property $\mathcal{P} \subseteq \{F : \mathbb{F}^n \rightarrow [R]\}$ is *degree-structural* if there exist integers $c, d \geq 1$ and a set of tuples

$$S \subset \{(c, \mathbf{d}, \Gamma) \mid \mathbf{d} \in [0, d]^c, \Gamma : \mathbb{F}^c \rightarrow [R]\},$$

such that a function $F \in \mathcal{P}$ if and only if F is (c, \mathbf{d}, Γ) -structured for some $(c, \mathbf{d}, \Gamma) \in S$. We call R the *range*, c the *scope* and d the *max-degree* of the degree-structural property \mathcal{P} .

Remark 16.2. We could allow for an even more general definition of degree-structural properties, by allowing the polynomials P_i above to be nonclassical with some bound on the depth. Everything below can be extended to this setting as well.

It is straightforward to see that the examples described above all satisfy this definition. By definition, any degree structural property is affine invariant. In this section, we present a result of Bhattacharyya *et al.* [14] that shows that any degree-structural property with bounded scope and max-degree has a local characterization.

Theorem 16.3. Fix a prime finite field \mathbb{F} . Every degree-structural property \mathcal{P} with range R , scope c and max-degree d is q -locally characterized for some $q = q(\mathbb{F}, R, c, d)$.

An immediate corollary of [Theorem 16.3](#) and [Theorem 14.3](#) is that any degree-structural property is proximity-oblivious locally testable.

Corollary 16.4. Fix a prime finite field \mathbb{F} . Every degree-structural property \mathcal{P} with range R , scope c and max-degree d is q -locally testable for some $q = q(\mathbb{F}, R, c, d)$.

The proof of [Theorem 16.3](#) will rely on [Theorem 7.32](#). In the original paper, Bhattacharyya *et al.* [14], the proof utilized a special version of [Theorem 7.32](#) for affine constraints, as these are the systems of linear forms which arise in affine invariant property testing. Here, we present a slightly different version, which avoids the need to introduce this specialized form of [Theorem 7.32](#).

16.1 Proof of [Theorem 16.3](#)

We proceed to give the proof of [Theorem 16.3](#). The proof would utilize the following lemma due to Bhattacharyya *et al.* [12], which shows that locally characterized properties cannot distinguish between polynomials of high rank, as long as they have the same degree and depth.

Lemma 16.5. Let $\mathbb{F} = \mathbb{F}_p$ be a prime finite field, and fix $d, c, q \geq 1$. Let \mathcal{P} be a q -locally characterized property of functions $\mathbb{F}^n \rightarrow [R]$. Let \mathcal{B} be a polynomial factor of rank $> r_{16.5}(\mathbb{F}, d, c, q)$ defined by homogeneous nonclassical polynomials $P_1, \dots, P_c : \mathbb{F}^n \rightarrow \mathbb{T}$ of degrees $\leq d$, and define $F(x) = \Gamma(P_1(x), \dots, P_c(x))$ for some function $\Gamma : \mathbb{T}^c \rightarrow [R]$.

Assume that $F \in \mathcal{P}$. Then, for any homogeneous polynomials $Q_1, \dots, Q_c : \mathbb{F}^n \rightarrow \mathbb{T}$ for which $\deg(Q_i) = \deg(P_i)$ and $\text{depth}(Q_i) =$

depth(P_i), it holds that the function $G(x) = \Gamma(Q_1(x), \dots, Q_c(x))$ is also in \mathcal{P} .

Proof. Assume that $G \notin \mathcal{P}$. As \mathcal{P} is q -locally characterized, there must be q points $x_1, \dots, x_q \in \mathbb{F}^n$ such that the values of $G(x_1), \dots, G(x_q)$ witness the fact that $G \notin \mathcal{P}$. We will show that there exists points $x'_1, \dots, x'_q \in \mathbb{F}^n$ which witness the fact that $F \notin \mathcal{P}$, a contradiction.

Let e denote the dimension of the linear subspace spanned by x_1, \dots, x_q , and assume without loss of generality that x_1, \dots, x_e are linearly independent. Let $L_1, \dots, L_q : \mathbb{F}^e \rightarrow \mathbb{F}$ be linear forms such that $x_i = L_i(x_1, \dots, x_e)$. For every Q_i , we have that $Q_i(L_1(x_1, \dots, x_e)), \dots, Q_i(L_q(x_1, \dots, x_e))$ are (d_i, k_i) -consistent with L_1, \dots, L_m , where $d_i = \deg(P_i) = \deg(Q_i)$ and $k_i = \text{depth}(P_i) = \text{depth}(Q_i)$ (see Definition 7.37). Theorem 7.38 then implies that, as long as \mathcal{B} has high enough rank (concretely, rank at least $r_{7.10}(\mathbb{F}, d, \varepsilon)$ for $\varepsilon = 1/2|\mathcal{B}|^m$) then $\left((P_i(L_j(X))) : X \in (\mathbb{F}^n)^e \right)$ attain all values in \mathbb{T}^{mc} which are (\mathbf{d}, \mathbf{k}) -consistent with L_1, \dots, L_m , where $\mathbf{d} = (d_1, \dots, d_c)$ and $\mathbf{k} = (k_1, \dots, k_c)$. In particular, there are some $x'_1, \dots, x'_e \in \mathbb{F}^n$ such that $P_i(L_j(x'_1, \dots, x'_e)) = Q_i(L_j(x_1, \dots, x_e))$ for all $i \in [c], j \in [m]$. This implies $F(x_i) = G(x_i)$ for all $i \in [q]$, which in turn shows $F \notin \mathcal{P}$. \square

As a specific corollary, we obtain that under the definitions of Lemma 16.5, if $F : \mathbb{F}^n \rightarrow \mathbb{F}$ is a polynomial of degree D then $\deg(G) \leq D$.

Corollary 16.6. Let $\mathbb{F} = \mathbb{F}_p$ be a prime finite field, and fix $c, d \geq 1$. Let \mathcal{B} be a polynomial factor of rank $> r_{16.6}(\mathbb{F}, d, c)$ defined by homogeneous nonclassical polynomials $P_1, \dots, P_c : \mathbb{F}^n \rightarrow \mathbb{T}$ of degrees $\leq d$, and define $F(x) = \Gamma(P_1(x), \dots, P_c(x))$ for some function $\Gamma : \mathbb{T}^c \rightarrow \mathbb{F}$.

Assume that $\deg(F) = D$. Then, for any homogeneous polynomials $Q_1, \dots, Q_c : \mathbb{F}^n \rightarrow \mathbb{T}$ for which $\deg(Q_i) = \deg(P_i)$ and $\text{depth}(Q_i) = \text{depth}(P_i)$, it holds that the function $G(x) = \Gamma(Q_1(x), \dots, Q_c(x))$ has $\deg(G) \leq D$.

Proof. The corollary follows as the property of being a polynomial of degree $\leq D$ is q -locally characterized for $q \leq 2^D$. Moreover, if

$\deg(F) = D$ and F is a function of c nonclassical polynomials of degree $\leq d$, then $D \leq dcp^{\lceil d/(p-1) \rceil}$. \square

Proof of Theorem 16.3. Let \mathcal{P} be a degree-structural property with range R , scope c and max-degree d . Denote by S the set of tuples (c, \mathbf{d}, Γ) such that \mathcal{P} is the union over all $(c, \mathbf{d}, \Gamma) \in S$ of (c, \mathbf{d}, Γ) -structured functions. It is clear that \mathcal{P} is affine-invariant, as having degree bounded by some d_i is an affine-invariant property. It is also immediate that \mathcal{P} is closed under taking restrictions to subspaces, since if F is (c, \mathbf{d}, Γ) -structured, then F restricted to any hyperplane is also (c, \mathbf{d}, Γ) -structured. The non-trivial part of the theorem is to show that the locality dimension is bounded. In other words, we need to show that there is a constant $K = K(\mathbb{F}, R, c, d)$, such that for $n \geq K$, if $F : \mathbb{F}^n \rightarrow [R]$ is a function with $F|_H \in \mathcal{P}$ for every K -dimensional subspace $H \leq \mathbb{F}^n$, then this implies that $F \in \mathcal{P}$. To do that, it suffices (by induction) to prove a weaker statement: for $n \geq K$, if $F : \mathbb{F}^n \rightarrow [R]$ is a function with $F|_H \in \mathcal{P}$ for every hyperplane (namely, an $(n - 1)$ -dimensional subspace) $H \leq \mathbb{F}^n$, then this implies that $F \in \mathcal{P}$.

For each $t \in [R]$ let $F_t : \mathbb{F}^n \rightarrow \{0, 1\} \subset \mathbb{F}$ be given by $F_t(x) = \mathbb{1}_{F(x)=t}$. The first step will be bound the degrees of F_t . Recall that we assume that $F|_H \in \mathcal{P}$ for every hyperplane H . Thus

$$F|_H(x) = \Gamma_H(P_{H,1}(x), \dots, P_{H,c}(x))$$

for some polynomials $P_{H,i}$ of degrees $\deg(P_{H,i}) \leq d_i \leq d$ and some $\Gamma_H : \mathbb{F}^c \rightarrow [R]$. This implies that $(F_t)|_H(x) = \Gamma_{H,t}(P_{H,1}(x), \dots, P_{H,c}(x))$ where $\Gamma_{H,t}(z) = \mathbb{1}_{\Gamma_H(z)=t}$. In particular, $\deg((F_t)|_H) \leq D := |\mathbb{F}|cd$ for all $t \in [R]$. As the property of being low degree is locally characterized, by making sure that $K > D$ we obtain that also $\deg(F_t) \leq D$.

Let $r_1 : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ be a function to be determined later. Define $r_2 : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ so that $r_2(\cdot) > r_1(C_{7.30}^{(\mathbb{F}, r, D)}(\cdot + c)) + C_{7.30}^{(\mathbb{F}, r, D)}(\cdot + c) + |\mathbb{F}|$. Apply Theorem 7.30 to the polynomial factor $\{F_1, \dots, F_R\}$ to obtain an r_2 -regular polynomial factor \mathcal{B} of degree $\leq D$, defined by homogeneous nonclassical polynomials $R_1, \dots, R_C : \mathbb{F}^n \rightarrow \mathbb{T}$, where $C \leq C_{7.30}^{\mathbb{F}, r_2, d}(R)$. Since each F_t is measurable with respect to \mathcal{B} , so is F . So, there exists a function $\Sigma : \mathbb{T}^C \rightarrow [R]$ such that

$$F(x) = \Sigma(R_1(x), \dots, R_C(x)).$$

From each R_i pick a monomial with degree equal to $\deg(R_i)$ and a monomial (possibly the same one) with depth equal to $\text{depth}(R_i)$. By taking K to be sufficiently large, and as we assume that $n > K$, we can guarantee the existence of an $i_0 \in [n]$ such that x_{i_0} is not involved in any of these monomials. Let R'_1, \dots, R'_C be the restrictions of R_1, \dots, R_C , respectively, to the hyperplane $H = \{x_{i_0} = 0\}$. By the choice of i_0 we have that $\deg(R'_i) = \deg(R_i)$ and $\text{depth}(R'_i) = \text{depth}(R_i)$ for all $i \in [C]$. Also, by Lemma 7.2, R'_1, \dots, R'_C have rank $> r_2(C) - p$. Since $F|_{x_{i_0}=0} \in \mathcal{P}$ by our assumption, by definition of \mathcal{P} , there must exist $(c, \mathbf{d}, \Gamma) \in S$ such that

$$\Sigma(R'_1(x), \dots, R'_C(x)) = \Gamma(P_1(x), \dots, P_c(x)),$$

where $\deg(P_i) \leq d_i$ for all $i \in [c]$.

Next, apply Theorem 7.30 again to find an r_1 -regular refinement of the factor defined by the tuple of polynomials $(R'_1, \dots, R'_C, P_1, \dots, P_c)$. Because of our choice of r_2 and the last part of Theorem 7.30, we obtain a syntactic refinement of $\{R'_1, \dots, R'_C\}$. That is, we obtain a polynomial factor \mathcal{B}' defined by homogeneous nonclassical polynomials $R'_1, \dots, R'_C, S_1, \dots, S_E : \mathbb{F}^n \rightarrow \mathbb{T}$ such that it has degree $\leq D$, rank $> r_1(C + E)$ where $C + E \leq C_{7.30}^{\mathbb{F}, d, r_1}(C + \sigma)$. In particular, for each $i \in [c]$ we have

$$P_i(x) = \Gamma_i(R'_1(x), \dots, R'_C(x), S_1(x), \dots, S_E(x))$$

for some function $\Gamma_i : \mathbb{T}^{C+E} \rightarrow \mathbb{T}$. So for all $x \in \mathbb{F}^n$,

$$\begin{aligned} \Sigma(R'_1(x), \dots, R'_C(x)) = & \\ & \Gamma\left(\Gamma_1(R'_1(x), \dots, R'_C(x), S_1(x), \dots, S_E(x)), \right. \\ & \left. \dots, \Gamma_c(R'_1(x), \dots, R'_C(x), S_1(x), \dots, S_E(x)))\right). \end{aligned}$$

Applying Lemma 7.24, we see that if the rank of \mathcal{B}' is $> r_{7.24}(\mathbb{F}, D, \varepsilon)$ where $\varepsilon > 0$ is sufficiently small (say $\varepsilon = \frac{1}{2\|\mathcal{B}'\|}$), then $(R'_1(x), \dots, R'_C(x), S_1(x), \dots, S_E(x))$ acquires every value in its range. Thus, we have the identity

$$\begin{aligned} & \Sigma(a_1, \dots, a_C) \\ & = \Gamma\left(\Gamma_1(a_1, \dots, a_C, b_1, \dots, b_E), \dots, \Gamma_c(a_1, \dots, a_C, b_1, \dots, b_E)\right), \end{aligned}$$

for every $a_i \in \mathbb{U}_{\text{depth}(R'_i)+1}$ and $b_i \in \mathbb{U}_{\text{depth}(S_i)+1}$. Hence, we can substitute R_i for R'_i and 0 for S_i in the above equation and still retain the identity

$$\begin{aligned} F(x) &= \Sigma(R_1(x), \dots, R_C(x)) \\ &= \Gamma\left(\Gamma_1(R_1(x), \dots, R_C(x), 0, \dots, 0), \right. \\ &\quad \left. \dots, \Gamma_c(R_1(x), \dots, R_C(x), 0, \dots, 0)\right) \\ &= \Gamma(Q_1(x), \dots, Q_c(x)) \end{aligned}$$

where $Q_i : \mathbb{F}^n \rightarrow \mathbb{T}$ are defined as $Q_i(x) = \Gamma_i(R_1(x), \dots, R_C(x), 0, \dots, 0)$. Since for every i , $\text{deg}(R_i) = \text{deg}(R'_i)$ and $\text{depth}(R_i) = \text{depth}(R'_i)$, we apply [Corollary 16.6](#) to conclude that $\text{deg}(Q_i) \leq \text{deg}(P_i) \leq d_i$ for every $i \in [c]$, as long as the rank of \mathcal{B}' is $> r_{16.6}(\mathbb{F}, d, C)$.

Finally, we argue that Q_1, \dots, Q_c are classical polynomials. Indeed, since P_1, \dots, P_c are classical polynomials, $\Gamma_1, \dots, \Gamma_c$ must map to \mathbb{U}_1 on all of $\prod_{i=1}^C \mathbb{U}_{\text{depth}(R'_i)+1} \times \prod_{i=1}^E \mathbb{U}_{\text{depth}(S_i)+1} \supseteq \prod_{i=1}^C \mathbb{U}_{\text{depth}(R_i)+1} \times \{0\}^E$. Hence, Q_1, \dots, Q_c take values in \mathbb{U}_1 , and hence are classical polynomials. We conclude that $F \in \mathcal{P}$. □

17

Estimating the Distance from Algebraic Properties

In [Chapter 15](#) and [Chapter 16](#) we discussed one-sided error testable affine-invariant properties. In this section we discuss two different settings, testability with two-sided error and parameter estimation. First let us define the notion of testability with two-sided error.

Definition 17.1 (Testability with two-sided error). A property $\mathcal{P} \subset \{\mathbb{F}^n \rightarrow [R]\}$ is said to be *testable* with two-sided error if there is a function $q : (0, 1) \rightarrow \mathbb{N}$ and a randomized algorithm T that, given as input a parameter $\varepsilon > 0$ and oracle access to a function $f : \mathbb{F}^n \rightarrow [R]$, makes at most $q(\varepsilon)$ queries to the oracle for f , accepts with probability at least $2/3$ if $f \in \mathcal{P}$ and rejects with probability at least $2/3$ if f is ε -far from \mathcal{P} .

Parameter estimation. A *function parameter* is any map π which maps a function $f : \mathbb{F}^n \rightarrow [R]$ to $\pi(f) \in [0, 1]$. A function parameter is affine invariant if for any $f : \mathbb{F}^n \rightarrow [R]$ and any invertible affine map $A : \mathbb{F}^n \rightarrow \mathbb{F}^n$ it holds that $\pi(f) = \pi(f \circ A)$. An example of a function parameter is the distance from an affine invariant property. That is, if $\mathcal{P} \subset \{f : \mathbb{F}^n \rightarrow [R]\}$ is an affine invariant property, then $\pi(f) = \text{dist}(f, \mathcal{P})$ is an affine-invariant function parameter.

Definition 17.2 (Parameter estimation). A function parameter π of functions $f : \mathbb{F}^n \rightarrow [R]$ is said to be *estimable* if there is a function $q : (0, 1) \rightarrow \mathbb{N}$ and a randomized algorithm T that, given as input a parameter $\varepsilon > 0$ and oracle access to a function $f : \mathbb{F}^n \rightarrow [R]$, makes at most $q(\varepsilon)$ queries to the oracle for f , and outputs a value which is within ε of $\pi(f)$ with probability at least $2/3$.

Hatami and Lovett [49] showed that the distance to any constant-query testable (with two-sided error) hereditary affine-invariant property is constant-query estimable.

Theorem 17.3 (Hatami and Lovett [49]). For every two-sided error testable hereditary affine-invariant property \mathcal{P} the parameter $\text{dist}(f, \mathcal{P})$ is estimable.

We will present a sketch of the proof of [Theorem 17.3](#). We would like to prove that given an oracle access to a function $f : \mathbb{F}^n \rightarrow [R]$ and an error parameter $\varepsilon > 0$, one can query the function f on $q(\varepsilon)$ points, and output an estimate of $\text{dist}(f, \mathcal{P})$ that is, with probability $\geq 2/3$ (say), within ε of the correct value. The test we study is very natural. We show that there exists a constant $m = m(\mathcal{P}, \varepsilon)$ such that for a random affine subspace H of dimension m it holds that

$$\Pr_H [|\text{dist}(f, \mathcal{P}) - \text{dist}(f|_H, \mathcal{P})| \leq \varepsilon] \geq 2/3.$$

Note that crucially, m is independent of n .

The proof of [Theorem 17.3](#) combines higher-order Fourier analysis with the framework of Fischer and Newman [30] which obtained similar results for graph properties. At a high level, the approach for the graph case and the affine-invariant case are similar. One applies a regularization process, which allows one to represent a graph (or a function) by a small structure. Then, one argues that a large enough random sample of the graph or function should have a similar small structure representing it. Hence, properties of the main object can be approximated by properties of a large enough sample of it. Fischer and Newman [30] implemented this idea in the graph case. Adapting this to the algebraic case inevitably introduces some new challenges.

To simplify the presentation, we focus on the case $R = 2$ from now on. That is, we assume that all functions are $f : \mathbb{F}^n \rightarrow \{0, 1\}$. Also, we fix the required success probability at $2/3$ (any other constant strictly less than 1 would also work). The following theorem formalizes the discussion above.

Theorem 17.4. Let $\mathcal{P} \subset \{f : \mathbb{F}^n \rightarrow \{0, 1\} : n \in \mathbb{N}\}$ be an affine-invariant hereditary property which is two-sided testable. Then, for any $\varepsilon > 0$, there exists a constant $m = m(\mathcal{P}, \varepsilon)$ such that the following holds. Let $H \subset \mathbb{F}^n$ be a uniformly chosen m -dimensional affine subspace. Then for any $f : \mathbb{F}^n \rightarrow \{0, 1\}$,

$$\Pr_H [|\text{dist}(f, \mathcal{P}) - \text{dist}(f|_H, \mathcal{P})| \leq \varepsilon] \geq 2/3.$$

17.1 Proof sketch of Theorem 17.4

Let $\mathcal{P} \subset \{f : \mathbb{F}^n \rightarrow \{0, 1\} : n \in \mathbb{N}\}$ be an affine-invariant hereditary property which is two-sided testable. Let $f : \mathbb{F}^n \rightarrow \{0, 1\}$ be a function, and let \tilde{f} be the restriction of the function to a random m -dimensional affine subspace $H \subset \mathbb{F}^n$. We show that, if $m = m(\mathcal{P}, \varepsilon)$ is chosen large enough, then

- **Completeness:** If $\text{dist}(f, \mathcal{P}) \leq \delta$, then with high probability, $\text{dist}(\tilde{f}, \mathcal{P}) \leq \delta + \varepsilon$.
- **Soundness:** If $\text{dist}(f, \mathcal{P}) \geq \delta + \varepsilon$, then with high probability, $\text{dist}(\tilde{f}, \mathcal{P}) \geq \delta$.

Let us first fix notations. Let $A : \mathbb{F}^m \rightarrow \mathbb{F}^n$ be a random full rank affine transformation. Note that $Af : \mathbb{F}^m \rightarrow \{0, 1\}$ defined as $Af(x) = f(Ax)$ is the restriction of f to the affine subspace which is the image of A . Thus in the above discussion $\tilde{f} = Af$.

The proof of the completeness is simple. If $\text{dist}(f, \mathcal{P}) \leq \delta$ then there exists $g \in \mathcal{P}$ for which $\text{dist}(f, g) \leq \delta$. With high probability over a random restriction, the distance of Af and Ag is at most $\delta + o_m(1)$. This is because (i) $\mathbb{E}_A \text{dist}(Af, Ag) = \text{dist}(f, g)$, as each point in \mathbb{F}^n has equal probability to be in the image of A , and (ii) a random affine subspace is

pairwise independent with regards to whether an element is contained in it. Thus, by Chebychev's inequality,

$$\Pr_A[|\text{dist}(Af, Ag) - \text{dist}(f, g)| \geq \varepsilon] \leq \frac{1}{\varepsilon^2 |\mathbb{F}|^m}.$$

Clearly, choosing m large enough guarantees that the probability is bounded by $1/10$ (say).

The main challenge (as in nearly all results in property testing) is to establish soundness. That is, we wish to show that if a function f is far from \mathcal{P} then, with high probability, a random restriction of it is also far from \mathcal{P} as well. The main idea is to show that if for a typical restriction Af is δ -close to a function $h : \mathbb{F}^m \rightarrow \{0, 1\}$ which is in \mathcal{P} , then h can be “pulled back” to a function $g : \mathbb{F}^n \rightarrow \{0, 1\}$ which is both roughly δ -close to f and also very close to \mathcal{P} . This will contradict our initial assumption that f is $(\delta + \varepsilon)$ -far from \mathcal{P} . In order to do so we apply the machinery of higher order Fourier analysis. Our description in this overview subsection will hide various “cheats” but will present the correct general outline. For the full details we refer to the original paper [49].

First, we apply two-sided testability to deduce that a restriction to a low-dimensional subspace can distinguish between $f \in \mathcal{P}$ and f which is ε -far from \mathcal{P} .

Claim 17.5. For any $\varepsilon > 0$, there is $k = k(\mathcal{P}, \varepsilon)$ and a partition of the set of functions $\{\mathbb{F}^k \rightarrow \{0, 1\}\}$ to two disjoint sets F_+, F_- such that the following holds. Let $B : \mathbb{F}^k \rightarrow \mathbb{F}^n$ be a random full rank affine transformation. Then for any $f : \mathbb{F}^n \rightarrow \{0, 1\}$:

- If $f \in \mathcal{P}$ then $\Pr[Bf \in F_+] \geq 0.9$.
- If $\text{dist}(f, \mathcal{P}) \geq \varepsilon$ then $\Pr[Bf \in F_-] \geq 0.9$.

Proof. The assumption that \mathcal{P} is two-sided locally testable means that for some $q = q(\varepsilon)$, there is a local test which uses q queries, that can distinguish with probability 0.9 between $f \in \mathcal{P}$ and f for which $\text{dist}(f, \mathcal{P}) \geq \varepsilon$. As the property is affine-invariant, we may assume that the queries come from affine forms in some $k \leq q$ variables. In particular, all the queries are contained in an affine subspace of dimension k . The

sets F_+, F_- are determined by which restrictions make the tester accept and which ones make it reject. \square

Next, we apply the decomposition theorems discussed in Chapter 9. These allow us to decompose f to “structured” parts which we will study, and “pseudo-random” parts which do not affect the distribution of restrictions to k -dimensional subspaces. In order to do so, for a function $f : \mathbb{F}^n \rightarrow \{0, 1\}$ define by $\mu_{f,k}$ the distribution of its restriction to k -dimensional subspaces. That is, for any $v : \mathbb{F}^k \rightarrow \{0, 1\}$ let

$$\mu_{f,k}[v] = \Pr_B[Bf = v],$$

where to recall $B : \mathbb{F}^k \rightarrow \mathbb{F}^n$ is a random full rank affine transformation.

We will need to slightly generalize this definition to randomized functions. In our context, these can be modeled as $f : \mathbb{F}^n \rightarrow [0, 1]$. Such a function describes a distribution over (deterministic) functions $f' : \mathbb{F}^n \rightarrow \{0, 1\}$ as follows: for each $x \in \mathbb{F}^n$ independently, sample $f'(x) \in \{0, 1\}$ so that $\mathbb{E}[f'(x)] = f(x)$. We extend the definition of $\mu_{f,k}$ to functions $f : \mathbb{F}^n \rightarrow [0, 1]$ by setting

$$\mu_{f,k}[v] = \mathbb{E}_{f'}[\mu_{f',k}[v]].$$

Our definition implies that if two functions $f, g : \mathbb{F}^n \rightarrow [0, 1]$ have distributions $\mu_{f,k}$ and $\mu_{g,k}$ close in statistical distance, then random restrictions to k -dimensional affine subspaces cannot distinguish f from g . This will be useful in the analysis of the soundness.

We next decompose our function $f : \mathbb{F}^n \rightarrow \{0, 1\}$ based on the above intuition. Let d to be determined later. Theorem 9.1 gives a decomposition

$$f = f_1 + f_2$$

where $f_1 = \Gamma(P_1(x), \dots, P_C(x)) : \mathbb{F}^n \rightarrow [0, 1]$ for a high-rank polynomial factor $\{P_1, \dots, P_C\}$ and where $\|f_2\|_{U^d}$ is small enough. In the actual proof one has to use the strong decomposition theorem (Theorem 9.3) into three parts $f = f_1 + f_2 + f_3$. However for the sake of sketching the proof one can ignore this technicality. For an appropriately chosen d ($d > |\mathbb{F}|^k$ suffices) we can then replace f with f_1 for the purposes of

analyzing its restrictions to k -dimensional subspaces. That is,

$$\mu_{f,k} \approx \mu_{f_1,k}$$

where $f_1 : \mathbb{F}^m \rightarrow [0, 1]$ and where closeness is in statistical distance. Thus, from now on we restrict our attention to f_1 , namely the “structured part” of f .

The next step is to show that the same type of decomposition can be applied to the restriction Af of f , recalling that $A : \mathbb{F}^m \rightarrow \mathbb{F}^n$ is a random full rank affine transformation. We will later choose $m \gg k$ for the proof to work. Clearly, $Af = Af_1 + Af_2$. We next analyze the typical behavior of Af_1 and Af_2 .

First, note that $Af_1 = \Gamma(Q_1(x), \dots, Q_C(x))$ where $Q_i = AP_i$ are the restrictions of P_1, \dots, P_C . One can show (and we omit the details in this proof sketch) that as P_1, \dots, P_C is a high-rank polynomial factor, then Q_1, \dots, Q_C is also a high-rank polynomial factor. Thus f_1 and Af_1 have the same “high level” factorization Γ to high-rank polynomials.

Next, we analyze Af_2 . We claim that with high probability over the choice of A , if m is chosen large enough, then $\|Af_2\|_{U^d} \approx \|f_2\|_{U^d}$. This holds since

$$\mathbb{E}_A \|Af_2\|_{U^d}^{2^d} = \|f_2\|_{U^d}^{2^d} \pm O(2^d |\mathbb{F}|^{-m}).$$

So, Af and Af_1 also have similar distribution of their restrictions to random k -dimensional subspaces. That is,

$$\mu_{Af,k} \approx \mu_{Af_1,k}.$$

Next, assume that $\text{dist}(Af, \mathcal{P}) \leq \varepsilon$. By definition, there exists $h : \mathbb{F}^m \rightarrow \{0, 1\}$, where $h \in \mathcal{P}$, such that $\text{dist}(Af, h) \leq \varepsilon$. We apply the same decomposition process to h . Thus, we decompose

$$h = h_1 + h_2,$$

where h_1 is structured and $\|h_2\|_{U^d}$ is very small. Thus as before, $\mu_{h,k} \approx \mu_{h_1,k}$.

By choosing the exact parameters of “high rank” for h_1 to be lower than those for Af_1 , but still high enough (for exact details, see the original paper [49]), we may assume that the polynomials that appear

in the decomposition of h_1 extend Q_1, \dots, Q_C which appear in the decomposition of Af_1 . That is, we have

$$h_1(x) = \Gamma'(Q_1(x), \dots, Q_{C'}(x))$$

for some $C' > C$, such that the entire set of polynomials $Q_1, \dots, Q_{C'}$ is of high rank. The important aspect here is that the polynomials Q_1, \dots, Q_C which comprise Af_1 are part of the description of h_1 (however, it may be the case that Γ' ignores them; we will see soon that this is impossible). To summarize: both Af_1 and h_1 can be defined in terms of the same basic “building blocks”, namely high rank polynomials $Q_1, \dots, Q_{C'}$.

The next step is to “pull back” h to a function $\phi : \mathbb{F}^n \rightarrow \{0, 1\}$, such that ϕ is very close to \mathcal{P} , and such that $\text{dist}(f, \phi) \approx \text{dist}(Af, h)$. This will show that

$$\text{dist}(f, \mathcal{P}) \leq \text{dist}(f, \phi) + o_m(1) = \text{dist}(Af, h) + o_m(1) \leq \delta + o_m(1).$$

Setting $m = m(k, \varepsilon)$ large enough would show that $\text{dist}(f, \mathcal{P}) \leq \delta + \varepsilon$, which is our goal.

The first step is to pull back h_1 . To recall, $h_1(x) = \Gamma'(Q_1(x), \dots, Q_{C'}(x))$, and moreover $Q_i = AP_i$ for $1 \leq i \leq C$. So, for $C < i \leq C'$ we need to define pull-back polynomials $P_i : \mathbb{F}^n \rightarrow \{0, 1\}$ such that (i) $Q_i = AP_i$; and (ii) $P_1, \dots, P_{C'}$ are of high rank. This can be done for example by letting $P_i = DQ_i$ for any affine map $D : \mathbb{F}^n \rightarrow \mathbb{F}^m$ for which AD is the identity map on \mathbb{F}^m . So, define $\phi_1 : \mathbb{F}^n \rightarrow [0, 1]$ given by

$$\phi_1(x) = \Gamma'(P_1(x), \dots, P_{C'}(x)).$$

Note that $A\phi_1 = h_1$; that is, ϕ_1 is the pull-back of the “structured part” h_1 of h . However, it does not in general generate a function close to f . This makes sense, as we still have not used the finer “pseudo-random” structure of h_2 .

However, we can already show something about ϕ_1 : it is very close to \mathcal{P} . More concretely, as $P_1, \dots, P_{C'}$ are high rank polynomials, and also $Q_1, \dots, Q_{C'}$ are high rank polynomials, we have

$$\mu_{\phi_1, k} \approx \mu_{h_1, k}.$$

Recall that we have shown $\mu_{h_1, k} \approx \mu_{h, k}$, and that $h \in \mathcal{P}$. Thus, the tester which distinguishes functions in \mathcal{P} from those ε -far from \mathcal{P} cannot distinguish ϕ_1 from functions in \mathcal{P} . Hence, ϕ_1 must be ε -close to \mathcal{P} .

The final step is to define the more refined pull-back ϕ of h . Define an atom as a subset $\{x \in \mathbb{F}^n : P_1(x) = a_1, \dots, P_{C'}(x) = a_{C'}\}$ for values $a_1, \dots, a_{C'}$. Note that the functions f_1, h_1 are constant over atoms. We next define $\phi : \mathbb{F}^n \rightarrow [0, 1]$ by redefining ϕ_1 inside each atom, so that the average over each atom of ϕ and ϕ_1 is the same, but such that ϕ is as close as possible to f given this constraint. Concretely, we can consider three cases:

- An atom where the average of f over the atom equals the value ϕ_1 assigns to this atom. In this atom, we simply set $\phi(x) = f(x)$ for all points x in this atom.
- An atom where the average of f over the atom is larger than the value ϕ_1 assigns to this atom. In this atom, for any x , if $f(x) = 0$ then we set $\phi(x) = 0$, and if $f(x) = 1$ then we set $\phi(x) = \alpha$ where the value α is chosen so that the average of ϕ and ϕ_1 over the atom is the same.
- An atom where the average of f over the atom is lower than the value ϕ_1 assigns to this atom. This is analogous to the previous case.

One can show that under this choice, ϕ is indeed a proper pull-back of h , in the sense that

$$\text{dist}(f, \phi) = \text{dist}(f_1, \phi_1) \approx \text{dist}(Af, h_1) \approx \text{dist}(Af, h).$$

Moreover,

$$\text{dist}(\phi, \mathcal{P}) \approx \text{dist}(\phi_1, \mathcal{P}) = o_m(1).$$

We thus conclude that $\text{dist}(f, \mathcal{P}) \leq \text{dist}(Af, h) + o_m(1) \leq \delta + o_m(1)$.

Part IV

Open Problems

18

Open Problems

We conclude this survey with a quick overview of the main open problems in this area. Several of these have already been mentioned in the text, and we repeat them here for completeness.

18.1 Testability of hereditary properties

A beautiful and fundamental result of Alon and Shapira [4] says that every hereditary graph property is testable. In [16] the analogue of Alon and Shapira's result is conjectured for algebraic properties.

Conjecture 14.7 (restated). *Every affine invariant subspace hereditary property is testable with one-sided error.*

Resolving [Conjecture 14.7](#) would yield a combinatorial *characterization* of the one-sided testable affine-invariant properties, similar to the characterization for testable dense graph properties [4]. In [Theorem 14.9](#), we made major progress towards resolving [Conjecture 14.7](#) by proving the testability under an additional assumption of “bounded complexity”. The authors of the survey believe that it is likely that [Conjecture 14.7](#) is in fact false, and some assumption on the complexity of the property is necessary.

18.2 Testing correlation with classical polynomials

As we saw in [Section 6.2](#), estimating the Gowers norm $\|f\|_{U_{d+1}}$ can be used as a test for whether a function f has significant or negligible correlation with degree d nonclassical polynomials. We also saw in [Chapter 5](#) that $d = 3$ is the first case where the Gowers norm is not a test for correlation with classical polynomials. The $d = 3$ still remains a fascinating open problem.

Problem 18.1. Does there exist a tester which queries a function $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$ on a constant number of positions, and which can distinguish whether f has noticeable or negligible correlation with classical cubic polynomials?

More precisely, we would like a universal test with a constant number of queries for which the following holds. For every $\varepsilon > 0$, there exists a constant $0 < \delta(\varepsilon) < \varepsilon$, such that with probability at least $2/3$ the test correctly distinguishes between functions that have correlation at most $\delta(\varepsilon)$ with all classical cubic polynomials, and the ones that have correlation at least ε with some classical cubic polynomial.

As we saw in [Chapter 17](#), a tester for which the number of queries depends on the error parameter exists and was given in [49], however the above problem is left open. See [50] for more results and discussion on the subject.

18.3 Quantitative bounds for inverse theorems

Recall that the inverse theorem for Gowers norms, [Theorem 6.9](#), shows existence of $\varepsilon(\delta, d, \mathbb{F})$ such that the following holds. For every function $f : \mathbb{F}^n \rightarrow \mathbb{C}$ with $\|f\|_\infty \leq 1$ and $\|f\|_{U_{d+1}} \geq \delta$, there exists a polynomial $P \in \text{Poly}_{\leq d}(\mathbb{F}^n \rightarrow \mathbb{T})$ of degree $\leq d$ that is ε -correlated with f , meaning

$$\left| \mathbb{E}_{x \in \mathbb{F}^n} f(x) e(-P(x)) \right| \geq \varepsilon.$$

The asymptotics of the required dependence of ε on δ are not well understood and in fact it is a major open problem whether this dependence can be made polynomial, even in the special case of $d = 3$ and $|\mathbb{F}| = 2$.

Conjecture 18.2 (Polynomial inverse Gowers conjecture for U^3). Let $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$. If $\left\| (-1)^f \right\|_{U^3} \geq \varepsilon$, then there exists a quadratic classical polynomial q such that $\Pr[f(x) = q(x)] \geq \frac{1}{2} + \delta$, where $\delta = c\varepsilon^c$ for some absolute constant $c > 0$.

As evidence for the importance of understanding quantitative bounds for the inverse theorem, Green and Tao [44] and Lovett [58] proved that the above conjecture is equivalent to a very important conjecture in additive combinatorics, namely the well-known polynomial Freiman-Ruzsa conjecture. The best known lower-bound for δ is quasi-polynomial in ε which follows from the paper of Sanders on the Bogolyubov-Ruzsa conjecture [71].

18.4 Complexity of linear forms

In [Theorem 11.8](#) we proved that if $\mathcal{L} = \{L_1, \dots, L_m\}$ is a system of linear forms for which $L_1^{d+1}, L_2^{d+1}, \dots, L_m^{d+1}$ are linearly independent, then for every $\varepsilon > 0$, there exists $\delta > 0$ such that for any collection of functions $f_1, \dots, f_m : \mathbb{F}^n \rightarrow \mathbb{D}$ with $\|f_1\|_{U^{d+1}} \leq \delta$, we have

$$\left| \mathbb{E}_{X \in (\mathbb{F}^n)^k} \left[\prod_{i=1}^m f_i(L_i(X)) \right] \right| \leq \varepsilon.$$

Our proof of this fact goes through inverse theorem for Gowers norms and regularity lemmas for polynomials and as a result the dependence of δ on ε is really bad. It is left open whether this theorem can be proved with reasonable bounds on δ . We state an open problem due to Gowers and Wolf [40] that focuses on the polynomial regime.

Problem 18.3 (Gowers and Wolf [40], Problem 7.6 reformulated). Does there exist an integer $d \geq 1$ and a system of linear forms L_1, \dots, L_m with $L_1^{d+1}, \dots, L_m^{d+1}$ linearly independent, such that the following holds: For every positive real number r , there exists $\varepsilon > 0$ and functions $f_i : \mathbb{F}^n \rightarrow \mathbb{C}$ such that $\|f_i\|_{U^d} \leq \varepsilon^r$ for every i , and yet

$$\left| \mathbb{E}_{X \in (\mathbb{F}^n)^d} \prod_{i=1}^m f_i(L_i(X)) \right| > \varepsilon.$$

18.5 Norms defined by linear forms.

A possibly challenging open problem is to give a full characterization of collections of linear forms L_1, \dots, L_m for which

$$\left| \mathbb{E}_{X \in \mathbb{F}^k} \prod_{i=1}^m f(L_i(X)) \right|^{1/m},$$

defines a norm on the space $\{f : \mathbb{F}^n \rightarrow \mathbb{R}\}$. Note that both the Gowers norm as well as the L_p norms for even values of p are of this type. In fact, one can show a characterization when \mathbb{F} has characteristic larger than the true complexity of the linear forms. It turns out that in this case, all such norms are essentially either equivalent to an L_p norm for some $p > 0$ or equivalent to the Gowers norm of some order d . However, the techniques do not seem to extend to the general question which is the interesting regime.

References

- [1] Alon, N. and R. Beigel (2001). “Lower bounds for approximations by low degree polynomials over Z^m ”. In: *Computational Complexity, 16th Annual IEEE Conference on, 2001*. IEEE. 184–187.
- [2] Alon, N., E. Fischer, I. Newman, and A. Shapira (2009). “A combinatorial characterization of the testable graph properties: it’s all about regularity”. *SIAM J. Comput.* 39(1): 143–167. DOI: [10.1137/060667177](https://doi.org/10.1137/060667177).
- [3] Alon, N., T. Kaufman, M. Krivelevich, S. Litsyn, and D. Ron (2005). “Testing Reed-Muller codes”. *IEEE Trans. Inform. Theory*. 51(11): 4032–4039.
- [4] Alon, N. and A. Shapira (2008a). “A Characterization of the (Natural) Graph Properties Testable with One-Sided Error”. *SIAM J. on Comput.* 37(6): 1703–1727.
- [5] Alon, N. and A. Shapira (2008b). “Every Monotone Graph Property Is Testable”. *SIAM J. on Comput.* 38(2): 505–522.
- [6] Babai, L., L. Fortnow, and C. Lund (1992). “Addendum to: “Non-deterministic exponential time has two-prover interactive protocols” [Comput. Complexity 1 (1991), no. 1, 3–40; MR1113533 (92h:68031)]”. *Comput. Complexity*. 2(4): 374. DOI: [10.1007/BF01200430](https://doi.org/10.1007/BF01200430).

- [7] Babai, L., L. Fortnow, L. A. Levin, and M. Szegedy (1991). “Checking computations in polylogarithmic time”. In: *Proc. 23rd Annual ACM Symposium on the Theory of Computing*. New York: ACM Press. 21–32.
- [8] Balog, A. and E. Szemerédi (1994). “A statistical theorem of set addition”. *Combinatorica*. 14(3): 263–268. URL: <http://dx.doi.org/10.1007/BF01212974>.
- [9] Bellare, M., D. Coppersmith, J. Håstad, M. Kiwi, and M. Sudan (1996). “Linearity testing over characteristic two”. *IEEE Trans. Inform. Theory*. 42(6): 1781–1795.
- [10] Bergelson, V., T. Tao, and T. Ziegler (2010). “An inverse theorem for the uniformity seminorms associated with the action of \mathbb{F}_p^∞ ”. *Geom. Funct. Anal.* 19(6): 1539–1596. DOI: [10.1007/s00039-010-0051-1](https://doi.org/10.1007/s00039-010-0051-1).
- [11] Bhattacharyya, A. (2014). “Polynomial decompositions in polynomial time”. *Tech. rep.* <http://eccc.hpi-web.de/report/2014/018>. Electronic Colloquium on Computational Complexity (ECCC).
- [12] Bhattacharyya, A., A. Bhowmick, and C. Gupta (2016). “On Higher-Order Fourier Analysis over Non-Prime Fields”. In: *LIPICs-Leibniz International Proceedings in Informatics*. Vol. 60. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik.
- [13] Bhattacharyya, A., V. Chen, M. Sudan, and N. Xie (2011). “Testing linear-invariant non-linear properties”. *Theory Comput.* 7: 75–99. DOI: [10.4086/toc.2011.v007a006](https://doi.org/10.4086/toc.2011.v007a006).
- [14] Bhattacharyya, A., E. Fischer, H. Hatami, P. Hatami, and S. Lovett (2013a). “Every locally characterized affine-invariant property is testable”. In: *Proceedings of the 45th annual ACM symposium on Symposium on theory of computing. STOC '13*. Palo Alto, California, USA: ACM. 429–436. DOI: [10.1145/2488608.2488662](https://doi.org/10.1145/2488608.2488662).
- [15] Bhattacharyya, A., E. Fischer, and S. Lovett (2013b). “Testing Low Complexity Affine-Invariant Properties”. In: *Proc. 24th ACM-SIAM Symposium on Discrete Algorithms*. 1337–1355.
- [16] Bhattacharyya, A., E. Grigorescu, and A. Shapira (2010a). “A Unified Framework for Testing Linear-Invariant Properties”. In: *Proc. 51st Annual IEEE Symposium on Foundations of Computer Science*. 478–487.

- [17] Bhattacharyya, A., P. Hatami, and M. Tulsiani (2015). “Algorithmic regularity for polynomials and applications”. In: *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2015, San Diego, CA, USA, January 4-6, 2015*. 1870–1889. DOI: [10.1137/1.9781611973730.125](https://doi.org/10.1137/1.9781611973730.125).
- [18] Bhattacharyya, A., S. Kopparty, G. Schoenebeck, M. Sudan, and D. Zuckerman (2010b). “Optimal testing of Reed-Muller codes”. In: *2010 IEEE 51st Annual Symposium on Foundations of Computer Science FOCS 2010*. IEEE Computer Soc., Los Alamitos, CA. 488–497.
- [19] Bhowmick, A. and S. Lovett (2015). “Bias vs structure of polynomials in large fields, and applications in effective algebraic geometry and coding theory”. *arXiv preprint arXiv:1506.02047*.
- [20] Blum, M., M. Luby, and R. Rubinfeld (1993). “Self-testing/correcting with applications to numerical problems”. In: *Proceedings of the 22nd Annual ACM Symposium on Theory of Computing (Baltimore, MD, 1990)*. Vol. 47 (3). 549–595. DOI: [10.1016/0022-0000\(93\)90044-W](https://doi.org/10.1016/0022-0000(93)90044-W).
- [21] Bogdanov, A. and E. Viola (2007). “Pseudorandom Bits for Polynomials”. In: *Proc. 48th Annual IEEE Symposium on Foundations of Computer Science*. Washington, DC, USA: IEEE Computer Society. 41–51. DOI: <http://dx.doi.org/10.1109/FOCS.2007.56>.
- [22] Borgs, C., J. Chayes, L. Lovász, V. T. Sós, B. Szegedy, and K. Vesztergombi (2006). “Graph limits and parameter testing”. In: *STOC’06: Proceedings of the 38th Annual ACM Symposium on Theory of Computing*. ACM, New York. 261–270. DOI: [10.1145/1132516.1132556](https://doi.org/10.1145/1132516.1132556).
- [23] Cohen, H. (2008). *Number theory: Volume I: Tools and diophantine equations*. Vol. 239. Springer Science & Business Media.
- [24] Conlon, D., J. Fox, and B. Sudakov (2010). “Hypergraph Ramsey numbers”. *Journal of the American Mathematical Society*. 23(1): 247–266.
- [25] Erdős, P. and R. Rado (1952). “Combinatorial theorems on classifications of subsets of a given set”. *Proceedings of the London mathematical Society*. 3(1): 417–439.

- [26] Erdős, P. and G. Szekeres (1935). “A combinatorial problem in geometry”. *Compositio Mathematica*. 2: 463–470.
- [27] Erdős, P. and P. Turán (1936). “On Some Sequences of Integers”. *J. London Math. Soc.* S1-11(4): 261. DOI: [10.1112/jlms/s1-11.4.261](https://doi.org/10.1112/jlms/s1-11.4.261).
- [28] Feige, U., S. Goldwasser, L. Lovász, S. Safra, and M. Szegedy (1996). “Interactive proofs and the hardness of approximating cliques”. *J. ACM*. 43(2): 268–292.
- [29] Fischer, E. (2004). “The art of uninformed decisions: A primer to property testing”. In: *Current Trends in Theoretical Computer Science: The Challenge of the New Century*. Ed. by G. Paun, G. Rozenberg, and A. Salomaa. Vol. 1. World Scientific Publishing. 229–264.
- [30] Fischer, E. and I. Newman (2007). “Testing versus estimation of graph properties”. *SIAM J. Comput.* 37(2): 482–501 (electronic). DOI: [10.1137/060652324](https://doi.org/10.1137/060652324).
- [31] Furstenberg, H. (1977). “Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions”. *J. Analyse Math.* 31: 204–256.
- [32] Goldreich, O., S. Goldwasser, and D. Ron (1998). “Property testing and its connection to learning and approximation”. *J. ACM*. 45(4): 653–750. DOI: [10.1145/285055.285060](https://doi.org/10.1145/285055.285060).
- [33] Goldreich, O. and T. Kaufman (2011). “Proximity oblivious testing and the role of invariances”. In: *Approximation, randomization, and combinatorial optimization*. Vol. 6845. *Lecture Notes in Comput. Sci.* Heidelberg: Springer. 579–592. DOI: [10.1007/978-3-642-22935-0_49](https://doi.org/10.1007/978-3-642-22935-0_49).
- [34] Goldreich, O. and L. A. Levin (1989). “A hard-core predicate for all one-way functions”. In: *Proceedings of the twenty-first annual ACM symposium on Theory of computing*. ACM. 25–32.
- [35] Goldreich, O. and D. Ron (2011). “On proximity-oblivious testing”. *SIAM J. Comput.* 40(2): 534–566. DOI: [10.1137/100789646](https://doi.org/10.1137/100789646).
- [36] Gopalan, P., R. O’Donnell, R. A. Servedio, A. Shpilka, and K. Wimmer (2009). “Testing Fourier Dimensionality and Sparsity”. In: *Proc. 36th Annual International Conference on Automata, Languages, and Programming*. 500–512.

- [37] Gowers, W. T. (2001). “A new proof of Szemerédi’s theorem”. *Geom. Funct. Anal.* 11(3): 465–588. DOI: [10.1007/s00039-001-0332-9](https://doi.org/10.1007/s00039-001-0332-9).
- [38] Gowers, W. T. (2010). “Decompositions, approximate structure, transference, and the Hahn-Banach theorem”. *Bull. Lond. Math. Soc.* 42(4): 573–606. DOI: [10.1112/blms/bdq018](https://doi.org/10.1112/blms/bdq018).
- [39] Gowers, W. T. and J. Wolf (2010). “The true complexity of a system of linear equations”. *Proc. Lond. Math. Soc. (3)*. 100(1): 155–176. DOI: [10.1112/plms/pdp019](https://doi.org/10.1112/plms/pdp019).
- [40] Gowers, W. T. and J. Wolf (2011). “Linear forms and higher-degree uniformity for functions on \mathbb{F}_p^n ”. *Geom. Funct. Anal.* 21(1): 36–69. DOI: [10.1007/s00039-010-0106-3](https://doi.org/10.1007/s00039-010-0106-3).
- [41] Green, B. (2007). “Montréal notes on quadratic Fourier analysis”. In: *Additive combinatorics*. Vol. 43. *CRM Proc. Lecture Notes*. Providence, RI: Amer. Math. Soc. 69–102.
- [42] Green, B. and T. Tao (2008). “An inverse theorem for the Gowers U^3 -norm”. *Proc. Edin. Math. Soc.* 51: 73–153.
- [43] Green, B. and T. Tao (2009). “The distribution of polynomials over finite fields, with applications to the Gowers norms”. *Contrib. Discrete Math.* 4(2): 1–36.
- [44] Green, B. and T. Tao (2010a). “An equivalence between inverse sumset theorems and inverse conjectures for the U^3 norm”. In: *Mathematical Proceedings of the Cambridge Philosophical Society*. Vol. 149. No. 01. Cambridge Univ Press. 1–19.
- [45] Green, B. and T. Tao (2010b). “Linear equations in primes”. *Ann. of Math. (2)*. 171(3): 1753–1850. DOI: [10.4007/annals.2010.171.1753](https://doi.org/10.4007/annals.2010.171.1753).
- [46] Hatami, H., P. Hatami, and J. Hirst (2014a). “Limits of Boolean functions on \mathbb{F}_p^n ”. *Electron. J. Combin.* 21(4): Paper 4.2, 15.
- [47] Hatami, H., P. Hatami, and S. Lovett (2014b). “General systems of linear forms: equidistribution and true complexity”. *arXiv preprint arXiv:1403.7703*.
- [48] Hatami, H. and S. Lovett (2011). “Higher-order Fourier analysis of \mathbb{F}_p^n and the complexity of systems of linear forms”. *Geom. Funct. Anal.* 21(6): 1331–1357. DOI: [10.1007/s00039-011-0141-8](https://doi.org/10.1007/s00039-011-0141-8).

- [49] Hatami, H. and S. Lovett (2013). “Estimating the distance from testable affine-invariant properties”. In: *Foundations of Computer Science (FOCS), 2013 IEEE 54th Annual Symposium on*. IEEE. 237–242.
- [50] Hatami, H. and S. Lovett (2014). “Correlation Testing for Affine Invariant Properties on \mathbb{F}_p^n in the High Error Regime”. *SIAM Journal on Computing*. 43(4): 1417–1455.
- [51] Host, B. and B. Kra (2005). “Nonconventional ergodic averages and nilmanifolds”. *Ann. of Math. (2)*. 161(1): 397–488. DOI: [10.4007/annals.2005.161.397](https://doi.org/10.4007/annals.2005.161.397).
- [52] Jutla, C. S., A. C. Patthak, A. Rudra, and D. Zuckerman (2004). “Testing Low-Degree Polynomials over Prime Fields”. In: *Proc. 45th Annual IEEE Symposium on Foundations of Computer Science*. 423–432.
- [53] Kaufman, T. and S. Litsyn (2005). “Almost orthogonal linear codes are locally testable”. In: *46th Annual IEEE Symposium on Foundations of Computer Science (FOCS’05)*. IEEE. 317–326.
- [54] Kaufman, T. and S. Lovett (2008). “Worst Case to Average Case Reductions for Polynomials”. *Foundations of Computer Science, IEEE Annual Symposium on*: 166–175. URL: <http://doi.ieeecomputersociety.org/10.1109/FOCS.2008.17>.
- [55] Kaufman, T. and D. Ron (2006). “Testing Polynomials over General Fields”. *SIAM J. on Comput.* 36(3): 779–802.
- [56] Kaufman, T. and M. Sudan (2008). “Algebraic property testing: the role of invariance”. In: *Proc. 40th Annual ACM Symposium on the Theory of Computing*. 403–412.
- [57] Lovett, S. (2009). “Unconditional Pseudorandom Generators for Low Degree Polynomials”. *Theory of Computing*. 5(1): 69–82. DOI: [10.4086/toc.2009.v005a003](https://doi.org/10.4086/toc.2009.v005a003). URL: <http://www.theoryofcomputing.org/articles/v005a003>.
- [58] Lovett, S. (2012). “Equivalence of polynomial conjectures in additive combinatorics”. *Combinatorica*. 32(5): 607–618. DOI: [10.1007/s00493-012-2714-z](https://doi.org/10.1007/s00493-012-2714-z).

- [59] Lovett, S. (2015). *An Exposition of Sanders' Quasi-Polynomial Freiman-Ruzsa Theorem*. *Graduate Surveys*. No. 6. Theory of Computing Library. 1–14. DOI: [10.4086/toc.gs.2015.006](https://doi.org/10.4086/toc.gs.2015.006). URL: <http://www.theoryofcomputing.org/library.html>.
- [60] Lovett, S., R. Meshulam, and A. Samorodnitsky (2011). “Inverse conjecture for the Gowers norm is false”. *Theory Comput.* 7: 131–145. DOI: [10.4086/toc.2011.v007a009](https://doi.org/10.4086/toc.2011.v007a009).
- [61] Lucas, E. (1878). “Théorie des Fonctions Numériques Simplement Périodiques”. *American Journal of Mathematics*. 1(2): 184–196. URL: <http://www.jstor.org/stable/2369308>.
- [62] MacWilliams, F. J. and N. J. A. Sloane (1977). *The theory of error-correcting codes*. Elsevier.
- [63] O’Donnell, R. (2014). *Analysis of boolean functions*. Cambridge University Press.
- [64] Reingold, O., L. Trevisan, M. Tulsiani, and S. Vadhan (2008). “Dense subsets of pseudorandom sets”. In: *Foundations of Computer Science, 2008. FOCS’08. IEEE 49th Annual IEEE Symposium on*. IEEE. 76–85.
- [65] Ron, D. (2009). “Algorithmic and Analysis Techniques in Property Testing”. *Foundations and Trends in Theoretical Computer Science*. 5(2): 73–205.
- [66] Roth, K. F. (1953). “On certain sets of integers”. *J. London Math. Soc.* 28: 104–109.
- [67] Rubinfeld, R. (1999). “On the robustness of functional equations”. *SIAM Journal on Computing*. 28(6): 1972–1997.
- [68] Rubinfeld, R. (2006). “Sublinear time algorithms”. In: *Proceedings of International Congress of Mathematicians 2006*. Vol. 3. 1095–1110.
- [69] Rubinfeld, R. and M. Sudan (1996). “Robust characterizations of polynomials with applications to program testing”. *SIAM J. Comput.* 25(2): 252–271. DOI: [10.1137/S0097539793255151](https://doi.org/10.1137/S0097539793255151).
- [70] Samorodnitsky, A. (2007). “Low-degree tests at large distances”. In: *STOC’07—Proceedings of the 39th Annual ACM Symposium on Theory of Computing*. New York: ACM. 506–515. DOI: [10.1145/1250790.1250864](https://doi.org/10.1145/1250790.1250864).

- [71] Sanders, T. (2012). “On the Bogolyubov-Ruzsa lemma”. *Analysis & PDE*. 5(3): 627–655.
- [72] Sanders, T. (2013). “The structure theory of set addition revisited”. *Bulletin of the American Mathematical Society*. 50: 93–127.
- [73] Shapira, A. (2009). “Green’s conjecture and testing linear-invariant properties”. In: *STOC’09—Proceedings of the 2009 ACM International Symposium on Theory of Computing*. ACM, New York. 159–166.
- [74] Sudan, M. (2010). “Invariance in Property Testing”. *Tech. rep.* No. 10-051. Electronic Colloquium in Computational Complexity.
- [75] Szegedy, B. (2012). “On higher order Fourier analysis”. *arXiv preprint arXiv:1203.2260*.
- [76] Szemerédi, E. (1969). “On sets of integers containing no four elements in arithmetic progression”. *Acta Math. Acad. Sci. Hungar.* 20: 89–104.
- [77] Szemerédi, E. (1975). “On sets of integers containing no k elements in arithmetic progression”. *Acta Arith.* 27: 199–245. Collection of articles in memory of Juriui Vladimirovive Linnik.
- [78] Tao, T. (2007). “Structure and randomness in combinatorics”. In: *Foundations of Computer Science, 2007. FOCS’07. 48th Annual IEEE Symposium on*. IEEE. 3–15.
- [79] Tao, T. (2008). “Some notes on “non-classical” polynomials in finite characteristic”. *Blog post available at <https://terrytao.wordpress.com/2008/11/13/some-notes-on-non-classical-polynomials-in-finite-characteristic>*.
- [80] Tao, T. (2012). *Higher order Fourier analysis*. Vol. 142. American Mathematical Soc.
- [81] Tao, T. and V. H. Vu (2010). *Additive combinatorics*. Vol. 105. *Cambridge Studies in Advanced Mathematics*. Paperback edition [of MR2289012]. Cambridge University Press, Cambridge. xviii+512.
- [82] Tao, T. and T. Ziegler (2010). “The inverse conjecture for the Gowers norm over finite fields via the correspondence principle”. *Anal. PDE*. 3(1): 1–20. DOI: [10.2140/apde.2010.3.1](https://doi.org/10.2140/apde.2010.3.1).

- [83] Tao, T. and T. Ziegler (2012). “The inverse conjecture for the Gowers norm over finite fields in low characteristic”. *Ann. Comb.* 16(1): 121–188. DOI: [10.1007/s00026-011-0124-3](https://doi.org/10.1007/s00026-011-0124-3).
- [84] Viola, E. (2011). “Selected Results in Additive Combinatorics: An Exposition.” *Theory of Computing, Graduate Surveys*. 2: 1–15.
- [85] Viola, E. (2009). “The Sum of D Small-Bias Generators Fools Polynomials of Degree D”. *Computational Complexity*. 18(2): 209–217.
- [86] Yoshida, Y. (2014a). “A characterization of locally testable affine-invariant properties via decomposition theorems”. In: *Proceedings of the 46th Annual ACM Symposium on Theory of Computing*. ACM. 154–163.
- [87] Yoshida, Y. (2014b). “Gowers Norm, Function Limits, and Parameter Estimation”. *arXiv preprint arXiv:1410.5053*.
- [88] Ziegler, T. (2007). “Universal characteristic factors and Furstenberg averages”. *J. Amer. Math. Soc.* 20(1): 53–97 (electronic). DOI: [10.1090/S0894-0347-06-00532-7](https://doi.org/10.1090/S0894-0347-06-00532-7).