

UC Irvine

UC Irvine Previously Published Works

Title

Draft Genome Sequence of *Bacillus safensis* RP10, Isolated from Soil in the Atacama Desert, Chile

Permalink

<https://escholarship.org/uc/item/7894v1jd>

Journal

Microbiology Resource Announcements, 8(44)

ISSN

2576-098X

Authors

Strahsburger, Erwin
Zapata, Felipe
Pedroso, Inti
et al.

Publication Date

2019-10-31

DOI

10.1128/mra.01150-19

Peer reviewed



Draft Genome Sequence of *Bacillus safensis* RP10, Isolated from Soil in the Atacama Desert, Chile

 Erwin Strahsburger,^{a,c} Felipe Zapata,^b Inti Pedroso,^b Derie Fuentes,^b Paz Tapia,^b Raul Ponce,^a Michael McClelland,^c Jorge Valdes^b

^aDoctoral Program in Agronomy in Desert and Arid Zones, Molecular Biotechnology Laboratory, Faculty of Renewal Natural Resources, Universidad Arturo Prat, Iquique, Chile

^bBio-Computing and Applied Genetics Division, Center for Systems Biotechnology, Fraunhofer Chile Research Foundation, Santiago, Chile

^cDepartment of Microbiology & Molecular Genetics, University of California, Irvine, Irvine, California, USA

ABSTRACT Genome analysis of *Bacillus safensis* RP10, a strain from the soil of Atacama Desert in northern Chile, reflects a bacterium adapted to live in soil containing high levels of heavy metals, high salt conditions, and low carbon and energy sources.

The Atacama Desert of Chile is the driest desert in the world (1, 2) and has soils naturally enriched in heavy metals and salts, which affects local agricultural activities (3). These soils also harbor species of the genus *Bacillus* (4, 5). This genus is adapted to live in many different environments (6, 7), and it has been used for biocontrol of agricultural plagues (8), for phytoremediation (9, 10), and as a probiotic (11). With the aim of understanding how *Bacillus* species have adapted to conditions in the Atacama Desert, we isolated *Bacillus* strains by mixing 1 g of soil with 9 ml of NaCl 0.8% and incubating at 50°C for 1 h. The solution was decanted, and 1 ml of the supernatant was spread on Luria broth (LB) agar and incubated at 37°C for 4 days under aerobic conditions. One of the colonies recovered was named RP10, and its genome was sequenced. The genomic DNA was purified using a QuickExtract bacterial DNA extraction kit (Epibio), and the library was constructed using a Nextera XT kit (Illumina). Whole-genome shotgun sequencing was performed using 2 × 250-bp (paired-end) reads on an Illumina MiSeq platform. A total of 14.2 million reads was obtained. Reads were filtered for a Phred quality score of at least 20 and assembled using the A5 pipeline (2015 Linux, default parameters) (12). Open reading frame prediction and annotation were performed using Prokka software version 1.11 (13).

Bacillus species identification was achieved by using the JSpeciesWS server online with BLAST (14) average nucleotide identity (ANIb) and MUMmer average nucleotide identity (ANIm) analysis (15–17). The RP10 strain was identified as a member of the species *Bacillus safensis*, and the highest identities were with *B. safensis* FO-36b (16S rRNA, 99.2%; ANIb, 98.83%; ANIm, 99.07%) and *B. safensis* JPL-MERTA-8-2 (ANIb, 98.7%; ANIm, 99.00%). Both strains were isolated from clean rooms at the Jet Propulsion Laboratory at NASA (Pasadena, CA). (18–19). A genome comparison between these three strains by Mauve software (20) and the Rapid Annotation using Subsystem Technology (RAST)-National Microbial Pathogen Database Resource (NMPDR) server with SEED view indicated that the RP10 genome contained an island enriched in phage-related genes (60 kb), an arsenic resistance operon, and other uncharacterized metabolic operons not found in the other two genomes.

The draft genome of *Bacillus safensis* RP10 consists of 3,813,379 bp distributed in 102 contigs, with an average GC content of 41.7%. The draft genome comprises 75 tRNAs and 20 T box leader sequences that are probably involved in a riboswitch mechanism described for Gram-positive bacteria (21). The RAST-NMPDR server with SEED view

Citation Strahsburger E, Zapata F, Pedroso I, Fuentes D, Tapia P, Ponce R, McClelland M, Valdes J. 2019. Draft genome sequence of *Bacillus safensis* RP10, isolated from soil in the Atacama Desert, Chile. *Microbiol Resour Announc* 8:e01150-19. <https://doi.org/10.1128/MRA.01150-19>.

Editor John J. Dennehy, Queens College

Copyright © 2019 Strahsburger et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Erwin Strahsburger, estrahbs@unap.cl.

Received 24 September 2019

Accepted 14 October 2019

Published 31 October 2019

indicates that this strain carries genes and operons for resistance to arsenic (*arsRRBC*), lead (*epsABCDEFFHIKLMN* and *cadA*), copper (*copZA*), and manganese (*mntP*, *mntH*, and *mntABC*) and an operon with potential antimicrobial activity and 34% identity with the bacteriocin operon AS-48 of *Enterococcus faecalis*. Finally, the genome encodes diverse capabilities for synthesizing several vitamins, siderophores, and aromatic compounds, including tryptophan (*trpEDCFBA*) and several multidrug efflux pumps, which are potentially associated with the ability to survive under saline soil conditions with a low content of energy and carbon sources.

Data availability. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [MKXN00000000](https://doi.org/10.1093/bioinformatics/btu153) with contigs under accession numbers [MKXN01000001](https://doi.org/10.1093/bioinformatics/btu153) to [MKXN01000102](https://doi.org/10.1093/bioinformatics/btu153), and the raw reads are available in the SRA under the accession number [PRJNA345377](https://doi.org/10.1093/bioinformatics/btu153).

ACKNOWLEDGMENTS

This work was supported by funds from academic innovation project DE1549, sustained by the Academic Vicerectory, by internal grants VRIIP0123-15 and VRIIP0218-15, supported by the Vicerectory of Research, Innovation, and Postgraduate (VRIIP) of Universidad Arturo Prat, and by the program of InnovaChile CORFO for Attraction of International Centers of Excellence (grant FCR-CSB 09CEII-6991).

REFERENCES

- Olson DM, Dinerstein E. 1998. The global 200: a representation approach to conserving the Earth's most biologically valuable ecoregions. *Conserv Biol* 12:502–515. <https://doi.org/10.1046/j.1523-1739.1998.012003502.x>.
- Betancourt JL, Latorre C, Rech JA, Quade J, Rylander KA. 2000. A 22,000-year record of monsoonal precipitation from northern Chile's Atacama Desert. *Science* 289:1542–1546. <https://doi.org/10.1126/science.289.5484.1542>.
- De Gregori I, Fuentes E, Rojas M, Pinochet H, Potin-Gautier M. 2003. Monitoring of copper, arsenic and antimony levels in agricultural soils impacted and non-impacted by mining activities, from three regions in Chile. *J Environ Monit* 5:287–295. <https://doi.org/10.1039/b211469k>.
- Crits-Christoph A, Robinson CK, Barnum T, Fricke WF, Davila AF, Jedynek B, McKay CP, DiRuggiero J. 2013. Colonization patterns of soil microbial communities in the Atacama Desert. *Microbiome* 1:28. <https://doi.org/10.1186/2049-2618-1-28>.
- Drees KP, Neilson JW, Betancourt JL, Quade J, Henderson DA, Pryor BM, Maier RM. 2006. Bacterial community structure in the hyperarid core of the Atacama Desert, Chile. *Appl Environ Microbiol* 72:7902–7908. <https://doi.org/10.1128/AEM.01305-06>.
- Nicholson WL, McCoy LE, Kerney KR, Ming DW, Golden DC, Schuergler AC. 2012. Aqueous extracts of a Mars analogue regolith that mimics the Phoenix landing site do not inhibit spore germination or growth of model spacecraft contaminants *Bacillus subtilis* 168 and *Bacillus pumilus* SAFR-032. *Icarus* 220:904–910. <https://doi.org/10.1016/j.icarus.2012.06.033>.
- Cortês M, Fuchs FM, Commichau FM, Eichenberger P, Schuergler AC, Nicholson WL, Setlow P, Moeller R. 2019. *Bacillus subtilis* spore resistance to simulated Mars surface conditions. *Front Microbiol* 10:333. <https://doi.org/10.3389/fmicb.2019.00333>.
- Bhattacharya S, Das A, Samadder S, Rajan SS. 2016. Biosynthesis and characterization of a thermostable, alkali-tolerant chitinase from *Bacillus pumilus* JUBCH08 displaying antagonism against phytopathogenic *Fusarium oxysporum*. *3 Biotech* 6:1–8. <https://doi.org/10.1007/s13205-016-0406-x>.
- Tiwari S, Sarangi BK, Thul ST. 2016. Identification of arsenic resistant endophytic bacteria from *Pteris vittata* roots and characterization for arsenic remediation application. *J Environ Manage* 180:359–365. <https://doi.org/10.1016/j.jenvman.2016.05.029>.
- Wu T, Xu J, Liu J, Guo W, Li X, Xia J, Xie W, Yao Z, Zhang Y, Wang R. 2019. Characterization and initial application of endophytic *Bacillus safensis* strain ZY for improving phytoremediation of oil-contaminated saline soils. *Front Microbiol* 10:991. <https://doi.org/10.3389/fmicb.2019.00991>.
- Mu Y, Cong Y. 2019. *Bacillus coagulans* and its applications in medicine. *Benef Microbes* 10:679–688. <https://doi.org/10.3920/BM2019.0016>.
- Tritt A, Eisen JA, Facciotti MT, Darling AE. 2012. An integrated pipeline for *de novo* assembly of microbial genomes. *PLoS One* 7:e42304. <https://doi.org/10.1371/journal.pone.0042304>.
- Seemann T. 2014. Prokka: prokaryotic genome annotation system. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic Local Alignment Search Tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).
- Richter M, Rosselló-Móra R, Glöckner FO, Peplies J. 2016. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931. <https://doi.org/10.1093/bioinformatics/btv681>.
- Espariz M, Zuljan FA, Esteban L, Magni C. 2016. Taxonomic identity resolution of highly phylogenetically related strains and selection of phylogenetic markers by using genome-scale methods: the *Bacillus pumilus* group case. *PLoS One* 11:e0163098. <https://doi.org/10.1371/journal.pone.0163098>.
- Federhen S, Rossello-Mora R, Klenk HP, Tindall BJ, Konstantinidis KT, Whitman WB, Brown D, Labeda D, Ussery D, Garrity GM, Colwell RR, Hasan N, Graf J, Parte A, Yarza P, Goldberg B, Sichtig H, Karsch-Mizrachi I, Clark K, McVeigh R, Pruitt KD, Tatusova T, Falk R, Turner S, Madden T, Kitts P, Kimchi A, Klimke W, Agarwala R, DiCuccio M, Ostell J. 2016. Meeting report: GenBank microbial genomic taxonomy workshop (12–13 May, 2015). *Stand Genomic Sci* 11:15.
- Coil DA, Benardini JN, Eisen JA. 2015. Draft genome sequence of *Bacillus safensis* JPL-MERTA-8-2, isolated from a Mars-bound spacecraft. *Genome Announc* 3:e01360-15. <https://doi.org/10.1128/genomeA.01360-15>.
- Satomi M, La Duc MT, Venkateswaran K. 2006. *Bacillus safensis* sp. nov., isolated from spacecraft and assembly-facility surfaces. *Int J Syst Evol Microbiol* 56:1735–1740. <https://doi.org/10.1099/ijs.0.64189-0>.
- Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.
- Chopin A, Biauudet V, Ehrlich SD. 1998. Analysis of the *Bacillus subtilis* genome sequence reveals nine new T-box leaders. *Mol Microbiol* 29:662–664. <https://doi.org/10.1046/j.1365-2958.1998.00912.x>.