

UCSF

UC San Francisco Previously Published Works

Title

Correlates of lifetime blunt/spliff use among cigarette smokers in substance use disorders treatment

Permalink

<https://escholarship.org/uc/item/78h9m0qt>

Authors

Campbell, Barbara K

Le, Thao

Kapiteni, Kwinoja

et al.

Publication Date

2020-09-01

DOI

10.1016/j.jsat.2020.108064

Peer reviewed

# Correlates of Lifetime Blunt/Spliff Use among Cigarette Smokers in Substance Use Disorders Treatment

Barbara K. Campbell<sup>a</sup>, Thao Le<sup>b</sup>, Kwinoja Kapiteni<sup>b</sup>, Noah R. Gubner<sup>c</sup>, Joseph Guydish<sup>b</sup>

*<sup>a</sup>OHSU/PSU School of Public Health  
Oregon Health and Sciences University  
3181 SW Sam Jackson Park Road  
Portland, OR 97239-3098*

*<sup>b</sup>Philip R. Lee Institute for Health Policy Studies  
University of California San Francisco, CA, USA  
[Thao.Le@ucsf.edu](mailto:Thao.Le@ucsf.edu)  
[Kwinoja.Kapiteni@ucsf.edu](mailto:Kwinoja.Kapiteni@ucsf.edu)  
[Joseph.Guydish@ucsf.edu](mailto:Joseph.Guydish@ucsf.edu)*

*<sup>c</sup>Department of Psychiatry and Behavioral Sciences,  
University of Washington, WA, USA  
[ngubner@uw.edu](mailto:ngubner@uw.edu)*

## **Corresponding Author:**

Barbara K. Campbell, Ph.D.  
OHSU/PSU School of Public Health  
Oregon Health and Sciences University  
3181 SW Sam Jackson Park Road  
Portland, OR 97239-3098  
Phone: 503-221-7074  
Email: [drbarbaracampbell@earthlink.net](mailto:drbarbaracampbell@earthlink.net)

**Declarations of interest:** none

## Abstract

**Background:** Co-use of tobacco and cannabis has been associated with greater dependence on and lower quit rates for both substances. Tobacco/cannabis co-use among individuals with substance use disorders (SUDs), a population with high rates of cigarette smoking, may hinder the effectiveness of smoking cessation interventions. We examined rates of lifetime (i.e., ever vs. never) cannabis use among current cigarette smokers in SUD treatment, and we identified the subgroup who had used tobacco and cannabis together in the form of blunts and/or spliffs. We then examined variables associated with lifetime use of blunts and/or spliffs.

**Methods:** We surveyed 562 clients in 20 residential SUD treatment programs in California, USA in 2019. Measures included demographics, lifetime use of any cannabis product, lifetime use of blunt/spliffs, patterns of tobacco use and smoking cessation-related questions. We asked current cigarette smokers who also reported lifetime cannabis use whether they had ever used blunts and/or spliffs. We then assessed relationships of demographic, tobacco use, use of cannabis/tetrahydrocannabinol (THC) in e-cigarettes/vape pens, and smoking cessation-related variables with ever use versus never use of blunts/spliffs.

**Results:** Among 340 current cigarette smokers, 93.2% (n=317) reported lifetime use of any cannabis product. Among current cigarette smokers with lifetime cannabis use, 64.4% reported lifetime blunt/spliff use. Compared to those who had never used blunts/spliffs, lifetime blunt/spliff users were more likely to be younger (OR = 0.93, 95% CI 0.90- 0.95), more likely to report lifetime use of cigars/cigarillos (OR = 2.95, CI .1.37-6.32), and to have ever used cannabis/THC in e-cigarettes/vape pens (OR = 4.26, CI 1.54-11.80). They were less often ready to quit smoking within 30 days (OR =0.37, CI 0.23 – 0.60), but more likely to want help with smoking cessation (OR = 2.39, CI 1.52-3.77).

**Conclusion:** Current cigarette smokers in SUD treatment reported a high prevalence of lifetime cannabis use. Smokers with a history of blunt/spliff use were more likely to report lifetime use of e-cigarettes/vape pens for cannabis/THC delivery. They wanted help quitting smoking, but felt less prepared to quit in the next 30 days.

Attention to cannabis co-use may be warranted when providing smoking cessation interventions during SUD treatment.

**Keywords:** tobacco, cannabis, blunts/spliffs, substance use, smoking cessation

## 1. Introduction

Rates of co-use of cannabis and tobacco are high among smokers in the general population (Schauer, Berg, Kegler, Donovan, & Windle, 2015a) and have been increasing, particularly in states with legalized medical and/or recreational cannabis use (Goodwin et al., 2018; Wang & Cataldo, 2016). Co-use of tobacco and cannabis has been associated with higher rates of dependence, more difficulty quitting, and higher relapse rates relative to single use of each substance (McClure, Tomko, et al., 2018; Schauer, King, & McAfee, 2017; Voci, Zawertailo, Baliunas, Masood, & Selby, 2020; Weinberger, Platt, Copeland, & Goodwin, 2018). Co-use includes sequential use of cannabis and tobacco as well as co-administration in the form of blunts (i.e., cannabis rolled in cigar wrappers) or spliffs (i.e., cannabis and tobacco rolled together in a joint). Studies of co-administration have reported higher rates of use and poorer psychosocial functioning than use of either substance alone (Tucker et al., 2019), higher rates of dependence for both substances (Fairman, 2015; Ream, Benoit, Johnson, & Dunlap, 2008; Schauer, Rosenberry, & Peters, 2017; Timberlake, 2009) and greater lifetime use of tobacco and illicit drugs (Montgomery & Mantey, 2017).

Individuals with substance use disorders (SUDs) smoke tobacco at rates much higher than the general population. In a national sample of clients enrolled in SUD treatment programs in 2014-2015, 77.9% were current cigarette smokers (Guydish, Tajima, et al., 2016). Efforts to address smoking in SUD treatment over the past thirty years at regulatory, policy, and treatment program levels have achieved modest gains. (Apollonio, Philipps, & Bero, 2016; Skelton et al., 2018; Thurgood, McNeill, Clark-Carter, & Brose, 2016). Nevertheless, unlike the dramatic decrease in smoking within the general population, smoking rates among those with SUDs remain high (Weinberger, Gbedemah, et al., 2018). Co-use of tobacco and cannabis, particularly blunt/spliff use, may hinder the effectiveness of smoking cessation interventions delivered in SUD treatment. Yet, blunt/spliff use may not be addressed when assessing tobacco use in SUD treatment, missing both the additional tobacco/nicotine exposure that occurs through these methods of administration, and identification of a use pattern that may decrease the probability of successful smoking cessation. We did not identify any studies

examining cannabis/tobacco co-use among smokers in treatment for SUDs other than cannabis use disorder (CUD). Among cannabis treatment studies, findings indicate tobacco co-use is associated with poorer cannabis treatment outcomes (Gray et al., 2017), that moderate/high nicotine dependence is associated with increases in smoking after CUD treatment (McClure, Baker, et al., 2018), and that treatment interventions aimed at both cannabis and tobacco cessation may have some efficacy in reducing cannabis use, but evidence is weaker for tobacco cessation (Walsh, McNeill, Purssell, & Duaso, 2020).

The current study was designed to address the research gap regarding cannabis/tobacco co-use among persons with any SUDs. We examined rates of lifetime cannabis use and lifetime use of blunts and/or spliffs among smokers in residential treatment programs in California. We then examined differences between current cigarette smokers reporting lifetime blunt/spliff use and current cigarette smokers reporting never having used blunt/spliffs on variables including demographic characteristics, smoking severity, use of other tobacco products, use of e-cigarettes/vape pens, including e-cigarettes/vape pens for cannabis/THC delivery, and variables associated with smoking cessation.

## **2. Methods**

### *2.1 Program Selection and Participants*

This study surveyed 562 clients from 20 residential SUD treatment programs located in California, USA in 2019. Data from three ongoing studies were combined for this analysis. There were 7 programs in the first study, 9 in the second, and 4 in the third. Eligible programs in the first study had a minimum 20-bed capacity and had applied to participate in a policy development intervention provided by the University of California San Francisco (UCSF) Smoking Cessation Leadership Center. In study two, eligible programs were those expressing interest in creating tobacco-free grounds policies. The third study was designed to improve tobacco intervention in four residential SUD programs in San Francisco, CA. All participating programs were licensed, residential SUD treatment programs and all clients participated in SUD treatment; three program included beds

for prison post-release clients and all programs included some clients with comorbid SUD and mental health problems. Clients in all programs were surveyed during site visits using the same data collection procedures and a core set of tobacco-related measures. All clients present on site visit days were eligible to complete surveys.

## *2.2 Measures*

### *2.2.1. Demographics and Primary Reason for Treatment Entry*

Survey items included demographic information (age, gender, race/ethnicity, education) and primary reason for treatment entry. Clients were asked, “What are you currently in treatment for?” with response options including SUD, mental health disorder, both SUD and mental health disorder, and other. Only clients who responded SUD or both SUD and mental health disorder were asked their primary drug for treatment entry. Based on these two questions, we developed the variable, primary reason for treatment entry, coded as alcohol, stimulants, marijuana/cannabis, opioids, mental health, or other. Gender was coded as male, female, or something else. Race/ethnicity was categorized as non-Hispanic White, African American, Hispanic/ Latino, or other. Education was coded as less than a high school education, high school, or GED equivalent, or greater than a high school education.

### *2.2.2. Tobacco Use.*

Clients were categorized as current cigarette smokers if they responded yes to two survey questions “Have you smoked at least 100 cigarettes in your entire life?” and “Do you currently smoke cigarettes?” (CDC, 2017). Smoking severity was measured using the Heaviness of Smoking Index (HSI) (Heatherton, Kozlowski, Frecker, Rickert, & Robinson, 1989), a 6-point scale based on two self-report items, number of cigarettes smoked per day, and time to first cigarette after awakening. The HSI has demonstrated reliability and validity as a measure of nicotine dependence (Borland, Yong, O'connor, Hyland, & Thompson, 2010). The HSI categorizes level of nicotine dependence as, low (0-1), medium (2-4), or high (5-6; Diaz et al., 2005). Smokers were asked whether their usual cigarette was menthol or non-menthol. Participants also reported lifetime use of

other tobacco products (i.e., “Have you ever in your lifetime used any of the following products?”) coded as smokeless tobacco, cigarillos and/or cigars, and e-cigarettes.

### *2.2.3. Cannabis Use.*

Participants reported lifetime cannabis use (“Have you ever in your lifetime used marijuana/cannabis/hash?”). Those responding “yes” were asked about lifetime use of blunt/spliffs (“Have you ever in your lifetime mixed tobacco and marijuana and smoked them together, as part of the same blunt or spliff?”) Participants who had reported lifetime use of e-cigarettes were asked to report whether they had ever used an e-cigarette with a drug besides nicotine. Those responding yes were asked to select which drugs from a list including the option, marijuana/cannabis/THC/hash.

### *2.2.4. Smoking cessation-related questions*

Participants were asked whether they had quit smoking for at least 24 hours within the past year; those responding yes were asked the number of times they had done so. To identify lifetime use of smoking cessation products, participants were asked whether they had ever used any of the following products to try to quit smoking (coded as use of nicotine replacement therapy [NRT], use of non-NRT medication, and/or use of e-cigarette/vape pens), as well as whether they had ever called the California Smokers Helpline (yes/no). Smokers were also asked whether they were seriously thinking of quitting within 30 days, six months, but not the next 30 days, or not within the next six months (DiClemente et al., 1991). We used readiness to quit within the next 30 days (yes/no) for this analysis. Smokers were asked their primary reason for reducing or quitting smoking (i.e., program requirement, cost of a pack of cigarettes, health concerns, family pressure or encouragement, other, and did not try to quit). Finally, participants were asked whether they wanted help with quitting (yes/no) and whether quitting smoking was a part of their personal treatment plan (yes/no) in their current program.

## *2.3 Data collection procedures*



Data collection was conducted by two research staff members during 1-2-day site visits at each program. Research staff verbally reviewed study information sheets with clients and distributed iPads with unique participant IDs to each potential participant. Potential participants then used the iPads to review the study information page again, and clicked to agree or decline study participation. Those agreeing to participation then completed the online survey. The self-administered surveys, developed using Qualtrics™ (Provo, Utah) software, took approximately 30 minutes to complete. Clients received a \$20.00 gift card for study participation. Study procedures were approved by the institutional review board of the University of California, San Francisco.

#### *2.4 Data Analysis*

Categorical variables were summarized using frequencies and percentages. We identified lifetime blunt/spliff users and non-users among current cigarette smokers based on the lifetime cannabis question and the lifetime blunt/spliff question. See Figure 1. Among current cigarette smokers, lifetime blunt/spliff users vs. non-users were compared on: demographics; drug use (primary drug for treatment entry, cannabis as a primary or secondary drug for treatment entry, lifetime use of cannabis in e-cigarettes/vape pens); tobacco use (HSI, menthol use, lifetime use of smokeless, cigar/cigarillo, and/or e-cigarette/vape pens); and smoking cessation-related variables (lifetime use of smoking cessation products, lifetime use of the California Helpline, readiness to quit within 30 days, desire for help with quitting and quitting as part of their treatment plan). Descriptive statistics (mean, standard deviation) were used to summarize continuous variables. Statistical comparisons were computed using t-test for continuous variables and chi-square test for categorical variables. Last, multivariable logistic regression was used to assess differences between ever/never blunt spliff users, while controlling for covariates. All demographic variables were entered into the regression model. All other variables were included in the model if they significant at a  $p \leq 0.10$  in the univariate comparisons. The model accounted for nesting clients within treatment program. Generalized estimating equation (GEE) method was applied for correlated data. The GEE method was conducted in the SAS GENMOD procedure with a logit link function and repeated

statement. As the rate of missing data was low (3.6%), the multivariable model used complete case analysis. The variance inflation factor (VIF) and tolerance coefficient were used to assess multi-collinearity. We used SAS version 9.4 to conduct the analyses.

### **3. Results**

#### *3.1 Demographics and Drug Use*

Among all 562 clients surveyed, 340 reported current cigarette smoking. Among 340 current cigarette smokers, 317 (93.2 %) reported lifetime use of any cannabis product, and 23 (6.8%) reported never having used a cannabis product. Among current cigarette smokers who reported lifetime use of cannabis, 197 (64.4%) reported lifetime blunt/spliff use, and 109 reported never having used blunt/spliffs. In addition, 11 cases did not respond to the blunt/spliff use survey question and were removed from the analyses, as shown in Figure 1. Among current smokers comprising our blunt/spliff analyses (N=329), average age was 37.8, (s.d.=11.4) with the majority identifying as male, 33.7% identifying as Hispanic/Latino and 38.6% as non-Hispanic white (see Table 1). There were 117 (35.6%) reporting high school or GED diplomas and 119 (36.2%) reporting more than a high school education. Stimulants were reported as primary drug for treatment entry by 47.7%, followed by 21.7% reporting opioids. Cannabis was reported as either a primary or secondary drug for treatment entry by 12.7% of participants, while 23.4% reported using cannabis/THC in e-cigarettes/vape pens.

#### *3.2 Variables Associated with Blunt/Spliff Use*

Table 1 shows results of univariate analyses of demographic and drug use variables for ever vs never blunt/spliff users. Blunt/spliff users were younger ( $p < 0.0001$ ), were more often non-Hispanic white, and were less often African American, ( $p < 0.018$ ). Blunt/spliff users were significantly more likely to report use of cannabis/THC in e-cigarettes/vape pens ( $p < 0.0001$ ). Results of univariate comparisons for tobacco use and smoking cessation-related variables are shown in Table 2. Variables significant at the  $p \leq 0.10$  level and

included in the multivariate model were lifetime use of cigars/cigarillos, lifetime use of smokeless tobacco and lifetime use of e-cigarettes/vape pens, all of which were reported in higher numbers by blunt/spliff users. Smoking-cessation related variables significant at  $p \leq 0.10$  were lifetime use of NRT, lifetime use of e-cigarettes for cutting down or quitting smoking, thinking of quitting in the next 30 days and wanting help with quitting.

Results for the multivariate analysis are shown in Table 3. Participants reporting lifetime use of cannabis/THC in e-cigarettes/vape pens were 4.26 times more likely to report lifetime blunt/spliff use (95% CI: 1.54-11.80). Participants reporting lifetime use of cigars/cigarillos were 2.95 times more likely to report blunt/spliff use (CI: 1.37-6.32). Participants thinking of quitting smoking within the next 30 days were significantly less likely to report blunt/spliff use (OR= 0.37, CI: 0.23-0.60). However, those reporting that they wanted help with quitting smoking were significantly more likely to report lifetime blunt/spliff use (OR=2.39, CI: 1.52-3.77). Multi-collinearity was evaluated using VIF and tolerance coefficient. For the variables in the model, the VIF are small ( $< 2$ ) and the tolerance coefficients are greater than 0.5. These indicators suggest that multicollinearity did not threaten the model.

#### **4. Discussion**

Among clients surveyed in residential treatment in California, 60.5% self-reported current cigarette smoking, a finding indicative of continued high smoking prevalence among individuals with other SUDs. Smokers in this sample reported a very high rate of lifetime cannabis use (93.2%), compatible with other reports of high cannabis use rates among smokers in general population samples (Dierker, Braymiller, Rose, Goodwin, & Selya, 2018; Goodwin et al., 2018; Hindocha et al., 2015) and high rates of tobacco smoking among cannabis users (Schauer, Berg, Kegler, Donovan, & Windle, 2015b; Schauer, King, et al., 2017; Weinberger, Platt, et al., 2018). Additionally, a majority of those reporting lifetime cannabis use also reported lifetime blunt/spliff use. Lifetime blunt/spliff users in our sample were more likely to be younger than non-

blunt/spliff users, and more likely to report lifetime cigar/cigarillo use and use of cannabis/THC in e-cigarettes/vape pens. The association of cigar/cigarillo use with blunt/spliff use may be expected given that blunts are comprised of cannabis rolled in cigar wrappers. This is concerning because blunt use may lead to cigar use among adolescents (Audrain-McGovern, Rodriguez, Alexander, Pianin, & Sterling, 2019) and conversely, cigar use may lead to blunt/spliff use (Fairman & Anthony, 2018). Moreover, multiple tobacco product use has been associated with greater nicotine dependence among smokers in SUD treatment (Guydish, Tajima, et al., 2016). Blunt/spliff users in our sample were also more likely to report lifetime use of e-cigarettes/vape pens for cannabis/THC delivery. The emergence of e-cigarette/vaping product use-associated lung injury (EVALI) has raised alarm regarding acute health hazards associated with e-cigarettes/vape pen use, particularly for cannabis or THC-delivery (King, Jones, Baldwin, & Briss, 2020). Our survey did not ask for information regarding co-administration of cannabis and tobacco via e-cigarette/vape pens which has also been associated with EVALI. This should be addressed in future research.

Current cigarette smokers in our sample who expressed readiness to quit within 30 days were significantly less likely to report lifetime blunt/spliff use. However, those who wanted help with quitting were over two times more likely to report blunt/spliff use. It is possible that smokers with histories of blunt/spliff use in this high smoking population want to quit but feel less confident, thus less ready, to do so and want more help to achieve smoking cessation. This interpretation is supported by evidence showing that co-users of tobacco and cannabis have more difficulty quitting and sustaining smoking cessation (Voci et al., 2020; Weinberger, Platt, et al., 2018)

Quit-related findings in our study offer directions for research and may also inform clinical considerations for addressing cigarette smoking in SUD treatment. Research should examine the relationship between co-use and cigarette smoking cessation outcomes for those in SUD treatment. Readiness interventions that address co-use could be developed and tested , with the goal of facilitating motivation to address cannabis use as a part of quitting tobacco (Guydish, Gruber, et al., 2016). Additionally, consideration should be given to

simultaneous treatment of tobacco and cannabis use in outpatient SUD treatment settings, although this may not be relevant in residential treatment, where, presumably, smokers are not engaged in active cannabis use. Emerging evidence suggests that such interventions are feasible, although the evidence also suggests that simultaneous treatment may be more helpful for cannabis cessation than tobacco cessation (Becker, Haug, Kraemer, & Schaub, 2015; Lee et al., 2019; Walsh et al., 2020). Clinical trials examining simultaneous cannabis/tobacco treatment interventions utilizing evidence-based treatments for both should be studied among co-users in SUD treatment.

These findings should be interpreted with several limitations in mind. As a cross-sectional survey, causal interpretations are not possible. Self-reported cigarette smoking, cannabis, and blunt/spliff use may not have been fully accurate. For example, an additional 47 clients self-reported as non-smokers but blew expired carbon monoxide above the 9 parts per million (ppm) cutoff. These probable, current tobacco smokers were not surveyed about cigarette smoking behaviors, thus not included in our sample. Our measures of lifetime cannabis and blunt/spliff use preclude the ability to assess the relationship of frequency and heaviness of cannabis use with cigarette smoking and nicotine dependence. Additionally, although information regarding recent or current blunt/spliff use was likely to be confounded with length of time in residential treatment, our study's lack of findings regarding recent/current blunt/spliff use also precludes understanding the impact of that use on smoking cessation-related variables. There are generalizability limitations related to our sample selection, since it was a convenience sample restricted to residential programs within California. However, given the legalization of medical cannabis in California in 1996 and recreational cannabis in 2016, cross-sectional data regarding co-use among smokers in SUD treatment in California provides information on which to build regarding possible increases in co-use rates and consequences of co-use that may occur as recreational cannabis becomes legally available across states. The racial/ethnic composition of our sample may be another limitation. African Americans comprised 17.6% of our study sample, higher than the estimated 11.6% of adults in the United States with SUDS who are African American (McCance-Katz, 2019). However, reports indicate a higher prevalence

of blunt use among African Americans than other racial/ethnic groups, thus they may be underrepresented in our study sample (Cohn, Johnson, Ehlke, & Villanti, 2016; Montgomery & Mantey, 2017).

#### *4.1 Conclusion*

Co-use of tobacco and cannabis has been associated with more severe nicotine dependence and more difficulty quitting cigarette smoking, but has been understudied in the high numbers of smokers in SUD treatment. We found a high rate of lifetime cannabis use, including blunt/spliff use, among smokers in residential SUD treatment. Cigarette smokers expressing less readiness to quit, but requesting help with quitting were more likely to have a history of blunt/spliff use, suggesting blunt/spliff use may be a variable that affects the probability of smoking cessation success. Co-use of cannabis and tobacco should be addressed for smokers in SUD treatment.

#### **Funding Sources**

This work was supported by the California Tobacco Control Program (18-10025), by the California Tobacco Related Disease Research Program (TRDRP 27IR-0040, 28CP-0038), and by the NCI Cancer Center Support Grant (P30 CA082103). The content is solely the responsibility of the authors.

#### **Acknowledgments**

The authors would like to thank the directors, staff, and patients enrolled in the participating treatment programs.

## References

- Apollonio, D., Philipps, R., & Bero, L. (2016). Interventions for tobacco use cessation in people in treatment for or recovery from substance use disorders. *Cochrane Database of Systematic Reviews*(11).
- Audrain-McGovern, J., Rodriguez, D., Alexander, E., Pianin, S., & Sterling, K. L. (2019). Association Between Adolescent Blunt Use and the Uptake of Cigars. *JAMA Network Open*, 2(12), e1917001-e1917001.
- Becker, J., Haug, S., Kraemer, T., & Schaub, M. P. (2015). Feasibility of a group cessation program for co-smokers of cannabis and tobacco. *Drug and alcohol review*, 34(4), 418-426.
- Borland, R., Yong, H.-H., O'connor, R., Hyland, A., & Thompson, M. (2010). The reliability and predictive validity of the Heaviness of Smoking Index and its two components: findings from the International Tobacco Control Four Country study. *Nicotine & Tobacco Research*, 12(suppl 1), S45-S50.
- CDC. (2017). *National Center for Health Statistics. Adult Tobacco Use Information. National Health Interview Survey*. Retrieved from [https://www.cdc.gov/nchs/nhis/tobacco/tobacco\\_glossary.htm](https://www.cdc.gov/nchs/nhis/tobacco/tobacco_glossary.htm).
- Cohn, A., Johnson, A., Ehlke, S., & Villanti, A. C. (2016). Characterizing substance use and mental health profiles of cigar, blunt, and non-blunt marijuana users from the National Survey of Drug Use and Health. *Drug and alcohol dependence*, 160, 105-111.
- Diaz, F. J., Jané, M., Saltó, E., Pardell, H., Salleras, L., Pinet, C., & De Leon, J. (2005). A brief measure of high nicotine dependence for busy clinicians and large epidemiological surveys. *Australian & New Zealand Journal of Psychiatry*, 39(3), 161-168.
- DiClemente, C. C., Prochaska, J. O., Fairhurst, S. K., Velicer, W. F., Velasquez, M. M., & Rossi, J. S. (1991). The process of smoking cessation: an analysis of precontemplation, contemplation, and preparation stages of change. *Journal of consulting and clinical psychology*, 59(2), 295.
- Dierker, L., Braymiller, J., Rose, J., Goodwin, R., & Selya, A. (2018). Nicotine dependence predicts cannabis use disorder symptoms among adolescents and young adults. *Drug and alcohol dependence*, 187, 212-220.

- Fairman, B. J. (2015). Cannabis problem experiences among users of the tobacco–cannabis combination known as blunts. *Drug and alcohol dependence, 150*, 77-84.
- Fairman, B. J., & Anthony, J. C. (2018). Does starting to smoke cigars trigger onset of cannabis blunt smoking? *Nicotine and Tobacco Research, 20*(3), 355-361.
- Goodwin, R. D., Pacek, L. R., Copeland, J., Moeller, S. J., Dierker, L., Weinberger, A., . . . Hasin, D. S. (2018). Trends in daily cannabis use among cigarette smokers: United States, 2002–2014. *American journal of public health, 108*(1), 137-142.
- Gray, K. M., Sonne, S. C., McClure, E. A., Ghitza, U. E., Matthews, A. G., McRae-Clark, A. L., . . . Mooney, L. J. (2017). A randomized placebo-controlled trial of N-acetylcysteine for cannabis use disorder in adults. *Drug and alcohol dependence, 177*, 249-257.
- Guydish, J., Gruber, V. A., Le, T., Tajima, B., Andrews, K. B., Leo, H., . . . Tsoh, J. Y. (2016). A pilot study of a readiness group to increase initiation of smoking cessation services among women in residential addiction treatment. *Journal of Substance Abuse Treatment, 63*, 39-45.
- Guydish, J., Tajima, B., Pramod, S., Le, T., Gubner, N. R., Campbell, B., & Roman, P. (2016). Use of multiple tobacco products in a national sample of persons enrolled in addiction treatment. *Drug and alcohol dependence, 166*, 93-99.
- Heatherton, T. F., Kozlowski, L. T., Frecker, R. C., Rickert, W., & Robinson, J. (1989). Measuring the heaviness of smoking: using self-reported time to the first cigarette of the day and number of cigarettes smoked per day. *Addiction, 84*(7), 791-800.
- Hindocha, C., Shaban, N. D., Freeman, T. P., Das, R. K., Gale, G., Schafer, G., . . . Curran, H. V. (2015). Associations between cigarette smoking and cannabis dependence: a longitudinal study of young cannabis users in the United Kingdom. *Drug and alcohol dependence, 148*, 165-171.
- King, B. A., Jones, C. M., Baldwin, G. T., & Briss, P. A. (2020). The EVALI and Youth Vaping Epidemics—Implications for Public Health. *New England Journal of Medicine*.



- Lee, D. C., Walker, D. D., Hughes, J. R., Brunette, M. F., Scherer, E., Stanger, C., . . . Budney, A. J. (2019). Sequential and simultaneous treatment approaches to cannabis use disorder and tobacco use. *Journal of Substance Abuse Treatment*, 98, 39-46.
- McCance-Katz, E. F. (2019). The National Survey on Drug Use and Health: 2017. *Substance Abuse and Mental Health Services Administration*. <https://www.samhsa.gov/data/sites/default/files/nsduh-ppt-09-2018.pdf>. Accessed May, 7.
- McClure, E. A., Baker, N. L., Sonne, S. C., Ghitza, U. E., Tomko, R. L., Montgomery, L., . . . Gray, K. M. (2018). Tobacco use during cannabis cessation: Use patterns and impact on abstinence in a National Drug Abuse Treatment Clinical Trials Network study. *Drug and alcohol dependence*, 192, 59-66.
- McClure, E. A., Tomko, R. L., Salazar, C. A., Akbar, S. A., Squeglia, L. M., Herrmann, E., . . . Peters, E. N. (2018). Tobacco and cannabis co-use: Drug substitution, quit interest, and cessation preferences. *Experimental and clinical psychopharmacology*.
- Montgomery, L., & Mantey, D. S. (2017). Correlates of blunt smoking among African American, Hispanic/Latino, and white adults: results from the 2014 national survey on drug use and health. *Substance use & misuse*, 52(11), 1449-1459.
- Ream, G. L., Benoit, E., Johnson, B. D., & Dunlap, E. (2008). Smoking tobacco along with marijuana increases symptoms of cannabis dependence. *Drug and alcohol dependence*, 95(3), 199-208.
- Schauer, G. L., Berg, C. J., Kegler, M. C., Donovan, D. M., & Windle, M. (2015a). Assessing the overlap between tobacco and marijuana: Trends in patterns of co-use of tobacco and marijuana in adults from 2003–2012. *Addictive behaviors*, 49, 26-32.
- Schauer, G. L., Berg, C. J., Kegler, M. C., Donovan, D. M., & Windle, M. (2015b). Differences in tobacco product use among past month adult marijuana users and nonusers: findings from the 2003–2012 National Survey on Drug Use and Health. *Nicotine & Tobacco Research*, 18(3), 281-288.


- Schauer, G. L., King, B. A., & McAfee, T. A. (2017). Prevalence, correlates, and trends in tobacco use and cessation among current, former, and never adult marijuana users with a history of tobacco use, 2005–2014. *Addictive behaviors*, 73, 165-171.
- Schauer, G. L., Rosenberry, Z. R., & Peters, E. N. (2017). Marijuana and tobacco co-administration in blunts, spliffs, and mulled cigarettes: A systematic literature review. *Addictive behaviors*, 64, 200-211.
- Skelton, E., Tzelepis, F., Shakeshaft, A., Guillaumier, A., McCrabb, S., & Bonevski, B. (2018). Integrating smoking cessation care in alcohol and other drug treatment settings using an organizational change intervention: a systematic review. *Addiction*, 113(12), 2158-2172.
- Thurgood, S. L., McNeill, A., Clark-Carter, D., & Brose, L. S. (2016). A systematic review of smoking cessation interventions for adults in substance abuse treatment or recovery. *Nicotine & Tobacco Research*, 18(5), 993-1001.
- Timberlake, D. S. (2009). A comparison of drug use and dependence between blunt smokers and other cannabis users. *Substance use & misuse*, 44(3), 401-415.
- Tucker, J. S., Pedersen, E. R., Seelam, R., Dunbar, M. S., Shih, R. A., & D'Amico, E. J. (2019). Types of cannabis and tobacco/nicotine co-use and associated outcomes in young adulthood. *Psychol Addict Behav*, 33(4), 401-411. doi:10.1037/adb0000464
- Voci, S., Zawertailo, L., Baliunas, D., Masood, Z., & Selby, P. (2020). Is cannabis use associated with tobacco cessation outcome? An observational cohort study in primary care. *Drug and alcohol dependence*, 206, 107756.
- Walsh, H., McNeill, A., Purssell, E., & Duaso, M. (2020). A systematic review and Bayesian meta-analysis of interventions which target or assess co-use of tobacco and cannabis in single or multi-substance interventions. *Addiction*.
- Wang, J. B., & Cataldo, J. K. (2016). Medical marijuana legalization and co-use in adult cigarette smokers. *American journal of health behavior*, 40(2), 205-214.

Weinberger, A. H., Gbedemah, M., Wall, M. M., Hasin, D. S., Zvolensky, M. J., & Goodwin, R. D. (2018).

Cigarette use is increasing among people with illicit substance use disorders in the United States, 2002–14: emerging disparities in vulnerable populations. *Addiction*, 113(4), 719-728.

Weinberger, A. H., Platt, J., Copeland, J., & Goodwin, R. D. (2018). Is Cannabis Use Associated With Increased Risk of Cigarette Smoking Initiation, Persistence, and Relapse? Longitudinal Data From a Representative Sample of US Adults. *The Journal of clinical psychiatry*, 79(2).

**Figure 1. Selection of participants among current cigarette smokers in residential SUD treatment in California who report ever vs. never blunt/spliff use**



**Table 1. Demographics and drug use among current cigarette smokers in residential SUD treatment in California who report ever vs. never blunt/spliff use (N = 329)**

	Mean (SD) or n (%)			
	Total (N=329)	Smokers reporting ever blunt spliff use (n=197)	Smokers reporting never blunt spliff use (n=132)	p-value
Demographics				
Age	37.8 (11.4)	34.0 (9.0)	43.5 (12.1)	<0.0001
Gender <sup>1</sup>				
Male	242 (73.8%)	150 (76.1%)	92 (70.2%)	
Female	81 (24.7%)	44 (22.3%)	37 (28.2%)	
Something else	5 (1.5%)	3 (1.5%)	2 (1.5%)	
Race/ethnicity				0.018
Hispanic/Latino	111 (33.7%)	67 (34.0%)	44 (33.3%)	
African American	58 (17.6%)	25 (12.7%)	33 (25.0%)	
Non- Hispanic White	127 (38.6%)	86 (43.7%)	41 (31.1%)	
Other	33 (10.0%)	19 (9.6%)	14 (10.6%)	
Education				0.355
<HS	93 (28.3%)	50 (25.4%)	43 (32.6%)	
HS/GED	117 (35.6%)	72 (36.6%)	45 (34.1%)	
>HS	119 (36.2%)	75 (38.1%)	44 (33.3%)	
Primary reason for treatment entry <sup>1</sup>				0.156
Alcohol	51 (15.8%)	28 (14.6%)	23 (17.6%)	
Stimulant	154 (47.7%)	89 (46.4%)	65 (49.6%)	
Marijuana/Cannabis	11 (3.4%)	7 (3.7%)	4 (3.1%)	
Opioids	70 (21.7%)	50 (26.0%)	20 (15.3%)	
Mental health or other	37 (11.5%)	18 (9.4%)	19 (14.5%)	
Marijuana/cannabis as primary or secondary drug for treatment entry <sup>1,2</sup>	37 (12.7%)	25 (14.0%)	12 (10.6%)	0.402
Use of cannabis/THC in e-cigarettes/vape pens <sup>1</sup>	76 (23.4%)	69 (35.2%)	7 (5.4%)	<0.0001

<sup>1</sup>Missing data: Gender (2), Primary drug for treatment entry (6), Marijuana/cannabis as primary or secondary drug for treatment entry (1), Use of cannabis/THC in e-cigarettes/vape pens (4)

<sup>2</sup>Excluded 37 smokers whose reason for treatment entry was mental health or other

**Table 2. Tobacco use characteristics among current cigarette smokers in residential SUD treatment in California who report ever vs. never blunt/spliff use**

	Mean (SD) or n (%)		p-value
	Smokers reporting ever blunt spliff use (N=197)	Smokers reporting never blunt spliff use (N=132)	
<b>Tobacco/Nicotine Use</b>			
Heaviness of Smoking Index (HSI)	2.1 (1.3)	2.0 (1.5)	0.563
Menthol as usual cigarette	64 (32.5%)	42 (31.8%)	0.899
Lifetime cigars/cigarillos <sup>1</sup>	168 (85.7%)	68 (53.1%)	<b>&lt;0.0001</b>
Lifetime smokeless	106 (53.8%)	32 (24.2%)	<b>&lt;0.0001</b>
Lifetime e-cigarettes/vape pens <sup>1</sup>	150 (76.5%)	58 (45.0%)	<b>&lt;0.0001</b>
<b>Smoking cessation-related variables</b>			
Past year number of quit attempts	4.5 (10.1)	5.8 (13.2)	0.343
Main reason for quitting/cutting down <sup>1</sup>			0.373
Treatment program requirement	33 (16.8%)	14 (10.7%)	
Cost of a pack of cigarettes	33 (16.8%)	19 (14.5%)	
Health concerns	63 (32.1%)	51 (38.9%)	
Family (pressure, encouragement, being a good model)	15 (7.7%)	16 (12.2%)	
Other reason	26 (13.3%)	15 (11.5%)	
I have not tried to reduce my smoking/quitting	26 (13.3%)	16 (12.2%)	
<b>Methods Used for Quitting</b>			
Lifetime NRT use <sup>1</sup>	101 (51.3%)	48 (36.6%)	<b>0.009</b>
Lifetime non-NRT medication <sup>1</sup>	16 (8.5%)	11 (8.7%)	0.935
Lifetime e-cigarettes/vape pens use <sup>1</sup>	107 (55.7%)	40 (31.0%)	<b>&lt;0.0001</b>
Lifetime California Smokers Helpline use <sup>1</sup>	15 (7.7%)	12 (9.1%)	0.642
Readiness- Thinking of quitting next 30 days	53 (26.9%)	52 (39.4%)	<b>0.017</b>
Want help with quitting <sup>1</sup>	83 (42.1%)	41 (31.3%)	<b>0.048</b>
Quitting part of personal treatment plan <sup>1</sup>	60 (30.6%)	40 (30.3%)	0.952

<sup>1</sup>Missing data: Lifetime cigars/cigarillos (5), Lifetime e-cigarettes/vape pens (4), Main reason for quitting/cutting down (2), Lifetime NRT use (1), Lifetime non- NRT medication (14), Lifetime e-cigarettes/vape pens use (8), Lifetime California Smokers Helpline use (1), Want help with quitting (1), Quitting part of personal treatment plan (1)

**Table 3. Factors associated with blunt/spliff use among current cigarette smokers in residential SUD treatment in California**

	Adjusted OR (95% CI)	p <sup>1</sup>
<b>Use of cannabis/THC in e-cigarettes/vape pens</b>		
No (Ref)	1	
Yes	4.26 (1.54- 11.80)	<b>0.005</b>
<b>Tobacco/Nicotine Use</b>		
Lifetime cigars/cigarillos		
No (Ref)	1	
Yes	2.95 (1.37- 6.32)	<b>0.006</b>
Lifetime Smokeless		
No (Ref)	1	
Yes	1.71 (0.75- 3.87)	0.201
Lifetime e-cigarettes/vape pens		
No (Ref)	1	
Yes	1.01 (0.55- 1.84)	0.982
<b>Smoking cessation-related variables</b>		
Lifetime NRT use		
No (Ref)	1	
Yes	1.24 (0.65- 2.36)	0.513
Lifetime e-cigarette/vape pen use for quitting		
No (Ref)	1	
Yes	0.97 (0.53- 1.80)	0.929
Readiness- Thinking of quitting next 30 days		
No (Ref)	1	
Yes	0.37 (0.23- 0.60)	<b>&lt;0.0001</b>
Want help with quitting		
No (Ref)	1	
Yes	2.39 (1.52- 3.77)	<b>&lt;0.001</b>
<b>Demographics</b>		
Age	0.93 (0.90- 0.95)	<b>&lt;0.0001</b>
Gender		
Male	1	
Female	0.72 (0.38- 1.34)	0.298
Other	0.74 (0.49- 1.12)	0.156
Race/Ethnicity		
Hispanic (Ref)	1	
Non-Hispanic White	1.55 (0.78- 3.07)	0.208
Non-Hispanic Black	1.06 (0.39- 2.84)	0.913
Non-Hispanic Other	0.96 (0.42- 2.21)	0.924
Education		
<HS	1	
HS/GED	1.62 (0.91- 2.87)	0.102
>HS	1.44 (0.71- 2.92)	0.316

<sup>1</sup>Logistic regression with clients nested within clinic