UC San Diego

UC San Diego Previously Published Works

Title

Neutrino mass hierarchy and stepwise spectral swapping of supernova neutrino flavors.

Permalink

<https://escholarship.org/uc/item/78k0c86x>

Journal

Physical review letters, 99(24)

ISSN

0031-9007

Authors

Duan, Huaiyu
Fuller, George M
Carlson, J
et al.

Publication Date

2007-12-01

DOI

10.1103/physrevlett.99.241802

Peer reviewed

Neutrino Mass Hierarchy and Stepwise Spectral Swapping of Supernova Neutrino Flavors

Huaiyu Duan,¹ George M. Fuller,¹ J. Carlson,² and Yong-Zhong Qian³

¹*Department of Physics, University of California, San Diego, La Jolla, California 92093-0319, USA*

²*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

³*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

(Received 7 July 2007; published 14 December 2007)

We examine a phenomenon recently predicted by numerical simulations of supernova neutrino flavor evolution: the swapping of supernova ν_e and $\nu_{\mu,\tau}$ energy spectra below (above) energy E_C for the normal (inverted) neutrino mass hierarchy. We present the results of large-scale numerical calculations which show that in the normal neutrino mass hierarchy case, E_C decreases as the assumed effective 2×2 vacuum $\nu_e \rightleftharpoons \nu_{\mu,\tau}$ mixing angle ($\approx \theta_{13}$) is decreased. In contrast, these calculations indicate that E_C is essentially independent of the vacuum mixing angle in the inverted neutrino mass hierarchy case. With a good neutrino signal from a future galactic supernova, the above results could be used to determine the neutrino mass hierarchy even if θ_{13} is too small to be measured by terrestrial neutrino oscillation experiments.

DOI: 10.1103/PhysRevLett.99.241802

PACS numbers: 14.60.Pq, 97.60.Bw

In this Letter we point out how two grand themes in contemporary science, physics beyond the standard model of elementary particles and the physics of stars undergoing gravitational collapse, overlap in a way that could allow unique insight into the nature of neutrinos. Recent experiments have established that neutrinos have nonvanishing rest masses and that the flavor states ν_e , ν_μ , and ν_τ for these particles are mixtures of the vacuum mass eigenstates ν_1 , ν_2 , and ν_3 (see, e.g., Refs. [1,2] for recent reviews). However, key issues remain unresolved. Among these is the nature of the neutrino mass hierarchy: the sign of the mass-squared difference $\delta m_{31}^2 = m_3^2 - m_1^2 \approx \pm \delta m_{\text{atm}}^2$ remains unknown. Here δm_{atm}^2 is the neutrino mass-squared difference associated with atmospheric neutrino oscillations, and the plus (minus) sign corresponds to the normal (inverted) neutrino mass hierarchy. Conventional laboratory experimental resolution of the mass hierarchy issue is problematic, in part because neutrino rest masses are tiny and because θ_{13} , the mixing angle relating ν_e to ν_3 , is small. One possible way to probe the neutrino mass hierarchy is to analyze neutrino signals from galactic supernovae. Supernova neutrinos can experience significant flavor transformation through the Mikheyev-Smirnov-Wolfenstein (MSW) effect [3,4] as they stream out from the surface of a proto-neutron star (with very high matter density) into the vacuum. In addition, it has been pointed out that neutrino-neutrino forward scattering can provide an additional source for neutrino refractive indices [5–8]. This neutrino self-coupling is especially important in the supernova environment because neutrino fluxes are large. Therefore, previous studies of supernova neutrino oscillations based on the pure MSW effect [9–15] may not apply in some scenarios. Following suggestions in Refs. [16–18], in this Letter we present and analyze new large-scale numerical calculations of supernova neutrino flavor evolution that suggest a novel method to determine the neutrino

mass hierarchy. This method is independent of absolute neutrino masses and can work even for tiny θ_{13} .

Our method is based on a stunning feature revealed by recent numerical simulations of supernova neutrino flavor evolution [16,19]: (i) For the normal neutrino mass hierarchy case, ν_e and $\nu_{\mu,\tau}$ swap their energy spectra at energies below a transition energy E_C , but retain their original spectra at higher energies; (ii) for the inverted neutrino mass hierarchy case, the situation is exactly the opposite. This phenomenon is known as “stepwise spectral swapping” [16] or “spectral split” [18].

The stepwise swapping of the ν_e and $\nu_{\mu,\tau}$ energy spectra has its origin in nonlinear neutrino self-coupling. Assuming coherent neutrino propagation and the efficacy of the mean field approach [20,21], for 2×2 flavor evolution, it is possible to define the neutrino flavor isospin (NFIS) as [22]

$$\mathbf{s}_\nu \equiv \psi_\nu^\dagger \frac{\boldsymbol{\sigma}}{2} \psi_\nu \quad \text{and} \quad \mathbf{s}_{\bar{\nu}} \equiv (\sigma_y \psi_{\bar{\nu}})^\dagger \frac{\boldsymbol{\sigma}}{2} (\sigma_y \psi_{\bar{\nu}}) \quad (1)$$

for a neutrino (with flavor wave function ψ_ν) and an antineutrino ($\psi_{\bar{\nu}}$), respectively. Flavor evolution for neutrino or antineutrino mode i is described by precession of a corresponding NFIS $\mathbf{s}_i$ around an effective field:

$$\frac{d}{dt} \mathbf{s}_i = \mathbf{s}_i \times \left[\omega_i \mathbf{H}_V - \sqrt{2} G_F n_e \hat{\mathbf{e}}_z^f - 2\sqrt{2} G_F \sum_j (1 - \cos \vartheta_{ij}) n_j \mathbf{s}_j \right]. \quad (2)$$

Here $\hat{\mathbf{e}}_{x,y,z}^f$ are the flavor-basis unit vectors in flavor space, $\mathbf{H}_V \equiv -\sin 2\theta_v \hat{\mathbf{e}}_x^f + \cos 2\theta_v \hat{\mathbf{e}}_z^f$ generates vacuum mixing for a nonvanishing effective mixing angle θ_v , $\omega_i = \pm \delta m^2 / 2E_i$ is the vacuum precession angular velocity around $\mathbf{H}_V$ for a NFIS corresponding to a neutrino (plus sign) or antineutrino (minus sign) with energy E_i , G_F is the

Fermi constant, n_e is the net electron number density, ϑ_{ij} is the angle between the propagation directions of neutrinos in modes i and j , and n_j is the number density of neutrinos in mode j . Because flavor transformation in the $\nu_e \rightleftharpoons \nu_{\mu,\tau}$ and $\bar{\nu}_e \rightleftharpoons \bar{\nu}_{\mu,\tau}$ channels is the most important in supernovae (e.g., for shock reheating [23,24] and nucleosynthesis [24–27]), and because δm_{atm}^2 will give flavor transformation deeper in the supernova envelope than will $\delta m_{\odot}^2$, the mass-squared difference associated with solar neutrino oscillations, we take $\delta m^2 = \pm 3 \times 10^{-3} \text{ eV}^2 \simeq \pm \delta m_{\text{atm}}^2$ and $\theta_v \simeq \theta_{13} \ll 1$. For this 2×2 mixing we use ν_{τ^*} to designate the relevant linear combination of ν_μ and ν_τ [28].

In a stepwise-swapping scenario, the probability $P_{\nu\nu}$ for neutrinos to remain in their initial flavor states is

$$P_{\nu\nu}(\omega) \simeq \frac{1}{2} \left[1 - \text{sgn}(\omega - \omega_{\text{pr}}^0) \right], \quad (3)$$

where $\text{sgn}(\xi) = \xi/|\xi|$ is the sign of ξ , and $\omega_{\text{pr}}^0 = \delta m^2/2E_C$ specifies the transition energy E_C . Equation (3) is slightly different from Eq. (57b) in Ref. [29], with different conventions for θ_v and δm^2 . This stepwise spectral swapping feature has been demonstrated in two different approaches [16,19,30,31]: “multiangle” simulations, where flavor evolution on independently followed neutrino trajectories is self-consistently coupled, and “single-angle” simulations, where the evolution history of radially propagating neutrinos is assumed to apply to all trajectories. Single-angle calculations capture the qualitative features of stepwise spectral swapping, and they suggest the following generic explanation for this phenomenon.

Because neutrinos and antineutrinos are in flavor eigenstates when they leave the neutrino sphere, they naturally form a “bipolar system” in which the corresponding NFIS’s form two oppositely oriented groups [22]. (Note that $\nu_e/\bar{\nu}_{\tau^*}$ and $\bar{\nu}_e/\nu_{\tau^*}$ correspond to NFIS’s in the directions of $+\hat{\mathbf{e}}_z^f$ and $-\hat{\mathbf{e}}_z^f$, respectively.) This bipolar system behaves like a gyroscopic pendulum in flavor space [32], which can have both nutation and precession modes. These neutrinos and antineutrinos initially follow a quasistatic, MSW-like solution near the neutrino sphere before being driven away from this solution by the collective nutation of the gyroscopic pendulum [29]. Subsequently, the gyroscopic pendulum can execute regular precession around $\mathbf{H}_V$, corresponding to the collective precession of NFIS’s in flavor space [17]. This precession, although not perfectly regular, is indeed found in both single-angle and multiangle simulations [16,17]. If NFIS’s stay in the regular collective precession mode, a stepwise spectral swapping given by Eq. (3) will occur when the neutrino fluxes decrease toward 0 [16–18]. In this case, ω_{pr}^0 in Eq. (3) is just the precession angular velocity for vanishing neutrino fluxes.

Strictly speaking, the regular precession mode obtains for $n_e = 0$ where the “lepton number”

$$\mathcal{L} \equiv \int_0^\infty [f_{\nu_1}(E) - f_{\nu_3}(E) - f_{\bar{\nu}_1}(E) + f_{\bar{\nu}_3}(E)] dE \quad (4)$$

is conserved [32]. Here $f_{\nu_1(\bar{\nu}_1)}(E)$ and $f_{\nu_3(\bar{\nu}_3)}(E)$ are the distribution functions specifying the populations of the corresponding neutrino (antineutrino) vacuum mass eigenstates within energy interval dE . These are normalized to the total (summing over all states) neutrino number density n_ν^{tot} and in general evolve with time. The conservation of $\mathcal{L}$ is exact for $n_e = 0$ and holds even when neutrino number densities change [17]. Because the presence of the matter field does not change the collective precession qualitatively [17,29], the conservation of $\mathcal{L}$ can be used to compute E_C [18].

In the inverted neutrino mass hierarchy case, flavor transformation is suppressed when n_e and neutrino fluxes are high. As ν_e are dominant in supernovae, the bipolar system of neutrinos and antineutrinos resembles a gyroscopic pendulum near its highest point (displacement angle equal to π) in flavor space. For a simple bipolar system initially consisting of monoenergetic ν_e and $\bar{\nu}_e$, the analogy is exact, with the initial displacement angle being $\pi - 2\theta_v$. When the total neutrino flux decreases below some critical value, the flavor pendulum evolves away from its maximum displacement [17,32]. Its nutation then pushes neutrinos and antineutrinos into the collective precession mode. The presence of a matter field does not change this nutation qualitatively, but effectively reduces the mixing angle [22,32]. Therefore, $\mathcal{L}$ is essentially unchanged before the precession mode begins. As a result, when n_e and neutrino fluxes become small, $f_{\nu_1(\bar{\nu}_1)}(E)$ and $f_{\nu_3(\bar{\nu}_3)}(E)$ are related simply through Eq. (3) to the initial neutrino energy spectra at the neutrino sphere. Specifically, for $\theta_v \ll 1$ we have

$$\begin{aligned} \mathcal{L} &\simeq \int_0^{E_C} [f_{\nu_e}(E) - f_{\nu_{\tau^*}}(E)] dE \\ &\quad + \int_{E_C}^\infty [f_{\nu_{\tau^*}}(E) - f_{\nu_e}(E)] dE + \frac{n_{\bar{\nu}_e} - n_{\bar{\nu}_{\tau^*}}}{n_\nu^{\text{tot}}} \quad (5) \\ &\simeq \frac{n_{\nu_e} - n_{\nu_{\tau^*}} - n_{\bar{\nu}_e} + n_{\bar{\nu}_{\tau^*}}}{n_\nu^{\text{tot}}}, \end{aligned}$$

where, e.g., $f_{\nu_e}(E)$ and $n_{\nu_e}(E)$ are the initial spectrum (normalized to n_ν^{tot}) and number density, respectively, of ν_e at the neutrino sphere. The transition energy E_C can then be found from Eq. (5).

The conservation of $\mathcal{L}$ can also be used to find E_C for the normal mass hierarchy case. However, in this case, $\mathcal{L}$ cannot be related simply to the initial neutrino spectra at the neutrino sphere through Eq. (3). This is because there is a resonance in the quasistatic MSW-like solution initially followed by neutrinos and antineutrinos. For example, in the large neutrino luminosity limit, all neutrinos and antineutrinos are synchronized [33] and experience simultaneously an MSW-like resonance near the radius where a single ν_e with a representative energy E_{sync} would encoun-

ter a conventional MSW resonance [26]. During this MSW-like evolution, $\mathcal{L}$ is decreased. Using the initial spectra for supernova neutrinos and Eqs. (3) and (4), it can be shown that the less $\mathcal{L}$ is reduced, the smaller E_C becomes. If θ_v is tiny and/or the neutrino luminosities are not large enough, the MSW-like conversion will be non-adiabatic, and $\mathcal{L}$ will change very little. In this case, $E_C \rightarrow 0$ and the stepwise nature of the swapping of ν_e and $\nu_{\mu,\tau}$ spectra becomes unobservable. On the other hand, given sufficiently large values of θ_v and/or neutrino luminosities, MSW-like flavor conversion can be adiabatic and efficient for a large range of neutrino energies. This can engender nearly complete swapping of the entire neutrino and anti-neutrino spectra and, consequently, $\mathcal{L}$ can retain its magnitude but reverse its sign. In this case, E_C for the subsequent stepwise spectral swapping is roughly the same as in the inverted neutrino mass hierarchy case.

Though the transition energy E_C is sensitive to θ_v in the normal mass hierarchy case, it appears to be essentially independent of θ_v in the inverted mass hierarchy case. We have carried out multiangle simulations under the same conditions as those discussed in Refs. [16,19] except with smaller θ_v . The probability $P_{\nu_e\nu_e}(E, \vartheta_0)$ at radius $r = 250$ km is plotted in Fig. 1 for both a normal mass hierarchy case ($\theta_v = 0.01$, left panel) and an inverted mass hierarchy case ($\theta_v = 10^{-9}$, right panel). Here ϑ_0 is the angle between the propagation direction of the neutrino and the normal at its emission position on the neutrino sphere (see Fig. 1 of Ref. [19]). A comparison of the results shown in Fig. 1 with those for $\theta_v = 0.1$ shown in Fig. 3 of Ref. [19] is revealing. In the normal neutrino mass hierarchy case, E_C decreases from ~ 10 MeV to ~ 3 MeV as θ_v is reduced from 0.1 to 0.01. However, for the inverted

neutrino mass hierarchy case, E_C is essentially unchanged as θ_v is decreased by 8 orders of magnitude. The value $E_C \simeq 8.4$ MeV calculated from Eq. (5) for this case agrees very well with the numerical results. We note that E_C has a slight dependence on neutrino trajectory ($\cos\vartheta_0$) in the multiangle simulations for $\theta_v \ll 0.1$.

The apparent insensitivity of E_C to θ_v in the inverted neutrino mass hierarchy case requires discussion. We note that the neutrino system transitions from the MSW-like evolution to the collective precession mode through nutation. If the system does not develop significant nutation while it is in the collective flavor transformation regime, it will not enter the collective precession mode, and therefore, stepwise spectral swapping will not occur. For a uniform and isotropic gas of monoenergetic neutrinos initially in pure ν_e and $\bar{\nu}_e$ states, the nutation time scale is $T_{\text{nut}} \sim -\ln\theta_v$ in the inverted mass hierarchy case [32]. Estimating the nutation time scale for realistic supernova neutrino systems is problematic, partly because this quantity depends on time-varying electron and neutrino number densities [22,32,34]. In Fig. 2 we plot as a function of θ_v the radius r_X (as defined in Ref. [16]) where the energy-averaged value of $P_{\nu_e\nu_e}$ drops below 0.9 and significant nutation develops in our single-angle simulations for the inverted mass hierarchy. Our single-angle calculations suggest that the onset of significant nutation is nearly independent of θ_v as it is decreased from 0.1 to $\simeq 10^{-20}$. As θ_v is decreased further, r_X begins to increase. We expect that for sufficiently small θ_v , the onset of significant nutation is pushed to so large a radius that the corresponding neutrino number density becomes too low to generate any collective flavor evolution or stepwise spectral swapping.

FIG. 1 (color online). The neutrino survival probability $P_{\nu\nu}$ as a function of neutrino emission angle ϑ_0 (relative to the normal at the emission point on the neutrino sphere) and energy E . The left panel is calculated for a normal neutrino mass hierarchy with $\theta_v = 0.01$, and the right panel is for an inverted neutrino mass hierarchy with $\theta_v = 10^{-9}$. These results are taken at radius $r = 250$ km. Except for θ_v , all parameters are the same as those for Fig. 3 of Ref. [19].

FIG. 2 (color online). Single-angle simulation results for the dependence of r_X on θ_v in the inverted neutrino mass hierarchy case. Here θ_v is the effective 2×2 vacuum mixing angle ($\simeq \theta_{13}$), and r_X is the radius where the energy-averaged value of $P_{\nu_e \nu_e}$ drops below 0.9. Except for θ_v , all parameters are the same as those for Fig. 2 of Ref. [19].

Just after the bounce of the supernova core, when the supernova shock breaks through the neutrino sphere, there is a brief intense burst of neutrinos which are emitted predominantly as ν_e 's. Lacking in $\bar{\nu}_e$'s [35], this burst is not likely to be affected by stepwise spectral swapping at the δm_{atm}^2 scale (but see Ref. [36]). Later, both neutrinos and antineutrinos are emitted and they form a bipolar system. If θ_{13} is not too small, for the normal mass hierarchy case, stepwise spectral swapping at the δm_{atm}^2 scale can be observable at late times when matter has been sufficiently condensed toward the proto-neutron star. For the inverted mass hierarchy case, however, stepwise spectral swapping occurs even at early times [22,31] for essentially any nonvanishing θ_{13} . In light of the insensitivity of E_C to θ_{13} in the latter case, supernova neutrino signals can offer a unique probe of the neutrino mass hierarchy even for θ_{13} too small to be measured by conventional neutrino oscillation experiments.

This work was supported in part by NSF Grant No. PHY-04-00359, the TSI Collaboration's DOE SciDAC grant at UCSD, DOE Grant No. DE-FG02-87ER40328 at UMN, and an IGPP/LANL mini-grant. This work was also supported in part by the LDRD Program and Open Supercomputing at LANL, and by NERSC through the TSI Collaboration using Bassi. We thank A. Burrows, A. Mezzacappa, and A. Mirizzi for valuable conversations.

-
- [1] W.-M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
 [2] A. Strumia and F. Vissani, arXiv:hep-ph/0606054.
 [3] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978).

- [4] S.P. Mikheyev and A.Y. Smirnov, Yad. Fiz. **42**, 1441 (1985) [Sov. J. Nucl. Phys. **42**, 913 (1985)].
 [5] G.M. Fuller, R.W. Mayle, J.R. Wilson, and D.N. Schramm, Astrophys. J. **322**, 795 (1987).
 [6] D. Nötzold and G. Raffelt, Nucl. Phys. **B307**, 924 (1988).
 [7] J.T. Pantaleone, Phys. Rev. D **46**, 510 (1992).
 [8] G. Sigl and G. Raffelt, Nucl. Phys. **B406**, 423 (1993).
 [9] A.S. Dighe and A.Y. Smirnov, Phys. Rev. D **62**, 033007 (2000).
 [10] R.C. Schirato and G.M. Fuller, arXiv:astro-ph/0205390.
 [11] C. Lunardini and A.Y. Smirnov, J. Cosmol. Astropart. Phys. **06** (2003) 009.
 [12] J.P. Kneller and G.C. McLaughlin, Phys. Rev. D **73**, 056003 (2006).
 [13] S.-H. Chiu and T.-K. Kuo, Phys. Rev. D **73**, 033007 (2006).
 [14] T. Yoshida *et al.*, Phys. Rev. Lett. **96**, 091101 (2006).
 [15] J.P. Kneller, G.C. McLaughlin, and J. Brockman, arXiv:0705.3835.
 [16] H. Duan, G.M. Fuller, J. Carlson, and Y.-Z. Qian, Phys. Rev. D **74**, 105014 (2006).
 [17] H. Duan, G.M. Fuller, J. Carlson, and Y.-Z. Qian, Phys. Rev. D **75**, 125005 (2007).
 [18] G.G. Raffelt and A.Y. Smirnov, Phys. Rev. D **76**, 081301(R) (2007).
 [19] H. Duan, G.M. Fuller, J. Carlson, and Y.-Z. Qian, Phys. Rev. Lett. **97**, 241101 (2006).
 [20] A. Friedland and C. Lunardini, J. High Energy Phys. **10** (2003) 043.
 [21] A.B. Balantekin and Y. Pehlivan, J. Phys. G **34**, 47 (2007).
 [22] H. Duan, G.M. Fuller, and Y.-Z. Qian, Phys. Rev. D **74**, 123004 (2006).
 [23] G.M. Fuller, R.W. Mayle, B.S. Meyer, and J.R. Wilson, Astrophys. J. **389**, 517 (1992).
 [24] G.M. Fuller and Y.-Z. Qian, Phys. Rev. D **73**, 023004 (2006).
 [25] Y.-Z. Qian, G.M. Fuller, G.J. Mathews, R.W. Mayle, J.R. Wilson, and S.E. Woosley, Phys. Rev. Lett. **71**, 1965 (1993).
 [26] S. Pastor and G. Raffelt, Phys. Rev. Lett. **89**, 191101 (2002).
 [27] A.B. Balantekin and H. Yüksel, New J. Phys. **7**, 51 (2005).
 [28] A.B. Balantekin and G.M. Fuller, Phys. Lett. B **471**, 195 (1999).
 [29] H. Duan, G.M. Fuller, and Y.-Z. Qian, Phys. Rev. D **76**, 085013 (2007).
 [30] A. Esteban-Pretel, S. Pastor, R. Tomas, G.G. Raffelt, and G. Sigl, arXiv:0706.2498 [Phys. Rev. D (to be published)].
 [31] G.L. Fogli, E. Lisi, A. Marrone, and A. Mirizzi, arXiv:0707.1998.
 [32] S. Hannestad, G.G. Raffelt, G. Sigl, and Y.Y.Y. Wong, Phys. Rev. D **74**, 105010 (2006).
 [33] S. Pastor, G.G. Raffelt, and D.V. Semikoz, Phys. Rev. D **65**, 053011 (2002).
 [34] V.A. Kostelecky and S. Samuel, Phys. Rev. D **52**, 621 (1995).
 [35] M. Kachelriess *et al.*, Phys. Rev. D **71**, 063003 (2005).
 [36] H. Duan, G.M. Fuller, J. Carlson, and Y.-Z. Qian, arXiv:0710.1271 [Phys. Rev. Lett. (to be published)].