

# UC Office of the President

## Recent Work

### Title

Nectar-inhabiting microorganisms influence nectar volatile composition and attractiveness to a generalist pollinator

### Permalink

<https://escholarship.org/uc/item/78z355k3>

### Authors

Rering, Caitlin C  
Beck, John J  
Hall, Griffin W  
et al.

### Publication Date

2017-09-28

### DOI

10.1111/nph.14809

Peer reviewed

# Nectar-inhabiting microorganisms influence nectar volatile composition and attractiveness to a generalist pollinator

Caitlin C. Rering<sup>1</sup>, John J. Beck<sup>1</sup>, Griffin W. Hall<sup>2</sup>, Mitchell M. McCartney<sup>3</sup> and Rachel L. Vannette<sup>2</sup>

<sup>1</sup>Chemistry Research Unit, Center for Medical, Agricultural and Veterinary Entomology, Agricultural Research Service, United States Department of Agriculture, 1700 SW 23<sup>rd</sup> Dr., Gainesville, FL 32608, USA; <sup>2</sup>Department of Entomology and Nematology, University of California, Davis, One Shields Ave., Davis, CA 95616, USA; <sup>3</sup>Mechanical and Aerospace Engineering, University of California, Davis, One Shields Ave., Davis, CA 95616, USA

## Summary

Author for correspondence:

Rachel L. Vannette

Tel: +1 530 752 3379

Email: [rlvannette@ucdavis.edu](mailto:rlvannette@ucdavis.edu)

Received: 16 June 2017

Accepted: 21 August 2017

*New Phytologist* (2017)

doi: 10.1111/nph.14809

**Key words:** *Apis mellifera*, floral headspace, microbial volatile (MVOC), nectar microbes, pollination, semiochemical, volatile organic chemicals (VOCs), yeast.

- The plant microbiome can influence plant phenotype in diverse ways, yet microbial contribution to plant volatile phenotype remains poorly understood. We examine the presence of fungi and bacteria in the nectar of a coflowering plant community, characterize the volatiles produced by common nectar microbes and examine their influence on pollinator preference.
- Nectar was sampled for the presence of nectar-inhabiting microbes. We characterized the headspace of four common fungi and bacteria in a nectar analog. We examined electrophysiological and behavioral responses of honey bees to microbial volatiles. Floral headspace samples collected in the field were surveyed for the presence of microbial volatiles.
- Microbes commonly inhabit floral nectar and the common species differ in volatile profiles. Honey bees detected most microbial volatiles tested and distinguished among solutions based on volatiles only. Floral headspace samples contained microbial-associated volatiles, with 2-ethyl-1-hexanol and 2-nonanone – both detected by bees – more often detected when fungi were abundant.
- Nectar-inhabiting microorganisms produce volatile compounds, which can differentially affect honey bee preference. The yeast *Metschnikowia reukaufii* produced distinctive compounds and was the most attractive of all microbes compared. The variable presence of microbes may provide volatile cues that influence plant–pollinator interactions.

## Introduction

Plant volatile emissions are a key phenotype that mediate a myriad of ecological interactions in natural and managed systems (Paré & Tumlinson, 1999; Kessler & Baldwin, 2001; Pichersky & Gershenzon, 2002; Junker & Tholl, 2013), including pollination (Dobson, 1994; Raguso, 2001). Indeed, floral volatile emission can mediate pollinator choice among (Byers *et al.*, 2014) and within (Galen & Newport, 1988; Kessler *et al.*, 2011) plant species, and is often a target of pollinator-mediated selection (Parachnowitsch *et al.*, 2012; Gervasi & Schiestl, 2017).

Microorganisms can contribute to plant phenotype in diverse ways (Friesen, 2013) including by changing plant volatile emissions (Pineda *et al.*, 2010; Shapiro *et al.*, 2012). In some cases, microbial pathogens alter the host plant's volatile profile to attract the pathogen's insect vector (Jiménez-Martínez *et al.*, 2004; Mann *et al.*, 2012). That microbial volatiles contribute to plant–pollinator interactions has been hypothesized (Raguso, 2004, 2008; Pozo *et al.*, 2009), and implicated (Golonka *et al.*, 2014; Peñuelas *et al.*, 2014; Schaeffer *et al.*, 2017). However, whether volatiles produced by microbes, rather than their plant hosts, can alter ecological interactions remains poorly understood (but see Davis *et al.*, 2013), particularly for plant–pollinator

interactions. If microorganisms directly modify a plant's chemical phenotype (chemotype), their influence could extend not only to pollination, but possibly other plant–insect interactions as well (Beck & Vannette, 2017).

Yeasts and bacteria are common and abundant inhabitants of floral nectar (Herrera *et al.*, 2008, 2009), and often rely on pollinators to disperse among individual flowers (Brysch-Herzberg, 2004; Canto *et al.*, 2008; Herrera *et al.*, 2008; Vannette & Fukami, 2017). The microorganisms that specialize on the nectar environment must rapidly and repeatedly disperse and re-establish themselves in new flowers, which in some cases may require multiple pollinator visits (Mittelbach *et al.*, 2016a,b). Yeasts that inhabit other ephemeral habitats, including decaying fruit or plant material, tend to rely on other organisms for dispersal to specific habitats (reviewed by Mittelbach & Vannette, 2017) in a process called phoresis. For example, volatile compounds emitted by yeasts are attractive to insect vectors including *Drosophila* (Buser *et al.*, 2014). In nectar, microbial metabolism can influence sugar composition and concentration (Herrera *et al.*, 2013; Vannette *et al.*, 2013; Good *et al.*, 2014; Schaeffer *et al.*, 2014, 2015), ethanol concentration, amino acid composition (Peay *et al.*, 2012) and other metabolic products that contribute to flavor and scent of nectar (Vannette & Fukami, 2016).

Pollinators may respond innately or learn to associate such products with nectar availability or quality (Knauer & Schiestl, 2015). Volatile chemicals produced by microbes are suspected to play a role in pollination: while nectar-inhabiting yeasts have been reported to emit volatile compounds (Golonka *et al.*, 2014), volatile profiles of nectar-inhabiting microorganisms have not been compared and the influence of microbial volatiles on pollinator preference has not been previously examined.

Here, we test the hypothesis that nectar-inhabiting microorganisms vary in volatile emission and differentially influence attractiveness to a generalist insect pollinator. First, to establish the ecological relevance of microbes in nectar, we examined the presence of bacteria and fungi in the nectar of 28 species of coflowering plants in California. Second, we characterized volatile headspace of four commonly isolated microbial species in a synthetic nectar analog at multiple time points. Third, we used electroantennographic (EAG) bioassays to examine which microbial volatiles were detectable to honey bees (*Apis mellifera*), and proboscis extension response (PER) bioassays to determine honey bee response to microorganisms inoculated in a nectar analog. We also determined whether microbe-associated volatiles could be detected in floral headspace samples taken from the field. Our results provide evidence that nectar-inhabiting microorganisms are common but variable inhabitants of nectar, and that different microbial species produce distinct volatile blends detectable in field samples, which may influence pollinator detection and preference.

## Materials and Methods

### Study system and microbial isolation

Nectar standing crop was collected from 1170 individual flowers of 28 nectar-producing plant species native to California in October 2015 and March–June 2016 (Supporting Information Table S1). All available flowering plant species at Stebbins Cold Canyon that produced at least 0.1  $\mu\text{l}$  of nectar during the sampling period were sampled to examine frequency of bacterial and fungal colonization of nectar. Honey bees, used in later experiments, are common pollinators of many of the plant species sampled here, and also very common generalist pollinators of native plants and agricultural crops in California and world-wide.

Briefly, open flowers were collected, nectar was extracted in the laboratory and dilutions were plated. Yeasts and yeast-like fungi were cultured on yeast media agar (YMA: 0.3% malt extract, 0.5% peptone, 1% glucose, 0.3% yeast extract, 2% agar) and a subset of representative colony morphotypes were identified from over 1200 plates by sequencing the D1/D2 domain of the large subunit nuclear ribosomal RNA with primers NL1/NL4 (O'Donnell, 1993). Bacteria were cultured on R2A media supplemented with 20% sucrose (BD, Franklin Lakes, NJ, USA) and colonies identified by sequencing the 16S rRNA gene using primers 27F/1492R (Lane, 1991; Turner *et al.*, 1999). The presence of fungi or bacterial colonies on plates was recorded for each individual flower, with lower detection limits of *c.* 5 colony forming units (CFU) for fungi and 50 CFU for bacteria. Isolates of

identified species were stored at  $-80^{\circ}\text{C}$  as glycerol stocks until use, and propagated on the media described above until inoculation into synthetic nectar solutions.

For this study, four commonly isolated microorganisms were selected for further analyses: the fungi *Metschnikowia reukaufii* and *Aureobasidium pullulans*, and the bacteria *Neokomagataea* sp. and *Asaia astilbes* (GenBank IDs: MF319536, MF325803, MF340296 and KC677740). Isolates were taken from the nectar of plant species *Epilobium canum* and *Mimulus aurantiacus*, which have a diverse group of flower visitors including honey bees, hummingbirds and other native insects. These microbes are all commonly isolated from flowers in the current and previous studies, so are all ecologically relevant to the nectar environment. However, these taxa are also likely to vary in their ecological affinity to flowers and pollinators. Previous work has found that *M. reukaufii* is dependent on pollinators for transmission among flowers (e.g. Belisle *et al.*, 2012; Vannette & Fukami, 2017), while *A. pullulans* has a broader habitat range (e.g. Andrews *et al.*, 1994; Wehner *et al.*, 2017). The ecological affinities of the bacterial taxa profiled here are unknown, but strains from these genera have been found in floral nectar in other studies previously (Vannette *et al.*, 2013; Good *et al.*, 2014).

### Microbial growth conditions

Sterile synthetic nectar (0.3% w/v sucrose; 0.6% w/v each of glucose and fructose; 0.1 mM each of glycine, L-alanine, L-asparagine, L-aspartic acid, L-glutamic acid, L-proline and L-serine), designed to mimic floral nectar of bee-pollinated flowers (Baker & Baker, 1982; Gardener & Gillman, 2001), was inoculated with 20 ml of  $10^3$  cells  $\mu\text{l}^{-1}$  from actively growing subcultures and incubated at  $29^{\circ}\text{C}$  under aerobic conditions in sealed 118 ml Mason jars ( $n = 3$  replicates of each microbial species: *M. reukaufii*, *A. pullulans*, *Neokomagataea* and *Asaia*). Mason jar lids remained hermetically sealed for the duration of each experiment, except for just before volatile collections times when the lids were gently lifted under sterile conditions to allow venting of accumulated volatiles. Lids were modified to accommodate two sampling ports by fitting GC septa (high temperature low bleed septa, 11 mm; Agilent, Santa Clara, CA, USA) into pre-drilled holes. Glassware was autoclaved immediately before use and preparations were carried out under sterile conditions. Microbial growth was confirmed visually, by pH decrease (Vannette *et al.*, 2013) and optical density (for bacteria) or cell count (for yeast) increase.

### Volatile collection, analysis and identification

Headspace volatiles from inoculated microbes were collected onto solid-phase microextraction (SPME) fibers (Supelco, Bellefonte, PA, USA; 50/30  $\mu\text{m}$ , 2 cm, divinylbenzene/carboxen/polydimethylsiloxane) at intervals of 0, 48 and 96 h ( $\pm 0.4$  h). Two identical SPME fiber types were simultaneously inserted into the sampling ports installed on jar lids. Volatile collections used the following fiber parameters (Beck *et al.*, 2008): accumulation of volatiles in the freshly sealed container (see above for venting procedure to allow for collection of volatiles at the specific time

point), 15 min; exposure of fiber to adsorb volatiles, 15 min; storage time of volatiles on fiber,  $\leq 1$  min; and thermal desorption of volatiles in injector ports, 6 min. The adsorbed volatiles were thermally desorbed in splitless mode onto an Agilent 7890A gas chromatograph coupled to a quadrupole 5975C MSD detector in electron ionization mode (Palo Alto, CA, USA) outfitted with a J&W Scientific (Folsom, CA, USA) DB-Wax column (60 m  $\times$  320  $\mu\text{m}$   $\times$  0.25  $\mu\text{m}$ ), and an Agilent 7890B gas chromatograph coupled to a quadrupole 5977B MSD detector in electron ionization mode and equipped with a J&W Scientific DB-1 column (60 m  $\times$  320  $\mu\text{m}$   $\times$  0.25  $\mu\text{m}$ ). Volatiles were analyzed using parameters identical to those previously described (Beck *et al.*, 2016) with the following modifications: DB-1 had a final temperature of 190°C, and the DB-Wax had an adapted flow of 3 ml min<sup>-1</sup>. Data from the GC MSD device fitted with a DB-Wax column provided superior peak shape for polar analytes and was used for the qualitative comparison of compounds. Data from the DB-1-equipped instrument were used for additional identification and retention index (RI) calculations. RIs were

calculated using a homologous series of *n*-alkanes on both the DB-1 and the DB-Wax columns. RI values from both columns were used to assist with initial identification, and identities were further confirmed by comparison to retention times and fragmentation patterns of standards. Compound identities not verified on both instruments with a commercial or other available standard were marked as tentatively identified. Additionally, if peaks could not be authenticated and the library matches were poor, their identities were labeled as unknown (see Table S2). The most abundant ions for unknowns are tabulated in Table S2. Peaks identified as background from the containers, fibers, columns and synthetic nectar that were found in media controls and blanks were removed before statistical analysis.

### Electroantennographic bioassays

To examine if honey bee antennae respond to microbial volatiles, we tested all available and identified compounds that were produced by microorganisms (21 compounds, see Table 1). Honey

**Table 1** Volatile compounds produced by microorganisms grown in synthetic nectar

Class	Chemical	Retention indices <sup>a</sup>		Normalized EAG response (%; <i>n</i> = 6 bees)		Peak area <sup>l</sup> at 96 h ( $\times 10^5$ ; mean $\pm$ SE, <i>n</i> = 3 jars)			
		DB-Wax	DB-1	40 $\mu\text{mol}$	0.4 $\mu\text{mol}$	<i>Asaia</i>	<i>Neok</i>	<i>Metsch</i>	<i>Aureo</i>
1° Alcohol	ethanol <sup>c</sup>	933	–	1 $\pm$ 3	1 $\pm$ 3	23 $\pm$ 8	18 $\pm$ 3	6800 $\pm$ 200	3000 $\pm$ 100
	<i>n</i> -propanol <sup>c</sup>	1038	–	17 $\pm$ 4**	4 $\pm$ 6	0	0	30 $\pm$ 2	103 $\pm$ 4
	isobutanol <sup>e</sup>	1092	613	24 $\pm$ 14	–4 $\pm$ 4	1.5 $\pm$ 0.8	2.2 $\pm$ 0.2	614 $\pm$ 3	730 $\pm$ 30
	2-methyl-1-butanol <sup>c,g</sup>	1209	720	99 $\pm$ 16**	–6 $\pm$ 8	44 $\pm$ 2 <sup>g</sup>	46 $\pm$ 1 <sup>g</sup>	6990 $\pm$ 80 <sup>g</sup>	1500 $\pm$ 100 <sup>g</sup>
	3-methyl-1-butanol <sup>c</sup>	1209	718	143 $\pm$ 11**	1 $\pm$ 3	– <sup>h</sup>	0	5.5 $\pm$ 0.2	3 $\pm$ 0.1
	3-methyl-3-buten-1-ol <sup>c</sup>	1250	713	104 $\pm$ 34*	8 $\pm$ 12	– <sup>h</sup>	0	8.9 $\pm$ 0.6	8.9 $\pm$ 0.4
	4-penten-1-ol <sup>c</sup>	1304	735	69 $\pm$ 10**	0 $\pm$ 3	0	0	6 $\pm$ 2	4.6 $\pm$ 0.3
	<b><i>n</i>-hexanol<sup>c</sup></b>	1355	853	160 $\pm$ 16**	62 $\pm$ 21*	5.1 $\pm$ 0.3	3.38 $\pm$ 0.08	1.8 $\pm$ 0.4	0
	3-ethoxy-1-propanol <sup>c</sup>	1380	817	30 $\pm$ 7**	–4 $\pm$ 2	0	0	29 $\pm$ 2	90 $\pm$ 60
	<b>2-ethyl-1-hexanol<sup>c</sup></b>	1493	1014	60 $\pm$ 9**	59 $\pm$ 15**	77 $\pm$ 6	60 $\pm$ 2	260 $\pm$ 20	16 $\pm$ 8
	<b>2-phenylethanol<sup>c</sup></b>	1913	1082	62 $\pm$ 5**	50 $\pm$ 9**	4.7 $\pm$ 0.5	4.6 $\pm$ 0.2	10 $\pm$ 1	0
2° Alcohol	2-butanol <sup>d</sup>	1024	–	465 $\pm$ 86**	–3 $\pm$ 6	0	0	96 $\pm$ 7	130 $\pm$ 40
Aldehyde	acetaldehyde <sup>b</sup>	–	–	–	–	3 $\pm$ 2	1.7 $\pm$ 0.4	130 $\pm$ 10	18 $\pm$ 3
Ester	ethyl acetate <sup>c</sup>	885	–	4 $\pm$ 8	–2 $\pm$ 4	0	0	5.3 $\pm$ 0.6	0
	2-methylpropyl acetate <sup>c</sup>	1012	756	133 $\pm$ 26**	–24 $\pm$ 16	0	0	6 $\pm$ 1	0
	ethyl butyrate <sup>c</sup>	1035	783	68 $\pm$ 19**	–4 $\pm$ 3	0	0	41 $\pm$ 2	0
	<b>3-methylbutyl acetate<sup>c</sup></b>	1122	861	81 $\pm$ 16**	6 $\pm$ 7	0	0	0	0
Isoprenoid	Isoprene <sup>b</sup>	–	–	9 $\pm$ 7	9 $\pm$ 8	9 $\pm$ 1	10 $\pm$ 9	0	0
Ketone	4-methyl-2-pentanone <sup>c</sup>	717	1008	127 $\pm$ 20*	–7 $\pm$ 8	0	0	0	20 $\pm$ 10
	4-methyl-3-penten-2-one <sup>f</sup>	1132	776	–	–	1 $\pm$ 0.5	0	0	– <sup>h</sup>
	3-hydroxy-2-butanone <sup>c</sup>	1284	677	177 $\pm$ 93	21 $\pm$ 13	15 $\pm$ 1	12.3 $\pm$ 0.3	53 $\pm$ 0.9	20 $\pm$ 3
	<b>2-nonanone<sup>c</sup></b>	1389	1072	87 $\pm$ 4**	76 $\pm$ 10**	0	0	0	6 $\pm$ 1
Misc.	<b>2,5-dimethylfuran<sup>c</sup></b>	950	696	10 $\pm$ 5	–13 $\pm$ 13	16 $\pm$ 4	9 $\pm$ 1	0	0
	Unknown 1	1087	727	–	–	0	0	0	– <sup>h</sup>
	Unknown 2	1197	1022	–	–	1 $\pm$ 0.5	1.7 $\pm$ 0.2	20 $\pm$ 10	70 $\pm$ 70
	Unknown 3	1278	719	–	–	– <sup>h</sup>	0	19 $\pm$ 1	9 $\pm$ 1
	Unknown 4	1430	1094	–	–	0	0	0	5 $\pm$ 4

Fungal species included *Metschnikowia reukaufii* (*Metsch*) and *Aureobasidium pullulans* (*Aureo*); bacterial species included strains from the genera *Asaia* and *Neokomagataea* (*Neok.*). Full sampling details are found in the Materials and Methods section.

<sup>a</sup>Retention indices relative to *n*-alkanes on DB-1 and DB-Wax columns for compound identification. A dash indicates retention time was too low to calculate the retention index. <sup>b</sup>Sources were Fisher Scientific. <sup>c</sup>Sources were Sigma-Aldrich. <sup>d</sup>Sources were Mallinckrodt Baker. <sup>e</sup>Sources were J. T. Baker. <sup>f</sup>Compounds were tentatively identified due to lack of commercial standard. <sup>g</sup>Peak areas for the isomers 2-methyl-1-butanol and 3-methyl-1-butanol are summed as a result of co-elution. <sup>h</sup>Compound observed in only one replicate on day 4. <sup>l</sup>Peak areas gathered from DB-Wax-equipped GC-MS. \*\*Normalized electroantennographic (EAG) response is significantly different from 0 (false discovery rate < 0.05; \**P* < 0.05, \*\**P* < 0.01). \*Normalized EAG response is significantly different from 0 (false discovery rate < 0.05). Compounds in bold type could be detected using *in situ* methods.

bees ( $n=6$  bees tested for each identified compound) were collected each day from outside hives housed in an apiary located at the USDA-ARS Center for Medical, Agricultural and Veterinary Entomology campus in Gainesville (FL, USA). Before experiments, honey bees were fed a sugar solution (1 : 1 water/sucrose) and stored in the dark. Immediately before bioassays, bees were placed in a trimmed 15 ml centrifuge tube and secured from behind with cotton. Under a low power stereo-microscope, both antennae were excised at the scape using micro-scissors and mounted on a forked probe (Syntech, Kirchzarten, Germany; internal gain 10 $\times$ ) with electrode gel (Parker Laboratories, Fairfield, NJ, USA). The prepared probe was mounted in the humidified constant air and volatile sample tube and allowed to equilibrate to the air flow for 3–5 min (until signal reached  $c. 0 \mu\text{V}$ ). Sample puffs were delivered at 1-min intervals to allow antennae to re-equilibrate after exposure. Antennal responses, which indicate detection rather than attraction or repulsion, were recorded with the Autospike software (SYNTECH, v.3.9).

Odor samples were prepared in similar fashion to Beck *et al.* (2014) by loading 7 mm bioassay discs with 20  $\mu\text{l}$  of pentane containing each compound at each of two loading doses: 40 or 0.4  $\mu\text{mol}$ . Pilot experiments revealed these loading doses provided consistent antennal responses to known honey bee pheromones and have been used previously (Bhagavan & Smith, 1997). Slightly higher compound loadings were used to accommodate the 'wind tunnel' design of the EAG bioassay system used (Fig. S1). The pentane was allowed to evaporate for 2 min and then the filter paper was placed within a Pasteur pipette trimmed at the tapered end to a final length of 6.5 cm and sealed with Parafilm. A 0.5 s pulse flow (300 ml min $^{-1}$ ) and a humidified continuous flow (125 ml min $^{-1}$ ) directed odors through an air and volatile tube (1.5 cm diameter) containing the mounted antennae and probe. A Faraday cage protected against ambient electrical interference. To account for variability in response among individuals, responses to blanks (20  $\mu\text{l}$  pentane and bioassay disc) were subtracted from each sample and antennal response values were normalized to a standard stimulus (0.4  $\mu\text{mol}$  citral, a component of bee pheromone produced by the Nasanov gland), which was set at 100%.

### Proboscis extension response assay

To assay the effect of microbial scent on honey bee feeding preference, we used a PER assay, which has been previously used to examine how nectar solutes influence acceptability of a nectar source (Scheiner *et al.*, 2004) based on odor or solute detection by honey bee antennae. Synthetic nectar solutions, including a sterile control, were prepared as above, incubated for 68–74 h, then used to assess bee preference. Foraging bees were collected at the entrance of three hives at UC Davis Laidlaw Honey Bee Research Facility. Bees were restrained, screened for responsiveness ( $\sim 75\%$  success) and allowed to feed on the control solution, then starved for 2 h before initiation of the assay. Microcapillary tubes containing 2  $\mu\text{l}$  of the solution were wafted 5 mm from an individual bee's antennae for 6 s and proboscis extension was recorded. Solutions were introduced in two different sequences.

### *In situ* floral headspace analysis

Volatiles were collected from the floral headspace of five plant species (Table S3), a subset of those surveyed above, at the UC Davis Stebbins Cold Canyon Reserve (Winters, CA, USA) in May and June 2016 (a total of 24 samples across three sampling dates). The plant species *Delphinium nudicaule*, *M. aurantiacus*, *Clarkia unguiculata*, *Collinsia heterophylla* and *Delphinium hesperium* were chosen because all were found to host bacteria or fungi in floral nectar, often at high abundance. Pairs of flowers or an entire floral cluster were enclosed in a nylon oven bag (406  $\times$  444 mm; Reynolds, Richmond, VA, USA) secured using metal clips to minimize total headspace. A PDMS Twister bar (Gerstel Inc., Linthicum, MD, USA) was enclosed within each bag to collect volatiles for 60 min between 09:00 and 10:00 h. After headspace analysis, flowers were collected and culturable microorganisms were plated as described above for sampled flowers, and colonies counted to quantify microbial abundance (CFU). No-flower controls were collected by bagging a Twister bar in the field. Twisters were kept on ice for no more than 48 h, then were thermally desorbed in splitless mode using a thermal desorption unit (Gerstel), which ramped from 30 to 250 $^{\circ}\text{C}$ , holding for 3 min. Desorbed volatiles were cryofocused at  $-80^{\circ}\text{C}$ , then heated to 260 C for 3 min in a cooled injection unit (Gerstel), where they were splitlessly introduced to the GC column. Separation occurred on an Agilent 7890B gas chromatograph coupled to a single quadrupole 5977A MSD device outfitted with an HP-5MS column (30 m  $\times$  250  $\mu\text{m}$   $\times$  0.25  $\mu\text{m}$ ; Agilent) at a constant flow of 1.5 ml min $^{-1}$  with a temperature program as follows: initial temperature, 40 $^{\circ}\text{C}$ ; 4 $^{\circ}\text{C}$  min $^{-1}$  ramp to 200 $^{\circ}\text{C}$ ; 30 $^{\circ}\text{C}$  min $^{-1}$  ramp to 300 C; 4 min hold. MS detection scanned from 33 to 300  $m/z$ . A series of commercially available chemical standards (subset of the microbial volatiles identified above) was compared to identify compounds in floral headspace samples by similarity in retention times and mass spectrum fragmentation patterns. Samples were scored blind (without identifying sample information). Volatile presence in a sample was scored based on similarity in retention times and fragmentation patterns or a match to the NIST mass spectrum library (Table S2).

This analysis was intended to qualitatively report compounds that were detected in both floral microbes and the headspace of the same flower. Because of its affinity to apolar compounds, our Twister-DB5 system may have not recovered all of the polar compounds found in the headspace above the flowers (Table 1). Thus, we may be underreporting the number of compounds found in both isolated microbes and flower samples. Still, the experiment as executed allowed us to report that some volatiles produced by floral microbes were also found in the headspace of wild flowers.

### Statistical analyses

To compare volatile blends among microbial species and days following inoculation, peak areas were log-transformed. Microbial volatile composition was visualized using principal



coordinates analysis (PCoA) based on Bray–Curtis dissimilarities and the interactive effects of microbial strain and day were assessed using PerMANOVA implemented using the *adonis* function in the R package ‘VEGAN’ (Oksanen *et al.*, 2012). DESeq2 (Love *et al.*, 2013) was used to assess which compounds differed in relative abundance between prespecified groups. First, we examined which compounds differentiated bacteria from fungi (days 2 and 4 only) and, second, which differentiated the nectar-specialist yeast *M. reukaufii* from the generalist yeast *A. pullulans*. Because bacterial species did not differ significantly in volatile composition, we did not examine their differences further. We examined if microbial species differed in variance in volatile composition at each timepoint (days 2 and 4 separately) using the *betadisper* function in *vegan*.

To examine which compounds were detected by honey bees, we used a *t*-test to examine if normalized EAG responses were significantly different from zero (no detectable response), and used a false discovery rate (FDR) correction to control for multiple comparisons.


To examine if microbial species in nectar influenced honey bee proboscis extension (PER), we used a binomial regression to examine if the proportion of positive responses by honey bees varied among microbial species. The sequence of solution introduction was included in the model, but was not significant ( $P > 0.10$ ), so was dropped from the model.

For field volatile data, we examined if the presence of each validated microbial-associated volatile (Table 1) was predicted by fungal CFU abundance, bacterial CFU abundance or plant species identity of the sampled flowers using Pearson chi-squared tests.

## Results

Floral nectar often contained culturable nectar-inhabiting microorganisms, including bacteria and fungi (Fig. 1). Among plant species, the proportion of individual flowers that contained detectable microorganisms ranged from 20% to 86% of flowers sampled.


All focal microorganisms produced detectable volatile compounds when grown in artificial nectar, and species differed in the composition of the volatile blend emitted ( $F_{3,28} = 40.22$ ,  $P < 0.001$ ; Fig. 2). Surprisingly, species differences in volatile composition were detectable immediately after inoculation (Day = 0, 30 min after inoculation into synthetic nectar with 15 min each permeation and exposure time), and volatile blends further diverged after 2 and 4 d of growth (Species  $\times$  Day  $F_{3,28} = 6.86$ ,  $P < 0.001$ ; Fig. 2). Alcohols, esters and ketones were more abundant in fungal solutions, while the volatile 2,5-dimethylfuran was characteristic of bacteria (Table 1). All four microbes produced *n*-hexanol. When we compared which compounds differentiated the fungal taxa, the generalist fungi *A. pullulans* emitted the ketones 2-nonanone and 4-methyl-2-pentanone, the short chain alcohols isobutanol and *n*-propanol, and two additional unidentified compounds (Table 1, unknowns 1 and 4). By contrast, the specialist nectar yeast *M. reukaufii* was characterized by esters, including ethyl butyrate, 2-methylpropyl



**Fig. 1** Proportion of individual nectar samples from plant species hosting culturable (a) bacteria or (b) fungi (black) or no detectable microorganisms (light gray). Thresholds for detection were *c.* 5 colony-forming units (CFU) for fungi and 50 CFUs for bacteria. Some flowers contained both bacteria and fungi. Flower abbreviation indicates the first three letters of the genus and species for each plant species sampled with  $n = 1170$  individual flowers sampled with  $n = 13$ –133 samples per plant species. Microbial density within individual flowers ranged from 0 to  $10^6$  CFUs  $\mu\text{l}^{-1}$  nectar.

acetate and 3-methylbutyl acetate, the alcohols 2-butanol and 3-ethoxy-1-propanol, and a relatively greater abundance of carbon dioxide, ethyl acetate, and the alcohols 3-methyl-1-butanol, 2-methyl-1-butanol, 3-methyl-3-buten-1-ol, ethanol and 2-phenylethanol. Across all microbial species, the replicates of *M. reukaufii* showed the least variance in volatile composition (Fig. 2, *betadisper* Day 2  $P = 0.02$ ; Day 4  $P = 0.01$ ).

Honey bee antennae detected 14 of the 20 microbial volatiles tested using EAG at a loading of 40  $\mu\text{mol}$ , but only three compounds at 0.4  $\mu\text{mol}$ . Compounds detected at low concentrations included *n*-hexanol, 2-ethyl-1-hexanol and 2-phenylethanol, which were produced by all surveyed microbes but most abundantly by yeasts (Table 1). Additionally, 2-nonanone was also detected by antennae but was produced only by the fungi *A. pullulans*. Antennal response was strongest in response to the short chain secondary alcohol 2-butanol, which was only



**Fig. 2** Principal coordinates analysis showing variation in microbial volatile composition across microbial taxa and sampling days (days 0, 2, and 4) based on Bray–Curtis dissimilarity. Microbial species differ from each other (Species  $P < 0.001$ ) and diverge from each other over time (Species  $\times$  Day  $P < 0.001$ ). In panels for days 2 and 4, replicates of *Metschnikowia reukaufii* are largely overlapping.


produced by the yeast *M. reukaufii* ( $465 \pm 86\%$  normalized antennal response at  $40 \mu\text{mol}$  exposure dose, Table 1), while its structural isomer and primary alcohol isobutanol, produced by all microbes, did not elicit a significant response ( $24 \pm 14\%$  normalized antennal response at  $40 \mu\text{mol}$ ).

Honey bees varied in acceptance of microbial solutions based solely on volatile exposure: control and *M. reukaufii* solutions were accepted in over 70% of trials, whereas *A. pullulans* and *Asaia* were accepted 36% and 48% of the time, respectively. *Neokomagataea* received the fewest positive responses, with only 17% of bees accepting this solution (Fig. 3,  $P < 0.001$ ).

A subset of compounds that characterized microbial volatile emission in synthetic nectar were identified in floral headspace of naturally occurring plants. Specifically, *n*-hexanol, 2,5-dimethyl furan, 2-ethyl-1-hexanol, 3-hexen-1-ol and 2-nonanone were detected and either absent or present at low abundance in the no-flower controls (Table S3). Notably, floral samples with abundant fungal colony-forming units in nectar were more likely to contain some microbial-associated volatiles (Fig. 4), including 2-ethyl-1-hexanol ( $\chi^2 = 9.98$ ,  $P = 0.001$ ) and 2-nonanone ( $\chi^2 = 5.83$ ,  $P = 0.01$ ; Table S4) in the headspace sample. Plant species varied in the presence of *n*-hexanol ( $\chi^2 = 13.10$ ,  $P = 0.02$ ), but species identity did not predict the presence of other volatiles and no apparent relationship was found between bacterial abundance and focal volatiles detected in the field analyses (Table S4).


## Discussion

Here, we demonstrate that common microbial inhabitants of floral nectar differ in the volatile profiles emitted, and can influence acceptability of nectar to a generalist pollinator. The chemical compounds produced by the microorganisms assayed have been previously described in the floral headspace of plant species (Lemfack *et al.*, 2014). Some compounds, including



**Fig. 3** Proportion of positive proboscis extension responses by honey bee (*Apis mellifera*) following exposure to solutions inoculated with microorganisms. Acceptance differed among microbial species ( $P < 0.001$ ). Proboscis extension response (PER) indicates proboscis extension when bee was exposed to focal solution. Letters indicate treatments that differ at  $P < 0.05$  with Tukey's honest significant difference test. Number of bees tested for each treatment ranged between 40 and 82.

2-phenylethanol (Knudsen *et al.*, 2006; Galen *et al.*, 2011), are commonly documented and produced by many plant species, while others such as ethanol, 3-methyl-1-butanol, 3-hydroxy-2-butanone (acetoin) or ethyl acetate are described as major components of 'fetid' smelling flowers, including those that attract flies



**Fig. 4** Relationship between fungal abundance and the presence of microbially produced volatiles (Table 1) from floral headspace samples collected in the field. Points indicate individual samples containing flowers sampled from the field. A list of plant species sampled is provided in Supporting Information Table S3. Fungal abundance was quantified by plating nectar on YMA media. The line on each plot indicates the best-fit binomial regression model between fungal abundance (fungal colony forming units (CFU)) and presence of each individual compound ( $P < 0.05$ ). See also full results and statistics in Table S4.

or beetles (Dobson, 2006; Goodrich *et al.*, 2006). For example, the nectar of *Agave palmeri*, which smells of rotting fruit, overlaps to some degree with the chemicals described here, containing short-chain and aromatic alcohols including 3-methyl-1-butanol, *n*-hexanol and 2-phenylethanol (Raguso, 2004). By contrast, bee- or butterfly-pollinated species are typically not characterized by those fatty acid derivatives or fermentation volatiles (Dobson, 2006). Instead, these compounds are minor components of the volatile profile in some bee-pollinated plants, including field-collected clover (Buttery *et al.*, 1984), *Silene caroliniana* (Golonka *et al.*, 2014), and the current study. Although some compounds may be produced by both plants and microorganisms, if yeasts or other microbes contribute to floral scent, the presence of these components of the volatile profiles should be

dynamic over time, and vary with microbial presence and abundance in nectar, as our field survey suggests (Fig. 4). A recent meta-analysis indicated that floral volatile profiles are, on average, the most variable of all plant or animal volatile blends examined (Junker *et al.*, 2017). Our data suggest that nectar microbes can contribute to this variability.

Volatiles produced by nectar microbes are likely to be byproducts of microbial metabolism or fermentation, but may have diverse ecological functions. Nectar-inhabiting microbes often suppress the growth of late-arriving species in nectar (Peay *et al.*, 2012; Vannette & Fukami, 2014; Mittelbach *et al.*, 2016b), and the volatiles ethyl acetate, 2-butanol, isobutanol, ethanol, 2-ethyl-1-hexanol and 2-phenylethanol have been shown to inhibit microbial growth (Cruz *et al.*, 2012; Hua *et al.*, 2014; Pereira *et al.*, 2016). Alternatively, for microbes that rely on pollinator-mediated phoresis, volatiles may be attractive or aid in dispersal. The yeast *M. reukaufii* is a nectar specialist (Brysch-Herzberg, 2004; Pozo *et al.*, 2011) and is largely reliant on floral visitors for dispersal among flowers (Belisle *et al.*, 2012). Notably, *M. reukaufii* produced the volatile blend most attractive to honey bees among all microbes tested (Fig. 3). In field trials and feeding assays, *M. reukaufii* has been found to be either attractive (Herrera *et al.*, 2013; Schaeffer & Irwin, 2014; Schaeffer *et al.*, 2017) or not deterrent to bee pollinators (Good *et al.*, 2014). By contrast, the other microbes tested in this experiment have been isolated from a broad range of habitats, including plant surfaces, rotting fruits or pollinator-associated habitats (Swings & De Ley, 1981; Andrews *et al.*, 1994), so may be less reliant on pollinators for dispersal to appropriate habitats. Differences in honey bee attraction to microbial species support the hypothesis that microbes vary in their dependence on insect vectors or differ in the identity of dispersal vectors (Davis & Landolt, 2013; Vannette *et al.*, 2013).

Our study suggests that microbially produced volatiles have differential effects on honey bee physiology and behavior. For example, 2-butanol, a compound only emitted by *M. reukaufii*, elicited the strongest antennal response, over 400× greater than that of the 0.4 μmol citral control (e.g. Table 1). While EAG cannot reveal if this compound is attractive or deterrent and further studies should verify the ecological validity of chosen concentrations, PER assays indicated that the blend of compounds including 2-butanol was not deterrent. Yet other compounds emitted by *M. reukaufii* have been shown to act as insect honey bee semiochemicals. For example, 3-methylbutyl acetate (isoamyl acetate or isopentyl acetate) is the principal component of the honey bee alarm pheromone (Free, 1987; Hunt, 2007) and can influence *A. mellifera* behavior (Pastor & Seeley, 2005; Nieh, 2010; Urlacher *et al.*, 2010). Alternatively, it is possible that yeast-emitted volatiles could function, either individually or as a blend, as ‘honest signals’ of nectar rewards (Knauer & Schiestl, 2015). Our data cannot address if the identified microbial volatiles are attractive, honest signals or if *M. reukaufii* simply lacks deterrent volatile cues that may be produced by other microbes (Mittelbach *et al.*, 2016a). For example, scent produced by nectar-inhabiting microorganisms may indicate the presence of floral nectar and provide foraging cues for some pollinators.



Nevertheless, given that individual flowers vary in the presence and abundance of microorganisms (Fig. 1 and Herrera *et al.*, 2009), it is possible that pollinators could use these varying cues to distinguish among flowers depending on the presence or abundance of particular microorganisms.

The consequences of altered bee visitation or behavior for microbial dispersal and pollination remain unclear. If bacteria inhibit visitation by pollinators, their presence could reduce microbial dispersal and pollination, whereas decreased visit duration could instead increase dispersal and potential for outcrossing. Our results support the former (reduced attraction), but cannot resolve this question. Moreover, we only examined responses of the generalist pollinator *A. mellifera* in the current study so it is possible that other floral visitors respond either more or less strongly to the presence of microbial volatiles or differ in attraction to specific microbial taxa (Davis *et al.*, 2013), so the consequences for plant reproduction and microbial dispersal may depend on the specific combination of plant, microbe and floral visitor.

More generally, the finding that microbial volatiles can contribute to plant phenotype suggests a novel mechanism of microbial influence on ecological interactions between plants and animals. In addition to previously documented microbial effects on phenotype mediated by changes in nutrition, production of defensive compounds and altered hormonal signaling (Friesen, 2013), we present evidence that microbes can directly contribute to plant volatile chemotype, an understudied phenomenon. The fitness consequences for plants and microorganisms were not examined in this study, but our results suggest that when they align (as may be the case for *M. reukaufii* and plant reproduction), microbial volatile emission may enhance fitness benefits for both partners. Although more research is necessary to fully examine the prevalence, magnitude and consequences of microbial contribution to host semiochemicals, our results imply that microbial contribution to host volatile signaling may be an important but largely overlooked effect of the microbiome on host phenotype and subsequent ecological interactions.

## Acknowledgements

We thank W. S. Gee and S. D. Willms for laboratory assistance, the Niño lab for honey bees collected in Davis and C. Stuhl and B. Smith for ARS bees, the Insect Ecology group at UC Davis and Robert Schaeffer for helpful comments on previous versions of this work, and the Ebeler lab for assistance with GC-MS analyses of floral headspace from field samples. We acknowledge USDA-ARS research project no. 6036-22000-028 (J.J.B. and C.C.R.) and USDA Multistate Hatch grant NE1501 to R.L.V. (CA-D-ENM-2354-RR), and C.C.R. was supported by a USDA-ARS HQ-funded grant (0101-88888-016).

## Author contributions

R.L.V. and J.J.B. conceived the study, R.L.V. and G.W.H. performed field work, including nectar volatile collections and microbial isolations. G.W.H. performed PER assays. C.C.R.

performed volatile analyses of single strain inocula and EAG analyses. M.M.M. assisted with methods and GC-MS analyses of floral headspace samples. All authors contributed to writing and manuscript editing.

## References

- Andrews JH, Harris RF, Spear RN, Lau GW, Nordheim EV. 1994. Morphogenesis and adhesion of *Aureobasidium pullulans*. *Canadian Journal of Microbiology* 40: 6–17.
- Baker HG, Baker I. 1982. Chemical constituents of nectar in relation to pollination mechanisms and phylogeny. In: Nitecki HM, ed. *Biochemical aspects of evolutionary biology*. Chicago, IL, USA: Chicago University Press, 131–171.
- Beck JJ, Light DM, Gee WS. 2014. Electrophysiological responses of male and female *Amyeloid transitella* antennae to pistachio and almond host plant volatiles. *Entomologia Experimentalis et Applicata* 153: 217–230.
- Beck JJ, Smith L, Merrill GB. 2008. *In situ* volatile collection, analysis, and comparison of three *Centaurea* species and their relationship to biocontrol with herbivorous insects. *Journal of Agricultural and Food Chemistry* 56: 2759–2764.
- Beck JJ, Vannette RL. 2017. Harnessing insect–microbe chemical communications to control insect pests of agricultural systems. *Journal of Agricultural and Food Chemistry* 65: 23–28.
- Beck JJ, Willett DS, Gee WS, Mahoney NE, Higbee BS. 2016. Differentiation of volatile profiles from stockpiled almonds at varying relative humidity levels using benchtop and portable GC MS. *Journal of Agricultural and Food Chemistry* 64: 9286–9292.
- Belisle M, Peay KG, Fukami T. 2012. Flowers as islands: spatial distribution of nectar-inhabiting microfungi among plants of *Mimulus aurantiacus*, a hummingbird-pollinated shrub. *Microbial Ecology* 63: 711–718.
- Bhagavan S, Smith BH. 1997. Olfactory conditioning in the honey bee, *Apis mellifera*: effects of odor intensity. *Physiology & Behavior* 61: 107–117.
- Brysch-Herzberg M. 2004. Ecology of yeasts in plant-bumblebee mutualism in Central Europe. *FEMS Microbiology Ecology* 50: 87–100.
- Buser CC, Newcomb RD, Gaskett AC, Goddard MR. 2014. Niche construction initiates the evolution of mutualistic interactions. *Ecology Letters* 17: 1257–1264.
- Buttery RG, Kamm JA, Ling LC. 1984. Volatile components of red clover leaves, flowers, and seed pods: possible insect attractants. *Journal of Agricultural and Food Chemistry* 32: 254–256.
- Byers KJ, Bradshaw H, Riffell JA. 2014. Three floral volatiles contribute to differential pollinator attraction in monkeyflowers (*Mimulus*). *The Journal of Experimental Biology* 217: 614–623.
- Canto A, Herrera CM, Medrano M, Perez R, Garcia IM. 2008. Pollinator foraging modifies nectar sugar composition in *Helleborus foetidus* (Ranunculaceae): an experimental test. *American Journal of Botany* 95: 315–320.
- Cruz AF, Hamel C, Yang C, Matsubara T, Gan YT, Singh AK, Kuwada K, Ishii T. 2012. Phytochemicals to suppress *Fusarium* head blight in wheat-chickpea rotation. *Phytochemistry* 78: 72–80.
- Davis TS, Crippen TL, Hofstetter RW, Tomberlin JK. 2013. Microbial volatile emissions as insect semiochemicals. *Journal of Chemical Ecology* 39: 840–859.
- Davis TS, Landolt PJ. 2013. A survey of insect assemblages responding to volatiles from a ubiquitous fungus in an agricultural landscape. *Journal of Chemical Ecology* 39: 860–868.
- Dobson HE. 1994. Floral volatiles in insect biology. *Insect–Plant Interactions* 5: 47–81.
- Dobson HE. 2006. Relationship between floral fragrance composition and type of pollinator. In: Dudareva NA, Pichersky E, eds. *Biology of floral scent*. Boca Raton, FL, USA: CRC, 147–198.
- Free JB. 1987. *Pheromones of social bees*. London, UK: Chapman and Hall.
- Friesen ML. 2013. Microbially mediated plant functional traits. In: de Bruijn FJ, ed. *Molecular microbial ecology of the rhizosphere, two volume set*. Hoboken, NJ, USA: John Wiley & Sons, 87–102.
- Galen C, Kaczorowski R, Todd SL, Geib J, Raguso RA. 2011. Dose-dependent impacts of a floral volatile compound on pollinators, larcenists, and

- the potential for floral evolution in the Alpine Skypilot *Polemonium viscosum*. *American Naturalist* 177: 258–272.
- Galen C, Newport MEA. 1988. Pollination quality, seed set, and flower traits in *Polemonium viscosum*: complementary effects of variation in flower scent and size. *American Journal of Botany* 75: 900–905.
- Gardener MC, Gillman MP. 2001. Analyzing variability in nectar amino acids: composition is less variable than concentration. *Journal of Chemical Ecology* 27: 2545–2558.
- Gervasi DD, Schiestl FP. 2017. Real-time divergent evolution in plants driven by pollinators. *Nature Communications* 8: 14691.
- Golonka AM, Johnson BO, Freeman J, Hinson DW. 2014. Impact of nectarivorous yeasts on *Silene caroliniana*'s scent. *Eastern Biologist* 3: 1–26.
- Good AP, Gauthier M-PL, Vannette RL, Fukami T. 2014. Honey bees avoid nectar colonized by three bacterial species, but not by a yeast species, isolated from the bee gut. *PLoS ONE* 9: e86494.
- Goodrich KR, Zjhra ML, Ley CA, Raguso RA. 2006. When flowers smell fermented: the chemistry and ontogeny of yeasty floral scent in pawpaw (*Asimina triloba*: Annonaceae). *International Journal of Plant Sciences* 167: 33–46.
- Herrera CM, de Vega C, Canto A, Pozo MI. 2009. Yeasts in floral nectar: a quantitative survey. *Annals of Botany* 103: 1415–1423.
- Herrera CM, Garcia IM, Perez R. 2008. Invisible floral larcenies: microbial communities degrade floral nectar of bumble bee-pollinated plants. *Ecology* 89: 2369–2376.
- Herrera CM, Pozo MI, Medrano M. 2013. Yeasts in nectar of an early-blooming herb: sought by bumble bees, detrimental to plant fecundity. *Ecology* 94: 273–279.
- Hua SST, Beck JJ, Sarreal SBL, Gee W. 2014. The major volatile compound 2-phenylethanol from the biocontrol yeast, *Pichia anomala*, inhibits growth and expression of aflatoxin biosynthetic genes of *Aspergillus flavus*. *Mycotoxin Research* 30: 71–78.
- Hunt GJ. 2007. Flight and fight: a comparative view of the neurophysiology and genetics of honey bee defensive behavior. *Journal of Insect Physiology* 53: 399–410.
- Jiménez-Martínez ES, Bosque-Pérez NA, Berger PH, Zemetra RS, Ding H, Eigenbrode SD. 2004. Volatile cues influence the response of *Rhopalosiphum padi* (Homoptera: Aphididae) to Barley yellow dwarf virus-infected transgenic and untransformed wheat. *Environmental Entomology* 33: 1207–1216.
- Junker RR, Kuppler J, Amo L, Blande JD, Borges RM, Dam NM, Dicke M, Dötterl S, Ehlers BK, Etl F. 2017. Covariation and phenotypic integration in chemical communication displays: biosynthetic constraints and eco-evolutionary implications. *New Phytologist*. doi: 10.1111/nph.14505.
- Junker RR, Tholl D. 2013. Volatile organic compound mediated interactions at the plant–microbe interface. *Journal of Chemical Ecology* 39: 810–825.
- Kessler A, Baldwin IT. 2001. Defensive function of herbivore-induced plant volatile emissions in nature. *Science* 291: 2141–2144.
- Kessler A, Halitschke R, Poveda K. 2011. Herbivory-mediated pollinator limitation: negative impacts of induced volatiles on plant–pollinator interactions. *Ecology* 92: 1769–1780.
- Knauer A, Schiestl F. 2015. Bees use honest floral signals as indicators of reward when visiting flowers. *Ecology Letters* 18: 135–143.
- Knudsen JT, Eriksson R, Gershenzon J, Stahl B. 2006. Diversity and distribution of floral scent. *The Botanical Review* 72: 1–120.
- Lane DJ. 1991. 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M, eds. *Nucleic acid techniques in bacterial systematics*. New York, NY: John Wiley & Sons, 115–175.
- Lemfack MC, Nickel J, Dunkel M, Preissner R, Piechulla B. 2014. mVOC: a database of microbial volatiles. *Nucleic Acids Research* 42: D744–D748.
- Love M, Anders S, Huber W. 2013. *Differential analysis of count data, the DESeq2 package*. [WWW document] URL <http://bioconductor.org/packages/release/bioc/html/DESeq2.html>. v.1.16.1. [accessed 1 January 2017].
- Mann RS, Ali JG, Hermann SL, Tiwari S, Pelz-Stelinski KS, Alborn HT, Stelinski LL. 2012. Induced release of a plant–defense volatile ‘deceptively’ attracts insect vectors to plants infected with a bacterial pathogen. *PLoS Pathogens* 8: e1002610.
- Mittelbach M, Vannette RL. 2017. Mutualistic interactions of yeasts. Ecology and diversity. In: Buzzini P, Lachance M-A, Yurkov AM, eds. *Yeasts in natural ecosystems*. Berlin, Germany: Springer-Verlag.
- Mittelbach M, Yurkov AM, Begerow D. 2016a. Adaptive anonymity: crypsis as an evolutionary trait of floral yeasts? *bioRxiv*. doi: 10.1101/088179.
- Mittelbach M, Yurkov AM, Stoll R, Begerow D. 2016b. Inoculation order of nectar-borne yeasts opens a door for transient species and changes nectar rewarded to pollinators. *Fungal Ecology* 22: 90–97.
- Nieh JC. 2010. A negative feedback signal that is triggered by Peril Curbs Honey Bee recruitment. *Current Biology* 20: 310–315.
- O'Donnell K. 1993. Fusarium and its near relatives. In: Reynolds DR, Taylor JW, eds. *The fungal holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics*. Wallingford, UK: CAB International, 225–233.
- Oksanen J, Blanchet FG, Kindt R, Legendre P, O'Hara RB, Simpson GL, Solymos P, Stevens MH, Wagner H. 2012. *vegan: Community ecology package*. <https://cran.r-project.org/web/packages/vegan/index.html>. V 2.4-1.
- Parachnowitsch AL, Raguso RA, Kessler A. 2012. Phenotypic selection to increase floral scent emission, but not flower size or colour in bee-pollinated *Penstemon digitalis*. *New Phytologist* 195: 667–675.
- Paré PW, Tumlinson JH. 1999. Plant volatiles as a defense against insect herbivores. *Plant Physiology* 121: 325–332.
- Pastor KA, Seeley TD. 2005. The brief piping signal of the honey bee: begging call or stop signal? *Ethology* 111: 775–784.
- Peay KG, Belisle M, Fukami T. 2012. Phylogenetic relatedness predicts priority effects in nectar yeast communities. *Proceedings of the Royal Society B: Biological Sciences* 279: 749–758.
- Peñuelas J, Farré-Armengol G, Llusia J, Gargallo-Garriga A, Rico L, Sardans J, Terradas J, Filella I. 2014. Removal of floral microbiota reduces floral terpene emissions. *Scientific Reports* 4: 6727.
- Pereira JPC, Verheijen PJT, Straathof AJJ. 2016. Growth inhibition of *S. cerevisiae*, *B. subtilis*, and *E. coli* by lignocellulosic and fermentation products. *Applied Microbiology and Biotechnology* 100: 9069–9080.
- Pichersky E, Gershenzon J. 2002. The formation and function of plant volatiles: perfumes for pollinator attraction and defense. *Current Opinion in Plant Biology* 5: 237–243.
- Pineda A, Zheng S-J, van Loon JJA, Pieterse CMJ, Dicke M. 2010. Helping plants to deal with insects: the role of beneficial soil-borne microbes. *Trends in Plant Science* 15: 507–514.
- Pozo MI, de Vega C, Canto A, Herrera CM. 2009. Presence of yeasts in floral nectar is consistent with the hypothesis of microbial-mediated signaling in plant–pollinator interactions. *Plant Signal Behavior* 4: 1102–1104.
- Pozo MI, Herrera CM, Bazaga P. 2011. Species richness of yeast communities in floral nectar of southern Spanish plants. *Microbial Ecology* 61: 82–91.
- Raguso RA. 2001. Floral scent, olfaction, and scent-driven foraging behavior. In: Chittka L, Thomson JD, eds. *Cognitive ecology of pollination*. Cambridge, UK: Cambridge University Press, 83–105.
- Raguso RA. 2004. Why are some floral nectars scented? *Ecology* 85: 1486–1494.
- Raguso RA. 2008. Wake up and smell the roses: the ecology and evolution of floral scent. *Annual Review of Ecology, Evolution, and Systematics* 39: 549–569.
- Schaeffer R, Irwin RE. 2014. Yeasts in nectar enhance male fitness in a montane perennial herb. *Ecology* 95: 1792–1798.
- Schaeffer RN, Mei YZ, Andicoechea J, Manson JS, Irwin RE. 2017. Consequences of a nectar yeast for pollinator preference and performance. *Functional Ecology* 31: 613–621.
- Schaeffer RN, Phillips CR, Duryea MC, Andicoechea J, Irwin RE. 2014. Nectar yeasts in the tall larkspur *Delphinium barbeyi* (Ranunculaceae) and effects on components of pollinator foraging behavior. *PLoS One* 9: e108214.
- Schaeffer RN, Vannette RL, Irwin RE. 2015. Nectar yeasts in *Delphinium nuttallianum* (Ranunculaceae) and their effects on nectar quality. *Fungal Ecology* 18: 100–106.
- Scheiner R, Page R, Erber J. 2004. Sucrose responsiveness and behavioral plasticity in honey bees (*Apis mellifera*). *Apidologie* 35: 133–142.
- Shapiro L, De Moraes CM, Stephenson AG, Mescher MC. 2012. Pathogen effects on vegetative and floral odours mediate vector attraction and host exposure in a complex pathosystem. *Ecology Letters* 15: 1430–1438.

- Swings J, De Ley J. 1981. The genera *Gluconobacter* and *Acetobacter*. In: Starr MP, Stolp H, Truper HG, Balows A, Schlegel HG, eds. *The prokaryotes*. Berlin, Germany, New York, USA: Springer, 771–778.
- Turner S, Pryer KM, Miao VPW, Palmer JD. 1999. Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis. *Journal of Eukaryotic Microbiology* 46: 327–338.
- Urlacher E, Francés B, Giurfa M, Devaud J-M. 2010. An alarm pheromone modulates appetitive olfactory learning in the honeybee (*Apis Mellifera*). *Frontiers in Behavioral Neuroscience* 4: 1–10.
- Vannette RL, Fukami T. 2014. Historical contingency in species interactions: towards niche-based predictions. *Ecology Letters* 17: 115–124.
- Vannette RL, Fukami T. 2016. Nectar microbes can reduce secondary metabolites in nectar and alter effects on nectar consumption by pollinators. *Ecology* 97: 1410–1419.
- Vannette RL, Fukami T. 2017. Dispersal enhances beta diversity in nectar microbes. *Ecology Letters* 20: 901–910.
- Vannette RL, Gauthier M-PL, Fukami T. 2013. Nectar bacteria, but not yeast, weaken a plant–pollinator mutualism. *Proceedings of the Royal Society B: Biological Sciences* 280: 20122601.
- Wehner J, Mittelbach M, Rillig MC, Verbruggen E. 2017. Specialist nectar-yeasts decline with urbanization in Berlin. *Scientific Reports* 7: 45315.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information tab for this article:

**Fig. S1** The ‘wind tunnel’ volatile introduction system used for the electroantennographic bioassays.

**Table S1** Full names of plant species at Stebbins Cold Canyon reserve sampled for fungi and bacteria

**Table S2** Unknown compound retention indices, principal ions and peak areas in *ex situ* microbial headspace

**Table S3** Plant species names and other metadata for floral headspace samples

**Table S4** Statistical results of generalized linear model examining fungal abundance, bacterial abundance and plant species on the presence of microbially associated volatiles in floral headspace collected in the field

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



## About New Phytologist

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication ‘as ready’ via *Early View* – our average time to decision is <26 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**