

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Constraints on anomalous Higgs boson couplings from its production and decay using the WW channel in proton-proton collisions at $s = 13$ TeV.

Permalink

<https://escholarship.org/uc/item/7973x5f0>

Journal

European Physical Journal C: Particles and Fields, 84(8)

ISSN

1434-6044

Publication Date

2024

DOI

10.1140/epjc/s10052-024-12925-0

Peer reviewed



Constraints on anomalous Higgs boson couplings from its production and decay using the WW channel in proton–proton collisions at $\sqrt{s} = 13$ TeV

CMS Collaboration*

CERN, 1211 Geneva 23, Switzerland

Received: 1 March 2024 / Accepted: 16 May 2024
© CERN for the benefit of the CMS Collaboration 2024

Abstract A study of the anomalous couplings of the Higgs boson to vector bosons, including CP -violation effects, has been conducted using its production and decay in the WW channel. This analysis is performed on proton–proton collision data collected with the CMS detector at the CERN LHC during 2016–2018 at a center-of-mass energy of 13 TeV, and corresponds to an integrated luminosity of 138 fb^{-1} . The different-flavor dilepton ($e\mu$) final state is analyzed, with dedicated categories targeting gluon fusion, electroweak vector boson fusion, and associated production with a W or Z boson. Kinematic information from associated jets is combined using matrix element techniques to increase the sensitivity to anomalous effects at the production vertex. A simultaneous measurement of four Higgs boson couplings to electroweak vector bosons is performed in the framework of a standard model effective field theory. All measurements are consistent with the expectations for the standard model Higgs boson and constraints are set on the fractional contribution of the anomalous couplings to the Higgs boson production cross section.

1 Introduction

After the discovery of the Higgs boson (H) by the ATLAS and CMS Collaborations in 2012 [1–3], the CMS [4–11] and ATLAS [12–18] experiments set constraints on the spin-parity properties of the Higgs boson and its couplings with gluons and electroweak (EW) gauge bosons, denoted here as Hgg and HVV , respectively. The Higgs boson quantum numbers are consistent with the standard model (SM) expectation $J^{PC} = 0^{++}$, but the possibility of small, anomalous couplings is not yet ruled out. In beyond-the-SM (BSM) theories, interactions with the Higgs boson may occur through several anomalous couplings, which lead to new tensor structures in the interaction terms that can be both CP -even or CP -odd. The CP -odd anomalous couplings between the Higgs boson and BSM particles may generate CP violation in the interactions of the Higgs boson.

In this paper, we study the tensor structure of the Hgg and HVV couplings, and we search for several anomalous effects, including CP violation, using the different-flavor dilepton ($e\mu$) final state from $H \rightarrow WW$ decays. The Higgs boson production processes include gluon fusion (ggH), EW vector boson fusion (VBF), and associated production with a W or Z boson (VH). Higgs boson production and decay processes are sensitive to certain anomalous contributions, which can be described by higher-dimensional operators in an effective field theory (EFT) [19] that can modify the kinematic distributions of the Higgs boson decay products and the particles from associated production.

Each production process of the Higgs boson is identified using its kinematic features, and events are assigned to corresponding production categories. The matrix element likelihood approach (MELA) [20–24] is employed to construct observables that are optimal for the measurement of anomalous couplings, or EFT operators, at the production vertex. These and other decay-based variables are used to explore all kinematic features of the events, giving the analysis sensitivity to simultaneous anomalous effects at the Higgs boson production and decay vertices. Fully simulated signal samples that include anomalous couplings incorporate the detector response into the analysis.

The analysis is based on the proton–proton (pp) collision data collected at the CERN LHC from 2016 to 2018, at a center-of-mass energy of 13 TeV, corresponding to an integrated luminosity of 138 fb^{-1} . This paper builds on a previous analysis conducted by the CMS Collaboration in the $H \rightarrow WW$ channel [25], which focused on measuring the Higgs boson production cross sections and coupling parameters in the so-called κ framework [26]. We follow a formalism used in previous CMS analyses of anomalous couplings in Run 1 and Run 2 [4–11, 27, 28], focusing on the case where the Higgs boson is produced on-shell. The coupling parame-

* e-mail: cms-publication-committee-chair@cern.ch (corresponding author)

ters are extracted using the signal strength and the fractional contributions of the couplings to the cross section. A general study of the HVV interaction is performed with four anomalous couplings analyzed individually. Through $SU(2) \times U(1)$ symmetry considerations, the anomalous HVV couplings are reduced in number to three and analyzed simultaneously. The primary HVV coupling measurements are performed in terms of cross section fractions with additional interpretations in terms of EFT couplings included. A study of the Hgg interaction is also performed in terms of a CP -odd anomalous coupling cross section fraction.

This paper is organized as follows. The phenomenology of anomalous couplings is discussed in Sect. 2. Section 3 gives a brief overview of the CMS apparatus. Data sets and Monte Carlo (MC) simulation samples are discussed in Sect. 4. The event reconstruction and selection are outlined in Sects. 5 and 6, respectively. Methods to estimate backgrounds are given in Sect. 7. In Sect. 8, we discuss the kinematic variables associated with Higgs boson production and decay. Sources of systematic uncertainties are presented in Sect. 9. The results are presented and discussed in Sect. 10. Finally, a summary is given in Sect. 11. Tabulated results are provided in the HEPData record for this analysis [29].

2 Phenomenology

In this analysis, we investigate anomalous coupling effects in gluon fusion or electroweak Higgs boson production, as well as in its decay to WW pairs. A detailed discussion of the theoretical considerations can be found in Refs. [22, 24, 28]. The interaction of the spin-zero Higgs boson with two spin-one gauge bosons $V_1 V_2$, such as WW, ZZ, $Z\gamma$, $\gamma\gamma$, or gg, can be parametrized by the scattering amplitude

$$A(HV_1 V_2) \sim \left[a_1^{VV} + \frac{\kappa_1^{VV} q_{V1}^2 + \kappa_2^{VV} q_{V2}^2}{(\Lambda_1^{VV})^2} \right] m_{V1}^2 \epsilon_{V1}^* \epsilon_{V2}^* + \frac{1}{v} a_2^{VV} f_{\mu\nu}^{*(1)} f^{*(2),\mu\nu} + \frac{1}{v} a_3^{VV} f_{\mu\nu}^{*(1)} \tilde{f}^{*(2),\mu\nu}, \quad (1)$$

where q_{Vi} and ϵ_{Vi} are the spin-one gauge boson four-momentum and polarization vectors, m_{V1} is the pole mass of the boson, $f^{(i),\mu\nu} = \epsilon_{Vi}^\mu q_{Vi}^\nu - \epsilon_{Vi}^\nu q_{Vi}^\mu$ and $\tilde{f}_{\mu\nu}^{(i)} = \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} f^{(i),\rho\sigma}$ (with $\epsilon_{\mu\nu\rho\sigma}$ the Levi-Civita symbol), Λ_1^{VV} is the scale of BSM physics, and v is the Higgs field vacuum expectation value.

The only leading tree-level contributions in the scattering amplitude are $a_1^{ZZ} \neq 0$ and $a_1^{WW} \neq 0$; other a_1 coupling parameters ($Z\gamma$, $\gamma\gamma$, gg) do not contribute because the pole mass vanishes. Additional ZZ and WW couplings are considered anomalous contributions. Anomalous terms arising in the SM via loop effects are typically small and are

not yet accessible experimentally. The BSM contributions, however, could yield larger coupling parameters. Among the anomalous contributions, considerations of symmetry and gauge invariance require $\kappa_1^{ZZ} = \kappa_2^{ZZ}$, $\kappa_1^{\gamma\gamma} = \kappa_2^{\gamma\gamma} = 0$, $\kappa_1^{gg} = \kappa_2^{gg} = 0$, and $\kappa_1^{Z\gamma} = 0$ [24]. The presence of CP -odd a_3^{VV} couplings together with any of the other couplings (all of them CP -even), will result in CP violation. We reduce the number of independent parameters by assuming that $a_2^{\gamma\gamma}$, $a_3^{\gamma\gamma}$, $a_2^{Z\gamma}$ and $a_3^{Z\gamma}$ are constrained in direct decays of $H \rightarrow \gamma\gamma$ and $Z\gamma$, therefore fixing them to be zero. The a_2^{gg} term results from loop effects in the SM.

The relationship between the ZZ and WW couplings is mostly relevant for VBF production. There are no kinematic differences between the ZZ and WW fusion processes; therefore, it is not possible to disentangle the couplings. One possibility is to set the ZZ and WW couplings to be equal, $a_i^{ZZ} = a_i^{WW}$, leaving four HVV anomalous couplings to be measured: a_2 , a_3 , $\kappa_1/(\Lambda_1)^2$, and $\kappa_2^{Z\gamma}/(\Lambda_1^{Z\gamma})^2$. The $a_1^{ZZ} = a_1^{WW}$ relationship also appears under custodial symmetry. This approach provides a general test of the Higgs boson Lagrangian tensor structure and a search for CP violation in HVV interactions. In an alternative approach, the $SU(2) \times U(1)$ symmetry reduces the number of independent HVV anomalous couplings to three (a_2 , a_3 , and $\kappa_1/(\Lambda_1)^2$) through the introduction of the following coupling parameter relationships [19] :

$$a_1^{WW} = a_1^{ZZ}, \quad (2)$$

$$a_2^{WW} = c_w^2 a_2^{ZZ}, \quad (3)$$

$$a_3^{WW} = c_w^2 a_3^{ZZ}, \quad (4)$$

$$\frac{\kappa_1^{WW}}{(\Lambda_1^{WW})^2} = \frac{1}{c_w^2 - s_w^2} \left(\frac{\kappa_1^{ZZ}}{(\Lambda_1^{ZZ})^2} - 2s_w^2 \frac{a_2^{ZZ}}{m_Z^2} \right), \quad (5)$$

$$\frac{\kappa_2^{Z\gamma}}{(\Lambda_1^{Z\gamma})^2} = \frac{2s_w c_w}{c_w^2 - s_w^2} \left(\frac{\kappa_1^{ZZ}}{(\Lambda_1^{ZZ})^2} - \frac{a_2^{ZZ}}{m_Z^2} \right), \quad (6)$$

where c_w and s_w are the cosine and sine of the weak mixing angle, respectively, and m_Z is the Z boson mass. With this approach, there is a linear relationship between the scattering amplitude couplings and the SM EFT (SMEFT) couplings in the Higgs basis [19]:

$$\delta c_z = \frac{1}{2} a_1^{ZZ} - 1, \quad (7)$$

$$c_{zz} = -\frac{2s_w^2 c_w^2}{e^2} a_2^{ZZ}, \quad (8)$$

$$\tilde{c}_{zz} = -\frac{2s_w^2 c_w^2}{e^2} a_3^{ZZ}, \quad (9)$$

$$c_{z\Box} = \frac{m_Z^2 s_w^2}{e^2} \frac{\kappa_1^{ZZ}}{(\Lambda_1^{ZZ})^2}, \quad (10)$$

where e is the electron charge. The amplitude couplings may also be related to the SMEFT Warsaw basis [19, 30] couplings through the following translation [28, 31]:

$$\delta a_1^{ZZ} = \frac{v^2}{\Lambda^2} \left(2c_{H\Box} + \frac{6e^2}{s_w^2} c_{HWB} + \left(\frac{3c_w^2}{2s_w^2} - \frac{1}{2} \right) c_{HD} \right), \quad (11)$$

$$\kappa_1^{ZZ} = \frac{v^2}{\Lambda^2} \left(-\frac{2e^2}{s_w^2} c_{HWB} + \left(1 - \frac{1}{2s_w^2} \right) c_{HD} \right), \quad (12)$$

$$a_2^{ZZ} = -2 \frac{v^2}{\Lambda^2} \left(s_w^2 c_{HB} + c_w^2 c_{HW} + s_w c_w c_{HWB} \right), \quad (13)$$

$$a_3^{ZZ} = -2 \frac{v^2}{\Lambda^2} \left(s_w^2 c_{H\tilde{B}} + c_w^2 c_{H\tilde{W}} + s_w c_w c_{H\tilde{WB}} \right), \quad (14)$$

where Λ is the UV cutoff of the theory (set to 1 TeV), and δa_1^{ZZ} is a correction to the SM value of a_1^{ZZ} . Further discussion on the EFT operators corresponding to the couplings considered here may be found in Chapter 2.2 of Ref. [19]. The assumed constraints on $a_2^{\gamma\gamma}$, $a_3^{\gamma\gamma}$, $a_2^{Z\gamma}$ and $a_3^{Z\gamma}$ imply that only one of the three coupling parameters c_{HW} , c_{HWB} , and c_{HB} is independent; the same is also true for their CP -odd counterparts $c_{H\tilde{W}}$, $c_{H\tilde{WB}}$, and $c_{H\tilde{B}}$. Therefore, we have four independent HVV couplings in both the Higgs and Warsaw basis. All the EFT couplings are expected to be zero in the SM.

We thus adopt two approaches to the HVV coupling study. In Approach 1, we use the $a_i^{ZZ} = a_i^{WW}$ relationship and individually analyze each of the four anomalous couplings. In Approach 2, we enforce the $SU(2) \times U(1)$ relationships from Eqs. (2–6) and analyze the three independent anomalous couplings both individually and simultaneously. Approach 1 may be considered to follow the relationships from Eqs. (2–5) in the limiting case $c_w = 1$.

It is convenient to measure the fractional contribution of the anomalous couplings to the Higgs boson cross section rather than the anomalous couplings themselves. For the anomalous HVV couplings, the effective fractional cross sections f_{ai} are defined as

$$f_{ai} = \frac{|a_i|^2 \sigma_i}{\sum_j |a_j|^2 \sigma_j} \text{sign} \left(\frac{a_i}{a_1} \right), \quad (15)$$

where \sum_j sums over all the coupling parameters considered, including a_1 , and σ_i is the cross section for the process corresponding to $a_i = 1$ and $a_{j \neq i} = 0$. Many systematic uncertainties cancel out in the ratio, and the physical range is conveniently bounded between -1 and $+1$. Our primary measurements are performed in terms of cross section frac-

Table 1 The cross sections (σ_i) of the anomalous contributions (a_i) relative to the SM value (σ_1) used to define the fractional cross sections f_{ai} for the Approach 1 and 2 coupling relationships. For the κ_1 and $\kappa_2^{Z\gamma}$ couplings, the numerical values $\Lambda_1 = \Lambda_1^{Z\gamma} = 100$ GeV are chosen to keep all coefficients of similar order of magnitude

f_{ai}	a_i	Approach 1 σ_i/σ_1	Approach 2 σ_i/σ_1
f_{a2}	a_2	0.361	6.376
f_{a3}	a_3	0.153	0.153
f_{A1}	κ_1	0.682	5.241
$f_{A1}^{Z\gamma}$	$\kappa_2^{Z\gamma}$	1.746	—

tions, with additional interpretations in terms of the SMEFT Higgs and Warsaw basis couplings also included. For consistency with previous CMS measurements, the σ_i coefficients used to define the fractional cross sections correspond to the $gg \rightarrow H \rightarrow VV \rightarrow 2e2\mu$ process [28]. The numerical values are given in Table 1 as calculated using the JHUGEN simulation [20–23]. Two sets of values are shown corresponding to the different coupling relationships adopted in Approach 1 and 2.

It has been shown that the angular correlations of the associated jets in the $ggH + 2$ jets process are sensitive to anomalous Hgg coupling effects at the production vertex [32]. The quark-quark initiated process, $qq \rightarrow qqH$, corresponds to the gluon scattering topology sensitive to anomalous effects. For the anomalous Hgg coupling, the effective fractional cross section can be defined as

$$f_{a3}^{ggH} = \frac{|a_3^{gg}|^2 \sigma_3^{gg}}{|a_2^{gg}|^2 \sigma_2^{gg} + |a_3^{gg}|^2 \sigma_3^{gg}} \text{sign} \left(\frac{a_3^{gg}}{a_2^{gg}} \right). \quad (16)$$

The σ_3^{gg} and σ_2^{gg} cross sections correspond to $a_3^{gg} = 1$, $a_2^{gg} = 0$ and $a_2^{gg} = 1$, $a_3^{gg} = 0$, respectively, and are equal. With this analysis it is not possible to distinguish the top quark, bottom quark, and heavy BSM particle contributions in the gluon fusion loop. As such, the Hgg coupling is treated as an effective coupling with heavy degrees of freedom integrated out.

3 The CMS detector

The CMS apparatus [33] is a multipurpose, nearly hermetic detector, designed to identify electrons, muons, photons, and (charged and neutral) hadrons [34–37]. A global reconstruction “particle-flow” (PF) algorithm [38] combines the information provided by the all-silicon inner tracker and by the crystal electromagnetic and brass-scintillator hadron calorimeters, operating inside a 3.8 T superconducting solenoid, with data from gas-ionization muon detectors interleaved with the solenoid return yoke, to build τ lep-

tons, jets, missing transverse momentum, and other physics objects [39–41].

Events of interest are selected using a two-tiered trigger system [42,43]. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4 μs [42]. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [43]. A more detailed description of the CMS detector, together with a definition of the coordinate system and kinematic variables, can be found in Ref. [33].

4 Data sets and simulation

The data sets included in this analysis were recorded with the CMS detector in 2016, 2017, and 2018, and correspond to integrated luminosities of 36.3, 41.5, and 59.7 fb^{-1} , respectively [44–46]. The collision events must fulfill HLT selection criteria that require the presence of one or two leptons satisfying isolation and identification requirements. For the 2016 data set, the single-electron trigger has a transverse momentum (p_{T}) threshold of 25 GeV for electrons with pseudorapidity $|\eta| < 2.1$ and 27 GeV for $2.1 < |\eta| < 2.5$, whereas the single-muon trigger has a p_{T} threshold of 24 GeV for $|\eta| < 2.4$. For the 2017 (2018) data set, the p_{T} threshold is 35 (32) GeV for the single-electron trigger (covering $|\eta| < 2.5$) and 27 (24) GeV for the single-muon trigger ($|\eta| < 2.4$). The dilepton $e\mu$ trigger has p_{T} thresholds of 23 and 12 GeV for the leading and subleading leptons, respectively, with the same coverage in pseudorapidity for electrons and muons as above. During the first part of data taking in 2016, a lower p_{T} threshold of 8 GeV for the subleading muon was used.

Monte Carlo event generators are used to model the signal and background processes. For each process, three independent sets of simulated events, corresponding to the three years of data taking, are used. This approach includes year-dependent effects in the CMS detector, data taking, and event reconstruction. All simulated events corresponding to a given data set share the same set of parton distribution functions (PDFs), underlying event (UE) tune, and parton shower (PS) configuration. The PDF sets used are NNPDF 3.0 [47,48] for 2016 and NNPDF 3.1 [49] for 2017 and 2018. The CUETP8M1 [50] tune is used to describe the UE in 2016 simulations, whereas the CP5 [51] tune is adopted in 2017 and 2018 simulated events. The MC samples are interfaced with PYTHIA 8.226 [52] in 2016, and 8.230 in 2017 and 2018, for the modeling of UE, PS, and hadronization. Standard Model Higgs boson production through ggH, VBF, and VH is simulated at next-to-leading order (NLO) accuracy

in quantum chromodynamics (QCD), including finite quark mass effects, using POWHEG v2 [53–59]. The MINLO HVJ [58] extension of POWHEG v2 is used for the simulation of WH and quark-induced ZH production, providing NLO accuracy for the VH + 0- and 1-jet processes. For ggH production, the simulated events are weighted to match the NNLOPS [60,61] prediction in the hadronic jet multiplicity (N_{jet}) and Higgs boson p_{T} distributions. The weighting is based on p_{T} and N_{jet} as computed in the simplified template cross section scheme 1.0 [62]. The MINLO HHJ [63] generator, which provides NLO accuracy for $N_{\text{jet}} \geq 2$, is also used for ggH production. The associated production processes with top quarks ($t\bar{t}\text{H}$) and bottom quarks ($b\bar{b}\text{H}$) are simulated with POWHEG v2 and MADGRAPH5_aMC@NLO v2.2.2 [64], respectively, and have a negligible contribution in the analysis phase space. All SM Higgs boson samples are normalized to the cross sections recommended in Ref. [19]. The Higgs boson mass in the event generation is assumed to be 125 GeV, while a value of 125.38 GeV [65] is used for the calculation of cross sections and branching fractions. The decay to a pair of W bosons and subsequently to leptons or hadrons is performed using the JHUGEN v5.2.5 generator in 2016, and v7.1.4 in 2017 and 2018, for ggH, VBF, and quark-induced ZH samples. The Higgs boson and W boson decays are performed using PYTHIA 8.212 for the other signal simulations.

The ggH, VBF, and VH Higgs boson events with HVV anomalous couplings are generated with JHUGEN at LO accuracy. With respect to the $\kappa_2^{Z\gamma}/(\Lambda_1^{Z\gamma})^2$ coupling parameter discussed in Sect. 2, the sign convention of the photon field is determined by the sign in front of the gauge fields in the covariant derivative. In this analysis, we define the covariant derivative $D_{\mu} = \partial_{\mu} - ie\sigma^i W_{\mu}^i/(2s_w) + ieB_{\mu}/(2c_w)$ following the convention in JHUGEN [31]. The JHUGEN and POWHEG SM Higgs boson simulations were compared after parton showering and no significant differences in the distributions of kinematic observables were found. We adopt the JHUGEN simulation to describe the kinematic features in all production modes with HVV anomalous couplings. The expected yields are scaled to match the SM theoretical predictions [19] for inclusive cross sections and the POWHEG SM prediction of relative event yields in the event categorization based on associated particles. Simulation of the ggH + 2 jets process with Hgg anomalous couplings is done using MINLO X0JJ [66] at NLO in QCD. A large number of signal samples with various anomalous couplings were generated. The MELA package [20–24] contains a library of matrix elements from JHUGEN for different Higgs boson signal hypotheses. Matrix elements from different coupling signal hypotheses, but with the same production mechanism, are used to reweight the generated signal events. This procedure is used in the construction of the predictions for the different coupling components and their interference, allowing

ing us to cover all points in the signal model phase space with sufficient statistical precision.

Background events are produced using several simulations. The quark-initiated nonresonant WW process is simulated with POWHEG v2 [67] at NLO accuracy for inclusive production. A reweighting is performed to match the diboson p_T spectrum computed at NNLO+NNLL QCD accuracy [68, 69]. The MCFM v7.0 [70–72] generator is used to simulate gluon-induced WW production at LO accuracy, with the normalization chosen to match the NLO cross section [73]. Nonresonant EW production of WW pairs with two additional jets is simulated at LO accuracy with MADGRAPH5_amc@NLO v2.4.2 using the MLM matching and merging scheme [74]. Top quark pair production ($t\bar{t}$) and single top quark processes, including tW, s- and t-channel contributions, are simulated with POWHEG v2 [75–77]. A reweighting of the top quark and antiquark p_T spectrum at parton level is performed for the $t\bar{t}$ simulation in order to match the NNLO and next-to-next-to-leading logarithm (NNLL) QCD predictions, including also the NLO EW contribution [78].

The Drell–Yan (DY) production of a charged-lepton pair is simulated with MADGRAPH5_amc@NLO v2.4.2 at NLO accuracy with up to two additional partons, using the FxFx matching and merging scheme [79]. Production of a W boson associated with an initial state radiation photon ($W\gamma$) is simulated with MADGRAPH5_amc@NLO v2.4.2 at NLO accuracy with up to 1 additional parton, using the FxFx jet merging. Diboson processes containing at least one Z boson or a virtual photon (γ^*) with a mass as low as 100 MeV are generated with POWHEG v2 [67] at NLO accuracy. Production of a W boson in association with a γ^* ($W\gamma^*$) for masses below 100 MeV is simulated by PYTHIA 8.212 in the parton showering of $W\gamma$ events. Triboson processes with inclusive decays are also simulated at NLO accuracy with MADGRAPH5_amc@NLO v2.4.2.

For all processes, the detector response is simulated using a detailed description of the CMS detector, based on the GEANT4 toolkit [80]. The distribution of additional pp interactions within the same or nearby bunch crossings (pileup) in the simulation is reweighted to match that observed in data. The efficiency of the trigger system is evaluated in data on a per lepton basis using dilepton events consistent with the Z boson decay. The overall efficiencies of the trigger selections used in the analysis are obtained as the average of the per-lepton efficiencies weighted by their probability. The resulting efficiencies are applied directly on simulated events.

5 Event reconstruction

The identification and measurement of the properties of individual particles (PF candidates) in an event is achieved in the PF algorithm by combining information from various subde-

tectors. Electrons are identified and their momenta are measured in the pseudorapidity interval $|\eta| < 2.5$ by combining tracks in the silicon tracker with spatially compatible energy deposits in the electromagnetic calorimeter. Muons are identified and their momenta are measured in the pseudorapidity range $|\eta| < 2.4$ by matching tracks in the muon system and the silicon tracker. For better rejection of nonprompt leptons, increasing the sensitivity of the analysis, leptons are required to be isolated and well reconstructed using a set of criteria based on the quality of the track reconstruction, shape of calorimetric deposits, and energy flux in the vicinity of the particle's trajectory [34, 35]. In addition, a selection based on a dedicated multivariate analysis (MVA) tagger developed for the CMS $t\bar{t}H$ analysis [81] is added in all channels for muon candidates.

Multiple pp interaction vertices are identified from tracking information by use of the adaptive vertex fitting algorithm [82]. The primary pp interaction vertex is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in Section 9.4.1 of Ref. [83]. Leptons are required to be associated to the primary vertex using transverse and longitudinal impact parameter criteria [34, 35].

Hadronic jets are clustered from PF candidates using the infrared- and collinear-safe anti- k_T algorithm with distance parameters of 0.4 (AK4) and 0.8 (AK8). The jet momentum is determined as the vectorial sum of all particle momenta in the jet. The AK8 jets considered are required to be reconstructed within the silicon tracker acceptance ($|\eta| < 2.4$), whereas AK4 jets are reconstructed in the range $|\eta| < 4.7$. For AK4 jets, contamination from pileup is suppressed using charged-hadron subtraction which removes charged PF candidates originating from vertices other than the primary interaction vertex. The residual contribution from neutral particles originating from pileup vertices is removed by means of an event-by-event jet-area-based correction to the jet four-momentum [84]. For AK8 jets, the pileup-per-particle identification algorithm (PUPPI) [85] is used to mitigate the effect of pileup at the reconstructed-particle level, making use of local shape information, event pileup properties, and tracking information. Additional selection criteria are applied to remove jets potentially dominated by instrumental effects or reconstruction failures [84].

The AK8 jets are used to reconstruct hadronic V boson decays in a single merged jet when the decay products are highly collimated. This approach targets boosted W or Z bosons originating from the VH production mode. Such Lorentz-boosted V decays are identified using the ratio of the 2- to 1-subjettiness [86], τ_2/τ_1 , and the groomed jet mass m_J . The groomed mass is calculated after applying a modified mass drop algorithm [87, 88], known as the soft-drop algorithm [89], with parameters $\beta = 0$, $z_{\text{cut}} = 0.1$, and

$R_0 = 0.8$. The algorithm also identifies two hard subjets within the AK8 jet.

We refer to the identification of jets likely originating from bottom quarks as b tagging [90,91]. For each AK4 jet in the event, a score is calculated through a multivariate combination of different jet properties, making use of boosted decision trees and deep neural networks. A jet is considered b-tagged if its associated score exceeds a threshold, tuned to achieve a certain tagging efficiency as measured in $t\bar{t}$ events. The chosen working point corresponds to about 90% efficiency for bottom quark jets and to a mistagging rate of about 10% for light-flavor or gluon jets and of about 50% for charm quark jets.

The missing transverse momentum vector \vec{p}_T^{miss} is computed as the negative vector sum of the transverse momenta of all the PF candidates in an event, and its magnitude is denoted as p_T^{miss} [41]. The PUPPI algorithm is applied to reduce the pileup dependence of the \vec{p}_T^{miss} observable by computing the \vec{p}_T^{miss} from the PF candidates weighted by their probability to originate from the primary interaction vertex [41].

6 Event selection

The analysis is performed using $H \rightarrow WW$ candidate events in the $e\mu$ final state. For an event to be selected, the transverse momenta of the leading lepton $p_T^{\ell 1}$ and the subleading lepton $p_T^{\ell 2}$ must be greater than 25 and 13 GeV, respectively. The $p_T^{\ell 2}$ threshold in the case of a muon is lowered to 10 GeV for the 2016 data set because of the lower threshold in the corresponding HLT algorithm. Events containing additional leptons with $p_T > 10$ GeV are discarded. The dilepton system is required to have an invariant mass $m_{\ell\ell}$ greater than 12 GeV and transverse momentum $p_T^{\ell\ell}$ above 30 GeV. A requirement on the missing transverse momentum of $p_T^{\text{miss}} > 20$ GeV is implemented. We define transverse mass discriminating variables m_T^H and $m_T^{\ell 2}$ as

$$m_T^H = \sqrt{2 p_T^{\ell\ell} p_T^{\text{miss}} [1 - \cos \Delta\Phi(\vec{p}_T^{\ell\ell}, \vec{p}_T^{\text{miss}})]}, \quad (17)$$

$$m_T^{\ell 2} = \sqrt{2 p_T^{\ell 2} p_T^{\text{miss}} [1 - \cos \Delta\Phi(\vec{p}_T^{\ell 2}, \vec{p}_T^{\text{miss}})]}, \quad (18)$$

and select events with $m_T^H > 60$ GeV and $m_T^{\ell 2} > 30$ GeV. The m_T^H requirement suppresses the $DY \rightarrow \tau\tau$ background process and avoids overlap with the $H \rightarrow \tau\tau$ analysis [11]. To ensure orthogonality with a future off-shell $H \rightarrow WW$ analysis we require $m_T^H < 125$ GeV. In addition, the region $76.2 < m_{\ell\ell} < 106.2$ GeV is excluded to avoid overlap with the off-shell $H \rightarrow ZZ \rightarrow 2\ell 2\nu$ analysis [10]. These requirements will simplify a future combination of Higgs boson decay final states. Finally, events with any b-tagged jets with

Table 2 Summary of the base selection criteria

Variable	Selection
Number of leptons	2 ($e\mu$ of opposite charge)
$p_T^{\ell 1}$	> 25 GeV
$p_T^{\ell 2}$	> 13 GeV (10 GeV for 2016 data)
$m_{\ell\ell}$	12–76.2 GeV or > 106.2 GeV
$p_T^{\ell\ell}$	> 30 GeV
p_T^{miss}	> 20 GeV
$m_T^{\ell 2}$	> 30 GeV
m_T^H	60–125 GeV
$N_{\text{jet}}(\text{bjets})$	0

Table 3 Summary of the ggH, VBF, and VH production channels used for the HVV coupling study

Variable	ggH	VBF	Resolved VH	Boosted VH
$N_{\text{jet}}(\text{V jets})$	0	0	0	> 0
$N_{\text{jet}}(\text{AK4 jets})$	0 and 1	2	2	—
m_{jj}	—	> 120 GeV	60–120 GeV	—

$p_T > 20$ GeV are vetoed. These base selection criteria are summarized in Table 2.

For the HVV coupling analysis, exclusive selection criteria, which are based on the associated jet activity in the event, are applied that target the ggH, VBF, and VH production processes. The AK4 (AK8) jets considered are required to have $p_T > 30$ (200) GeV. In the ggH channel, zero or one AK4 jet is required in the event. For the VBF and Resolved VH channels, we require two AK4 jets with dijet masses of $m_{jj} > 120$ GeV and $60 < m_{jj} < 120$ GeV, respectively. The Boosted VH channel requires the presence of a V-tagged AK8 jet (V jet); such jets have a groomed mass in the region $65 < m_j < 105$ GeV and satisfy the requirement $\tau_2/\tau_1 < 0.4$. In the other channels, a V jet veto is implemented to ensure orthogonality. These production channels for the HVV coupling study are summarized in Table 3.

As the production vertex of the ggH + 2 jets process is sensitive to anomalous Hgg coupling effects, we use a 2-jet ggH channel that follows the VBF selection described above for the Hgg coupling analysis. The $HW\bar{W}$ decay vertex is not sensitive to anomalous Hgg effects, and so decay-based variables are not studied in this channel. This permits a relatively tight selection of $m_{\ell\ell} < 55$ GeV which is beneficial for background suppression. The 0- and 1-jet ggH channels are also included to constrain the ggH signal strength. All channels included for the Hgg coupling study are summarized in Table 4.

Control regions (CRs) are defined using the base selection criteria together with a set of alternative requirements sum-

Table 4 Summary of ggH channel selections used for the Hgg coupling study

Variable	ggH	2-jet ggH
N_{jet} (AK4 jets)	0 and 1	2
m_{jj}	—	$> 120 \text{ GeV}$
$m_{\ell\ell}$	—	$< 55 \text{ GeV}$

Table 5 Summary of the $\tau\tau$, top quark, and WW control region requirements

Variable	$\tau\tau$	Top quark	WW
$m_{\ell\ell}$	40–80 GeV	$> 50 \text{ GeV}$	$> 106.2 \text{ GeV}$
m_T^H	$< 60 \text{ GeV}$	—	60–125 GeV
$m_T^{\ell 2}$	—	$> 30 \text{ GeV}$	$> 30 \text{ GeV}$
N_{jet} (b jets)	0	> 0	0

marized in Table 5. They are used to validate the background description and to estimate the number of background events in the signal region (SR). A dedicated $\tau\tau$ CR targets events from the DY process $Z \rightarrow \tau\tau$ with τ leptons decaying leptonically to produce the $e\mu$ final state. Also a top quark CR is defined to enhance events with one or more top quarks decaying to a W boson and bottom quark. Splitting events according to the number of associated jets, separate $\tau\tau$ and top quark CRs are defined for the 0-, 1- and 2-jet SRs. An additional CR with an enhanced contribution from the nonresonant WW background is used in the 2-jet SR. All CRs are used in the final data fit to constrain the DY, top quark, and WW background normalizations.

Additional $\tau\tau$, top quark, and WWCRs are defined requiring a V jet. These CRs are used to validate the background description in the Boosted VH channel. However, they generally do not have a sufficient number of events to significantly constrain the background normalizations in the final fit to the data. As such, we rely on the 2-jet CRs to determine the normalizations to be used in the Boosted VH channel. Agreement between data and the background prediction in the V jet CRs is observed when using normalizations determined in the 2-jet CRs.

7 Background estimation

The nonprompt-lepton backgrounds originating from leptonic decays of heavy quarks, hadrons misidentified as leptons, and electrons from photon conversions are suppressed by identification and isolation requirements imposed on electrons and muons. In this analysis, the nonprompt-lepton background primarily originates from W+jets events and is estimated from data, as described in detail in Ref. [92]. The procedure involves measuring the rate at which a nonprompt

lepton passing a loose selection further passes a tight selection (misidentification rate) and the corresponding rate for a prompt lepton to pass this selection (prompt rate). The misidentification rate is measured in a data sample enriched in multijet events, whereas the prompt rate is measured using a tag-and-probe method [93] in a data sample enriched in DY events. The nonprompt-lepton background estimation is validated with data in a CR enriched with W+jets events, in which a pair of same-sign leptons is required.

The backgrounds from top quark processes and nonresonant WW production are estimated using a combination of MC simulations and the dedicated CRs described in the previous section. The normalisations of these backgrounds are left as free parameters in the fit, keeping different parameters for each jet multiplicity region. The top quark background normalization is measured from the observed data in the top quark enriched CRs. A separate normalization parameter is included for the quark-induced and gluon-induced WW backgrounds. For the 2-jet regions, the WW enriched CR is used to constrain the WW background normalisation parameters. In the 0- and 1-jet channels, these parameters are constrained directly in the signal regions, which span the high $m_{\ell\ell}$ phase space enriched in WW events.

The DY $\rightarrow \tau\tau$ background process is estimated with a data-embedding technique [94]. As for the top quark and WW backgrounds, the DY normalization is left unconstrained in the data fit. The DY $\rightarrow \tau\tau$ enriched CR described in Sect. 6 is used to constrain the free normalization parameters in the 0-, 1-, 2-jet regions. The data-embedded samples cover the events that pass the $e\mu$ triggers, which represent the vast majority of the selected events. The remaining DY $\rightarrow \tau\tau$ events, which enter the analysis through the single-lepton triggers ($\approx 5\%$ of the total), are estimated using MC simulation.

The WZ and $W\gamma^*$ background contributions are simulated as described in Sect. 4, and a data-to-simulation scale factor is derived in a three-lepton CR, as described in Ref. [92]. The contribution of the $W\gamma$ process may also be a background because of photon conversions in the detector material. This process is estimated using MC simulation and validated using data in a CR requiring events with a leading μ and a trailing e with same sign and a separation in $\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2}$ (where ϕ is the azimuthal angle in radians) smaller than 0.5. Triple vector boson production is a minor background in all channels and is estimated using MC simulation.

8 Observables and kinematic discriminants

In this paper, we search for anomalous HVV and Hgg coupling effects by studying:




Fig. 1 Topologies of the Higgs boson production and decay for vector boson fusion $qq' \rightarrow qqH$ (left), $qq' \rightarrow VH$ (center), and gluon fusion with decay $gg \rightarrow H \rightarrow 2\ell 2\nu$ (right). For the electroweak production topologies, the intermediate vector bosons and their decays are shown in green and the $H \rightarrow WW$ decay is marked in red. For the

$gg \rightarrow H \rightarrow 2\ell 2\nu$ topology, the W boson leptonic decays are shown in green. In all cases, the incoming particles are depicted in brown and the angles characterizing kinematic distributions are marked in blue. Five angles fully characterize the orientation of the production and decay chain and are defined in the suitable rest frames

1. the two quark jets from VBF and VH production (HV coupling);
2. the $H \rightarrow WW$ decay products (HV coupling); and
3. the two quark jets from $ggH + 2$ jets production (Hgg coupling).

The VBF, VH, and ggH production and decay topologies relevant for the HVV coupling are illustrated in Fig. 1.

When combined with the momentum transfer of the vector bosons, the five angles illustrated for VBF/VH production provide complete kinematic information for production and decay of the Higgs boson. The illustration for Higgs boson production via ggH in association with two jets is identical to the VBF diagram, except for replacing the intermediate vector bosons by gluons. Full production kinematic information is extracted for VBF, VH, and $ggH + 2$ jets candidate events using discriminants built from the matrix element calculations of the MELA package. The MELA approach is designed to reduce the number of observables to a minimum, while retaining all essential information. To form the production-based MELA kinematic discriminants, we use jets to reconstruct the four-momentum of the associated production particles. The presence of two neutrinos in the final state means it is not possible to reconstruct the four-momentum of all the Higgs boson decay products. Therefore, decay-based kinematic discriminants built from matrix elements are not used in this analysis. Instead, we rely on kinematic variables related to the measured final state of the Higgs boson decay. The strategies used for each of the topologies listed above are now discussed in more detail.

8.1 Kinematic features of two quark jets in VBF and VH channels

Kinematic distributions of associated particles in VBF and VH production are sensitive to the anomalous HVV couplings of the Higgs boson.

As illustrated in Fig. 1, a set of seven observables can be defined for the VBF and VH production topologies: $\Omega = \{\theta_1^{(i)}, \theta_2^{(i)}, \theta^*, \Phi, \Phi_1, q_1^2, q_2^2\}$, with q_1^2 and q_2^2 the squared four-momenta of the vector bosons [22]. Three types of discriminants are defined using the full kinematic description characterized by Ω . The first type of discriminant is designed to separate signal and background Higgs boson production processes:

$$\mathcal{D}_{\text{sig}} = \frac{\mathcal{P}_{\text{sig}}(\Omega)}{\mathcal{P}_{\text{sig}}(\Omega) + \mathcal{P}_{\text{bkg}}(\Omega)}, \quad (19)$$

where the probability density \mathcal{P} for a specific process is calculated from the matrix elements provided by the MELA package. The second type of discriminant separates the anomalous coupling BSM process from that of the SM:

$$\mathcal{D}_{\text{BSM}} = \frac{\mathcal{P}_{\text{BSM}}(\Omega)}{\mathcal{P}_{\text{BSM}}(\Omega) + \mathcal{P}_{\text{SM}}(\Omega)}. \quad (20)$$

Throughout this document the generic BSM label is generally replaced by the specific anomalous coupling state targeted. For the a_3 CP -odd and a_2 CP -even coupling parameters, we use, respectively, \mathcal{D}_{0-} and \mathcal{D}_{0+} , whereas for the Λ_1 coupling parameters we use \mathcal{D}_{Λ_1} and $\mathcal{D}_{\Lambda_1}^{Z\gamma}$. The third type of discriminant isolates the interference contribution:

$$\mathcal{D}_{\text{int}} = \frac{\mathcal{P}_{\text{SM-BSM}}^{\text{int}}(\Omega)}{\mathcal{P}_{\text{SM}}(\Omega) + \mathcal{P}_{\text{BSM}}(\Omega)}, \quad (21)$$

where $\mathcal{P}_{\text{SM-BSM}}^{\text{int}}$ is the interference part of the probability distribution for a process with a mixture of the SM and BSM contributions. The CP label is generally used for the a_3 coupling parameter, as the BSM signal in this case is a pseudoscalar and the interference discriminant is a CP -sensitive observable. The \mathcal{P} values are normalized to give the same integrated cross sections in the relevant phase space of each process. Such normalization leads to a balanced distribution of events in the range between 0 and 1 for \mathcal{D}_{sig} and \mathcal{D}_{BSM} , and between -1 and $+1$ for \mathcal{D}_{int} .

The selected events are split into three main production channels: VBF, Resolved VH, and Boosted VH. In the first two channels, the four-momenta of the two AK4 jets assigned as the associated particles are used in the MELA probability calculation. For the Boosted VH category, we use the four-momentum of the two subjets of the V-tagged AK8 jet. An estimate of the Higgs boson four-momentum is also required for the probability calculation. This can not be measured directly since the final state contains two neutrinos. As such, we construct a proxy Higgs boson four-momentum in the following manner. The p_x and p_y of the dineutrino system are estimated from the \vec{p}_T^{miss} in a given event. The corresponding p_z is then set to equal that of the dilepton system, which is based on the observed correlation between these variables at the generator level for simulated signals. Finally, the mass of the dineutrino system is set equal to the mean value of the generator-level dineutrino mass. The resulting four-momentum can then be combined with that of the measured dilepton system to create a proxy Higgs boson four-momentum. We note that the MELA probability calculation for the production vertices is largely based on the kinematic features of the associated particles, so the reconstruction of the proxy Higgs boson has a relatively small effect on the final discriminants. As an illustrative example of the MELA based discriminants used in this analysis, Fig. 2 shows the D_{0-} discriminant in the VBF and Resolved VH production channels for a number of different signal hypotheses. The discriminants are designed to target the dominant signal production process in a given channel.

In the VBF channel, a D_{VBF} discriminant is constructed, following Eq. (19), where \mathcal{P}_{sig} corresponds to the probability for the VBF production hypothesis, and \mathcal{P}_{bkg} corresponds to that of gluon fusion production in association with two jets. The discriminant is also suitable for separating SM backgrounds from the VBF signal process. In the Resolved and Boosted VH channels, the corresponding discriminants do not give a significant level of separation with respect to ggH production or SM backgrounds. This is due to the relatively tight selection criteria, which limit the phase space to VH-like events. Hence, these discriminants are not included in the VH channels.

The D_{CP} discriminant is sensitive to the sign of the interference between the CP -even SM and CP -odd BSM states. An asymmetry between the number of events detected with positive and negative D_{CP} values is expected for mixed CP states. Therefore, a forward-backward categorization (forward defined as $D_{CP} > 0$ and backward as $D_{CP} < 0$) is used to analyze the CP -odd couplings. Similarly, D_{int} gives sensitivity to the sign of the interference between the SM and a_2 HVV BSM states. A forward-backward D_{int} categorization is also included. The value of D_{int} used to define the categories is chosen to symmetrize the SM Higgs boson expectation. In




Fig. 2 The D_{0-} discriminant in the VBF (upper) and Resolved VH (lower) production channels for a number of VBF (upper) and VH (lower) signal hypotheses. Pure a_1 ($f_{a3} = 0$) and a_3 ($f_{a3} = 1$) HVV signal hypotheses are shown along with two mixed coupling hypotheses ($f_{a3} = 0.005$ and $f_{a3} = 0.01$). All distributions are normalized to unity

the case of the A_1 measurements, the interference discriminants were shown to be highly correlated with the D_{BSM} discriminants and so are not considered.

We now discuss the categorization and construction of the final multidimensional discriminants used for the two HVV coupling approaches defined in Sect. 2. The binning of the final discriminants was optimized to ensure sufficient statistical precision in the predictions of all bins, while retaining

the kinematic information required to discriminate between the SM and anomalous coupling signal hypotheses.

8.1.1 VBF/ VH analysis strategy for Approach 1

In Approach 1, each of the four anomalous HVV coupling parameters (a_2 , a_3 , $\kappa_1/(\Lambda_1)^2$, and $\kappa_2^{Z\gamma}/(\Lambda_1^{Z\gamma})^2$) are analyzed separately. For this purpose, we construct a multidimensional discriminant for each of the four anomalous couplings in the VBF, Resolved VH, and Boosted VH channels.

In the VBF channel, we use two bins of the production discriminant \mathcal{D}_{VBF} , corresponding to low and high purity, using a bin boundary of 0.75. The $m_{\ell\ell}$ variable, which is sensitive to anomalous effects at the $H \rightarrow WW$ decay vertex, is included with two bins in the range 12–76.2 GeV. A bin boundary of 45 GeV is chosen based on the expected signal shape changes induced by anomalous effects. Finally, one of the \mathcal{D}_{BSM} discriminants is included with ten equally sized bins. Depending on the anomalous coupling under study this discriminant may be \mathcal{D}_{0+} , \mathcal{D}_{0-} , \mathcal{D}_{A_1} or $\mathcal{D}_{A_1}^{Z\gamma}$.

For the VH channels, the $m_{\ell\ell}$ and \mathcal{D}_{BSM} observables are used to build 2D kinematic discriminants. The $m_{\ell\ell}$ bins are the same as for the VBF channel. In the Resolved VH channel, we use four \mathcal{D}_{BSM} bins of equal size. For the Boosted VH case, three variable bins with boundaries of 0.6 and 0.8 are used, a large first bin is chosen because relatively little signal is expected at low values of \mathcal{D}_{BSM} . A distinct multidimensional discriminant is constructed for each anomalous coupling hypothesis in the VH channels.

For the a_3 coupling parameter, a forward-backward categorization of events based on \mathcal{D}_{CP} is implemented. In the case of the a_2 coupling parameter, \mathcal{D}_{int} is largely correlated with \mathcal{D}_{0+} in the VH channels. Therefore, a forward-backward \mathcal{D}_{int} categorization is implemented only in the VBF channel. Figures 3, 4 and 5 show the discriminants used in the final fit to the data for the a_2 , a_3 , $\kappa_1/(\Lambda_1)^2$, and $\kappa_2^{Z\gamma}/(\Lambda_1^{Z\gamma})^2$ Approach 1 coupling studies in the VBF and VH channels. A summary of the observables used in the HVV Approach 1 analysis may be found in Table 6.

8.1.2 VBF/ VH analysis strategy for Approach 2

In Approach 2, we use one categorization strategy and build one multidimensional discriminant in each channel to target all the HVV coupling parameters (a_2 , a_3 , $\kappa_1/(\Lambda_1)^2$) simultaneously. In the VBF channel, the \mathcal{D}_{CP} and \mathcal{D}_{int} discriminants are used to create four interference categories. Both \mathcal{D}_{VBF} and $m_{\ell\ell}$ are used as for Approach 1. All three \mathcal{D}_{BSM} discriminants that target the a_2 , a_3 and $\kappa_1/(\Lambda_1)^2$ coupling parameters are included. However, the number of bins we implement is limited by the number of simulated events. Also the \mathcal{D}_{BSM} discriminants are significantly correlated and so have

similar performance for all couplings. Therefore, we use the CP -odd discriminant \mathcal{D}_{0-} and just one of the CP -even discriminants, \mathcal{D}_{0+} , both with three bins and bin boundaries of 0.1 and 0.9. A dedicated rebinning strategy is applied to the $[\mathcal{D}_{0-}, \mathcal{D}_{0+}]$ distribution merging bins dominated by the SM Higgs boson prediction or with low precision in the background prediction. In the VH channels, just two categories using \mathcal{D}_{CP} are defined and the discriminant is built using $m_{\ell\ell}$ as for Approach 1. Again, both \mathcal{D}_{0-} and \mathcal{D}_{0+} are chosen for the final discriminant. For the Resolved VH channel, we use three bins with boundaries of 0.25 and 0.75, whereas for the Boosted VH case we use two bins with a boundary of 0.8. The same rebinning strategy described for the VBF channel is applied to both Resolved and Boosted VH multidimensional discriminants. Table 6 includes a summary of the observables used in the HVV Approach 2 analysis.

8.2 Kinematic features of $H \rightarrow WW$ decay products in 0- and 1-jet ggH channels

Similar to the SM $H \rightarrow WW$ analysis [25], we use $m_{\ell\ell}$ and m_T to build 2D discriminants in the 0- and 1-jet ggH channels. The distributions have nine bins for $m_{\ell\ell}$ in the range 12–200 GeV and six bins for m_T in the range 60–125 GeV. The bin widths vary and are optimized to achieve good separation between the SM Higgs boson signal and backgrounds, as well as between the different anomalous coupling signal hypotheses. In particular, a finer binning with respect to the SM $H \rightarrow WW$ analysis is implemented in regions where anomalous effects are most significant. Figure 6 shows the $[m_T, m_{\ell\ell}]$ distributions in the 0- and 1-jet ggH channels. The same $[m_T, m_{\ell\ell}]$ discriminant is used to study all HVV anomalous couplings for both Approach 1 and 2.

8.3 Kinematic features of two quark jets in 2-jet ggH channel

For the Hgg coupling, we adopt a similar approach to the VBF CP study, where the CP -odd a_3 HVV coupling parameter is included. In this case, the optimal observables are $\mathcal{D}_{0-}^{\text{ggH}}$ and $\mathcal{D}_{CP}^{\text{ggH}}$, targeting the CP -odd a_3 Hgg coupling parameter. A forward-backward categorization is implemented using $\mathcal{D}_{CP}^{\text{ggH}}$, and the \mathcal{D}_{VBF} and $\mathcal{D}_{0-}^{\text{ggH}}$ observables are used to build 2D discriminants. The $m_{\ell\ell}$ variable is not considered in this case because it is not sensitive to anomalous Hgg effects. For \mathcal{D}_{VBF} , the bin boundary is relaxed to 0.5 to ensure sufficient ggH events are accepted in the more VBF-like bin. For \mathcal{D}_{0-} , eight (five) bins are used in the more (less) VBF-like bin with larger bin sizes at the extremes of the distribution to ensure sufficient precision in the background and signal predictions. The 0- and 1-jet channels discussed previously are also included in this study to constrain the ggH signal




Fig. 3 Observed and predicted distributions after fitting the data for $[\mathcal{D}_{\text{VBF}}, m_{ll}, \mathcal{D}_{0+}]$ in the VBF channel (upper), and for $[m_{ll}, \mathcal{D}_{0+}]$ in the Resolved VH (lower left) and Boosted VH (lower right) channels. For the VBF channel, the $\mathcal{D}_{\text{int}} < 0.4$ (left) and $\mathcal{D}_{\text{int}} > 0.4$ (right) categories are shown. The predicted Higgs boson signal is shown stacked on top of the background distributions. For the fit, the a_1 and a_2 HVV coupling contributions are included. The corresponding pure a_1 ($f_{a2} = 0$)

and a_2 ($f_{a2} = 1$) signal hypotheses are also shown superimposed, their yields correspond to the predicted number of SM signal events scaled by an arbitrary factor to improve visibility. The uncertainty band corresponds to the total systematic uncertainty. The lower panel in each figure shows the ratio of the number of events observed to the total prediction

strength. The $[\mathcal{D}_{\text{VBF}}, \mathcal{D}_{0-}^{\text{ggH}}]$ distributions used to analyze the Hgg anomalous coupling in the 2-jet ggH channel are shown in Fig. 7. A summary of the observables used in the Hgg analysis is given in Table 6.

9 Systematic uncertainties

The signal extraction is performed using binned templates to describe the various signal and background processes. Sys-

tematic uncertainties that change the normalization or shape of the templates are included. All the uncertainties are modeled as nuisance parameters that are profiled in the maximum likelihood fit described in Sect. 10. The systematic uncertainties arise from both experimental or theoretical sources.

9.1 Experimental uncertainties

The following experimental systematic uncertainties are included in the final fit to data:




Fig. 4 Observed and predicted distributions after fitting the data for $[\mathcal{D}_{\text{VBF}}, m_{\ell\ell}, \mathcal{D}_{0-}]$ in the VBF channel (upper), and for $[m_{\ell\ell}, \mathcal{D}_{0-}]$ in the Resolved VH (middle) and Boosted VH (lower) channels. For each

channel, the $\mathcal{D}_{CP} < 0$ (left) and $\mathcal{D}_{CP} > 0$ (right) categories are shown. For the fit, the a_1 and a_3 HVV coupling contributions are included. More details are given in the caption of Fig. 3




Fig. 5 Observed and predicted distributions after fitting the data for $[D_{VBF}, m_\ell, D_{\Lambda_1}]$ (upper left) and $[D_{VBF}, m_\ell, D_{\Lambda_1}^{Z\gamma}]$ (upper right) in the VBF channel, and for $[m_\ell, D_{\Lambda_1}]$ (left) and $[m_\ell, D_{\Lambda_1}^{Z\gamma}]$ (right) in the Resolved VH (middle) and Boosted VH (lower) channels. For the

fits, the a_1 and $\kappa_1/(\Lambda_1)^2$ (left) or a_1 and $\kappa_2^{Z\gamma}/(\Lambda_1^{Z\gamma})^2$ (right) HVV coupling contributions are included. More details are given in the caption of Fig. 3

Table 6 The kinematic observables used for the interference based categorization and for the final discriminants used in the fits to data to study the HVV and Hgg couplings. For each of the anomalous HVV couplings in Approach 1, we have a dedicated analysis in the VBF and VH channels. In Approach 2, we use one analysis to target all anomalous HVV couplings simultaneously

Analysis	Channel	Categorization	Final discriminant
HVV	VBF (a_2)	\mathcal{D}_{int}	$[\mathcal{D}_{\text{VBF}}, m_{\ell\ell}, \mathcal{D}_{0+}]$
Approach 1	VBF (a_3)	\mathcal{D}_{CP}	$[\mathcal{D}_{\text{VBF}}, m_{\ell\ell}, \mathcal{D}_{0-}]$
	VBF (κ_1)	—	$[\mathcal{D}_{\text{VBF}}, m_{\ell\ell}, \mathcal{D}_{A1}]$
	VBF ($\kappa_2^{Z\gamma}$)	—	$[\mathcal{D}_{\text{VBF}}, m_{\ell\ell}, \mathcal{D}_{A1}^{Z\gamma}]$
	VH (a_2)	—	$[m_{\ell\ell}, \mathcal{D}_{0+}]$
	VH (a_3)	\mathcal{D}_{CP}	$[m_{\ell\ell}, \mathcal{D}_{0-}]$
	VH (κ_1)	—	$[m_{\ell\ell}, \mathcal{D}_{A1}]$
	VH ($\kappa_2^{Z\gamma}$)	—	$[m_{\ell\ell}, \mathcal{D}_{A1}^{Z\gamma}]$
	0- and 1-jet ggH	—	$[m_T, m_{\ell\ell}]$
HVV	VBF	$\mathcal{D}_{CP}, \mathcal{D}_{\text{int}}$	$[\mathcal{D}_{\text{VBF}}, m_{\ell\ell}, \mathcal{D}_{0-}, \mathcal{D}_{0+}]$
Approach 2	VH	\mathcal{D}_{CP}	$[m_{\ell\ell}, \mathcal{D}_{0-}, \mathcal{D}_{0+}]$
	0- and 1-jet ggH	—	$[m_T, m_{\ell\ell}]$
Hgg	2-jet ggH	$\mathcal{D}_{CP}^{\text{ggH}}$	$[\mathcal{D}_{\text{VBF}}, \mathcal{D}_{0-}^{\text{ggH}}]$
	0- and 1-jet ggH	—	$[m_T, m_{\ell\ell}]$

- The total uncertainty associated with the measurement of the integrated luminosity for 2016, 2017, and 2018 is 1.2% [44], 2.3% [45], and 2.5% [46], respectively. This uncertainty is partially correlated among the three data sets, resulting in an overall uncertainty of 1.6%.
- The systematic uncertainty in the trigger efficiency is determined by varying the tag lepton selection criteria and the Z boson mass window used in the tag-and-probe method. It affects both the normalization and the shape of the signal and background distributions, and is kept uncorrelated among data sets. The total normalization uncertainty is less than 1%.
- The tag-and-probe method is also used to determine the lepton identification and isolation efficiency. Corrections are applied to account for any discrepancy in the efficiencies measured in data and simulation. The corresponding systematic uncertainty is about 1% for electrons and 2% for muons.
- The uncertainties in the determination of the lepton momentum scale mainly arise from the limited data sample used for their estimation. The impact on the normalization of the signal and background templates ranges between 0.6–1.0% for the electron momentum scale and is about 0.2% for the muon momentum scale. They are treated as uncorrelated among the three data-taking years.
- The jet energy scale uncertainty is modeled by implementing eleven independent nuisance parameters corresponding to different jet energy correction sources, six of which are correlated among the three data sets. Their effects vary in the range of 1–10%, mainly depending on the jet multiplicity in the analysis phase space. Another source of uncertainty arises from the jet energy resolution smearing applied to simulated samples to match the

p_T resolution measured in data. The effect varies in a range of 1–5%, depending on the jet multiplicity and is uncorrelated among the data sets. These uncertainties are included for both AK4 and AK8 jets. In addition, the m_j scale and resolution, and V tagging corrections with their corresponding uncertainties are included for V-tagged AK8 jets. These variables are calibrated in a top quark–antiquark sample enriched in hadronically decaying W bosons [95].

- The effects of the unclustered energy scale, jet energy scale, and lepton p_T scales are included for the calculation of the missing transverse momentum. The resulting normalization systematic uncertainty is 1–10% and is treated as uncorrelated among the years.
- Both the normalization and shape of the signal and background templates are affected by the jet pileup identification uncertainty. The effect is below 1%.
- The uncertainty associated with the b tagging efficiency is modeled by seventeen nuisance parameters out of which five are of a theoretical origin and are correlated among the three data sets. The remaining set of four parameters per data set are treated as uncorrelated as they arise from the statistical accuracy of the efficiency measurement [90].
- Estimation of the nonprompt-lepton background is affected by the limited size of the data sets used for the misidentification rate measurements. It is also affected by the difference in the flavor composition of jets misidentified as leptons between the misidentification rate measurement region (enriched in multijet events) and the signal phase space. The effects on the nonprompt-lepton background estimation range between a few percent to about 10% depending on the SR and are treated as nui-




Fig. 6 Observed and predicted distributions after fitting the data for $[m_T, m_{\ell\ell}]$ in the 0- (upper) and 1-jet (lower) ggH channels. For the fit, the a_1 and a_3 HH coupling contributions are included. More details are given in the caption of Fig. 3

sance parameters uncorrelated between electrons and muons and among the three data sets. A normalization uncertainty of 30% [92] is assigned to fully cover for any discrepancies with respect to data in a W+jets CR and is treated as uncorrelated among data sets.

- The statistical uncertainties due to the limited number of simulated events are also included for all bins of the background distributions used to extract the results [96].




Fig. 7 Observed and predicted distributions after fitting the data for $[D_{VBF}, D_{0-}^{ggH}]$ in the 2-jet ggH channel. Both the $D_{CP}^{ggH} < 0$ (upper) and $D_{CP}^{ggH} > 0$ (lower) categories are shown. In this case, the VBF and ggH signals are shown separately. For the fit, the a_2^{gg} and a_3^{gg} coupling contributions are included. The corresponding pure a_2^{gg} ($f_{a_3}^{ggH} = 0$) and a_3^{gg} ($f_{a_3}^{ggH} = 1$) signal hypotheses are also shown superimposed, their yields correspond to the predicted number of SM signal events. More details are given in the caption of Fig. 3

9.2 Theoretical uncertainties

Multiple theoretical uncertainties are considered and are correlated among data sets, unless stated otherwise:

- The uncertainties related to the choice of PDF and α_S have a minor effect on the shape of the distributions. Therefore, only normalization effects related to the event acceptance and to the cross section are included. However, these uncertainties are not considered for the backgrounds that have their normalization constrained through data in dedicated CRs. For the Higgs boson signal processes, these uncertainties are calculated by the LHC Higgs cross section working group [19].
- The theoretical uncertainties arising from missing higher-order corrections in the cross section calculations are also included. Background simulations are reweighted to the alternative scenarios corresponding to renormalization μ_R and factorization μ_F scales varied by factors 0.5 or 2 and the envelopes of the varied templates are taken as the one standard deviations. For background processes that have their normalization constrained through data in dedicated CRs, we consider only the shape effect of the uncertainties coming from the missing higher-order corrections. The WWnonresonant background has the uncertainties derived by varying μ_R , μ_F , and the resummation scale. For the ggH and VBF signal processes, the effects of the missing higher-order corrections on the overall cross section are decoupled into multiple sources according to the recipes described in Ref. [19].
- The uncertainty due to the pileup modeling was included for the main simulated background processes (DY, WW, top quark) as well as the ggH and VBF signals. The effect is determined by varying the total inelastic pp cross section (69.2 mb [97, 98]) within the assigned 5% uncertainty.
- The PS modeling mainly affects the jet multiplicity, causing migration of events between categories that results in template shape changes. Associated uncertainties are evaluated by reweighting events with varied PS weights computed with PYTHIA 8.212. The effect on the signal strength is found to be below 1%.
- Uncertainties associated with UE modeling are evaluated by varying the UE tune parameters used in the MC sample generation. Systematic uncertainties are correlated between the 2017 and 2018 data sets since they share the same UE tunes, whereas for 2016 the uncertainty is considered uncorrelated. The UE uncertainty has a minimal effect on the template shapes and affects the normalization by about 1.5%.
- A 15% uncertainty is applied to the relative fraction of the gg-induced component in nonresonant WW production [99]. The relative fraction between single top quark and $t\bar{t}$ processes is assigned a systematic uncertainty of 8% [100]. Additional process-specific (DY, VZ, $V\gamma$, $V\gamma^*$) uncertainties, related to corrections to account for possible discrepancies between data and simulation, are assigned and are correlated among data sets.

10 Results

The optimization and validation of the analysis were performed using simulation and data in CRs. The data in the SRs were examined once all details of the analysis were finalized. For the final results, we perform a binned maximum likelihood fit to the data combining all channels and data-taking periods. The statistical approach was developed by the ATLAS and CMS Collaborations in the context of the LHC Higgs Combination Group [101]. The likelihood function is defined for candidate events as:

$$\begin{aligned} \mathcal{L}(\text{data} | \mu_{\text{ggH}}, \mu_{\text{EW}}, f_{ai}, \theta) \\ = \prod_j \text{Poisson}(n_j | s_j(\mu_{\text{ggH}}, \mu_{\text{EW}}, f_{ai}, \theta) + b_j(\theta)) p(\tilde{\theta} | \theta), \end{aligned} \quad (22)$$

where j runs over all bins and n_j is the observed number of data events in each bin. Total signal and background expectations in each bin are represented by s_j and b_j , respectively. The individual signal and background processes considered in each category are described using binned templates of multidimensional discriminants as described in Sect. 8. Each signal process is parametrized as a linear combination of terms originating from the SM, and anomalous couplings and their interference. The signal expectation depends on the parameters μ_{ggH} , μ_{EW} , and f_{ai} , and is constrained by the data fit. Both the signal and background expectations are functions of θ , which represents the full set of nuisance parameters corresponding to the systematic uncertainties. The CRs described in Sect. 6 are included in the fit in the form of single bins, representing the number of events in each CR.

The μ_{ggH} and μ_{EW} parameters correspond to the Higgs boson signal strength modifiers for the ggH and VBF/VH signals, respectively. Signal yields for the VBF and VH processes are related to each other because the same HVV couplings enter both in production and decay of the Higgs boson. The ggH signal is initiated predominantly by the top fermion couplings and is unrelated to the VBF and VH production mechanisms. As the signal strength modifiers are free parameters in the fit, the overall signal event yield is not used to discriminate between alternative signal hypotheses. The f_{ai} parameter corresponds to the anomalous coupling cross section fraction and determines the shape of the signal expectation. The cross section fraction for the SM coupling is simply taken as $1 - |f_{a1}|$. In Approach 1, the SM and just one anomalous HVV coupling are included, and each f_{ai} is thus studied independently. Depending on the particular anomalous coupling under investigation, f_{ai} may represent f_{a2} , f_{a3} , f_{A1} , or $f_{A1}^{Z\gamma}$. For Approach 2, the SM and three anomalous HVV couplings are included. In this case, f_{ai} represents f_{a2} , f_{a3} and f_{A1} , which are studied simultaneously. It is explicitly required that $|f_{a2}| + |f_{a3}| + |f_{A1}| \leq 1$ to avoid probing an




Fig. 8 Expected (dashed) and observed (solid) profiled likelihood on f_{a2} (upper left), f_{A1} (upper right), f_{a3} (lower left), and $f_{A1}^{Z\gamma}$ (lower right) using Approach 1. In each case, the signal strength modifiers are

treated as free parameters. The dashed horizontal lines show the 68 and 95% CL regions. Axis scales are varied for f_{a2} and f_{A1} to improve the visibility of important features

unphysical parameter space. Finally, there is just one anomalous coupling corresponding to f_{a3}^{ggH} to consider for the Hgg vertex. For this study, we also include the effect of the CP -odd HVV anomalous coupling on the VBF process. This is achieved by including f_{a3} as a free parameter in the fit. The $p(\tilde{\theta}|\theta)$ are the probability density functions (PDFs) for the

observed values of the nuisance parameters, $\tilde{\theta}$, obtained from calibration measurements. The systematic uncertainties that affect only the normalizations of the signal and background processes are treated as PDFs following a log-normal distribution, whereas shape-altering systematic uncertainties are treated as Gaussian PDFs [101].




Fig. 9 Expected (dashed) and observed (solid) profiled likelihood on f_{a2} (upper left), f_{A1} (upper right) and f_{a3} (bottom) using Approach 2. The other two anomalous coupling cross section fractions are either fixed to zero (blue) or left floating in the fit (red). In each case, the signal

strength modifiers are treated as free parameters. The dashed horizontal lines show the 68 and 95% CL regions. Axis scales are varied for f_{a2} and f_{A1} to improve the visibility of important features

Additional interpretations in terms of the SMEFT Higgs and Warsaw basis coupling parameters are also considered using Eqs. (7–10) and Eqs. (11–14), respectively. In each case, four independent couplings are studied simultaneously and the effect of the couplings on the total width of the Higgs boson is taken into account. For the f_{ai} measurements, this effect is absorbed by the signal strength modifiers. A parameterization of the partial widths of the main Higgs boson decay

modes as a function of the couplings is used to determine the effect on the Higgs boson width [24, 28].

The likelihood is maximized with respect to the signal modifier parameters and with respect to the nuisance parameters. Confidence level (CL) intervals are determined from profile likelihood scans of the respective parameters. The allowed 68% and 95% CL intervals are defined using the set of parameter values at which the profile likelihood function

Table 7 Summary of constraints on the anomalous HVV and Hgg coupling parameters with the best fit values and allowed 68 and 95% CL (in square brackets) intervals. For Approach 1, each f_{ai} is studied inde-

pendently. For Approach 2, each f_{ai} is shown separately with the other two cross section fractions either fixed to zero or left floating in the fit. In each case, the signal strength modifiers are treated as free parameters

Analysis	f_{ai}		Observed ($\times 10^{-3}$)	Expected ($\times 10^{-3}$)
HVV	f_{a2}	Best fit	0.5	0.0
Approach 1		68% CL	[−0.8, 3.5]	[−1.4, 1.3]
		95% CL	[−5.7, 12.0]	[−5.2, 6.1]
	f_{a3}	Best fit	0.9	0.0
		68% CL	[−2.7, 4.1]	[−0.7, 0.7]
		95% CL	[−553.0, 561.0]	[−2.8, 2.9]
	f_{A1}	Best fit	−0.2	0.0
		68% CL	[−0.5, 0.0]	[−0.2, 0.5]
		95% CL	[−1.4, 0.7]	[−0.6, 1.4]
	$f_{A1}^{Z\gamma}$	Best fit	3.0	0.0
		68% CL	[−11.0, 9.1]	[−5.0, 3.8]
		95% CL	[−55.0, 42.0]	[−14.0, 11.0]
HVV	f_{a2}	Best fit	38.0	0.0
Approach 2		68% CL	[−112.2, 129.3]	[−30.9, 37.5]
(Fix others)		95% CL	[−376.6, 430.0] ∪ [−989.2, −826.3]	[−126.1, 136.8]
	f_{a3}	Best fit	0.8	0.0
		68% CL	[−0.8, 3.5]	[−0.8, 1.1]
		95% CL	[−7.6, 58.8]	[−3.4, 4.3]
	f_{A1}	Best fit	−0.15	0.0
		68% CL	[−1.21, 0.16]	[−0.4, 0.4]
		95% CL	[−19.5, 118.5] ∪ [909.9, 964.1]	[−1.7, 18.9]
HVV	f_{a2}	Best fit	−1.0	0.0
Approach 2		68% CL	[−104.1, 139.9]	[−31.1, 39.8]
(Float others)		95% CL	[−986.4, 981.2]	[−127.5, 148.7]
	f_{a3}	Best fit	0.34	0.0
		68% CL	[−0.69, 3.4]	[−1.0, 1.2]
		95% CL	[−201.3, 361.5]	[−4.3, 5.3]
	f_{A1}	Best fit	−0.1	0.0
		68% CL	[−1.08, 3.78] ∪ [7.2, 20.7]	[−0.4, 0.9]
		95% CL	[−994.8, 993.9]	[−1.9, 21.4]
Hgg	f_{a3}^{ggH}	Best fit	−34	0
		68% CL	[−721, 383]	[−1000, 1000]
		95% CL	[−1000, 1000]	[−1000, 1000]

$-2\Delta \ln \mathcal{L} = 1.00$ and 3.84 [102], respectively, for which exact coverage is expected in the asymptotic limit [103]. The likelihood value at a given f_{ai} is determined by the shape of the signal hypothesis and the relative signal event yields between categories. Expected results are obtained using the Asimov data set [104] constructed using the SM values of the signal modifier parameters.

For Approach 1, where we assume $a_i^{\text{ZZ}} = a_i^{\text{WW}}$, the expected and observed f_{a2} , f_{a3} , f_{A1} , and $f_{A1}^{Z\gamma}$ likelihood

scans are shown in Fig. 8. Significant interference effects for negative values of f_{a2} , around -0.25 , and positive values of f_{A1} , around 0.5 , are evident. Relatively large changes in the signal shape with respect to the SM are predicted at these values. Also evident are narrow minima around $f_{ai} = 0$. The anomalous coupling terms in Eq. (1) have a q_i^2 dependence, which can be larger at the VBF/VH production vertex than at the Higgs decay vertex. This causes the cross section and the shape of the VBF/VH signal hypothesis to change rapidly




Fig. 10 The observed correlation coefficients between HVV anomalous coupling cross section fractions and signal strength modifiers (left) and between SMEFT Higgs basis coupling parameters (right)

with f_{ai} . For $f_{A1}^{Z\gamma}$, there are no anomalous effects at the Higgs decay vertex and so the only structure present is the narrow minimum related to the VBF/VH production vertex. The axis scales are varied to improve the visibility of important features for f_{a2} and f_{A1} . For Approach 2, where the SU(2) \times U(1) coupling relationships from Eqs. (2–6) are adopted, the expected and observed f_{a2} , f_{a3} and f_{A1} likelihood scans are shown in Fig. 9. The results are shown for each f_{ai} separately with the other two f_{ai} either fixed to zero or left floating in the fit. The measured values of the signal strength parameters correspond to $\mu_{\text{EW}} = 0.9^{+0.19}_{-0.24}$ and $\mu_{\text{ggH}} = 0.9^{+0.38}_{-0.20}$ when all parameters float simultaneously. It is notable that the observed $-2\Delta \ln \mathcal{L}$ profile values are generally lower than expected. This is consistent with a downward statistical fluctuation in the number of VBF and VH events. The lowest μ_{EW} value measured is 0.82 for the Approach 1 f_{a3} fit which can be compared with the highest value of 0.97 for the corresponding f_{A1} fit. In each case, the uncertainty in μ_{EW} is about 20% and as such all fitted values are consistent with both the SM and each other. More generally, all anomalous HVV coupling parameter measurements are consistent with the expectations for the SM Higgs boson. The p-value compatibility of the full Approach 2 fit, where all signal parameters float simultaneously, with the SM is 91%. A summary of constraints on the anomalous HVV coupling parameters with the best fit values and allowed 68% and 95% CL intervals are shown in Table 7. The most stringent constraints on the HVV anomalous coupling cross section fractions are at the per mille level. Some constraints are less stringent than

expected due to the fitted values of μ_{EW} being lower than the SM expectation. The observed correlation coefficients between HVV anomalous coupling cross section fractions and signal strength modifiers are displayed in Fig. 10.

For the SMEFT Higgs basis interpretation, the expected and observed constraints on the δc_z , c_{zD} , c_{zz} , and \tilde{c}_{zz} coupling parameters are shown in Fig. 11. Table 8 presents a summary of the constraints on the couplings whereas Fig. 10 reports the observed correlation coefficients between them. For the Warsaw basis interpretation, the expected and observed constraints on the c_{\square} , c_{HD} , c_{HW} , c_{HWB} , c_{HB} , $c_{\text{H}\tilde{\text{W}}}$, $c_{\text{H}\tilde{\text{W}}\text{B}}$, and $c_{\text{H}\tilde{\text{B}}}$ coupling parameters are presented in Table 9. To cover all the Warsaw basis coupling parameters, three independent fits to the data were performed with a different choice of four independent couplings in each. A summary of the constraints on the SMEFT Higgs and Warsaw basis coupling parameters is presented in Fig. 12.

Finally, the expected and observed f_{a3}^{ggH} likelihood scans are shown in Fig. 13. The result is consistent with the expectation for a SM Higgs boson. Excluding the effect of the CP -odd HVV anomalous coupling, by fixing f_{a3} to zero, has a negligible effect. For $|f_{a3}^{\text{ggH}}|$ approaching unity, the observed $-2\Delta \ln \mathcal{L}$ profile values are larger than expected. This is consistent with downward statistical fluctuations in the data for a couple of bins where sensitivity to the a_3 Hgg coupling contribution is enhanced (Fig. 7 upper). The constraint on the anomalous Hgg coupling parameter with the best fit value and allowed 68% CL interval is shown in Table 7.




Fig. 11 Expected (dashed) and observed (solid) profiled likelihood on the δc_z (upper left), $c_{z\square}$ (upper right), c_{zz} (lower left), and \tilde{c}_{zz} (lower right) couplings of the SMEFT Higgs basis. All four couplings are studied simultaneously. The dashed horizontal lines show the 68 and 95% CL regions

Table 8 Summary of constraints on the SMEFT Higgs basis coupling parameters with the best fit values and 68% CL uncertainties. All four couplings are studied simultaneously

Coupling	Observed	Expected
δc_z	$-0.06^{+0.09}_{-0.16}$	$0.00^{+0.08}_{-0.10}$
$c_{z\square}$	$0.01^{+0.02}_{-0.06}$	$0.00^{+0.02}_{-0.02}$
c_{zz}	$0.03^{+0.30}_{-0.52}$	$0.00^{+0.23}_{-0.29}$
\tilde{c}_{zz}	$-0.17^{+0.42}_{-0.30}$	$0.00^{+0.29}_{-0.32}$

Table 9 Summary of constraints on the SMEFT Warsaw basis coupling parameters with the best fit values and 68% CL uncertainties. Only one of c_{HW} , c_{HWB} , and c_{HB} is independent, the same is also true for $c_{H\tilde{W}}$, $c_{H\tilde{W}B}$, and $c_{H\tilde{B}}$. Three independent fits to the data were performed with a different choice of four independent couplings in each

Coupling	Observed	Expected
$c_{H\square}$	$-0.76^{+1.43}_{-3.43}$	$0.00^{+1.37}_{-1.84}$
c_{HD}	$-0.12^{+0.93}_{-0.32}$	$0.00^{+0.43}_{-0.30}$
c_{HW}	$0.08^{+0.43}_{-0.87}$	$0.00^{+0.37}_{-0.48}$
c_{HWB}	$0.17^{+0.88}_{-1.79}$	$0.00^{+0.77}_{-0.96}$
c_{HB}	$0.03^{+0.13}_{-0.26}$	$0.00^{+0.11}_{-0.14}$
$c_{H\tilde{W}}$	$-0.26^{+0.67}_{-0.50}$	$0.00^{+0.48}_{-0.52}$
$c_{H\tilde{W}B}$	$-0.54^{+1.37}_{-1.03}$	$0.00^{+0.99}_{-1.07}$
$c_{H\tilde{B}}$	$-0.08^{+0.20}_{-0.15}$	$0.00^{+0.15}_{-0.16}$

11 Summary

This paper presents a study of the anomalous couplings of the Higgs boson (H) with vector bosons, including CP violating effects, using its associated production with hadronic jets in gluon fusion, electroweak vector boson fusion, and associated production with a W or Z boson, and its subsequent decay to a pair of W bosons. The results are based on the proton–proton collision data set collected by the CMS detector at the LHC during 2016–2018, corresponding to an integrated luminosity of 138 fb^{-1} at a center-of-mass energy of 13 TeV . The analysis targets the different-flavor dilepton ($e\mu$) final state, with kinematic information from associated jets combined using matrix element techniques to increase sensitivity to anomalous effects at the production vertex. Dedicated Monte Carlo simulation and matrix element reweighting provide modeling of all kinematic features in the production and decay of the Higgs boson with full simulation of detector effects. A simultaneous measurement of four Higgs boson

couplings to electroweak vector bosons has been performed in the framework of a standard model effective field theory. All measurements are consistent with the expectations for the standard model Higgs boson and constraints are set on the fractional contribution of the anomalous couplings to the Higgs boson cross section. The most stringent constraints on the HVV anomalous coupling cross section fractions are at the per mille level. These results are in agreement with those obtained in the $H \rightarrow ZZ$ and $H \rightarrow \tau\tau$ channels, and also significantly surpass those of the previous $H \rightarrow WW$ anomalous coupling analysis from the CMS experiment in both scope and precision.





Fig. 12 Summary of constraints on the SMEFT Higgs (upper) and Warsaw (lower) basis coupling parameters with the best fit values and 68% CL uncertainties. For the Warsaw basis, only one of c_{HW} , c_{HWB} , and c_{HB} is independent, the same is also true for $c_{H\tilde{W}}$, $c_{H\tilde{W}B}$, and $c_{H\tilde{B}}$




Fig. 13 Expected (dashed) and observed (solid) profiled likelihood on f_{a3}^{ggH} . The signal strength modifiers and the CP -odd HVV anomalous coupling cross section fraction are treated as free parameters. The crossing of the observed likelihood with the dashed horizontal line shows the observed 68% CL region

Acknowledgements We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia); BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMELT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA). Rachada-pisek Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and

FWO (Belgium) under the “Excellence of Science – EOS” – be.h project n. 30820817; the Beijing Municipal Science and Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), under Germany’s Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources and Institutional Development, Research and Innovation, grant B37G660013 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

Data Availability Statement Data cannot be made available for reasons disclosed in the data availability statement. [Author’s comment: Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the CMS policy as stated in the “[CMS data preservation, re-use and open access policy](#).”]

Code Availability Statement This manuscript has no associated code/software. [Author’s comment: There is no code availability statement.]

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³.

References

- ATLAS Collaboration, Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett. B* **716**, 1 (2012). <https://doi.org/10.1016/j.physletb.2012.08.020>. <arXiv:1207.7214>
- C.M.S. Collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett. B* **716**, 30 (2012). <https://doi.org/10.1016/j.physletb.2012.08.021>. <arXiv:1207.7235>
- C.M.S. Collaboration, Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV. *JHEP* **06**, 081 (2013). [https://doi.org/10.1007/JHEP06\(2013\)081](https://doi.org/10.1007/JHEP06(2013)081). <arXiv:1303.4571>
- C.M.S. Collaboration, On the mass and spin-parity of the Higgs boson candidate via its decays to Z boson pairs. *Phys. Rev. Lett.* **110**, 081803 (2013). <https://doi.org/10.1103/PhysRevLett.110.081803>. <arXiv:1212.6639>
- C.M.S. Collaboration, Measurement of the properties of a Higgs boson in the four-lepton final state. *Phys. Rev. D* **89**, 092007 (2014). <https://doi.org/10.1103/PhysRevD.89.092007>. <arXiv:1312.5353>
- C.M.S. Collaboration, Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at 7 and 8 TeV. *Phys. Rev. D* **92**, 012004 (2015). <https://doi.org/10.1103/PhysRevD.92.012004>. <arXiv:1411.3441>
- CMS Collaboration, Limits on the Higgs boson lifetime and width from its decay to four charged leptons. *Phys. Rev. D* **92**, 072010 (2015). <https://doi.org/10.1103/PhysRevD.92.072010>. <arXiv:1507.06656>
- CMS Collaboration, Combined search for anomalous pseudoscalar HVV couplings in VH production and H \rightarrow VV decay. *Phys. Lett. B* **759**, 672 (2016). <https://doi.org/10.1016/j.physletb.2016.06.004>. <arXiv:1602.04305>
- CMS Collaboration, Constraints on anomalous Higgs boson couplings using production and decay information in the four-lepton final state. *Phys. Lett. B* **775**, 1 (2017). <https://doi.org/10.1016/j.physletb.2017.10.021>. <arXiv:1707.00541>
- CMS Collaboration, Measurements of the Higgs boson width and anomalous HVV couplings from on-shell and off-shell production in the four-lepton final state. *Phys. Rev. D* **99**, 112003 (2019). <https://doi.org/10.1103/PhysRevD.99.112003>. <arXiv:1901.00174>
- CMS Collaboration, Constraints on anomalous HVV couplings from the production of Higgs bosons decaying to τ lepton pairs. *Phys. Rev. D* **100**, 112002 (2019). <https://doi.org/10.1103/PhysRevD.100.112002>. <arXiv:1903.06973>
- ATLAS Collaboration, Evidence for the spin-0 nature of the Higgs boson using ATLAS data. *Phys. Lett. B* **726**, 120 (2013). <https://doi.org/10.1016/j.physletb.2013.08.026>. <arXiv:1307.1432>
- ATLAS Collaboration, Study of the spin and parity of the Higgs boson in diboson decays with the ATLAS detector. *Eur. Phys. J. C* **75**, 476 (2015). <https://doi.org/10.1140/epjc/s10052-015-3685-1>. <arXiv:1506.05669>
- ATLAS Collaboration, Test of CP invariance in vector-boson fusion production of the Higgs boson using the Optimal Observable method in the ditau decay channel with the ATLAS detector. *Eur. Phys. J. C* **76**, 658 (2016). <https://doi.org/10.1140/epjc/s10052-016-4499-5>. <arXiv:1602.04516>
- ATLAS Collaboration, Measurement of inclusive and differential cross sections in the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *JHEP* **10**, 132 (2017). [https://doi.org/10.1007/JHEP10\(2017\)132](https://doi.org/10.1007/JHEP10(2017)132). <arXiv:1708.02810>
- ATLAS Collaboration, Measurement of the Higgs boson coupling properties in the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channel at $\sqrt{s} = 13$ TeV with the ATLAS detector. *JHEP* **03**, 095 (2018). [https://doi.org/10.1007/JHEP03\(2018\)095](https://doi.org/10.1007/JHEP03(2018)095). <arXiv:1712.02304>
- ATLAS Collaboration, Measurements of Higgs boson properties in the diphoton decay channel with 36 fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Rev. D* **98**, 052005 (2018). <https://doi.org/10.1103/PhysRevD.98.052005>. <arXiv:1802.04146>
- ATLAS Collaboration, Test of CP invariance in vector-boson fusion production of the Higgs boson in the $H \rightarrow \tau\tau$ channel in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett. B* **805**, 135426 (2020). <https://doi.org/10.1016/j.physletb.2020.135426>. <arXiv:2002.05315>
- LHC Higgs Cross Section Working Group, Handbook of LHC Higgs cross sections: 4. Deciphering the nature of the Higgs sector. CERN Report CERN-2017-002-M (2016). <https://doi.org/10.23731/CYRM-2017-002>. <arXiv:1610.07922>
- Y. Gao et al., Spin determination of single-produced resonances at hadron colliders. *Phys. Rev. D* **81**, 075022 (2010). <https://doi.org/10.1103/PhysRevD.81.075022>. <arXiv:1001.3396>
- S. Bolognesi et al., On the spin and parity of a single-produced resonance at the LHC. *Phys. Rev. D* **86**, 095031 (2012). <https://doi.org/10.1103/PhysRevD.86.095031>. <arXiv:1208.4018>
- I. Anderson et al., Constraining anomalous HVV interactions at proton and lepton colliders. *Phys. Rev. D* **89**, 035007 (2014). <https://doi.org/10.1103/PhysRevD.89.035007>. <arXiv:1309.4819>
- A.V. Gritsan, R. Röntsch, M. Schulze, M. Xiao, Constraining anomalous Higgs boson couplings to the heavy flavor fermions using matrix element techniques. *Phys. Rev. D* **94**, 055023 (2016). <https://doi.org/10.1103/PhysRevD.94.055023>. <arXiv:1606.03107>
- A.V. Gritsan et al., New features in the JHU generator framework: constraining Higgs boson properties from on-shell and off-shell production. *Phys. Rev. D* **102**, 056022 (2020). <https://doi.org/10.1103/PhysRevD.102.056022>. <arXiv:2002.09888>
- CMS Collaboration, Measurements of the Higgs boson production cross section and couplings in the W boson pair decay channel in proton–proton collisions at $\sqrt{s} = 13$ TeV. *Eur. Phys. J. C* **83**, 667 (2023). <https://doi.org/10.1140/epjc/s10052-023-11632-6>. <arXiv:2206.09466>
- LHC Higgs Cross Section Working Group, Handbook of LHC Higgs cross sections: 3. Higgs properties: report of the LHC Higgs Cross Section Working Group. CERN Report CERN-2013-004 (2013). <https://doi.org/10.5170/CERN-2013-004>. <arXiv:1307.1347>
- CMS Collaboration, Measurements of $t\bar{t}H$ production and the CP structure of the Yukawa interaction between the Higgs boson and top quark in the diphoton decay channel. *Phys. Rev. Lett.* **125**, 061801 (2020). <https://doi.org/10.1103/PhysRevLett.125.061801>. <arXiv:2003.10866>
- CMS Collaboration, Constraints on anomalous Higgs boson couplings to vector bosons and fermions in its production and decay using the four-lepton final state. *Phys. Rev. D* **104**, 052004 (2021). <https://doi.org/10.1103/physrevd.104.052004>. <arXiv:2104.12152>
- HEPData record for this analysis (2024). <https://doi.org/10.17182/hepdata.146013>
- B. Grzadkowski, M. Iskrzynski, M. Misiak, J. Rosiek, Dimension-six terms in the standard model Lagrangian. *JHEP* **10**, 085 (2010). [https://doi.org/10.1007/jhep10\(2010\)085](https://doi.org/10.1007/jhep10(2010)085). <arXiv:1008.4884>
- J. Davis et al., Constraining anomalous Higgs boson couplings to virtual photons. *Phys. Rev. D* **105**, 096027 (2022). <https://doi.org/10.1103/PhysRevD.105.096027>. <arXiv:2109.13363>

32. A.V. Gritsan et al., New features in the JHU generator framework: constraining Higgs boson properties from on-shell and off-shell production. *Phys. Rev. D* **102** (2020). <https://doi.org/10.1103/physrevd.102.056022>
33. CMS Collaboration, The CMS experiment at the CERN LHC. *JINST* **3**, S08004 (2008). <https://doi.org/10.1088/1748-0221/3/08/S08004>
34. CMS Collaboration, Performance of electron reconstruction and selection with the CMS detector in proton–proton collisions at $\sqrt{s} = 8$ TeV. *JINST* **10**, P06005 (2015). <https://doi.org/10.1088/1748-0221/10/06/P06005>. arXiv:1502.02701
35. CMS Collaboration, Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV. *JINST* **13**, P06015 (2018). <https://doi.org/10.1088/1748-0221/13/06/P06015>. arXiv:1804.04528
36. CMS Collaboration, Performance of photon reconstruction and identification with the CMS detector in proton–proton collisions at $\sqrt{s} = 8$ TeV. *JINST* **10**, P08010 (2015). <https://doi.org/10.1088/1748-0221/10/08/P08010>. arXiv:1502.02702
37. CMS Collaboration, Description and performance of track and primary-vertex reconstruction with the CMS tracker. *JINST* **9**, P10009 (2014). <https://doi.org/10.1088/1748-0221/9/10/P10009>. arXiv:1405.6569
38. CMS Collaboration, Particle-flow reconstruction and global event description with the CMS detector. *JINST* **12**, P10003 (2017). <https://doi.org/10.1088/1748-0221/12/10/P10003>. arXiv:1706.04965
39. CMS Collaboration, Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV. *JINST* **13**, P10005 (2018). <https://doi.org/10.1088/1748-0221/13/10/P10005>. arXiv:1809.02816
40. CMS Collaboration, Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV. *JINST* **12**, P02014 (2017). <https://doi.org/10.1088/1748-0221/12/02/P02014>. arXiv:1607.03663
41. CMS Collaboration, Performance of missing transverse momentum reconstruction in proton–proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector. *JINST* **14**, P07004 (2019). <https://doi.org/10.1088/1748-0221/14/07/P07004>. arXiv:1903.06078
42. CMS Collaboration, Performance of the CMS Level-1 trigger in proton–proton collisions at $\sqrt{s} = 13$ TeV. *JINST* **15**, P10017 (2020). <https://doi.org/10.1088/1748-0221/15/10/P10017>. arXiv:2006.10165
43. CMS Collaboration, The CMS trigger system. *JINST* **12**, P01020 (2017). <https://doi.org/10.1088/1748-0221/12/01/P01020>. arXiv:1609.02366
44. CMS Collaboration, Precision luminosity measurement in proton–proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS. *Eur. Phys. J. C* **800**, 81 (2021). <https://doi.org/10.1140/epjc/s10052-021-09538-2>. arXiv:2104.01927
45. CMS Collaboration, CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV. CMS Physics Analysis Summary CMS-PAS-LUM-17-004 (2017). <https://cds.cern.ch/record/2621960>
46. CMS Collaboration, CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV. CMS Physics Analysis Summary CMS-PAS-LUM-18-002 (2019). <https://cds.cern.ch/record/2676164>
47. NNPDF Collaboration, Parton distributions with QED corrections. *Nucl. Phys. B* **877**, 290 (2013). <https://doi.org/10.1016/j.nuclphysb.2013.10.010>. arXiv:1308.0598
48. NNPDF Collaboration, Unbiased global determination of parton distributions and their uncertainties at NNLO and at LO. *Nucl. Phys. B* **855**, 153 (2012). <https://doi.org/10.1016/j.nuclphysb.2011.09.024>. arXiv:1107.2652
49. NNPDF Collaboration, Parton distributions from high-precision collider data. *Eur. Phys. J. C* **77**, 663 (2017). <https://doi.org/10.1140/epjc/s10052-017-5199-5>. arXiv:1706.00428
50. CMS Collaboration, Event generator tunes obtained from underlying event and multiparton scattering measurements. *Eur. Phys. J. C* **76**, 155 (2016). <https://doi.org/10.1140/epjc/s10052-016-3988-x>. arXiv:1512.00815
51. CMS Collaboration, Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements. *Eur. Phys. J. C* **80**, 4 (2020). <https://doi.org/10.1140/epjc/s10052-019-7499-4>. arXiv:1903.12179
52. T. Sjöstrand et al., An introduction to PYTHIA 8.2. *Comput. Phys. Commun.* **191**, 159 (2015). <https://doi.org/10.1016/j.cpc.2015.01.024>. arXiv:1410.3012
53. P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms. *JHEP* **11**, 040 (2004). <https://doi.org/10.1088/1126-6708/2004/11/040>. arXiv:hep-ph/0409146
54. S. Frixione, P. Nason, C. Oleari, Matching NLO QCD computations with parton shower simulations: the POWHEG method. *JHEP* **11**, 070 (2007). <https://doi.org/10.1088/1126-6708/2007/11/070>. arXiv:0709.2092
55. S. Alioli, P. Nason, C. Oleari, E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX. *JHEP* **06**, 043 (2010). [https://doi.org/10.1007/JHEP06\(2010\)043](https://doi.org/10.1007/JHEP06(2010)043). arXiv:1002.2581
56. E. Bagnaschi, G. Degrassi, P. Slavich, A. Vicini, Higgs production via gluon fusion in the POWHEG approach in the SM and in the MSSM. *JHEP* **02**, 088 (2012). [https://doi.org/10.1007/JHEP02\(2012\)088](https://doi.org/10.1007/JHEP02(2012)088). arXiv:1111.2854
57. P. Nason, C. Oleari, NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG. *JHEP* **02**, 037 (2010). [https://doi.org/10.1007/JHEP02\(2010\)037](https://doi.org/10.1007/JHEP02(2010)037). arXiv:0911.5299
58. G. Luisoni, P. Nason, C. Oleari, F. Tramontano, $HW^\pm/HZ + 0$ and 1 jet at NLO with the POWHEG BOX interfaced to GoSam and their merging within MiNLO. *JHEP* **10**, 083 (2013). [https://doi.org/10.1007/JHEP10\(2013\)083](https://doi.org/10.1007/JHEP10(2013)083). arXiv:1306.2542
59. H.B. Hartanto, B. Jager, L. Reina, D. Wackerlo, Higgs boson production in association with top quarks in the POWHEG BOX. *Phys. Rev. D* **91**, 094003 (2015). <https://doi.org/10.1103/PhysRevD.91.094003>. arXiv:1501.04498
60. K. Hamilton, P. Nason, E. Re, G. Zanderighi, NNLOPS simulation of Higgs boson production. *JHEP* **10**, 222 (2013). [https://doi.org/10.1007/JHEP10\(2013\)222](https://doi.org/10.1007/JHEP10(2013)222). arXiv:1309.0017
61. K. Hamilton, P. Nason, G. Zanderighi, Finite quark-mass effects in the NNLOPS POWHEG+MiNLO Higgs generator. *JHEP* **05**, 140 (2015). [https://doi.org/10.1007/JHEP05\(2015\)140](https://doi.org/10.1007/JHEP05(2015)140). arXiv:1501.04637
62. N. Berger et al., Simplified template cross sections—stage 1.1 (2019). arXiv:1906.02754
63. R. Frederix, K. Hamilton, Extending the MINLO method. *JHEP* **05**, 042 (2016). [https://doi.org/10.1007/JHEP05\(2016\)042](https://doi.org/10.1007/JHEP05(2016)042). arXiv:1512.02663
64. J. Alwall et al., The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP* **07**, 079 (2014). [https://doi.org/10.1007/JHEP07\(2014\)079](https://doi.org/10.1007/JHEP07(2014)079). arXiv:1405.0301
65. CMS Collaboration, A measurement of the Higgs boson mass in the diphoton decay channel. *Phys. Lett. B* **805**, 135425 (2020). <https://doi.org/10.1016/j.physletb.2020.135425>. arXiv:2002.06398
66. P. Nason, C. Oleari, M. Rocco, M. Zaro, An interface between the POWHEG BOX and MadGraph5_aMC@NLO. *Eur. Phys. J. C* **80**, 10 (2020). <https://doi.org/10.1140/epjc/s10052-020-08559-7>. arXiv:2008.06364

67. P. Nason, G. Zanderighi, $W^+ W^-$, WZ and ZZ production in the POWHEG-BOX-V2. Eur. Phys. J. C **74**, 2702 (2014). <https://doi.org/10.1140/epjc/s10052-013-2702-5>. arXiv:1311.1365
68. P. Meade, H. Ramani, M. Zeng, Transverse momentum resummation effects in $W^+ W^-$ measurements. Phys. Rev. D **90**, 114006 (2014). <https://doi.org/10.1103/PhysRevD.90.114006>. arXiv:1407.4481
69. P. Jaiswal, T. Okui, Explanation of the WW excess at the LHC by jet-veto resummation. Phys. Rev. D **90**, 073009 (2014). <https://doi.org/10.1103/PhysRevD.90.073009>. arXiv:1407.4537
70. J.M. Campbell, R.K. Ellis, An update on vector boson pair production at hadron colliders. Phys. Rev. D **60**, 113006 (1999). <https://doi.org/10.1103/PhysRevD.60.113006>. arXiv:hep-ph/9905386
71. J.M. Campbell, R.K. Ellis, C. Williams, Vector boson pair production at the LHC. JHEP **07**, 018 (2011). [https://doi.org/10.1007/JHEP07\(2011\)018](https://doi.org/10.1007/JHEP07(2011)018). arXiv:1105.0020
72. J.M. Campbell, R.K. Ellis, W.T. Giele, A multi-threaded version of MCFM. Eur. Phys. J. C **75**, 246 (2015). <https://doi.org/10.1140/epjc/s10052-015-3461-2>. arXiv:1503.06182
73. F. Caola et al., QCD corrections to vector boson pair production in gluon fusion including interference effects with off-shell Higgs at the LHC. JHEP **07**, 087 (2016). [https://doi.org/10.1007/JHEP07\(2016\)087](https://doi.org/10.1007/JHEP07(2016)087). arXiv:1605.04610
74. J. Alwall et al., Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions. Eur. Phys. J. C **53**, 473 (2008). <https://doi.org/10.1140/epjc/s10052-007-0490-5>. arXiv:0706.2569
75. S. Frixione, P. Nason, G. Ridolfi, A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction. JHEP **09**, 126 (2007). <https://doi.org/10.1088/1126-6708/2007/09/126>. arXiv:0707.3088
76. S. Alioli, P. Nason, C. Oleari, E. Re, NLO single-top production matched with shower in POWHEG: s - and t -channel contributions. JHEP **09**, 111 (2009). arXiv:0907.4076. [Erratum: [https://doi.org/10.1007/JHEP02\(2010\)011](https://doi.org/10.1007/JHEP02(2010)011)
77. E. Re, Single-top Wt -channel production matched with parton showers using the POWHEG method. Eur. Phys. J. C **71**, 1547 (2011). <https://doi.org/10.1140/epjc/s10052-011-1547-z>. arXiv:1009.2450
78. M. Czakon et al., Top-pair production at the LHC through NNLO QCD and NLO EW. JHEP **10**, 186 (2017). [https://doi.org/10.1007/JHEP10\(2017\)186](https://doi.org/10.1007/JHEP10(2017)186). arXiv:1705.04105
79. R. Frederix, S. Frixione, Merging meets matching in MC@NLO. JHEP **12**, 061 (2012). [https://doi.org/10.1007/JHEP12\(2012\)061](https://doi.org/10.1007/JHEP12(2012)061). arXiv:1209.6215
80. GEANT4 Collaboration, GEANT4—a simulation toolkit. Nucl. Instrum. Methods A **506**, 250 (2003). [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8)
81. C.M.S. Collaboration, Muon identification using multivariate techniques in the CMS experiment in proton-proton collisions at $\sqrt{s} = 13$ TeV. INST **19**, P02031 (2024). <https://doi.org/10.1088/1748-0221/19/02/P02031>
82. W. Waltenberger, R. Fröhwirth, P. Vanlaer, Adaptive vertex fitting. J. Phys. G **34**, N343 (2007). <https://doi.org/10.1088/0954-3899/34/12/N01>
83. CMS Collaboration, Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid. CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02 (2015). <http://cds.cern.ch/record/2020886>
84. CMS Collaboration, Pileup mitigation at CMS in 13 TeV data. JINST **15**, P09018 (2020). <https://doi.org/10.1088/1748-0221/15/09/p09018>. arXiv:2003.00503
85. D. Bertolini, P. Harris, M. Low, N. Tran, Pileup per particle identification. JHEP **10**, 059 (2014). [https://doi.org/10.1007/JHEP10\(2014\)059](https://doi.org/10.1007/JHEP10(2014)059). arXiv:1407.6013
86. J. Thaler, K. Van Tilburg, Identifying boosted objects with N -subjettiness. JHEP **03**, 015 (2011). [https://doi.org/10.1007/JHEP03\(2011\)015](https://doi.org/10.1007/JHEP03(2011)015). arXiv:1011.2268
87. M. Dasgupta, A. Fregoso, S. Marzani, G.P. Salam, Towards an understanding of jet substructure. JHEP **09**, 029 (2013). [https://doi.org/10.1007/JHEP09\(2013\)029](https://doi.org/10.1007/JHEP09(2013)029). arXiv:1307.0007
88. J.M. Butterworth, A.R. Davison, M. Rubin, G.P. Salam, Jet substructure as a new Higgs search channel at the LHC. Phys. Rev. Lett. **100**, 242001 (2008). <https://doi.org/10.1103/PhysRevLett.100.242001>. arXiv:0802.2470
89. A.J. Larkoski, S. Marzani, G. Soyez, J. Thaler, Soft drop. JHEP **05**, 146 (2014). [https://doi.org/10.1007/JHEP05\(2014\)146](https://doi.org/10.1007/JHEP05(2014)146). arXiv:1402.2657
90. CMS Collaboration, Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV. JINST **13**, P05011 (2018). <https://doi.org/10.1088/1748-0221/13/05/P05011>. arXiv:1712.07158
91. CMS Collaboration, CMS Phase 1 heavy flavour identification performance and developments. CMS Detector Performance Summary CMS-DP-2020-019 (2017). <http://cds.cern.ch/record/2263802>
92. CMS Collaboration, Measurements of properties of the Higgs boson decaying to a W boson pair in pp collisions at $\sqrt{s} = 13$ TeV. Phys. Lett. B **791**, 96 (2019). <https://doi.org/10.1016/j.physlettb.2018.12.073>. arXiv:1806.05246
93. C.M.S. Collaboration, Measurements of inclusive W and Z cross sections in pp collisions at $\sqrt{s} = 7$ TeV. JHEP **01**, 080 (2011). [https://doi.org/10.1007/JHEP01\(2011\)080](https://doi.org/10.1007/JHEP01(2011)080). arXiv:1012.2466
94. CMS Collaboration, An embedding technique to determine $\tau\tau$ backgrounds in proton–proton collision data. JINST **14**, P06032 (2019). <https://doi.org/10.1088/1748-0221/14/06/P06032>. arXiv:1903.01216
95. C.M.S. Collaboration, Identification techniques for highly boosted W bosons that decay into hadrons. JHEP **12**, 017 (2014). [https://doi.org/10.1007/JHEP12\(2014\)017](https://doi.org/10.1007/JHEP12(2014)017). arXiv:1410.4227
96. R. Barlow, C. Beeston, Fitting using finite Monte Carlo samples. Comput. Phys. Commun. **77**, 219 (1993). [https://doi.org/10.1016/0010-4655\(93\)90005-W](https://doi.org/10.1016/0010-4655(93)90005-W)
97. ATLAS Collaboration, Measurement of the inelastic proton–proton cross section at $\sqrt{s} = 13$ TeV with the ATLAS detector at the LHC. Phys. Rev. Lett. **117**, 182002 (2016). <https://doi.org/10.1103/PhysRevLett.117.182002>. arXiv:1606.02625
98. CMS Collaboration, Measurement of the inelastic proton–proton cross section at $\sqrt{s} = 13$ TeV. JHEP **07**, 161 (2018). [https://doi.org/10.1007/JHEP07\(2018\)161](https://doi.org/10.1007/JHEP07(2018)161). arXiv:1802.02613
99. G. Passarino, Higgs CAT. Eur. Phys. J. C **74**, 2866 (2014). <https://doi.org/10.1140/epjc/s10052-014-2866-7>. arXiv:1312.2397
100. C.M.S. Collaboration, Measurement of Higgs boson production and properties in the WW decay channel with leptonic final states. JHEP **01**, 096 (2014). [https://doi.org/10.1007/JHEP01\(2014\)096](https://doi.org/10.1007/JHEP01(2014)096). arXiv:1312.1129
101. The ATLAS Collaboration, The CMS Collaboration, The LHC Higgs Combination Group, Procedure for the LHC Higgs boson search combination in Summer 2011. Technical Report ATL-PHYS-PUB 2011-11, CMS NOTE 2011/005 (2011). <http://cds.cern.ch/record/1379837>
102. CMS Collaboration, Combined measurements of Higgs boson couplings in proton–proton collisions at $\sqrt{s} = 13$ TeV. Eur. Phys. J. C **79**, 421 (2019). <https://doi.org/10.1140/epjc/s10052-019-6909-y>. arXiv:1809.10733
103. S.S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses. Ann. Math. Stat. **9**, 60 (1938). <https://doi.org/10.1214/aoms/1177732360>

104. G. Cowan, K. Cranmer, E. Gross, O. Vitells, Asymptotic formulae for likelihood-based tests of new physics. *Eur. Phys. J. C* **71**, 1554 (2011). <https://doi.org/10.1140/epjc/s10052-011-1554-0>.
[arXiv:1007.1727](https://arxiv.org/abs/1007.1727)

CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J. W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , P. S. Hussain , M. Jeitler , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck , R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger , C.-E. Wulz

Universiteit Antwerpen, Antwerp, Belgium

M. R. Darwish , T. Janssen , P. Van Mechelen

Vrije Universiteit Brussel, Brussels, Belgium

E. S. Bols , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , H. El Faham , S. Lowette , I. Makarenko , D. Müller , A. R. Sahasransu , S. Tavernier , M. Tytgat , G. P. Van Onsem , S. Van Putte , D. Vannerom

Université Libre de Bruxelles, Brussels, Belgium

B. Clerbaux , A. K. Das, G. De Lentdecker , L. Favart , P. Gianneios , D. Hohov , J. Jaramillo , A. Khalilzadeh, K. Lee , M. Mahdavikhorrami , A. Malara , S. Paredes , L. Pétré , N. Postiau, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , Y. Hong , J. Knolle , L. Lambrecht , G. Mestdach, K. Mota Amarilo , C. Rendón, A. Samalan, K. Skovpen , N. Van Den Bossche , J. van der Linden , L. Wezenbeek

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , C. Delaere , I. S. Donertas , A. Giammanco , K. Jaffel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , K. Mondal , T. T. Tran , S. Wertz

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G. A. Alves , E. Coelho , C. Hensel , T. Menezes De Oliveira , A. Moraes , P. Rebello Teles , M. Soeiro

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W. L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato , E. M. Da Costa , G. G. Da Silveira , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza, J. Martins , C. Mora Herrera , L. Mundim , H. Nogima , J. P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel , A. Vilela Pereira

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C. A. Bernardes , L. Calligaris , T. R. Fernandez Perez Tomei , E. M. Gregores , P. G. Mercadante , S. F. Novaes , B. Orzari , Sandra S. Padula

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , S. Thakur

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan

Department of Physics, Tsinghua University, Beijing, ChinaZ. Hu , J. Liu, K. Yi  ^{8,9}**Institute of High Energy Physics, Beijing, China**G. M. Chen , H. S. Chen , M. Chen , F. Iemmi , C. H. Jiang, A. Kapoor , H. Liao , Z.-A. Liu , R. Sharma , J. N. Song , J. Tao , C. Wang , J. Wang , Z. Wang , H. Zhang **State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China**A. Agapitos , Y. Ban , A. Levin , C. Li , Q. Li , Y. Mao, S. J. Qian , X. Sun , D. Wang , H. Yang, L. Zhang , C. Zhou **Sun Yat-Sen University, Guangzhou, China**Z. You **University of Science and Technology of China, Hefei, China**N. Lu **Nanjing Normal University, Nanjing, China**G. Bauer **Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE)-Fudan University, Shanghai, China**X. Gao , D. Leggat, H. Okawa **Zhejiang University, Hangzhou, Zhejiang, China**Z. Lin , C. Lu , M. Xiao **Universidad de Los Andes, Bogotá, Colombia**C. Avila , D. A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J. A. Reyes Vega**Universidad de Antioquia, Medellín, Colombia**J. Mejia Guisao , F. Ramirez , M. Rodriguez , J. D. Ruiz Alvarez **Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia**D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac **Faculty of Science, University of Split, Split, Croatia**M. Kovac , T. Sculac **Institute Rudjer Boskovic, Zagreb, Croatia**P. Bargassa , V. Brigljevic , B. K. Chitroda , D. Ferencek , S. Mishra , A. Starodumov , T. Susa  ¹⁶**University of Cyprus, Nicosia, Cyprus**A. Attikis , K. Christoforou , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P. A. Razis , H. Rykaczewski, H. Saka , A. Stepennov **Charles University, Prague, Czech Republic**M. Finger , M. Finger Jr. , A. Kveton **Escuela Politecnica Nacional, Quito, Ecuador**E. Ayala **Universidad San Francisco de Quito, Quito, Ecuador**E. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**S. Elgammal , A. Ellithi Kamel  ¹⁸**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**M. A. Mahmoud , Y. Mohammed 

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht ID, M. Kadastik, T. Lange ID, S. Nandan ID, C. Nielsen ID, J. Pata ID, M. Raidal ID, L. Tani ID, C. Veelken ID

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann ID, K. Osterberg ID, M. Voutilainen ID

Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar ID, E. Brücken ID, F. Garcia ID, K. T. S. Kallonen ID, R. Kinnunen, T. Lampén ID, K. Lassila-Perini ID, S. Lehti ID, T. Lindén ID, L. Martikainen ID, M. Myllymäki ID, M. M. Rantanen ID, H. Siikonen ID, E. Tuominen ID, J. Tuomiemi ID

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka ID, H. Petrow ID

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, FranceM. Besancon ID, F. Couderc ID, M. Dejardin ID, D. Denegri, J. L. Faure, F. Ferri ID, S. Ganjour ID, P. Gras ID, G. Hamel de Monchenault ID, V. Lohezic ID, J. Malcles ID, J. Rander, A. Rosowsky ID, M. Ö. Sahin ID, A. Savoy-Navarro ID¹⁹, P. Simkina ID, M. Titov ID, M. Tornago ID**Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**

C. Baldenegro Barrera ID, F. Beaudette ID, A. Buchot Perraguin ID, P. Busson ID, A. Cappati ID, C. Charlot ID, M. Chiusi ID, F. Damas ID, O. Davignon ID, A. De Wit ID, B. A. Fontana Santos Alves ID, S. Ghosh ID, A. Gilbert ID, R. Granier de Cassagnac ID, A. Hakimi ID, B. Harikrishnan ID, L. Kalipoliti ID, G. Liu ID, J. Motta ID, M. Nguyen ID, C. Ochando ID, L. Portales ID, R. Salerno ID, J. B. Sauvan ID, Y. Sirois ID, A. Tarabini ID, E. Vernazza ID, A. Zabi ID, A. Zghiche ID

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, FranceJ.-L. Agram ID²⁰, J. Andrea ID, D. Apparu ID, D. Bloch ID, J.-M. Brom ID, E. C. Chabert ID, C. Collard ID, S. Falke ID, U. Goerlach ID, C. Grimault, R. Haeberle ID, A.-C. Le Bihan ID, M. Meena ID, G. Saha ID, M. A. Sessini ID, P. Van Hove ID**Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**S. Beauceron ID, B. Blancon ID, G. Boudoul ID, N. Chanon ID, J. Choi ID, D. Contardo ID, P. Depasse ID, C. Dozen ID²¹, H. El Mamouni, J. Fay ID, S. Gascon ID, M. Gouzevitch ID, C. Greenberg, G. Grenier ID, B. Ille ID, I. B. Laktineh, M. Lethuillier ID, L. Mirabito, S. Perries, A. Purohit ID, M. Vander Donckt ID, P. Verdier ID, J. Xiao ID**Georgian Technical University, Tbilisi, Georgia**G. Adamov, I. Lomidze ID, Z. Tsamalaidze ID¹⁶**I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany**

V. Botta ID, L. Feld ID, K. Klein ID, M. Lipinski ID, D. Meuser ID, A. Pauls ID, N. Röwert ID, M. Teroerde ID

RWTH Aachen University, III. Physikalisches Institut A, Aachen, GermanyS. Diekmann ID, A. Dodonova ID, N. Eich ID, D. Eliseev ID, F. Engelke ID, J. Erdmann ID, M. Erdmann ID, P. Fackeldey ID, B. Fischer ID, T. Hebbeker ID, K. Hoepfner ID, F. Ivone ID, A. Jung ID, M. Y. Lee ID, L. Mastrolorenzo, F. Mausolf ID, M. Merschmeyer ID, A. Meyer ID, S. Mukherjee ID, D. Noll ID, F. Nowotny, A. Pozdnyakov ID, Y. Rath, W. Redjeb ID, F. Rehm, H. Reithler ID, U. Sarkar ID, V. Sarkisovi ID, A. Schmidt ID, A. Sharma ID, J. L. Spah ID, A. Stein ID, F. Torres Da Silva De Araujo ID²², L. Vigilante, S. Wiedenbeck ID, S. Zaleski**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**C. Dziwok ID, G. Flügge ID, W. Haj Ahmad ID²³, T. Kress ID, A. Nowack ID, O. Pooth ID, A. Stahl ID, T. Ziemons ID, A. Zottz ID**Deutsches Elektronen-Synchrotron, Hamburg, Germany**H. Aarup Petersen ID, M. Aldaya Martin ID, J. Alimena ID, S. Amoroso, Y. An ID, S. Baxter ID, M. Bayatmakou ID, H. Becerril Gonzalez ID, O. Behnke ID, A. Belvedere ID, S. Bhattacharya ID, F. Blekman ID²⁴, K. Borras ID²⁵, A. Campbell ID, A. Cardini ID, C. Cheng, F. Colombina ID, S. Consuegra Rodríguez ID, G. Correia Silva ID, M. De Silva ID, G. Eckerlin, D. Eckstein ID, L. I. Estevez Banos ID, O. Filatov ID, E. Gallo ID²⁴, A. Geiser ID, A. Giraldi ID, G. Greau, V. Guglielmi ID, M. Guthoff ID, A. Hinzmman ID, A. Jafari ID²⁶, L. Jeppe ID, N. Z. Jomhari ID, B. Kaech ID, M. Kasemann ID,

C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁷, W. Lohmann ²⁸, R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A. B. Meyer , G. Milella , A. Mussgiller , L. P. Nair , A. Nürnberg , Y. Otarid, J. Park , D. Pérez Adán , E. Ranken , A. Rasperera , B. Ribeiro Lopes , J. Rübenach, A. Saggio , M. Scham ^{25, 29}, S. Schnake ²⁵, P. Schütze , C. Schwanenberger ²⁴, D. Selivanova , K. Sharko , M. Shchedrolosiev , R. E. Sosa Ricardo , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²⁵, C. Wissing , Y. Yang , A. Zimermann Castro Santos

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , S. Bollweg, M. Bonanomi , P. Connor , M. Eich, K. El Morabit , Y. Fischer , A. Fröhlich, C. Garbers , E. Garutti , A. Grohsjean , M. Hajheidari, J. Haller , H. R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari , T. Kramer , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K. J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoiu , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ³⁰, R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , R. Koppenhöfer , J. M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H. J. Simonis , M. Toms , N. Trevisani , R. Ulrich , R. F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Agia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis, A. Papadopoulos ³⁰, A. Stakia

National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, A. Panagiotou, I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis, A. Zacharopoulou

University of Ioánnina, Ioannina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P. G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

M. Bartók ³¹, C. Hajdu , D. Horvath ^{32, 33}, K. Márton, F. Sikler , V. Veszpremi

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csand , K. Farkas , M. M. A. Gadallah ³⁴, Á. Kadlecik , P. Major , K. Mandal , G. Pasztor , A. J. Radl ³⁵, G. I. Veres

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

P. Raics, B. Ujvari , G. Zilizi

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

MATE Institute of Technology, Karoly Robert Campus, Gyongyos, Hungary

T. Csorgo ³⁵, F. Nemes ³⁵, T. Novak

Punjab University, Chandigarh, India

J. Babbar , S. Bansal , S. B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra ³⁶, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , K. Sandeep , T. Sheokand, J. B. Singh , A. Singla

University of Delhi, Delhi, India

A. Ahmed ID, A. Bhardwaj ID, A. Chhetri ID, B. C. Choudhary ID, A. Kumar ID, A. Kumar ID, M. Naimuddin ID, K. Ranjan ID, S. Saumya ID

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia ID, S. Barman ID³⁷, S. Bhattacharya ID, S. Dutta ID, S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Chennai, India

M. M. Ameen ID, P. K. Behera ID, S. C. Behera ID, S. Chatterjee ID, P. Jana ID, P. Kalbhor ID, J. R. Komaragiri ID³⁸, D. Kumar ID³⁸, L. Panwar ID³⁸, P. R. Pujahari ID, N. R. Saha ID, A. Sharma ID, A. K. Sikdar ID, S. Verma ID

Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, M. Kumar ID, G. B. Mohanty ID, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala ID, S. Banerjee ID, R. M. Chatterjee, R. K. Dewanjee ID³⁹, M. Guchait ID, Sh. Jain ID, A. Jaiswal, S. Karmakar ID, S. Kumar ID, G. Majumder ID, K. Mazumdar ID, S. Parolia ID, A. Thachayath ID

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati ID⁴⁰, C. Kar ID, D. Maity ID⁴¹, P. Mal ID, T. Mishra ID, V. K. Muraleedharan Nair Bindhu ID⁴¹, K. Naskar ID⁴¹, A. Nayak ID⁴¹, P. Sadangi, P. Saha ID, S. K. Swain ID, S. Varghese ID⁴¹, D. Vats ID⁴¹

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya ID⁴², A. Alpana ID, S. Dubey ID, B. Gomber ID⁴², B. Kansal ID, A. Laha ID, B. Sahu ID⁴², S. Sharma ID, K. Y. Vaish

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi ID⁴³, E. Khazaie ID⁴⁴, M. Zeinali ID⁴⁵

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Chenarani ID⁴⁶, S. M. Etesami ID, M. Khakzad ID, M. Mohammadi Najafabadi ID

University College Dublin, Dublin, Ireland

M. Grunewald ID

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia ID^{a,b}, R. Aly ID^{a,c,47}, A. Colaleo ID^{a,b}, D. Creanza ID^{a,c}, B. D'Anzi ID^{a,b}, N. De Filippis ID^{a,c}, M. De Palma ID^{a,b}, A. Di Florio ID^{a,c}, W. Elmetenawee ID^{a,b,47}, L. Fiore ID^a, G. Iaselli ID^{a,c}, M. Louka ID^{a,b}, G. Maggi ID^{a,c}, M. Maggi ID^a, I. Margjeka ID^{a,b}, V. Mastrapasqua ID^{a,b}, S. My ID^{a,b}, S. Nuzzo ID^{a,b}, A. Pellecchia ID^{a,b}, A. Pompili ID^{a,b}, G. Pugliese ID^{a,c}, R. Radogna ID^a, G. Ramirez-Sanchez ID^{a,c}, D. Ramos ID^a, A. Ranieri ID^a, L. Silvestris ID^a, F. M. Simone ID^{a,b}, Ü. Sözbilir ID^a, A. Stamerra ID^a, R. Venditti ID^a, P. Verwilligen ID^a, A. Zaza ID^{a,b}

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi ID^a, C. Battilana ID^{a,b}, D. Bonacorsi ID^{a,b}, L. Borgonovi ID^a, R. Campanini ID^{a,b}, P. Capiluppi ID^{a,b}, A. Castro ID^{a,b}, F. R. Cavallo ID^a, M. Cuffiani ID^{a,b}, G. M. Dallavalle ID^a, T. Diotalevi ID^{a,b}, F. Fabbri ID^a, A. Fanfani ID^{a,b}, D. Fasanella ID^{a,b}, P. Giacomelli ID^a, L. Giommi ID^{a,b}, C. Grandi ID^a, L. Guiducci ID^{a,b}, S. Lo Meo ID^{a,48}, L. Lunerti ID^{a,b}, G. Masetti ID^a, F. L. Navarria ID^{a,b}, A. Perrotta ID^a, F. Primavera ID^a, A. M. Rossi ID^{a,b}, T. Rovelli ID^{a,b}, G. P. Siroli ID^{a,b}

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa ID^{a,b,49}, A. Di Mattia ID^a, R. Potenza^{a,b}, A. Tricomi ID^{a,b,49}, C. Tuve ID^{a,b}

INFN Sezione di Firenze^a, Università di Firenze^b, Florence, Italy

P. Assiouras ID^a, G. Barbagli ID^a, G. Bardelli ID^{a,b}, B. Camaiani ID^{a,b}, A. Cassese ID^a, R. Ceccarelli ID^a, V. Ciulli ID^{a,b}, C. Civinini ID^a, R. D'Alessandro ID^{a,b}, E. Focardi ID^{a,b}, T. Kello ID^a, G. Latino ID^{a,b}, P. Lenzi ID^{a,b}, M. Lizzo ID^a, M. Meschini ID^a, S. Paoletti ID^a, A. Papanastassiou ID^{a,b}, G. Sguazzoni ID^a, L. Viliani ID^a

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi ID, S. Bianco ID, S. Meola ID⁵⁰, D. Piccolo ID

INFN Sezione di Genova^a, Università di Genova^b, Genoa, Italy

P. Chatagnon , F. Ferro , E. Robutti , S. Tosi

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milan, Italy

A. Benaglia , G. Boldrini , F. Brivio , F. Cetorelli , F. De Guio , M. E. Dinardo , P. Dini , S. Gennai , R. Gerosa , A. Ghezzi , P. Govoni , L. Guzzi , M. T. Lucchini , M. Malberti , S. Malvezzi , A. Massironi , D. Menasce , L. Moroni , M. Paganoni , D. Pedrini , B. S. Pinolini , S. Ragazzi , T. Tabarelli de Fatis , D. Zuolo

INFN Sezione di Napoli^a, Università di Napoli ‘Federico II’^b, Naples, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Naples, Italy

S. Buontempo , A. Cagnotta , F. Carnevali , N. Cavallo , F. Fabozzi , A. O. M. Iorio , L. Lista , P. Paolucci , B. Rossi , C. Sciacca

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino , P. Azzi , N. Bacchetta , D. Bisello , P. Bortignon , A. Bragagnolo , R. Carlin , P. Checchia , T. Dorigo , F. Gasparini , U. Gasparini , F. Gonella , E. Lusiani , M. Margoni , F. Marini , M. Migliorini , J. Pazzini , P. Ronchese , R. Rossin , F. Simonetto , G. Strong , M. Tosi , A. Triossi , S. Ventura , H. Yarar , M. Zanetti , P. Zotto , A. Zucchetta , G. Zumerle

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

S. Abu Zeid , C. Aimè , A. Braghieri , S. Calzaferri , D. Fiorina , P. Montagna , V. Re , C. Riccardi , P. Salvini , I. Vai , P. Vitulo

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal , G. M. Bilei , D. Ciangottini , L. Fanò , M. Magherini , G. Mantovani , V. Mariani , M. Menichelli , F. Moscatelli , A. Rossi , A. Santocchia , D. Spiga , T. Tedeschi

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

P. Asenov , P. Azzurri , G. Bagliesi , R. Bhattacharya , L. Bianchini , T. Boccali , E. Bossini , D. Bruschini , R. Castaldi , M. A. Ciocci , M. Cipriani , V. D’Amante , R. Dell’Orso , S. Donato , A. Giassi , F. Ligabue , D. Matos Figueiredo , A. Messineo , M. Musich , F. Palla , A. Rizzi , G. Rolandi , S. Roy Chowdhury , T. Sarkar , A. Scribano , P. Spagnolo , R. Tenchini , G. Tonelli , N. Turini , A. Venturi , P. G. Verdini

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Rome, Italy

P. Barria , M. Campana , F. Cavallari , L. Cunqueiro Mendez , D. Del Re , E. Di Marco , M. Diemoz , F. Errico , E. Longo , P. Meridiani , J. Mijuskovic , G. Organtini , F. Pandolfi , R. Paramatti , C. Quaranta , S. Rahatlou , C. Rovelli , F. Santanastasio , L. Soffi

INFN Sezione di Torino^a, Università di Torino^b, Turin, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane , R. Arcidiacono , S. Argiro , M. Arneodo , N. Bartosik , R. Bellan , A. Bellora , C. Biino , C. Borca , N. Cartiglia , M. Costa , R. Covarelli , N. Demaria , L. Finco , M. Grippo , B. Kiani , F. Legger , F. Luongo , C. Mariotti , L. Markovic , S. Maselli , A. Mecca , E. Migliore , M. Monteno , R. Mulargia , M. M. Obertino , G. Ortona , L. Pacher , N. Pastrone , M. Pelliccioni , M. Ruspa , F. Siviero , V. Sola , A. Solano , A. Staiano , C. Tarricone , D. Trocino , G. Umoret , E. Vlasov

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte , V. Candelise , M. Casarsa , F. Cossutti , K. De Leo , G. Della Ricca

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , C. Huh , B. Kim , D. H. Kim , J. Kim, H. Lee, S. W. Lee , C. S. Moon , Y. D. Oh , M. S. Ryu , S. Sekmen , Y. C. Yang 

Department of Mathematics and Physics, GWNU, Gangneung, Korea

M. S. Kim 

Institute for Universe and Elementary Particles, Chonnam National University, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D. H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , D. Kim , T. J. Kim , J. A. Merlin

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K. S. Lee , S. Lee , J. Park, S. K. Park, J. Yoo 

Department of Physics, Kyung Hee University, Seoul, Korea

J. Goh , S. Yang 

Sejong University, Seoul, Korea

H. S. Kim , Y. Kim, S. Lee

Seoul National University, Seoul, Korea

J. Almond, J. H. Bhyun, J. Choi , W. Jun , J. Kim , S. Ko , H. Kwon , H. Lee , J. Lee , J. Lee , B. H. Oh , S. B. Oh , H. Seo , U. K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D. Y. Kang, Y. Kang , S. Kim , B. Ko, J. S. H. Lee , Y. Lee , I. C. Park , Y. Roh, I. J. Watson 

Department of Physics, Yonsei University, Seoul, Korea

S. Ha , H. D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M. R. Kim , H. Lee, Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy, Y. Maghrbi 

Riga Technical University, Riga, Latvia

K. Dreimanis , A. Gaile , G. Pikurs, A. Potrebko , M. Seidel , V. Veckalns  ⁵⁵

University of Latvia (LU), Riga, Latvia

N. R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

N. Bin Norjoharuddeen , I. Yusuff , Z. Zolkapli  ⁵⁶

Universidad de Sonora (UNISON), Hermosillo, Mexico

J. F. Benitez , A. Castaneda Hernandez , H. A. Encinas Acosta, L. G. Gallegos Maríñez, M. León Coello , J. A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo , I. Heredia-De La Cruz , R. Lopez-Fernandez , C. A. Mondragon Herrera, A. Sánchez Hernández  ⁵⁷

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , I. Pedraza , H. A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, MontenegroI. Bubanja, N. Raicevic **University of Canterbury, Christchurch, New Zealand**P. H. Butler **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M. I. Asghar, A. Awais , M. I. M. Awan, H. R. Hoorani , W. A. Khan **Faculty of Computer Science, Electronics and Telecommunications, AGH University of Krakow, Kraków, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad **Warsaw University of Technology, Warsaw, Poland**K. Pozniak , W. Zabolotny **Laboratório de Instrumentação e Física Experimental de Partículas, Lisbon, Portugal**M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J. W. Wulff**Faculty of Physics, University of Belgrade, Belgrade, Serbia**P. Adzic , P. Milenovic **VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**M. Dordevic , J. Milosevic , V. Rekovic**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**M. Aguilar-Benitez, J. Alcaraz Maestre , Cristina F. Bedoya , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J. P. Fernández Ramos , J. Flix , M. C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J. M. Hernandez , M. I. Josa , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , D. D. Redondo Ferrero , L. Romero, S. Sánchez Navas , L. Urda Gómez , J. Vazquez Escobar , C. Willmott**Universidad Autónoma de Madrid, Madrid, Spain**J. F. de Trocóniz **Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Universidad de Oviedo, Oviedo, Spain**B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J. R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia **Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**S. Bhowmik , S. Blanco Fernández , J. A. Brochero Cifuentes , I. J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasosa García , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J. M. Vizan Garcia **University of Colombo, Colombo, Sri Lanka**M. K. Jayananda , B. Kailasapathy  ⁵⁸, D. U. J. Sonnadara , D. D. C. Wickramarathna **Department of Physics, University of Ruhuna, Matara, Sri Lanka**W. G. D. Dharmaratna , K. Liyanage , N. Perera , N. Wickramage 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , B. Bilin , A. A. Bin Anuar , A. Bocci , C. Botta , E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M. M. Defranchis , M. Deile , M. Dobson , L. Forthomme , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , L. Gouskos , M. Haranko , J. Hegeman , B. Huber, V. Innocente , T. James , P. Janot , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri , M. Mannelli , A. C. Marini , M. Matthewman, F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , H. Qu , D. Rabady , G. Reales Gutierrez, M. Rovere , H. Sakulin , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas ⁶⁰, A. G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , A. Tsirou, D. Walter , J. Wanczyk ⁶¹, J. Wang, S. Wuchterl , P. Zehetner , P. Zejdl , W. D. Zeuner

Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua ⁶², L. Caminada ⁶², A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H. C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶², L. Noehte ⁶², T. Rohe

ETH Zurich-Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T. K. Arrestad , K. Androsov ⁶¹, M. Backhaus , A. Calandri , C. Cazzaniga , K. Datta , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedra , F. Glessgen , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R. A. Manzoni , M. Marchegiani , L. Marchese , C. Martin Perez , A. Mascellani ⁶¹, F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, R. Seidita , J. Steggemann ⁶¹, D. Valsecchi , R. Wallny , C. Amsler ⁶³

Universität Zürich, Zurich, Switzerland

P. Bärtschi , D. Brzhechko, M. F. Canelli , K. Cormier , J. K. Heikkilä, M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S. P. Liechti , A. Macchiolo , P. Meiring , U. Molinatti , A. Reimers , P. Robmann, S. Sanchez Cruz , M. Senger , Y. Takahashi , R. Tramontano

National Central University, Chung-Li, Taiwan

C. Adloff ⁶⁴, D. Bhowmik, C. M. Kuo, W. Lin, P. K. Rout , P. C. Tiwari ³⁸, S. S. Yu

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao , K. F. Chen , P. S. Chen, Z. G. Chen, A. De Iorio , W.-S. Hou , T. H. Hsu, Y. W. Kao, R. Khurana, G. Kole , Y. Y. Li , R.-S. Lu , E. Paganis , X. F. Su , J. Thomas-Wilsker , L. S. Tsai, H. Y. Wu, E. Yazgan

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand

Physics Department, Science and Art Faculty, Çukurova University, Adana, Turkey

D. Agyel , F. Boran , Z. S. Demiroglu , F. Dolek , I. Dumanoglu ⁶⁵, E. Eskut , Y. Guler ⁶⁶, E. Gurpinar Guler ⁶⁶, C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir ⁶⁷, A. Polatoz , B. Tali ⁶⁸, U. G. Tok , S. Turkcapar , E. Uslan , I. S. Zorbakir

Physics Department, Middle East Technical University, Ankara, Turkey

M. Yalvac ⁶⁹

Bogazici University, Istanbul, Turkey

B. Akgun , I. O. Atakisi , E. Gülmез , M. Kaya ⁷⁰, O. Kaya ⁷¹, S. Tekten ⁷²

Istanbul Technical University, Istanbul, Turkey

A. Cakir ^{65,73}, K. Cankocak , Y. Komurcu , S. Sen ⁷⁴

Istanbul University, Istanbul, Turkey

O. Aydilek , S. Cerci ⁶⁸, V. Epshteyn , B. Hacisahinoglu , I. Hos ⁷⁵, B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez

Yildiz Technical University, Istanbul, Turkey

B. Isildak , D. Sunar Cerci

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk

University of Bristol, Bristol, UK

D. Anthony , J. J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H. F. Heath , L. Kreczko , S. Paramesvaran , S. Seif El Nasr-Storey, V. J. Smith , N. Stylianou , K. Walkingshaw Pass, R. White

Rutherford Appleton Laboratory, Didcot, UKA. H. Ball, K. W. Bell , A. Belyaev ⁷⁸, C. Brew , R. M. Brown , D. J. A. Cockerill , C. Cooke , K. V. Ellis, K. Harder , S. Harper , M.-L. Holmberg ⁷⁹, J. Linacre , K. Manolopoulos, D. M. Newbold , E. Olaiya, D. Petyt , T. Reis , G. Salvi , T. Schuh, C. H. Shepherd-Themistocleous , I. R. Tomalin , T. Williams **Imperial College, London, UK**R. Bainbridge , P. Bloch , C. E. Brown , O. Buchmuller, V. Cacchio, C. A. Carrillo Montoya , G. S. Chahal ⁸⁰, D. Colling , J. S. Dancu, I. Das , P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M. H. Hassanshahi , A. Howard, G. Iles , M. Knight , J. Langford , J. León Holgado , L. Lyons , A.-M. Magnan , S. Malik, M. Mieskolainen , J. Nash ⁸¹, M. Pesaresi , B. C. Radburn-Smith , A. Richards, A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G. P. Uttley , L. H. Vage, T. Virdee ³⁰, M. Vojinovic , N. Wardle , D. Winterbottom **Brunel University, Uxbridge, UK**

K. Coldham, J. E. Cole , A. Khan, P. Kyberd , I. D. Reid

Baylor University, Waco, TX, USA

S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , J. Hiltbrand , B. McMaster , M. Saunders , S. Sawant , C. Sutantawibul , J. Wilson

Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , C. Huerta Escamilla, A. E. Simsek , R. Uniyal , A. M. Vargas Hernandez

The University of Alabama, Tuscaloosa, AL, USAB. Bam , R. Chudasama , S. I. Cooper , S. V. Gleyzer , C. U. Perez , P. Rumerio ⁸², E. Usai , R. Yi **Boston University, Boston, MA, USA**

A. Akpinar , D. Arcaro , C. Cosby , Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , S. Yuan , A. G. Zecchinelli

Brown University, Providence, RI, USAG. Benelli , X. Coubez ²⁵, D. Cutts , M. Hadley , U. Heintz , J. M. Hogan ⁸³, T. Kwon , G. Landsberg , K. T. Lau , D. Li , J. Luo , S. Mondal , M. Narain [†], N. Pervan , S. Sagir ⁸⁴, F. Simpson , M. Stamenkovic , W. Y. Wong, X. Yan , W. Zhang**University of California, Davis, Davis, CA, USA**

S. Abbott , J. Bonilla , C. Brainerd , R. Breedon , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P. T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , D. Pellett , W. Wei , Y. Yao , F. Zhang

University of California, Los Angeles, CA, USA

M. Bachtis , R. Cousins , A. Datta , G. Flores Avila, J. Hauser , M. Ignatenko , M. A. Iqbal , T. Lam , E. Manca , A. Nunez Del Prado, D. Saltzberg , V. Valuev

University of California, Riverside, Riverside, CA, USAR. Clare ID, J. W. Gary ID, M. Gordon, G. Hanson ID, W. Si ID, S. Wimpenny [†]**University of California, San Diego, La Jolla, CA, USA**

J. G. Branson ID, S. Cittolin ID, S. Cooperstein ID, D. Diaz ID, J. Duarte ID, L. Giannini ID, J. Guiang ID, R. Kansal ID, V. Krutelyov ID, R. Lee ID, J. Letts ID, M. Masciovecchio ID, F. Mokhtar ID, S. Mukherjee ID, M. Pieri ID, M. Quinnan ID, B. V. Sathia Narayanan ID, V. Sharma ID, M. Tadel ID, E. Vourliotis ID, F. Würthwein ID, Y. Xiang ID, A. Yagil ID

Department of Physics, University of California, Santa Barbara, Santa Barbara, CA, USA

A. Barzdukas ID, L. Brennan ID, C. Campagnari ID, A. Dorsett ID, J. Incandela ID, J. Kim ID, A. J. Li ID, P. Masterson ID, H. Mei ID, J. Richman ID, U. Sarica ID, R. Schmitz ID, F. Setti ID, J. Sheplock ID, D. Stuart ID, T. Á. Vámi ID, S. Wang ID

California Institute of Technology, Pasadena, CA, USA

A. Bornheim ID, O. Cerri, A. Latorre, J. Mao ID, H. B. Newman ID, M. Spiropulu ID, J. R. Vlimant ID, C. Wang ID, S. Xie ID, R. Y. Zhu ID

Carnegie Mellon University, Pittsburgh, PA, USA

J. Alison ID, S. An ID, M. B. Andrews ID, P. Bryant ID, M. Cremonesi, V. Dutta ID, T. Ferguson ID, A. Harilal ID, C. Liu ID, T. Mudholkar ID, S. Murthy ID, P. Palit ID, M. Paulini ID, A. Roberts ID, A. Sanchez ID, W. Terrill ID

University of Colorado Boulder, Boulder, CO, USA

J. P. Cumalat ID, W. T. Ford ID, A. Hart ID, A. Hassani ID, G. Karathanasis ID, E. MacDonald, N. Manganelli ID, A. Perloff ID, C. Savard ID, N. Schonbeck ID, K. Stenson ID, K. A. Ulmer ID, S. R. Wagner ID, N. Zipper ID

Cornell University, Ithaca, NY, USA

J. Alexander ID, S. Bright-Thonney ID, X. Chen ID, D. J. Cranshaw ID, J. Fan ID, X. Fan ID, D. Gadkari ID, S. Hogan ID, P. Kotamnives, J. Monroy ID, M. Oshiro ID, J. R. Patterson ID, J. Reichert ID, M. Reid ID, A. Ryd ID, J. Thom ID, P. Wittich ID, R. Zou ID

Fermi National Accelerator Laboratory, Batavia, IL, USAM. Albrow ID, M. Alyari ID, O. Amram ID, G. Apollinari ID, A. Apresyan ID, L. A. T. Bauerdtick ID, D. Berry ID, J. Berryhill ID, P. C. Bhat ID, K. Burkett ID, J. N. Butler ID, A. Canepa ID, G. B. Cerati ID, H. W. K. Cheung ID, F. Chlebana ID, G. Cummings ID, J. Dickinson ID, I. Dutta ID, V. D. Elvira ID, Y. Feng ID, J. Freeman ID, A. Gandrakota ID, Z. Gecse ID, L. Gray ID, D. Green, A. Grummer ID, S. Grünendahl ID, D. Guerrero ID, O. Gutsche ID, R. M. Harris ID, R. Heller ID, T. C. Herwig ID, J. Hirschauer ID, L. Horyn ID, B. Jayatilaka ID, S. Jindariani ID, M. Johnson ID, U. Joshi ID, T. Klijnsma ID, B. Klma ID, K. H. M. Kwok ID, S. Lammel ID, D. Lincoln ID, R. Lipton ID, T. Liu ID, C. Madrid ID, K. Maeshima ID, C. Mantilla ID, D. Mason ID, P. McBride ID, P. Merkel ID, S. Mrenna ID, S. Nahm ID, J. Ngadiuba ID, D. Noonan ID, V. Papadimitriou ID, N. Pastika ID, K. Pedro ID, C. Pena ID⁸⁵, F. Ravera ID, A. Reinsvold Hall ID⁸⁶, L. Ristori ID, E. Sexton-Kennedy ID, N. Smith ID, A. Soha ID, L. Spiegel ID, S. Stoynev ID, J. Strait ID, L. Taylor ID, S. Tkaczyk ID, N. V. Tran ID, L. Uplegger ID, E. W. Vaandering ID, I. Zoi ID**University of Florida, Gainesville, FL, USA**

C. Aruta ID, P. Avery ID, D. Bourilkov ID, L. Cadamuro ID, P. Chang ID, V. Cherepanov ID, R. D. Field, E. Koenig ID, M. Kolosova ID, J. Konigsberg ID, A. Korytov ID, K. H. Lo, K. Matchev ID, N. Menendez ID, G. Mitselmakher ID, K. Mohrman ID, A. Muthirakalayil Madhu ID, N. Rawal ID, D. Rosenzweig ID, S. Rosenzweig ID, K. Shi ID, J. Wang ID

Florida State University, Tallahassee, FL, USA

T. Adams ID, A. Al Kadhim ID, A. Askew ID, S. Bower ID, R. Habibullah ID, V. Hagopian ID, R. Hashmi ID, R. S. Kim ID, S. Kim ID, T. Kolberg ID, G. Martinez, H. Prosper ID, P. R. Prova, M. Wulansatiti ID, R. Yohay ID, J. Zhang

Florida Institute of Technology, Melbourne, FL, USAB. Alsufyani, M. M. Baarmand ID, S. Butalla ID, T. Elkafrawy ID⁵³, M. Hohlmann ID, R. Kumar Verma ID, M. Rahmani, E. Yanes**University of Illinois Chicago, Chicago, USA**

M. R. Adams ID, A. Baty ID, C. Bennett, R. Cavanaugh ID, R. Escobar Franco ID, O. Evdokimov ID, C. E. Gerber ID, D. J. Hofman ID, J. H. Lee ID, D. S. Lemos ID, A. H. Merrit ID, C. Mills ID, S. Nanda ID, G. Oh ID, B. Ozek ID, D. Pilipovic ID, R. Pradhan ID, T. Roy ID, S. Rudrabhatla ID, M. B. Tonjes ID, N. Varelas ID, Z. Ye ID, J. Yoo ID

The University of Iowa, Iowa City, IA, USA

M. Alhusseini , D. Blend, K. Dilsiz ⁸⁷, L. Emediato , G. Karaman , O. K. Köseyan , J.-P. Merlo, A. Mestvirishvili ⁸⁸, J. Nachtman , O. Neogi, H. Ogul ⁸⁹, Y. Onel , A. Penzo , C. Snyder, E. Tiras ⁹⁰

Johns Hopkins University, Baltimore, MD, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A. V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz

The University of Kansas, Lawrence, KS, USA

A. Abreu , L. F. Alcerro Alcerro , J. Anguiano , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , C. Lindsey, J. Marquez , N. Minafra , M. Murray , M. Nickel , M. Pitt , S. Popescu ⁹¹, C. Rogan , C. Royon , R. Salvatico , S. Sanders , C. Smith , Q. Wang , G. Wilson

Kansas State University, Manhattan, KS, USA

B. Allmond , A. Ivanov , K. Kaadze , A. Kalogeropoulos , D. Kim, Y. Maravin , K. Nam, J. Natoli , D. Roy , G. Sorrentino

Lawrence Livermore National Laboratory, Livermore, CA, USA

F. Rebassoo , D. Wright

University of Maryland, College Park, MD, USA

A. Baden , A. Belloni , Y. M. Chen , S. C. Eno , N. J. Hadley , S. Jabeen , R. G. Kellogg , T. Koeth , Y. Lai , S. Lascio , A. C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M. M. Paranjpe, L. Wang

Massachusetts Institute of Technology, Cambridge, MA, USA

J. Bendavid , I. A. Cali , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grossos, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , L. Lavezzi , Y.-J. Lee , K. Long , C. Mironov , A. Novak , C. Paus , D. Rankin , C. Roland , G. Roland , S. Rothman , G. S. F. Stephanos , Z. Wang , B. Wyslouch , T. J. Yang

University of Minnesota, Minneapolis, MN, USA

B. Crossman , B. M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , S. Pandey , M. Revering , R. Rusack , R. Saradhy , N. Schroeder , N. Strobbe , M. A. Wadud

University of Mississippi, Oxford, MS, USA

L. M. Cremaldi

University of Nebraska-Lincoln, Lincoln, NE, USA

K. Bloom , D. R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , J. E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu

State University of New York at Buffalo, Buffalo, NY, USA

H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams

Northeastern University, Boston, MA, USA

G. Alverson , E. Barberis , J. Dervan, Y. Haddad , Y. Han , A. Krishna , J. Li , M. Lu , G. Madigan , R. McCarthy , D. M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , A. Tishelman-Charny , B. Wang , D. Wood

Northwestern University, Evanston, IL, USA

S. Bhattacharya , J. Bueghly, Z. Chen , S. Dittmer , K. A. Hahn , Y. Liu , Y. Miao , D. G. Monk , M. H. Schmitt , A. Taliercio , M. Velasco

University of Notre Dame, Notre Dame, IN, USA

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K. W. Ho , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. McAlister, T. McCauley , C. McGrady , C. Moore , Y. Musienko ¹⁶, H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygalas

The Ohio State University, Columbus, OH, USA

A. Basnet , B. Bylsma, M. Carrigan , L. S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, B. L. Winer , B. R. Yates 

Princeton University, Princeton, NJ, USA

F. M. Addesa , H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , A. Shevelev , D. Stickland , C. Tully

University of Puerto Rico, Mayagüez, PR, USA

S. Malik 

Purdue University, West Lafayette, IN, USA

A. S. Bakshi , V. E. Barnes , S. Chandra , R. Chawla , S. Das , A. Gu , L. Gutay, M. Jones , A. W. Jung , D. Kondratyev , A. M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer, J. F. Schulte , M. Stojanovic , J. Thieman , A. K. Virdi , F. Wang , W. Xie

Purdue University Northwest, Hammond, IN, USA

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, TX, USA

D. Acosta , T. Carnahan , K. M. Ecklund , P. J. Fernández Manteca , S. Freed, P. Gardner, F. J. M. Geurts , W. Li , O. Miguel Colin , B. P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, NY, USA

A. Bodek , P. de Barbaro , R. Demina , J. L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar, P. Parygin  ⁹², E. Popova  ⁹², R. Taus 

The Rockefeller University, New York, NY, USA

K. Goulianatos 

Rutgers, The State University of New Jersey, Piscataway, NJ, USA

B. Chiarito, J. P. Chou , Y. Gershtein , E. Halkiadakis , M. Heindl , D. Jaroslawski , O. Karacheban  ²⁸, I. Laflotte , A. Lath , R. Montalvo, K. Nash, H. Routray , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S. A. Thayil , S. Thomas, J. Vora , H. Wang

University of Tennessee, Knoxville, TN, USA

H. Acharya, D. Ally , A. G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A. R. Kanuganti , N. Karunaratna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, TX, USA

D. Aebi , M. Ahmad , O. Bouhali  ⁹³, R. Eusebi , J. Gilmore , T. Huang , T. Kamon  ⁹⁴, H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, TX, USA

N. Akchurin , J. Damgov , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane , S. W. Lee , A. Mankel , T. Peltola , I. Volobouev , A. Whitbeck 

Vanderbilt University, Nashville, TN, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, VA, USA

B. Cardwell , B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Neu , C. E. Perez Lara 

Wayne State University, Detroit, MI, USA

P. E. Karchin 

University of Wisconsin-Madison, Madison, WI, USA

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni, H. He ,

M. Herndon , A. Herve , C. K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , D. Pinna, A. Savin, V. Shang , V. Sharma , W. H. Smith , D. Teague, H. F. Tsoi , W. Vetens , A. Warden 

Authors Affiliated with an Institute or an International Laboratory Covered by a Cooperation Agreement with CERN, Geneva, Switzerland

S. Afanasiev , V. Andreev , Yu. Andreev , T. Aushev , M. Azarkin , A. Babaev , A. Belyaev , V. Blinov ⁹⁵, E. Boos , V. Borshch , D. Budkouski , V. Chekhovsky, R. Chistov ⁹⁵, M. Danilov ⁹⁵, A. Dermenev , T. Dimova ⁹⁵, D. Druzhkin ⁹⁶, M. Dubinin ⁸⁵, L. Dudko , A. Ershov , G. Gavrilov , V. Gavrilov , S. Gninenco , V. Golovtcov , N. Golubev , I. Golutvin , I. Gorbunov , A. Gribushin , Y. Ivanov , V. Kachanov , V. Karjavine , A. Karneyeu , V. Kim ⁹⁵, M. Kirakosyan, D. Kirpichnikov , M. Kirsanov , V. Klyukhin , O. Kodolova ⁹⁷, V. Korenkov , A. Kozyrev ⁹⁵, N. Krasnikov , A. Lanev , P. Levchenko ⁹⁸, N. Lychkovskaya , V. Makarenko , A. Malakhov , V. Matveev ⁹⁵, V. Murzin , A. Nikitenko ^{97,99}, S. Obraztsov , V. Oreshkin , V. Palichik , V. Perelygin , S. Petrushanko , S. Polikarpov ⁹⁵, V. Popov , O. Radchenko ⁹⁵, M. Savina , V. Savrin , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen ⁹⁵, S. Slabospitskii , V. Smirnov , A. Snigirev , D. Sosnov , V. Sulimov , E. Tcherniaev , A. Terkulov , O. Teryaev , I. Tlisova , A. Toropin , L. Uvarov , A. Uzunian , A. Vorobyev †, N. Voytishin , B. S. Yuldashev ¹⁰⁰, A. Zarubin , I. Zhizhin , A. Zhokin

[†] Deceased

- 1: Also at Yerevan State University, Yerevan, Armenia
- 2: Also at TU Wien, Vienna, Austria
- 3: Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt
- 4: Also at Ghent University, Ghent, Belgium
- 5: Also at Universidade Estadual de Campinas, Campinas, Brazil
- 6: Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
- 7: Also at UFMS, Nova Andradina, Brazil
- 8: Also at Nanjing Normal University, Nanjing, China
- 9: Now at The University of Iowa, Iowa City, IA, USA
- 10: Also at University of Chinese Academy of Sciences, Beijing, China
- 11: Also at China Center of Advanced Science and Technology, Beijing, China
- 12: Also at University of Chinese Academy of Sciences, Beijing, China
- 13: Also at China Spallation Neutron Source, Dongguan, Guangdong, China
- 14: Now at Henan Normal University, Xinxiang, China
- 15: Also at Université Libre de Bruxelles, Brussels, Belgium
- 16: Also at an Institute or an International Laboratory Covered by a Cooperation Agreement with CERN, Geneva, Switzerland
- 17: Now at British University in Egypt, Cairo, Egypt
- 18: Now at Cairo University, Cairo, Egypt
- 19: Also at Purdue University, West Lafayette, IN, USA
- 20: Also at Université de Haute Alsace, Mulhouse, France
- 21: Also at Department of Physics, Tsinghua University, Beijing, China
- 22: Also at The University of the State of Amazonas, Manaus, Brazil
- 23: Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- 24: Also at University of Hamburg, Hamburg, Germany
- 25: Also at III. Physikalisch Institut A, RWTH Aachen University, Aachen, Germany
- 26: Also at Isfahan University of Technology, Isfahan, Iran
- 27: Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- 28: Also at Brandenburg University of Technology, Cottbus, Germany
- 29: Also at Forschungszentrum Jülich, Jülich, Germany
- 30: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- 31: Also at Institute of Physics, University of Debrecen, Debrecen, Hungary

- 32: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
33: Now at Universitatea Babes-Bolyai-Facultatea de Fizica, Cluj-Napoca, Romania
34: Also at Physics Department, Faculty of Science, Assiut University, Asyût, Egypt
35: Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
36: Also at Punjab Agricultural University, Ludhiana, India
37: Also at University of Visva-Bharati, Santiniketan, India
38: Also at Indian Institute of Science (IISc), Bangalore, India
39: Also at Birla Institute of Technology, Mesra, Mesra, India
40: Also at IIT Bhubaneswar, Bhubaneswar, India
41: Also at Institute of Physics, Bhubaneswar, India
42: Also at University of Hyderabad, Hyderabad, India
43: Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
44: Also at Department of Physics, Isfahan University of Technology, Isfahan, Iran
45: Also at Sharif University of Technology, Tehran, Iran
46: Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
47: Also at Helwan University, Cairo, Egypt
48: Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
49: Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
50: Also at Università degli Studi Guglielmo Marconi, Rome, Italy
51: Also at Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Naples, Italy
52: Also at Fermi National Accelerator Laboratory, Batavia, IL, USA
53: Also at Ain Shams University, Cairo, Egypt
54: Also at Consiglio Nazionale delle Ricerche-Istituto Officina dei Materiali, Perugia, Italy
55: Also at Riga Technical University, Riga, Latvia
56: Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
57: Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
58: Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
59: Also at Saegis Campus, Nugegoda, Sri Lanka
60: Also at National and Kapodistrian University of Athens, Athens, Greece
61: Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
62: Also at Universität Zürich, Zurich, Switzerland
63: Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
64: Also at Laboratoire d’Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
65: Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
66: Also at Konya Technical University, Konya, Turkey
67: Also at Izmir Bakircay University, Izmir, Turkey
68: Also at Adiyaman University, Adiyaman, Turkey
69: Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
70: Also at Marmara University, Istanbul, Turkey
71: Also at Milli Savunma University, Istanbul, Turkey
72: Also at Kafkas University, Kars, Turkey
73: Now at Istanbul Okan University, Istanbul, Turkey
74: Also at Hacettepe University, Ankara, Turkey
75: Also at Faculty of Engineering, Istanbul University-Cerrahpasa, Istanbul, Turkey
76: Also at Yildiz Technical University, Istanbul, Turkey
77: Also at Vrije Universiteit Brussel, Brussels, Belgium
78: Also at School of Physics and Astronomy, University of Southampton, Southampton, UK
79: Also at University of Bristol, Bristol, UK
80: Also at IPPP Durham University, Durham, UK
81: Also at Faculty of Science, Monash University, Clayton, Australia
82: Also at Università di Torino, Turin, Italy
83: Also at Bethel University, St. Paul, MN, USA

- 84: Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
85: Also at California Institute of Technology, Pasadena, CA, USA
86: Also at United States Naval Academy, Annapolis, MD, USA
87: Also at Bingöl University, Bingöl, Turkey
88: Also at Georgian Technical University, Tbilisi, Georgia
89: Also at Sinop University, Sinop, Turkey
90: Also at Erciyes University, Kayseri, Turkey
91: Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
92: Now at an Institute or an International Laboratory Covered by a Cooperation Agreement with CERN,
Geneva, Switzerland
93: Also at Texas A&M University at Qatar, Doha, Qatar
94: Also at Kyungpook National University, Daegu, Korea
95: Also at Another Institute or International Laboratory Covered by a Cooperation Agreement with CERN,
Geneva, Switzerland
96: Also at Universiteit Antwerpen, Antwerp, Belgium
97: Also at Yerevan Physics Institute, Yerevan, Armenia
98: Also at Northeastern University, Boston, MA, USA
99: Also at Imperial College, London, UK
100: Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan