UC Irvine

UC Irvine Previously Published Works

Title

Molecular genetic analysis of human alcohol dehydrogenase.

Permalink

<https://escholarship.org/uc/item/7b9324pt>

Journal

Alcohol (Fayetteville, N.Y.), 2(1)

ISSN

0741-8329

Authors

Duester, G
Hatfield, GW
Smith, M

Publication Date

1985

DOI

10.1016/0741-8329(85)90015-1

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Molecular Genetic Analysis of Human Alcohol Dehydrogenase

GREGG DUESTER,* G. WESLEY HATFIELD* AND MOYRA SMITH*†

*Department of Microbiology and Molecular Genetics and †Department of Pediatrics
College of Medicine, University of California, Irvine, CA 92717

DUESTER, G., G. W. HATFIELD AND M. SMITH. *Molecular genetic analysis of human alcohol dehydrogenase*. ALCOHOL 2(1) 53-56, 1985.—Human alcohol dehydrogenase (ADH) consists of a complex group of isozymes encoded by at least five non-identical genes, two of which have previously been shown through enzymatic analysis to possess polymorphic variants. Using a cDNA probe the *ADH2* gene encoding the β subunit of human ADH was mapped to human chromosome 4. The cDNA probe for *ADH2* was also used to detect a restriction fragment length polymorphism present in human populations. This polymorphism may help establish whether certain ADH allelic variants are linked with certain types of altered alcohol tolerance observed in various individuals. The restriction fragment length polymorphism may also be of use in genetic linkage studies of other genes located near ADH on human chromosome 4.

Chromosome mapping Restriction fragment length polymorphism Gene family cDNA

IN humans three different classes of ADH isozymes have been described. Isozymes of the three classes differ from each other in terms of electrophoretic mobility, substrate specificities, inhibition characteristics and stability [6, 8, 10]. Peptide analyses have revealed that the peptide profiles of Class I ADH (homo- or heterodimers of α , β , and γ subunits), Class II ADH (dimers of π subunits), and Class III ADH (dimers of X subunits) differ significantly from each other while α , β , and γ subunits within Class I have very similar profiles [14]. Also, subunits from one class do not form heterodimers with subunits from the other two classes [6, 8, 10]. Class I ADH isozymes are the products of three gene loci, *ADH1*, *ADH2*, and *ADH3*, which encode the three different polypeptide chains designated α , β , and γ , respectively [10]. Amino acid sequence analysis of human β and γ ADH (H. Jornvall, personal communication), and hybridization analysis of human ADH genomic clones (G. Duester, unpublished data) indicate that the Class I ADH genes (*ADH1*, *ADH2*, and *ADH3*) are similar enough to be considered a gene family. Class II ADH π subunits are encoded by *ADH4* and Class III ADH X subunits by *ADH5* [1]. The relationship of *ADH4* and *ADH5* to the class I ADH genes is unknown.

Studies on the gene products of human Class I ADH genes have revealed genetic polymorphisms. Data on the gene frequencies of polymorphic variants have been derived from studies on post-mortem tissues [11]. Both the *ADH2* and *ADH3* gene loci have been shown to be polymorphic. About 10% of Caucasians express a variant form of *ADH2* which differs in its electrophoretic mobility, pH optimum, stability and kinetic properties from the usual *ADH2* isozyme. This variant form was called atypical ADH [13]. Studies on Oriental populations have revealed that the atypical *ADH2* phenotype occurs in 85% of Japanese [5] and 89% of Chinese individuals [15]. The atypical phenotype is, however, infrequent in the Asian Indian population [15]. It has

been suggested that the incidence of this atypical variant at the *ADH2* locus in Oriental populations parallels the incidence of alcohol intolerance in these populations [16]. In the English population two common *ADH3* alleles have been described. The *ADH3*₁ allele occurs with a frequency of 0.60 while the *ADH3*₂ allele occurs with a frequency of 0.40 [9]. In the Chinese and Japanese populations the frequency of the *ADH3*₁ allele is 0.91 [15].

Genetic studies on Class I ADH and studies aimed at determining the physiological consequences of the different Class I ADH genotypes have been hampered by the fact that these genotypes are not expressed in accessible tissues such as blood cells or cultured fibroblasts. In order to facilitate genetic studies in individuals with altered alcohol tolerance and to define the genomic organization and regulation of the human Class I ADH genes, we have isolated a cDNA probe for the *ADH2* gene [4]. Nucleotide sequencing revealed that this cDNA (pADH12) contained a 593 base pair (bp) 3'-untranslated region in addition to a stretch of 273 nucleotides encoding 91 amino-acids at the carboxy terminal region of β ADH [4]. Here, we have used the β ADH cDNA to analyze the chromosomal location and polymorphism of the β ADH gene.

METHOD

Isolation of Plasmid DNA

Plasmid DNA was isolated by using an alkaline lysis procedure [2] and further purified by CsCl/ethidium bromide density gradient centrifugation. The 1.0 kilobase pair (kb) cDNA insert of pADH12 (Fig. 1) was excised with PstI and isolated by polyacrylamide gel electrophoresis. This cDNA insert was uniformly labeled by incorporation of [α -³²P] dGTP in a nick-translation reaction [7] to a specific activity of approximately 10⁶ cpm per μ g.

FIG. 1. Map of plasmid containing the β ADH cDNA. The plasmid pADH12 was isolated from an adult human liver cDNA library [4]. A cDNA insert of approximately 1.0 kilobase pairs (kb) containing the region of *ADH2* encoding amino acids 283–373 plus a 3'-untranslated region of 593 base pairs is indicated.

Hybridization Analysis of Human Genomic DNA

Genomic DNA was isolated from fresh human leucocytes, cultured Chinese hamster ovary (CHO) cells, and CHO-human hybrid cell lines by methods described elsewhere [7]. DNA was cleaved with either *EcoRI* or *MspI*, subjected to agarose gel electrophoresis, and transferred to nitrocellulose by the method of Southern [12]. A *HindIII* digest of λ bacteriophage DNA provided molecular size markers. Southern blots were hybridized to the β ADH cDNA probe, washed and subjected to autoradiography as described previously [7]. A high stringency wash was performed in $0.1\times$ SSC ($1\times$ SSC is 0.15 M sodium chloride/0.015 M sodium citrate, pH 7.0) at 66°C for 1.0 to 2.5 hours.

RESULTS

Since knowledge of the chromosomal location of the Class I ADH genes will aid human genetic studies on ADH we undertook the mapping of *ADH2* to a specific chromosome. Definitive information on the assignment was obtained using the cDNA probe pADH12 (Fig. 1). Genomic DNA was isolated from fresh human leucocytes, cultured Chinese hamster ovary (CHO) cells, and 2 different CHO-human hybrid cell lines. DNA was restricted with *EcoRI* and then subjected to agarose gel electrophoresis and Southern transfer [12]. Following hybridization to pADH12, nitrocellulose filters were washed under high stringency conditions and then autoradiographed. Using *EcoRI* six fragments were detected in human leucocyte DNA with molecular sizes of 9.6, 7.4, 4.2, 3.2, 2.9, and 2.2 kb. (Fig. 2). The multiple bands

FIG. 2. Southern blots of genomic DNA hybridized to the β ADH cDNA probe (pADH12). All lanes contain 10 μ g of genomic DNA digested with *EcoRI*. Lane 1: human leucocyte, Lane 2: CHO-human hybrid HHW366 (containing only human chromosomes 4 and 5), Lane 3: CHO-human hybrid HHW 416 (containing only human chromosome 4), Lane 4: CHO.

observed in DNA cleaved with *EcoRI* suggest that the β ADH cDNA probe cross-hybridizes with the genes encoding α and γ ADH. Analysis of genomic clones containing human ADH genes has revealed that the six *EcoRI* fragments observed in the human genomic blots are derived from the α , β , and γ ADH genes, each possessing two of the *EcoRI* fragments (G. Duester, unpublished results). Since all six *EcoRI* fragments are accounted for by the three Class I ADH genes, there is no evidence as yet for cross-hybridization of the β ADH cDNA to the Class II or Class III ADH genes, and no evidence for the existence of ADH pseudogenes.

Assignment of the *ADH2* locus to chromosome 4 was obtained from Southern blot analysis of two CHO-human hybrids which contain human chromosome 4. The CHO-human hybrid HHW 366 contains only human chromosomes 4 and 5. This hybrid cell line has locked in human chromo-

FIG. 3. Southern blot to detect polymorphism. Ten μ g of leucocyte genomic DNA from three unrelated individuals (lanes 1 and 2, Caucasian, lane 3 Oriental) was restricted with MspI, and subjected to Southern blot analysis using the β ADH cDNA probe (pADH12). The sizes of the fragments are indicated in kilobase pairs (kb).

some 5 since this chromosome is required to complement a leucyl-tRNA synthetase deficiency in the CHO mutant line thus allowing the hybrid cells to grow under selective conditions (i.e., in leucine deficient medium at 36° C) [3]. By growing the HHW 366 cell line under non-selective conditions and then recloning this cell line, it was possible to derive a CHO-human hybrid HHW 416, which contained only human chromosome 4. It is of interest to note that with a high stringency wash all of the EcoRI fragments detected by pADH12 in human leucocyte genomic DNA could be demonstrated in EcoRI digests of genomic DNA from the HHW 366 and HHW 416 cell lines (Fig. 2). These findings allow us to assign the *ADH2* gene to human chromosome 4. A more detailed description of the mapping of *ADH2* to chromosome 4 using a panel of 20 human-rodent cell lines will be reported elsewhere (M. Smith, manuscript in preparation).

Since genetic linkage studies require the availability of polymorphic chromosomally assigned gene markers, we have undertaken studies to determine whether the pADH12 cDNA probe can be used to detect restriction fragment length polymorphisms in human genomic DNA. Using leucocyte DNA from 3 unrelated individuals we have been able to demonstrate a restriction fragment length polymorphism with the enzyme MspI (Fig. 3). The 8.5 kb MspI fragment is common to all three individuals (lanes 1–3, Fig. 3). On the other hand the 9.6 kb MspI fragment is present only in the two Caucasian DNA samples (lanes 1 and 2, Fig. 3),

being replaced by a 10.3 kb and a 7.4 kb fragment in the Oriental DNA sample (lane 3, Fig. 3). Since the pADH12 probe hybridizes to all three Class I ADH genes, the particular ADH gene involved with the polymorphism is as yet unknown. This issue can be investigated using DNA probes specific for α , β , or γ ADH.

DISCUSSION

The results presented here indicate that the *ADH2* gene is located on chromosome 4. In addition, based upon genomic cloning studies (G. Duester, unpublished results), we conclude that the other two Class I ADH genes *ADH1* and *ADH3* are also located on chromosome 4. Interestingly, studies on the *ADH5* gene encoding Class III ADH indicate that this gene is also located on human chromosome 4 (S. Hiroshige and M. Smith, unpublished data). There is as yet no information on the chromosomal location of the *ADH4* gene encoding Class II ADH. Additional regional mapping and genomic cloning studies will be required to determine if all the Class I, II, and III ADH genes form a gene family, or whether only certain members therein, such as the three Class I genes, form a gene family.

An important finding here is that the β ADH cDNA is useful probe for detecting a restriction fragment length polymorphism on human chromosome 4. Interestingly, the two Caucasian DNA samples exhibited a common hybridization pattern, whereas the Oriental sample was of a different pattern. More individuals, both Caucasian and Oriental, will of course need to be examined, but it is interesting to speculate that the restriction fragment length polymorphism observed here might correlate to either the β_1/β_2 or γ_1/γ_2 polymorphisms that were discussed above. This question can be approached using DNA probes specific for particular ADH genes, such as DNA fragments that contain only 3'-untranslated regions or intron sequences that do not exhibit cross-hybridization to other ADH genes. In conclusion, DNA fragments derived from human ADH cDNA and genomic clones constitute useful probes for detecting polymorphisms on chromosome 4, and may facilitate genetic studies of various alcohol-related syndromes.

ACKNOWLEDGEMENTS

We wish to thank Drs. Leon Carlock and John Wasmuth for providing the HHW 366 and HHW 416 cell lines. This work was supported by research grants to M.S. from the National Institute of Alcohol Abuse and Alcoholism (AA-05781) and the March of Dimes (1-811), and to G.W.H. from the National Institute of General Medical Sciences (GM-32112). G.D. was supported by a National Institutes of Health Postdoctoral Fellowship (GM-08733).

REFERENCES

- Adinolfi, A., M. Adinolfi and D. A. Hopkinson. Immunological and biochemical characterization of the human alcohol dehydrogenase X-ADH isozyme. *Ann Hum Gen* 48: 1–10, 1984.
- Birnboim, H. C. and J. Doly. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* 7: 1513–1523, 1979.
- Dana, S. and J. J. Wasmuth. Selective linkage disruption in human-Chinese hamster cell hybrids: Deletion mapping of the *leuS*, *hexB*, *emtB*, and *chr* genes on human chromosome 5. *Mol Cell Biol* 2: 1220–1228, 1982.
- Duester, G., G. W. Hatfield, R. Buhler, J. Hempel, H. Jornvall and M. Smith. Molecular cloning and characterization of a c-DNA for the β subunit of human alcohol dehydrogenase. *Proc Natl Acad Sci* 81: 4055–4059, 1984.
- Harada, S., S. Misawa, D. P. Agarwal and H. W. Goede. Liver alcohol dehydrogenase and aldehyde dehydrogenases in the Japanese. *Am J Hum Gen* 32: 8–15, 1980.
- Li, T.-K. and L. J. Magnes. Identification of a distinctive molecular form of alcohol dehydrogenase in livers with high activity. *Biochem Biophys Res Commun* 63: 202–208, 1975.

7. Maniatis, T., E. F. Fritsch and J. Sambrook, *Molecular Cloning*. Cold Spring Harbor: Cold Spring Harbor Laboratory, 1982.
8. Pares, X. and B. L. Vallee. New human liver alcohol dehydrogenase forms with unique kinetic characteristics. *Biochem Biophys Res Commun* **98**: 122-130, 1981.
9. Smith, M. The genetics of alcoholism. In *Advances in Alcohol and Substance Abuse*, vol 1, edited by B. Stimmel. New York: Haworth Press, 1982, pp. 127-146.
10. Smith M. and D. A. Hopkinson. Studies on the subunit structure and molecular size of the human alcohol dehydrogenase isozymes determined by the different loci ADH1, ADH2, ADH3. *Ann Hum Genet* **36**: 401-414, 1973.
11. Smith, M., D. A. Hopkinson and H. Harris. Alcohol dehydrogenase isozymes in adult human stomach and liver: evidence for activity of the ADH3 locus. *Ann Hum Genet* **35**: 243-253, 1972.
12. Southern, E. M. Detection of specific sequences among DNA fragments separated by gel electrophoreses. *J Mol Biol* **98**: 503-518, 1975.
13. Stamatoyannopoulos, G., S. H. Chen and M. Fukui. Liver alcohol dehydrogenase: High population frequency of atypical form and its possible role in alcohol sensitivity. *Am J Hum Genet* **27**: 789-796, 1975.
14. Strydom, D. J. and B. L. Vallee. Characterization of human alcohol dehydrogenase isoenzymes by high-performance liquid chromatographic peptide mapping. *Anal Biochem* **123**: 422-429, 1982.
15. Teng, Y. S., S. Jehan and L. E. Lie-Injo. Human Alcohol dehydrogenase ADH2 and ADH3 polymorphisms in ethnic Chinese and Indians of West Malaysia. *Hum Genet* **53**: 87-90, 1979.
16. Yoshida, A., C. C. Impraim and I.-Y. Huang. Enzymatic and structural differences between usual and atypical human liver alcohol dehydrogenases. *J Biol Chem* **256**: 12430-12436, 1981.