

# UCSF

## UC San Francisco Previously Published Works

### Title

Bringing population mobility into focus to achieve HIV prevention goals.

### Permalink

<https://escholarship.org/uc/item/7bf066cx>

### Journal

Journal of the International AIDS Society, 21 Suppl 4(Suppl Suppl 4)

### ISSN

1758-2652

### Authors

Camlin, Carol S  
Cassels, Susan  
Seeley, Janet

### Publication Date

2018-07-01

### DOI

10.1002/jia2.25136

Peer reviewed

## EDITORIAL

# Bringing population mobility into focus to achieve HIV prevention goals

Carol S Camlin<sup>1§</sup> , Susan Cassels<sup>2</sup>  and Janet Seeley<sup>3</sup> 

§Corresponding author: Carol S. Camlin, Department of Obstetrics, Gynecology & Reproductive Sciences, University of California, San Francisco, ANSIRH Program, 1330 Broadway, Suite 1100, Oakland, CA, USA 94612. Tel:+1(510) 986 8981 ([carol.camlin@ucsf.edu](mailto:carol.camlin@ucsf.edu))

**Keywords:** Population mobility; Migration; Universal testing and treatment; Structural drivers; Prevention; Key and vulnerable populations

Received 18 May 2018; Accepted 22 May 2018

**Copyright** © 2018 The Authors. *Journal of the International AIDS Society* published by John Wiley & sons Ltd on behalf of the International AIDS Society. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

The ambitious UNAIDS ‘90-90-90’ targets, aiming to end the AIDS epidemic by the year 2030 [1], are a response to the profound therapeutic and preventive benefits of HIV treatment. Universal voluntary HIV counselling and testing followed by prompt initiation of antiretroviral therapy (ART) for all those diagnosed HIV-infected, an approach known as Universal Test and Treat (UTT), is now seen as the primary means through which the 90-90-90 targets can be achieved. Beyond the targets, the recently emergent concept of a prevention cascade [2] also recognizes the imperative of “coverage” for achieving population-level effects of HIV prevention interventions, which are needed together with the expansion of treatment to end the epidemic [3].

Perhaps nowhere else in the world is more momentum required to bring HIV treatment *and* prevention to all who need it than in Africa. The HIV epidemic on the continent has been concentrated in eastern and southern Africa, where an estimated 19.4 million adults are living with HIV [4]. Despite substantial progress in recent years, only approximately 60% of people living with HIV (PLHIV) in the region were receiving ART in 2016 [4]; an estimated 45% had a suppressed viral load as a result of their successful engagement in care in 2015 [5].

These estimates reflect tremendous successes of HIV prevention and care initiatives towards accelerating ART coverage for sub-Saharan African populations, and many successful efforts to reach key populations—yet also the sobering reality that what has been achieved to date has been achieved in easier to reach populations. A focus by UNAIDS and global leadership on harder to reach, key populations, that are particularly important for further scale-up, is laudable; but to date, inadequate attention has been paid to population mobility, a major force that challenges the HIV care cascade and threatens the promise of “treatment as prevention” particularly in high HIV prevalence, resource-limited regions. Ultimately, strategies to attain the 90-90-90 targets that do not account

for the complex dynamics of mobility in specific settings will fail to engage successfully with the magnitude of populations necessary to end the epidemic [6].

There is an urgent need to deepen our understanding of population mobility, its heterogeneous forms and gender dimensions in high HIV prevalence areas, its effects on sexual behaviour and sexual networks, HIV testing and engagement in HIV care and treatment, and on HIV acquisition and onward transmission, even as widespread HIV prevention interventions are underway—indeed, *especially* in contexts in which interventions such as UTT are underway. To understand whether and how mobility attenuates the effectiveness of HIV prevention and care initiatives is a vital first step towards identifying the full set of solutions, inclusive of mobile populations, that will be needed to end the epidemic.

Historically, the spread of HIV in sub-Saharan Africa followed the corridors of population movement, as people moved to expanding urban areas and other places of employment opportunity [7-10]. Today, mobility continues to place individuals at great risk of HIV acquisition as well as onward transmission [11-13]. The highly gendered nature of that linkage is well-documented, yet rarely commented upon. Forms of mobility in sub-Saharan Africa are diverse and complex relative to those of other regions, and men’s and women’s patterns of movement significantly differ, with women increasingly participating in mobility [14-18], as has been seen elsewhere [19-22]. While transcontinental migration from Africa is low [23], the intra-sub-Saharan African emigration rate represents the largest south to south movement of people in the world [24]. Data for the study of internal migration have been limited; however, data from Demographic Surveillance Sites show that 7% to 20% of local populations in surveillance areas, often over 30% of young adults, have migrated annually in recent years [14,25]. Rural to urban migration flows do not predominate in all settings [26]; rather, counter-urbanization [27] and circulation between rural areas, semi-urban towns, and rural

perimeters of cities are common [14]. Mobility has positive benefits for development and is a key driver of economic growth in southern and eastern Africa: from 50% to 80% of rural households have at least one migrant member [24], and the “sending” of a female migrant has been particularly advantageous to the poorest households [23,28,29].

Despite its importance, research on the effects of mobility in sub-Saharan Africa on HIV care engagement, and on effectiveness of ART as treatment and as prevention, is in its nascent. Existing literature mostly from the global north has shown that mobility can break bonds between individuals and HIV care systems [30], thus leading to disengagement from care, treatment interruption and poor health outcomes. Research on mobility and care engagement in sub-Saharan African settings is relatively scarce; thus there are key questions about the extent to which mobility is contributing to the gap that exists between the remarkable promise of ART for large-scale prevention [31–33], and actual engagement in care [34].

Recent meta-analyses in sub-Saharan African populations have shown that 30% to 60% of people living with HIV are lost to follow-up at each step after HIV diagnosis [35–37]. People repeatedly exit and re-enter the care cascade at various points [38,39]. Greater attention to the contextual factors that shape HIV treatment experiences, engagement and outcomes has recently emerged in social science research on HIV [40], revealing the complex social realities underlying “bottle-necks” in the cascade [40]. Yet data have been limited for investigating the role of mobility in these care cascade shortfalls.

Several meta-analyses and systematic reviews of the literature examining factors associated with care entry, engagement and retention [35,37,41–43] show that optimal lifelong engagement in HIV care can be threatened by a range of factors at the individual, social, and structural levels. Mobility affects many of the factors found to contribute to delayed entry or lapses in care, including psychological factors (e.g. seeking care away from home because of stigma) [39,44,45], clinic characteristics (e.g. waiting times) [46,47] and structural barriers [44,48] such as distance to clinic and transportation costs [44,47,49,50]. Yet the direct impact of mobility, and the pathways for this impact, have not been examined in depth [30,51]. Better measures and methods are needed to understand how mobility affects the ability of individuals to successfully navigate the HIV care cascade. This is especially urgent for sub-Saharan Africa, where levels of mobility are high and forms of mobility are complex and dynamic. The effects of mobility on initiatives designed to extend the reach of primary HIV prevention interventions in these settings are, to date, unknown—a glaring gap in the literature.

The articles in this supplement of the *Journal of the International AIDS Society* focus on crucial questions regarding the forms and dimensions of mobility and the impact on HIV prevention and care initiatives—particularly in Southern and Eastern Africa, but also in the ‘receiving’ nations in the global north of sub-Saharan African migrants. This volume includes contributions from each of the large-scale UTT studies in five African countries [52], which include HPTN 071 Population Effects of Antiretroviral Therapy to Reduce HIV Transmission (PopART), Sustainable East Africa Research in Community Health (SEARCH); the MaxART (Early Access to ART for All)

implementation study; and the ANRS 12249 Antiretroviral Treatment as Prevention (TasP) trial. It also includes contributions from studies outside UTT trial settings on the impacts of mobility on care engagement. The authors of the papers use different methods and approaches to identify emergent themes, key problems and gaps in the study of mobility and HIV prevention and care and point to potential solutions. The collection includes quantitative research that advances approaches towards measurement of diverse forms of mobility and its impacts, and also highlights the value of qualitative research for understanding the meaning of factors driving linkage to HIV treatment and prevention in the context of migration and mobility.

We observe four major themes across the contributions to this volume: first, the spatial *and* temporal dimensions of mobility are important, because mobility is inherently linked to time exposed to interventions and services within geographies; secondly, types of mobility in different populations and settings are diverse and highly gendered, and this heterogeneity across contexts should inform HIV prevention and treatment approaches in specific settings; thirdly, while mobility is intertwined with other factors that affect engagement in HIV care and prevention, it often sets into motion a “chain of events” that leads to disengagement; and fourth, the period following resettlement in new destinations is one of instability, in which risk is heightened, disrupting engagement and leading to ‘missed opportunities’ for progress along the prevention and care cascades. Such missed opportunities must be addressed through health systems- and policy-level actions. In sum, this collection speaks to a need for reconfiguration of programs and services to respond to the challenges to HIV care and prevention engagement that are presented by the mobility of populations.

Speaking to the dynamic nature of mobility and the importance of temporality, Larmarange and colleagues [53] present findings from the ANRS 12249 Antiretroviral Treatment as Prevention (TasP) trial, that highlight the effect of population mobility on temporality of and geographic exposure to the UTT intervention in rural KwaZulu-Natal, South Africa. The TasP cluster-randomized trial failed to show a reduction in HIV incidence at population level when ART was proposed regardless of CD4 count. The authors describe a dynamic cascade of various degrees of exposure to the trial interventions, using both calendar (population) and exposure (individual) time approaches in their analyses, and show that the structural effects of mobility diluted the impact of the UTT strategy. In a context of high HIV incidence, the circulation of newly infected individuals in and out of communities slowed down TasP efforts to increase ART coverage and population viral suppression, ultimately attenuating any population-level impact on HIV incidence.

In their study of population mobility embedded within the SEARCH trial in rural populations in Kenya and Uganda, Camlin and colleagues [54] describe significant heterogeneity in mobility across regions in Kenya and Uganda, which correlated with heterogeneous levels of risk behaviour and HIV prevalence observed across the regions: communities with higher proportions of mobile residents tended to also have higher HIV prevalence. Mobility may undermine UTT interventions in high prevalence areas if mobile populations are spending time exposed outside of their communities and indeed they were

as follows: both migration and also localized short-term mobility were associated with higher risk sexual behaviour, especially among women. Livelihoods requiring mobility for women, such as market trading, and the fish trade in particular, were strongly associated with their higher risk sexual behaviour. In contrast, men's labour-related mobility was not associated with higher risk behaviour; their travel for other purposes (e.g. attending funerals, seeking care, visiting family), was. This work highlights the need for gender-specific interventions among mobile populations.

The links between mobility and livelihoods that affect engagement in HIV care and prevention services and interventions also figure prominently in two contributions from the PopART trial. Bond and colleagues [55] used longitudinal qualitative data focused on the lives of six people living with HIV in urban Zambia to show the challenges of juggling household responsibility, livelihood mobility and HIV management. For five of the six, ongoing engagement in treatment could not be sustained because of travel. The authors highlight the need for differential care options which can adjust more to clients' temporal and spatial realities. Hoddinott and colleagues [56] used ethnographic and participatory research to describe patterns of 'household fluidity' in the Western Cape of South Africa, and explored how movements in and out of households to ensure livelihoods and social support shape HIV service access. With their conceptualization of 'fluidity', the authors challenge conventional sociological concepts of households, that is, they are not static, contained or bounded as the typical measures would suggest. Echoing the recommendations of Bond and colleagues, the authors call for "responsive, flexible health service delivery systems designed to support continuity in care across many shifts in client circumstances."

Shabalala and colleagues [57] present findings from a mixed methods study to improve understanding of care retention among ART clients in the MaxART implementation study in Swaziland. The authors examined associations between socio-demographic characteristics and retention, and conducted in-depth interviews with clients who were lost-to-follow-up from the study to explore their reasons for leaving care. Mobility, particularly relocations to another community far from the facility where the client initially obtained ART, was described by study participants as the first step in a complex chain of events that affected retention in care. Several often-intersecting reasons for discontinuing ART also mattered, including "harsh treatment by health care workers", ART side effects, stigma, and food insecurity. These findings provide further evidence in support of Ware's conceptualization of care disengagement as a process through which missed visits, and ensuing reluctance to return, can erode patients' feelings of connectedness to care over time [39]. They also expand upon this conceptualization by documenting how migration events can trigger the chain of events that leads to disengagement.

Expanding upon this theme, studies conducted outside of UTT trial contexts highlight how the period following resettlement in new destinations is a period of instability in which behavioural risks of HIV acquisition are heightened, and engagement in care and prevention is disrupted, leading to missed opportunities for improving the prevention and care cascades. Fakoya and colleagues [58] present findings from a study of care engagement among migrants living with HIV in

Europe. Using data from 57 HIV clinics in nine countries, the authors investigated the barriers and facilitators to HIV testing, and assessed the current treatment and healthcare needs of migrants living with HIV in Europe. Noting that immigrants are overrepresented in the European HIV epidemic, the authors present evidence suggesting that exposure to HIV after migrating accounts for a substantial proportion of infections among migrants, and that opportunities for HIV prevention are being missed. They call for a greater attention to the HIV prevention needs of immigrants to Europe, including interventions to reduce HIV acquisition among migrant gay and bisexual men, and to expand HIV testing opportunities for migrant heterosexual men and women.

Two articles in this volume examine women's migration in the context of pregnancy. Clouse and colleagues [59] present findings from a mixed methods study to identify drivers and types of mobility among pregnant and postpartum women living with HIV in the Johannesburg area in South Africa. They examine long-distance travel of mothers and infants before and after delivery, finding that the frequent mobility in the peripartum period "underscores the challenge of ensuring a continuity of HIV care in a fragmented health care system that is not adapted for a mobile population." In another setting in South Africa (Cape Town), Phillips and colleagues [60] also examined the mobility of women attending an integrated antenatal-ART clinic. Using routine electronic health data (including laboratory testing, ART dispensing and clinic visits) to measure the movements and access points of pregnancy and postpartum women, the authors show that a substantial proportion of women do not link to postpartum care, and among those who do, long-term retention remains a challenge as women move to a wide variety of facilities locally and nationally.

Vearey [61] offers a Commentary that speaks to a need for improving the policy environment to address better the HIV prevention and care needs of mobile populations. She explores challenges and strategic opportunities for "re-setting" the policy agenda on migration and HIV in southern Africa, specifically in the policy environment of the Southern African Development Community (SADC) – a region associated with high levels of migration, and home to the largest population of people living with HIV globally. Drawing upon policy review, empirical data, and on-going participant observation within local, regional and global policy processes, Vearey shows how current policy processes have the potential to undermine efforts to improve the global responses to migration and HIV. She argues that, "without mainstreaming migration, HIV programmes will continue to struggle, and key health targets will not be met."

Lastly, in our Viewpoint, the editors have expanded on two specific organizing concepts that have emerged from our work on this volume: the time scales of mobility, and migrants' sexual networks. Temporal scales of population mobility are complex, sexual risk behaviour can change in relation to the timing of migration, and timing of migration can interact with timing of, and exposure to, HIV prevention efforts. Additionally, the timing, sequence and spatial scale of migrant's sexual networks can mitigate intervention effectiveness. Cassels, Camlin and Seeley [62] argue that focusing on temporal patterns of mobility and network characteristics will not only help to explain why population mobility presents a challenge for HIV

prevention and care, but can be leveraged to improve future interventions.

The collection of articles in this volume contribute to framing a future policy and research agenda. Treatment access is a key area of concern: how can migrants sustain their care in the myriad destinations to which they travel? What interventions, policies, or health systems improvements are needed to maximize the engagement of mobile individuals in HIV care? Recognising the need for approaches that are responsive to diversity, that risk and opportunities for care may be affected by gender, age, sexual identity and ethnicity, is essential if hard to reach populations are to be reached by UTT. What possibilities exist for reconceptualising care delivery to address these challenges?

Mobile populations may be among those who stand to benefit the most from new models of differentiated care or differentiated service delivery, which aim to simplify and adapt HIV services across the cascade to better meet the needs of PLHIV and reduce burdens on health systems [63]. These models include patient-led community adherence groups, healthcare worker-managed groups known as adherence clubs, fast-track or multi-month scripting, mobile outreach, and community drug distribution points. To the extent that these models can be informed by an understanding of the needs of mobile women and men living with HIV, they hold promise for engaging and retaining these populations who struggle to fit their needs to the requirements of clinic-based HIV care systems.

In addition, mobile populations stand to benefit from improved therapeutic technologies such as long-acting ART as well as longer-acting formulations of biomedical prevention technologies such as Pre-Exposure Prophylaxis (PrEP), and expansion of the delivery of these technologies beyond clinic settings into communities and key migration destinations and transit hubs. Structural and behavioural interventions to facilitate demand are needed to complement these 'supply side' interventions [2]. Without such innovations, migrants will continue to be left behind in the quest to end the AIDS epidemic. We hope that this collection stimulates focus and commitment towards meeting this critical public health challenge.

## AUTHORS' AFFILIATIONS

<sup>1</sup>Department of Obstetrics, Gynecology and Reproductive Sciences, Department of Medicine, University of California, San Francisco, USA; <sup>2</sup>Department of Geography, University of California, Santa Barbara, USA; <sup>3</sup>Department of Global Health and Development, London School of Hygiene and Tropical Medicine, London, UK

## COMPETING INTERESTS

The authors have no competing interests to declare.

## AUTHORS' CONTRIBUTIONS

CC, SC, and JS conceptualized the main messages. CC drafted the manuscript; SC and JS provided critical reviews. All authors read and approved of the final manuscript.

## REFERENCES

1. UNAIDS. 90-90-90: an ambitious treatment target to help end the AIDS epidemic. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS (UNAIDS); 2014.
2. Hargreaves JR, Delany-Moretwe S, Hallett TB, Johnson S, Kapiga S, Bhat-tacharjee P, et al. The HIV prevention cascade: integrating theories of

- epidemiological, behavioural, and social science into programme design and monitoring. *Lancet HIV*. 2016;3(7):e318–22.
3. Isbell MT, Kilonzo N, Mugurungi O, Bekker L-G. We neglect primary HIV prevention at our peril. *Lancet HIV*. 2016;3(7):e284–5.
4. UNAIDS. Fact sheet - Latest global and regional statistics on the status of the AIDS epidemic. Geneva: UNAIDS; 2017.
5. UNAIDS. Prevention Gap Report. Geneva: UNAIDS; 2016.
6. Akullian A, Bershteyn A, Jewell B, Camlin CS. The missing 27%. *AIDS*. 2017;31(17):2427–9.
7. Glynn JR, Ponnighaus J, Crampin AC, Sibande F, Sichali L, Nkhosa P, et al. The development of the HIV epidemic in Karonga District, Malawi. *AIDS*. 2001;15(15):2025–9.
8. Jochelson K, Mothibeli M, Leger JP. Human immunodeficiency virus and migrant labor in South Africa. *Int J Health Serv*. 1991;21(1):157–73.
9. Garin B, Jeannel D, Kazadi K, Combe P, Singa L, De The G. Introduction of HIV-1 in a rural city of Zaire. *Ann Soc Belg Med Trop*. 1993;73(2):143–7.
10. Coffee MP, Garnett GP, Mlilo M, Voeten HA, Chandiwana S, Gregson S. Patterns of movement and risk of HIV infection in rural Zimbabwe. *J Infect Dis*. 2005;191(Suppl 1):S159–67.
11. Olawore O, Tobian AAR, Kagaayi J, Bazaale JM, Nantume B, Kigozi G, et al. Migration and risk of HIV acquisition in Rakai, Uganda: a population-based cohort study. *Lancet HIV*. 2018;5(4):e181–9.
12. McGrath N, Eaton JW, Newell ML, Hosegood V. Migration, sexual behaviour, and HIV risk: a general population cohort in rural South Africa. *Lancet HIV*. 2015;2(6):e252–9.
13. Cassels S, Jenness SM, Khanna AS. Conceptual framework and research methods for migration and HIV transmission dynamics. *AIDS Behav*. 2014;18(12):2302–13.
14. Camlin CS, Snow RC, Hosegood V. Gendered patterns of migration in Rural South Africa. *Popul Space Place*. 2014;20(6):528–51.
15. Agesa RU, Agesa J. Sources of gender difference in rural to urban migration in Kenya: does human capital matter? *Appl Econ Lett*. 2005;12(11):705–9.
16. Reed HE, Andrzejewski CS, White MJ. Men's and women's migration in coastal Ghana: an event history analysis. *Demogr Res*. 2010;22:771–812.
17. Thadani VN. Social Relations and Geographic Mobility: Male and Female Migration in Kenya. New York: Center for Policy Studies, The Population Council; 1982. Contract No.: 85.
18. Posel D, Casale D. What has been happening to internal labour migration in South Africa, 1993-1999? *South Afr J Econ*. 2003;71(3):455–79.
19. Curran SR, Rivero-Fuentes E. Engendering migrant networks: the case of Mexican migration. *Demography*. 2003;40(2):289–307.
20. Bilsborrow RE. Preliminary report of the United Nations expert group meeting on the feminization of internal migration. *Int Migr Rev*. 1992;26(1):138–61.
21. Gaetano AM, Jacka T (eds). On the Move: women in Rural-to-Urban Migration in Contemporary China. New York: Columbia University Press; 2004.
22. Zlotnick H. The Global Dimensions of Female Migration. Washington, D.C.: Migration Policy Institute; 2003 [Available from: <http://www.migrationinformation.org/Feature/display.cfm?ID=109>].
23. Zlotnick H. The Dimensions of Migration in Africa. In: Africa on the move: African migration and urbanisation in comparative perspective. Tienda M, Findley S, Tollman S, Preston-Whyte E (eds). Johannesburg: Wits University Press; 2006.
24. IOM. Internal Migration and Development: a Global Perspective. Geneva, Switzerland: International Organization for Migration; 2005. Contract No.: 19.
25. Collinson M. Age-Sex Profiles of Migration: who is a Migrant? In: The dynamics of migration, health and livelihoods: INDEPTH network perspectives. Collinson MA, Adazu K, White M, Findley S (eds). Surrey, England: Ashgate; 2009.
26. Potts D. Whatever happened to Africa's rapid urbanisation? *World Economics*. 2012;13(2):17–29.
27. Potts D. Counter-urbanisation on the Zambian copperbelt? interpretations and implications. *Urban Stud*. 2005;42(4):583–609.
28. Collinson M, Gerritsen A, Clark S, Kahn K, Tollman S. Migration and socio-economic change in rural South Africa, 2000-2007. In: The Dynamics of Migration, Health and Livelihoods: INDEPTH Network Perspectives. Collinson MA, Adazu K, White M, Findley S (eds). Surrey, England: Ashgate; 2009.
29. Kok P, Gelderblom D, Oucho JO, van Zyl J (eds). Migration in South and Southern Africa: dynamics and Determinants. Human Sciences Research Council: Cape Town; 2006.
30. Taylor BS, Garduno LS, Reyes EV, Valino R, Rojas R, Donastorg Y, et al. HIV care for geographically mobile populations. *Mt Sinai J Med*. 2011;78(3):342–51.
31. Granich RM, Gilks CF, Dye C, De Cock KM, Williams BG. Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: a mathematical model. *Lancet*. 2009;373(9657):48–57.

32. Granich R, Gupta S, Suthar AB, Smyth C, Hoos D, Vitoria M, et al. Antiretroviral therapy in prevention of HIV and TB: Update on current research efforts. *Curr HIV Res.* **2011**;9(6):446–69.
33. Charlebois ED, Havlir DV. "A bird in the hand...": a commentary on the test and treat approach for HIV. *Arch Intern Med.* **2010**;170(15):1354–6.
34. Gardner EM, McLees MP, Steiner JF, del Rio C, Burman WJ. The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. *Clin Infect Dis.* **2011**;52: 793–800. (6%U <http://cid.oxfordjournals.org/content/52/6/793.abstract>):
35. Rosen S, Fox MP, Gill CJ. Patient retention in antiretroviral therapy programs in sub-Saharan Africa: a systematic review. *PLoS Med.* **2007**;4(10):e298.
36. Kranzer K, Govindasamy D, Ford N, Johnston V, Lawn SD. Quantifying and addressing losses along the continuum of care for people living with HIV infection in sub-Saharan Africa: a systematic review. *J Int AIDS Soc.* **2012**;15(2):17383.
37. Mugglin C, Estill J, Wandeler G, Bender N, Egger M, Gsponer T, et al. Loss to programme between HIV diagnosis and initiation of antiretroviral therapy in sub-Saharan Africa: systematic review and meta-analysis. *Trop Med Int Health.* **2012**;17(12):1509–20.
38. Hallett TB, Eaton JW. A side door into care cascade for HIV-infected patients? *J Acquir Immune Defic Syndr.* **2013**;63(Suppl 2):S228–32.
39. Ware NC, Wyatt MA, Geng EH, Kaaya SF, Agbaji OO, Muyindike WR, et al. Toward an understanding of disengagement from HIV treatment and care in sub-Saharan Africa: a qualitative study. *PLoS Med.* **2013**;10(1):e1001369; discussion e.
40. Skovdal M, Wringe A, Seeley J, Renju J, Papanini S, Wamoyi J, et al. Using theories of practice to understand HIV-positive persons varied engagement with HIV services: a qualitative study in six Sub-Saharan African countries. *Sex Transm Infect.* **2017**;93(Suppl 3):pii: e052977. <https://doi.org/10.1136/sextrans-2016-052977>.
41. Rosen S, Fox MP. Retention in HIV care between testing and treatment in sub-Saharan Africa: a systematic review. *PLoS Med.* **2011**;8(7):e1001056.
42. Geng EH, Bangsberg DR, Musinguzi N, Emenyonu N, Bwana MB, Yiannoutsos CT, et al. Understanding reasons for and outcomes of patients lost to follow-up in antiretroviral therapy programs in Africa through a sampling-based approach. *J Acquir Immune Defic Syndr.* **2010**;53(3):405–11.
43. Geng EH, Nash D, Kambugu A, Zhang Y, Braitstein P, Christopoulos KA, et al. Retention in care among HIV-infected patients in resource-limited settings: emerging insights and new directions. *Curr HIV/AIDS Rep.* **2010**;7(4):234–44.
44. Coetzee B, Kagee A, Vermeulen N. Structural barriers to adherence to antiretroviral therapy in a resource-constrained setting: the perspectives of health care providers. *AIDS Care.* **2011**;23(2):146–51.
45. Murray LK, Semrau K, McCurley E, Thea DM, Scott N, Mwiya M, et al. Barriers to acceptance and adherence of antiretroviral therapy in urban Zambian women: a qualitative study. *AIDS Care.* **2009**;21(1):78–86.
46. Lubega M, Nsabagasani X, Tumwesigye NM, Wabwire-Mangen F, Ekstrom AM, Pariyo G, et al. Policy and practice, lost in transition: Reasons for high drop-out from pre-antiretroviral care in a resource-poor setting of eastern Uganda. *Health Policy.* **2010**;95(2–3):153–8.
47. Hardon AP, Akurut D, Comoro C, Ekezie C, Irunde HF, Gerrits T, et al. Hunger, waiting time and transport costs: time to confront challenges to ART adherence in Africa. *AIDS Care.* **2007**;19(5):658–65.
48. Kagee A, Remien RH, Berkman A, Hoffman S, Campos L, Swartz L. Structural barriers to ART adherence in Southern Africa: challenges and potential ways forward. *Global Public Health.* **2011**;6(1):83–97.
49. Tuller DM, Bangsberg DR, Senkungu J, Ware NC, Emenyonu N, Weiser SD. Transportation costs impede sustained adherence and access to HAART in a clinic population in southwestern Uganda: a qualitative study. *AIDS Behav.* **2010**;14(4):778–84.
50. Lankowski AJ, Siedner MJ, Bangsberg DR, Tsai AC. Impact of geographic and transportation-related barriers on HIV outcomes in sub-saharan Africa: a systematic review. *AIDS Behav.* **2014**;18(7):1199–223. <https://doi.org/10.1007/s10461-014-0729-8>. Review.
51. Andrews JR, Wood R, Bekker LG, Middelkoop K, Walensky RP. Projecting the benefits of antiretroviral therapy for HIV prevention: the impact of population mobility and linkage to care. *J Infect Dis.* **2012**;206(4):543–51.
52. Perriat D, Balzer L, Hayes R, Lockman S, Walsh F, Ayles H, et al. Comparative assessment of five trials of universal HIV testing and treatment in sub-Saharan Africa. *J Int AIDS Soc.* **2018**;21(1):e25048.
53. Larmarange J, Diallo MH, McGrath N, Iwuji C, Plazy M, Thiébaud R, et al. The impact of population dynamics on the population HIV care cascade: results from the ANRS 12249 treatment as prevention trial in rural KwaZulu-Natal (South Africa). *J Int AIDS Soc.* **2018**;21(54):e25128.
54. Camlin CS, Akullian A, Neilands TB, Getahun M, Eyul P, Maeri I, et al. Population mobility associated with higher risk sexual behaviour in Eastern African communities participating in a Universal Testing and Treatment trial. *J Int AIDS Soc.* **2018**;21(54):e25115.
55. Bond V, Ngwenya F, Thomas A, Simuyaba M, Hoddinott G, Fidler S, et al. Spinning plates: livelihood mobility, household responsibility and anti-retroviral treatment in an 1 urban zambian community during the HPTN 071 (PopART) study. *J Int AIDS Soc.* **2018**;21(54):e25117.
56. Hoddinott G, Myburgh H, de Villiers L, Nduvani R, Mantantana J, Thomas A, et al. Households, fluidity, and HIV service delivery in Zambia and South Africa – an exploratory analysis of longitudinal qualitative data from the HPTN 071 (PopART) trial. *J Int AIDS Soc.* **2018**;21(54):e25135.
57. Shabalala FS, Vernooij E, Pell C, Simelane N, Masilela N, Spiegelman D, et al. Understanding reasons for discontinued antiretroviral treatment among clients in test and treat: a qualitative study in Swaziland. *J Int AIDS Soc.* **2018**;21(54):e25120.
58. Fakoya I, Álvarez-Del Arco D, Monge S, Copas AJ, Gennotte A-F, Volny-Anne A, et al. HIV testing history and access to treatment among migrants living with HIV in Europe. *J Int AIDS Soc.* **2018**;21(54):e25123.
59. Clouse K, Fox MP, Mongwenya C, Motlathledi M, Buthelezi S, Bokaba D, et al. "I will leave the baby with my mother": Long-distance travel and follow-up care among HIV-positive pregnant and postpartum women in South Africa. *J Int AIDS Soc.* **2018**;21(54):e25123.
60. Phillips TK, Clouse K, Zerbe A, Orrell C, Abrams EJ, Myer L. Linkage to care, mobility and retention of HIV-positive postpartum women in antiretroviral therapy services in South Africa. *J Int AIDS Soc.* **2018**;21(54):e25114.
61. Vearey J. Moving forward: why responding to migration, mobility and HIV in South(ern) Africa is a public health priority. *J Int AIDS Soc.* **2018**;21(54):e25137.
62. Cassels S, Camlin CS, Seeley J. One step ahead: timing and sexual networks in population mobility and HIV prevention and care. *J Int AIDS Soc.* **2018**;21(54):e25140.
63. Grimsrud A, Barnabas RV, Ehrenkranz P, Ford N. Evidence for scale up: the differentiated care research agenda. *J Int AIDS Soc.* **2017**;20(54):22024.