

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### **Title**

Nuclear Reactions of Arsenic with 190 Mev Deuterons

### **Permalink**

<https://escholarship.org/uc/item/7db85775>

### **Author**

Hopkins Jr, H.H.

### **Publication Date**

2010-02-04

Peer reviewed

UCRL 47  
cy 74/A  
c. 2

UNIVERSITY OF  
CALIFORNIA

Radiation  
Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-47  
c. 2

UCRL


DF

**Special Review of  
Declassified Reports**

Authorized by USDOE JK Bratton

Unclassified TWX P182206Z May 79

**REPORT PROPERLY DECLASSIFIED**



AUG 15 1979

Date

Derivative Classifier

**Lawrence Radiation Laboratory Library**  
University of California, Berkeley



UCRL-47  
Physics-General  
CLASSIFICATION CONTROLLED BY AUTHORITY OF THE DEPARTMENT OF ENERGY BY THE DECLASSIFICATION COMMITTEE

NUCLEAR REACTIONS OF ARSENIC WITH 190 Mev DEUTERONS

By

H. H. Hopkins, Jr., and B. B. Cunningham

Department of Chemistry  
Radiation Laboratory  
University of California  
Berkeley, California

Contract No. W-7405-eng-48

January 21, 1948

Special Review of Declassified Reports  
Authorized by USDOE JK Bratton  
Unclassified TWX P182206Z May 79

REPORT PROPERLY DECLASSIFIED

\_\_\_\_\_  
Special Review Classifier  
Date 8-15-79  
\_\_\_\_\_  
By  
Date 8-20-79

Standard DistributionNo. of Copies

Argonne National Laboratory	1-10
Atomic Energy Commission, Washington	11-13
Brookhaven National Laboratories	14-21
Carbide & Carbon Chemicals Corp. (K-25 Area)	22-23
Carbide & Carbon Chemicals Corp. (Y-12 Area)	24-25
Clinton Laboratories	26-33
General Electric Company	34-37
Hanford Engineer Works	38-39
Iowa State College	40
Los Alamos	41-43
Madison Square Area	44
Monsanto Chemical Company, Dayton	45
National Bureau of Standards	46-47
Patent Advisor	48
Research Division (for NEPA), Oak Ridge	49
Research Division, Oak Ridge	50-64
University of Rochester	65-66
Chemistry Department, Radiation Laboratory	67-71
University of California, Radiation Laboratory	72-77
Declassification Procedure	78-87

---

 Total

87

University of California  
 Radiation Laboratory  
 Berkeley, California

## NUCLEAR REACTIONS OF ARSENIC WITH 190 Mev DEUTERONS

H. H. Hopkins, Jr., and B. B. Cunningham

Department of Chemistry and Radiation Laboratory  
University of California, Berkeley, California

January 21, 1948

Abstract

Isotopes formed in the bombardment of  ${}_{33}\text{As}^{75}$  with 190 Mev deuterons range in atomic number up to twenty-four (or more) mass units lighter than  $\text{As}^{75}$ . Identification of these isotopes was based on chemical behavior and half-life determination. Relative yields have been calculated and show that 30% of the observed reactions produce isotopes within 8 mass units of  $\text{As}^{75}$ .

Three new isotopes have been observed: 9.5 d.  $\text{Se}^{72}$  (K), 44 m.  $\text{Se}^{71}$  ( $\beta^+$ ), and 52 m.  $\text{As}^{71}$  ( $\beta^+$ ).

Contract No. W-7405-eng-48

To be published as a Letter to the Editor of Physical Review.

## NUCLEAR REACTIONS OF ARSENIC WITH 190 Mev DEUTERONS

H. H. Hopkins, Jr., and B. B. Cunningham

Department of Chemistry and Radiation Laboratory  
University of California, Berkeley, California

January 21, 1948


Bombardment of pure arsenic ( ${}_{33}\text{As}^{75}$ ) with 190 Mev deuterons has led to the observation of nuclear reactions in which some of the product nuclei are more than 20 mass units lighter than the target nucleus. Identification of these and of other reaction products was made by chemical separation of the various radioisotopes into elemental fractions, followed by an investigation of the radiations associated with each fraction.

The gross rate of decay of the radioisotopes in each elemental fraction was determined with argon-filled (10 cm. pressure) Geiger-Müller counting tubes of the thin window type (ca. 3 mg/cm.<sup>2</sup> of mica). The half-lives obtained from the resolved decay curves formed the chief basis for identification of known isotopes. In those cases in which sufficient activity was available samples were placed in a crude beta-ray spectrometer and tested for positive or negative beta-particles.

Most of the radiations observed could be assigned to known isotopes; these isotopes are listed in the table below. The identifications of  $\text{Zn}^{72}$ ,  $\text{Cu}^{60}$ ,  $\text{Ni}^{57}$ , and  $\text{Co}^{55}$  are uncertain as they are based on half-lives obtained with low accuracy due to the high level of accompanying activities. The assignments of  $\text{Mn}^{52}$  and  $\text{Cr}^{51}$  are based on reliable half-life determinations from measurements of a small amount of activity in the Mn and Cr fractions. The identification of the remaining isotopes is beyond doubt.

Several new radioactivities have been observed. Mass assignments have been made by demonstration of the following decay chains.





"Milking" of  $\text{As}^{73}$  from  $\text{Se}^{73}$  ( $\beta^+$ , 6.7 h.) have failed to produce measurable activity. This behavior is consistent with the assignment of the 100 d. activity<sup>(2)</sup> to  $\text{As}^{73}$ .

Rough values for the yields of the isotopes have been determined. The figures presented in the table are ratios relative to the yield of  $\text{As}^{72}$ , and are average figures for several bombardments. The calculations are only approximate because of errors due to (a) self absorption and absorption in air, (b) scattering, and (c) unknown counting efficiencies for orbital electron capturing isotopes.

In the column "Reaction  $\Delta$ " are given the differences in mass and charge between the product isotopes and the target nucleus,  ${}_{33}\text{As}^{75}$ . Due to the time required for chemical separations, no activities of less than about 15 minutes half-life would have been seen. Isotopes of half-life greater than about 200 days formed in yield less than 0.05 that of  $\text{As}^{72}$  would not have been detected.

Salient features of the data are: (a) the low yield of selenium isotopes as compared with arsenic isotopes (b) the predominance of neutron deficient isotopes from selenium to gallium and (c) the presence of proton deficient isotopes of the elements below zinc in yields comparable to those of the neutron deficient isotopes.

It will be noted that over 80% of the reactions observed produce isotopes within 8 mass units of  $\text{As}^{75}$ . Since these reactions require excitation of less than  $\sim 75$  Mev, it appears that the high energy deuteron gives up only part of its energy in most of the reactions. This behavior is consistent with the picture of

high energy nuclear reactions recently proposed by Serber. (3)

This work was performed under Contract No. W-7405-Eng-48, with the Atomic Energy Commission in connection with the Radiation Laboratory of the University of California, Berkeley, California. The bombardments were conducted by Dr. Duane Sewell and the 184-inch cyclotron group.

ISOTOPES PRODUCED BY  ${}_{33}\text{As}^{75} + 200 \text{ Mev D}^+$ 

Isotope	Type of Radiation (4)	Half-Life		Yield <sup>(6)</sup> Rel As <sup>72</sup> (7)	Reaction $\Delta$	
		Observed	Literature <sup>(5)</sup>		$\Delta Z$	$\Delta A$
${}_{34}\text{Se}^{75}$	(K, $\gamma$ , $e^-$ )	120 d	160 d <sup>(8)</sup>	0.11 (5%)	+1	+0
${}_{34}\text{Se}^{73}$	$\beta^+$	6.7 h	—	0.09	+1	-2
${}_{34}\text{Se}^{72}$	K	9.5 d	—	0.1 (100%)	+1	-3
${}_{34}\text{Se}^{71}$	$\beta^+$	44 m	—	0.01	+1	-4
${}_{33}\text{As}^{74}$	$\beta^-, \beta^+, \gamma$	19.0 d	16 d	1.1	0	-1
${}_{33}\text{As}^{72}$	$\beta^+$	26 h	26 h	1.00	0	-3
${}_{33}\text{As}^{71}$	$\beta^+$	52 m	—	0.3	0	-4
${}_{32}\text{Ge}^{71}$	$\beta^+, K$	{ 38 h 11.4 d	40 h 11 d	2 ~2 (3%)	-1	-4
${}_{32}\text{Ge}^{68}$	K	250 d	195 d	~5 (100%)	-1	-6
${}_{31}\text{Ga}^{68}$	$\beta^+$	68 m	68 m	0.2	-2	-7
${}_{31}\text{Ga}^{67}$	(K, $\gamma$ , $e^-$ )	83 h	83 h	~3 (5%)	-2	-8
${}_{31}\text{Ga}^{66}$	$\beta^+$	10 h	9.4 h	~0.1	-2	-9
${}_{30}\text{Zn}^{72}$	( $\beta^-, \gamma$ )	~50 h	49 h <sup>(9)</sup>	$\leq 0.001$	-3	-3
${}_{30}\text{Zn}^{69}$	( $\beta^-, \text{I.T.}, \gamma$ )	{ 51 m 14 h	57 m 13.8 h	0.07 0.05	-3	-6
${}_{29}\text{Cu}^{67}$	$\beta^-$	61 h	56 h <sup>(10)</sup>	0.02	-4	-8
${}_{29}\text{Cu}^{64}$	( $\beta^-, \beta^+, K, \gamma$ )	15 h	12.8 h	0.1	-4	-11
${}_{29}\text{Cu}^{61}$	( $\beta^+, K$ )	3.3 h	3.4 h	0.1 (100%)	-4	-14
${}_{29}\text{Cu}^{60}$	( $\beta^+$ )	~20 m	24.5 m <sup>(11)</sup>	0.06	-4	-15
${}_{28}\text{Ni}^{66}$	$\beta^-$	56 h	56 h <sup>(10)</sup>	0.002	-5	-9
${}_{28}\text{Ni}^{65}$	( $\beta^-, \gamma$ )	2.6 h	2.6 h	0.001	-5	-10
${}_{28}\text{Ni}^{57}$	$\beta^+$	34 h	36 h	0.0002	-5	-18
${}_{27}\text{Co}^{61}$	( $\beta^-$ )	1.8 h	1.75 h <sup>(12)</sup>	0.003	-6	-14
${}_{27}\text{Co}^{56-58}$	( $\beta^+, \gamma, K$ )	~80 d	72 d	0.06 (35%)	-6	{ -19 -17
${}_{27}\text{Co}^{55}$	$\beta^+, \gamma$	~16 h	18.2 h	0.003	-6	-20
${}_{26}\text{Fe}^{59}$	( $\beta^-, \gamma$ )	43 d	47 d	0.005	-7	-16
${}_{25}\text{Mn}^{56}$	( $\beta^-, \gamma$ )	2.6 h	2.59 h	0.002	-8	-19
${}_{25}\text{Mn}^{52}$	( $\beta^+, \gamma, K$ )	6 d	6.5 d	0.0002 (40%)	-8	-23
${}_{24}\text{Cr}^{51}$	(K, $\gamma$ , $e^-$ )	26 d	26.5 d	0.005 (5%)	-9	-24

References

- (1) Mann, W. B., Phys. Rev. 54, 649 (1938).
- (2) Elliot, L. G. and M. Deutsch, Phys. Rev. 63, 457 (1943).
- (3) Serber, R., Phys. Rev. 72, 1114 (1947).
- (4) Parentheses signify identification based on half-life determination only.
- (5) Seaborg, G. T., Rev. Mod. Phys., 16, 1-32 (1944).
- (6) The yield of As<sup>72</sup> corresponds to a cross-section of about  $0.02 \times 10^{-24}$  cm.<sup>2</sup>
- (7) Parenthesized figures signify assumed counting efficiencies.
- (8) Kent, C. V., J. M. Cork, and W. G. Wadey, Phys. Rev. 61, 339 (1942).
- (9) Plutonium Project compilation, "Nuclei Formed in Fission", J. Am. Chem. Soc. 68, 2411 (1946).
- (10) Goeckermann, R. H. and I. Perlman, Private Communication (January, 1948).
- (11) Leith, C. E., A. Bratenahl, and J. B. Mayer, Phys. Rev. 72, 732 (1947).
- (12) Farnley, T. J. and B. S. Moyer, Phys. Rev. 72, 82 (1947).



