

Lawrence Berkeley National Laboratory

Recent Work

Title

COMPRESSION EFFECTS IN RELATIVISTIC NUCLEUS-NUCLEUS COLLISIONS

Permalink

<https://escholarship.org/uc/item/7dv7v2m6>

Author

Stock, R.

Publication Date

1982-06-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY

JUN 5 1982

LIBRARY AND
DOCUMENTS SECTION

To be published in Physical Review Letters


COMPRESSION EFFECTS IN RELATIVISTIC NUCLEUS-NUCLEUS COLLISIONS

R. Stock, R. Bock, R. Brockmann, A. Dacal,
J.W. Harris, M. Maier, M.E. Ortiz, H.G. Pugh,
R.E. Renfordt, A. Sandoval, L.S. Schroeder,
H. Stroebele, and K.L. Wolf

June 1982

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, [REDACTED]*



*LBL-14583
c2*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

COMPRESSION EFFECTS IN RELATIVISTIC NUCLEUS-NUCLEUS COLLISIONS*

R. Stock,¹ R. Bock,¹ R. Brockmann,¹ A. Dacal,⁴
J. W. Harris,¹ M. Maier,³ M. E. Ortiz,⁴ H. G. Pugh,²
R. E. Renfordt,³ A. Sandoval,¹ L. S. Schroeder,²
H. Stroebele,¹ and K. L. Wolf¹

Abstract

The negative pion multiplicity is measured for central collisions of ^{40}Ar with KCl at eight energies from 0.36 to 1.8 GeV/nucleon and for ^4He on KCl and ^{40}Ar on BaI_2 at 977 and 772 MeV/nucleon, respectively. A systematic discrepancy with a cascade model calculation which fits proton- and pion-nucleus cross sections but omits potential energy effects is used to derive the energy going into bulk compression of the system. A value of the incompressibility constant of $K = 240$ MeV is extracted in a parabolic form of the nuclear matter equation of state.

*This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

¹Gesellschaft für Schwerionenforschung, D-6100 Darmstadt, West Germany

²Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720

³Universität Marburg, D-355 Marburg, West Germany

⁴Instituto de Fisica, UNAM, Mexico City 21D.F, Mexico

Central collisions of nuclei at relativistic energies are predicted by dynamical models^{1,2,3} to proceed via a compression-expansion cycle of hadronic bulk matter, with a first stage of interpenetration and pile-up of the nucleon densities from target and projectile followed by expansion towards a final freeze-out stage. While these models predict densities several times the nuclear ground state density to be reached at the end of the compression stage there has been no direct experimental evidence of such densities or of any new physical effects resulting from them. The reason is twofold: firstly, the compression is accompanied by heating manifested by chaotic kinetic effects which mask the collective motion⁴; secondly, a great deal of information about the compression stage is lost during expansion through final state interactions, approach to chemical equilibrium, etc. Various approaches are being made to overcome these difficulties but none has so far yielded definitive results. These approaches include study of penetrating particles produced in the early stages of the collision,⁵ and exclusive studies of all produced particles in the hope that analysis in terms of global variables may reduce chaotic effects relative to collective ones.⁶ In the present work we propose the total multiplicity of produced pions as an observable linked to the high density stage of the collisions. We present data on this variable for several colliding nuclear systems, demonstrate that the analysis in terms of nuclear density is consistent, extract values for the bulk compressional energy, and deduce a nuclear matter equation of state.

The reason why such a simple variable as total pion multiplicity can be a good measure of the compression stage is an interplay of three considerations. Firstly, the primary production yield in nucleon-nucleon collisions in the Bevalac energy range (up to 2.1 GeV/nucleon) is a rapidly rising function of energy. Secondly, the relative nucleon-nucleon energy in the nucleus-nucleus collision is degraded during the compression stage and is low by the time expansion begins. Thus, pion production is heavily weighted towards the compression stage. Finally, even though complex interactions during the expansion and freeze-out stages strongly affect differential pion observables such as angular distributions, spectra, etc.⁷, it appears that the total number of pions and delta-resonances, and hence the eventual pion yield, remains approximately constant.

These statements are best understood in the framework of an intranuclear cascade calculation. The cascade code used, that of Cugnon, et al.,³ was chosen because it is extensively described in recent publications and has input data in good agreement with pp, np and pn data that have recently been gathered⁸ for the energy range of the Bevalac. Figure 1 shows the most important results for central collisions ($b_{\max} \leq 2.4$ fm) of $^{40}\text{Ar} + \text{KCl}$ at 977 MeV/nucleon. Figure 1a shows the baryon density in a sphere of 3 fm diameter about the origin of the center-of-mass coordinate system of the participant nucleons, expressed as a ratio to the ground state nuclear density $\rho_0 = 0.17 \text{ fm}^{-3}$. It peaks at about 7 fm/c of elapsed reaction time (in the laboratory system), corresponding to the end of the

compression stage. Fig. 1b shows that about half the baryon-baryon collisions occur during the compression and expansion stages, respectively. Fig. 1c shows that by the end of the compression stage the total number of pions and delta resonances $N_{\Delta+\pi}$ reaches a plateau where it remains approximately constant through expansion and freeze-out. The three elementary processes, Δ decay into $\pi+N$, Δ absorption by the $\Delta+N \rightarrow N+N$ process, and Δ formation both by $N+N \rightarrow N+\Delta$ and $\pi+N \rightarrow \Delta$ stabilize the value of $N_{\Delta+\pi}$ and hence the observed number of pions. Fig. 1d shows the maximum density reached as a function of bombarding energy and also the mean density weighted according to the rate of $\pi+\Delta$ production. The latter, which reflects closely the maximum density, will be used in our subsequent analysis.

In our experiment, we used the Bevalac and the LBL Streamer Chamber facility to study the interaction of ^{40}Ar with KCl at bombarding energies from 0.36 to 1.8 GeV/nucleon. Between 4,000 and 10,000 events were accumulated at each of the energies 360, 566, 722, 977, 1180, 1385, 1609 and 1808 MeV/nucleon in both inelastic and central trigger modes. In addition data were obtained for $^4\text{He} + \text{KCl}$ at 977 MeV/nucleon and $^{40}\text{Ar} + \text{BaI}_2$ at 772 MeV/nucleon. Techniques and a part of the data have been presented elsewhere⁹. For $^{40}\text{Ar} + \text{KCl}$ the central trigger corresponded to a reaction cross section of 180 ± 20 mb, or impact parameters up to $b_{\text{max}} = 2.4$ fm in a geometrical model. This value, when used in the cascade code, enables us to predict successfully many experimental quantities such as the proton participant multiplicity distribution.

Figure 2a shows the negative pion multiplicity $\langle n_{\pi^-}(E) \rangle$ observed in $^{40}\text{Ar} + \text{KCl}$ collisions as a function of laboratory and cm energy. The cascade model prediction is also shown: it is systematically too high, with overestimates ranging from a factor of 4 at 360 MeV/nucleon to 1.35 at 1.8 GeV/nucleon. The overestimate also appears for the other systems studied. For example, consider a sequence of data at approximately constant incident energy, namely $p + ^{48}\text{Ti}$ (730 MeV), $^4\text{He} + \text{KCl}$ (977 MeV/nucleon), $^{40}\text{Ar} + \text{KCl}$ (772 MeV/nucleon) and $^{40}\text{Ar} + \text{BaI}_2$ (772 MeV/nucleon), where the proton data are taken from Cochran, et al.,¹⁰ and the other results from our own measurements. The ratios of cascade model predictions to the data are 1.2, 1.4, 2.1 and 2.4, respectively.

This discrepancy is not due to an inability of the cascade model to deal with pions. Several studies¹¹ have found it to work well for pion production in proton-nucleus collisions, for pion-nucleus scattering and for pion absorption on nuclei. Despite the many ways in which the model appears as a rather crude approach to the real physics it takes into account many extremely important features of the collision process¹². It appears to be a reasonable first order approach to high energy nuclear collisions in a normal nuclear matter medium. The factor that is present in nucleus-nucleus collisions but not important in p-nucleus or π -nucleus collisions, where the cascade model was successful, is the feature of density increase, or compression^{13,14}. We therefore examine, for the four interacting nuclear systems discussed above, the density reached as defined in

Fig. 1d. Such a definition is not useful for the $p + {}^{40}\text{Ti}$ system, but for the other three systems the calculated density is given by $\rho/\rho_0 = 1.9, 3.0, 3.4$, respectively. The overestimate ratios are almost exactly proportional to these densities, at roughly constant incident energy.

Now at a given cm energy E_i consider the measured multiplicity $\langle m_\pi(E_i) \rangle$. The same multiplicity is reached in the cascade model at a lower energy $E_i' < E_i$, as indicated in Fig. 2a. As a crude approximation let us interpret the difference $(E_i - E_i')$ as that part of the cm internal energy per nucleon which goes into potential (compressional) degrees of freedom. This energy becomes inactive as far as pion production is concerned. By the time that it reappears in kinetic energy of the baryons, overall thermalization has reduced the mean energy of binary collisions to a point where pion production is no longer important. Reading the compressional energy per nucleon $E_i^C = E_i - E_i'$ from the graph of Fig. 2a at each experimental point and plotting it against the mean compression ρ/ρ_0 derived from Fig. 1d at the energy E_i , we obtain the nuclear matter equation of state graph¹⁵ shown in Fig. 2b. It is noteworthy that above 1.2 GeV/nucleon where the density in $\text{Ar} + \text{KCl}$ becomes constant, so are the values of E_i^C . Furthermore, the reaction ${}^{40}\text{Ar} + \text{BaI}_2$ at 772 MeV/nucleon, for which the density prediction is $\rho = 3.4 \rho_0$, leads to a value of E_i^C in agreement with that derived from ${}^{40}\text{Ar} + \text{KCl}$ at 977 MeV/nucleon, for which the density is also $3.4 \rho_0$. The values of E_i^C plotted in Fig. 2b are offset by 10 MeV in order to

allow for the ground state binding energy of mass 40 nuclei. The dashed curve is a parabola representing an equation of state without phase transitions, corresponding to a compressibility constant $K = 250$ MeV, i.e., a "hard" equation of state. The value of 200 MeV extracted from excited nuclear energy levels yields the partial curve shown for comparison. The best fit to the data is found at $K = 240$ MeV. The horizontal and vertical error bars shown are statistical only. No estimate of the systematic errors implicit in our treatment has been made, as this should be the subject of more penetrating future theoretical studies, transcending the present simplistic approach.

In summary, we argue that the total pion multiplicity reflects the maximum density reached in central nucleus-nucleus collisions at Bevalac energies. We find that this assumption, taken with densities derived from a cascade calculation, serves to correlate data for different interacting systems at different energies. A systematic discrepancy between cascade calculations and experiment depends on the density and we attribute it to a bulk compressional effect not present in the calculation. A simple analysis of this effect yields an equation of state for nuclear matter which is somewhat harder than expected from low energy nuclear excitations.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

References


1. H. Stöcker, et al., Phys. Rev. Lett. 44, 725 (1980); J. R. Nix and D. Strottman, Phys. Rev. C23, 2548 (1981).
2. Y. Yariv and Z. Fraenkel, Phys. Rev. C20, 2227 (1979) and C24, 488 (1981).
3. J. Cugnon, et al., Nucl. Phys. A352, 505 (1981) and A379, 553 (1982).
4. H. Stöcker, M. Gyulassy, and J. Boguta, Phys. Lett. 103B, 269 (1981).
5. J. W. Harris, et al., Phys. Rev. Lett. 47, 229 (1981); J. Randrup and C. M. Ko, Nucl. Phys. A343, 519 (1980); S. Nagamiya, Proc. 5th High Energy Heavy Ion Study, ed., L. S. Schroeder, LBL-12652, 144 (1981).
6. R. Stock, Proc. 5th High Energy Heavy Ion Study, ed., L. S. Schroeder, LBL-12652, 284 (1981); J. Kapusta and D. Strottman, Phys. Lett. 106B, 33 (1981); M. Gyulassy, K. A. Frankel, and H. Stocker, Phys. Lett. 110B, 185 (1982).
7. K. L. Wolf, et al., submitted to Phys. Rev. C.
8. F. Shimizu, et al., University of Tokyo preprint (1982); G. Alexander, et al., Nucl. Phys. B52, 221 (1973); V. Flaminio, et al., CERN-HERA 79-03 (1979).
9. S. Y. Fung, et al., Phys. Rev. Lett. 40, 292 (1978); A. Sandoval, et al., Phys. Rev. Lett. 45, 874 (1980).
10. D. R. F. Cochran, et al., Phys. Rev. D6, 3085 (1972).

11. M. Sternheim and R. Silbar, Phys. Rev. D6, 3117 (1972); Y. Yariv and Z. Fraenkel, to be published. In calculations with the Cugnon code for pion absorption, quantitative agreement is obtained with the data of K. Nakai, et al., Phys. Rev. Lett. 44, 1446 (1980).
12. J. R. Nix, Prog. Part. Nucl. Phys. 2, 237 (1979).
13. I. Montvay and J. Zimanyi, Nucl. Phys. A316, 490 (1979).
14. H. Stöcker, et al., Phys. Lett. 81B, 303 (1979).
15. A. Bohr and B. R. Mottelson, Nuclear Structure, Vol. I, Benjamin Publ. 1969, p. 257.

Figure Captions


Fig. 1 Results of a cascade calculation for near-central collisions ($b \leq 2.4$ fm) of $^{40}\text{Ar} + \text{KCl}$ at a laboratory bombarding energy of 977 MeV/nucleon. The time dependence of the reactions is shown for (a) the baryon density, relative to the ground state value, (b) the integrated number of baryon-baryon collisions, and (c) the instantaneous number of pions and Δ -resonances. Part (d) shows the maximum baryon density attained as a function of bombarding energy and the mean density weighted by the rate of $\pi^+\Delta$ production.

Fig. 2 (a) The mean π^- multiplicity as a function of bombarding energy for near-central collisions of $^{40}\text{Ar} + \text{KCl}$. The triangles show the data. Open circles show the results of cascade calculations, where vertical lines at $E_{\text{LAB}} \geq 1.4$ GeV/nucleon estimate the uncertainty due to multiple pion production from single NN collisions, not included in the calculation. Horizontal arrows are the values of E_i^C , the compressional energy per nucleon determined at each experimental point. (b) The values of E_i^C as a function of the calculated mean baryon density. Points determined at 1.6 and 1.8 GeV/nucleon are not shown since values of E_C and ρ/ρ_0 are nearly identical to the results at 1.2 GeV/nucleon. The dashed lines represent equations of state with an incompressibility constant K of 250 and 200 MeV, respectively.




XBL 826-1434

Figure 1a,b,c




XBL 826-815

Figure 1d



XBL 826-1436

Figure 2a



XBL 826-1435

Figure 2b

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720