

UC Davis

UC Davis Previously Published Works

Title

Enantioselective Si-H Insertion Reactions of Diarylcarbenes for the Synthesis of Silicon-Stereogenic Silanes.

Permalink

<https://escholarship.org/uc/item/7gp5b5pv>

Journal

Journal of the American Chemical Society, 142(27)

ISSN

0002-7863

Authors

Jagannathan, Jake R
Fettinger, James C
Shaw, Jared T
[et al.](#)

Publication Date

2020-07-01

DOI

10.1021/jacs.0c04533

Peer reviewed



Published in final edited form as:

J Am Chem Soc. 2020 July 08; 142(27): 11674–11679. doi:10.1021/jacs.0c04533.

Enantioselective Si–H Insertion Reactions of Diarylcarbenes for the Synthesis of Silicon-Stereogenic Silanes

Jake R. Jagannathan, James C. Fettinger, Jared T. Shaw*, Annaliese K. Franz*

Department of Chemistry, University of California, One Shields Avenue, Davis, California 95616, United States

Abstract

We report the first example of enantioselective, intermolecular diarylcarbene insertion into Si–H bonds for synthesis of silicon-stereogenic silanes. Dirhodium(II) carboxylates catalyze an Si–H insertion using carbenes derived from diazo compounds where selective formation of an enantioenriched silicon center is achieved using prochiral silanes. Fourteen prochiral silanes were evaluated with symmetrical and prochiral diazo reactants to produce a total of 25 novel silanes. Adding an ortho substituent on one phenyl ring of a prochiral diazo enhances enantioselectivity up to 95:5 er with yields up to 98 %. Using *in situ* IR spectroscopy, the impact of the off-cycle azine formation is supported based on the structural dependence for relative rates of diazo decomposition. A catalytic cycle is proposed where the Si–H insertion step is rate-determining, supported by kinetic isotope effect experiments. Transformations of an enantioenriched silane derived from this method, including selective synthesis of a novel sila-indane, are demonstrated.

The potential utility of chiral-at-silicon compounds incorporated into more complex structures has not been fully understood due to a shortage of synthetic methods. Silicon-stereogenic molecules are rare in number and diversity of structures as compared to carbon. Selected examples to generate silicon-stereogenic silanes include dehydrocouplings,^{1–3} arylation,^{4,5} hydrosilylation,^{6–9} Si–C activation,^{10,11} and reactions controlled by chiral auxiliaries.^{12–14} Brief explorations of the effect of silicon chirality on reaction outcome to produce more complex molecules have occurred,^{15–17} yet remain limited.

The catalytic insertion of carbenes into Si–H bonds to generate organosilicon compounds has been intermittently explored since Doyle's original work in 1988.^{18,19} Methods to date have focused on generation of stereogenic carbon centers using donor/acceptor carbenes (Figure 1A).^{20–23} Si–H insertion to generate stereogenic silicon centers has been demonstrated by Katsuki²⁴ and Iwasa²⁵ using donor/acceptor carbenes (Figure 1A). Donor/donor carbenes (also referred to as diarylcarbenes) are less reactive, with few reports of intermolecular Si–H insertion, and one report of an enantioselective variant using functionalized alkynes as precursors (Figure 1B).^{26–29}

*Corresponding Authors: akfranz@ucdavis.edu or jtshaw@ucdavis.edu.

Author Contributions

All authors have given approval to the final version of the manuscript.

The authors declare no competing financial interest.

Donor/donor carbenes have recently emerged as useful substrates for highly selective C–H insertion reactions.^{30–34} Rhodium carbene complexes demonstrate sufficient reactivity at the insertion carbon despite the presence of two aryl rings for potential stabilization.^{35,36} The Franz group has a long-standing interest in organosilicon chemistry and expertise synthesizing prochiral dihydrosilanes with variation of steric and electronic factors.^{37–39} We envisioned that the additional aryl ring could accomplish an enantioselective intermolecular Si–H insertion process with prochiral silanes (Figure 1C). Herein, we communicate the first enantioselective diarylcarbene Si–H insertion to produce silicon-stereogenic organosilanes.

We began our studies screening metal catalysts [Ru(II), Ir(I), Fe(II), Rh(II) and Cu(II)] with diphenyldiazomethane (**2a**) and prochiral methylphenylsilane (**1a**). For all experiments, inverse addition of **2a** using a syringe pump over 1 hour increased yield of **3a** by preventing azine formation, as seen in previous studies with Si–H insertion methodologies.^{21,40,41} Insertion product **4a** was only observed using dirhodium tetraacetate (Table 1, entry 1).⁴² Based on this lead result, we proceeded to screen chiral dirhodium(II)-based catalysts to identify an enantioselective variant.

A screen of well-studied chiral dirhodium compounds highlighted the reactivity of dirhodium tetracarboxylates. Carboxylate ligands afforded higher yields compared to amido-containing ligands due to the increased electrophilicity of the metal center and resulting carbene (entry 2 vs. entries 4–9).⁴³ Of the catalysts studied, Rh₂(*S*-TCPTTL)₄ provided the highest levels of enantioselectivity when compared to others (entries 5–8 vs 9), which improved further using toluene (entry 10). When an insertion was tested using prochiral diazo compound **3a**, the enantioselectivity of silane product **5a** increased *from 82:18 to 93:7 er*, with a notable increase in yield (76% to 91% yield, entry 10 vs. 14). Manual slow addition of **3a** is seen to form **5a** in comparable yield and selectivity to use of a syringe pump (81%, entry 14 vs. 15), but the use of a syringe pump was continued for further studies. Reducing the reaction temperature below 23 °C did not increase selectivity and no insertion was observed below –30 °C. With optimized conditions in hand, we investigated the effect of substituents with both symmetrical and prochiral diazo compounds.

A series of sterically and electronically varied silanes and symmetrical diazo compounds were evaluated to study the effects on enantioselectivity (Scheme 1, **4a–o**). Electron-rich diazo **2b** was less reactive than **2a** and provides lower yield for **4b** (45%, 81:19 er). Yield improved using diazo **2c** (91%) and lower enantioselectivity was observed for silane **4c** (76:24 er). Electronwithdrawing groups do not strongly affect selectivity (**4d**, 80:20 er), while electron-donating groups on the silane proved deleterious to enantioselectivity (**4e** and **4f**, 50:50 er and 74:26 er respectively). Additional steric bulk on the aryl ring of the silane generally eroded enantioselectivity (**4g–j**) but maintained fair to good yields (55–69%). Selectivity similar to **4a** (82:18) was also observed using 2-naphthyl silane **1g**, with the yield also higher compared to **4h** (69 vs 60%). A slight recovery of enantioselectivity was also observed with **4k** (52%, 82:18 er) compared to **4h** (79:21 er), and comparable to **4a**. Studies with varied alkyl substitution on the silicon center were conducted with diazo **2a**.

Isobutyl-containing **4l** provided the highest enantioselectivity observed using **2a** (86:14 er). However, neopentyl substitution led to loss of enantioselectivity (**4m**, 50:50 er), and cyclohexyl substitution reduced enantioselectivity as well (**4n**, 70:30 er). Lastly, switching to a siloxane also deleteriously affected enantioselectivity while maintaining fair yield (**4o**, 0 60%, 61:39 er). We next turned our focus to insertion of prochiral diazo reactants.

The ability of the ortho substituent on one phenyl ring of the diazo compounds to control enantioselectivity was explored (Scheme 2). Electron-donating substituents lower diastereoselectivity (**5b**, 90:10 dr vs 93:7 dr), but slightly improve enantioselectivity (95:5 vs 93:7 er). With an electron-withdrawing group (**5c**), excellent yield and enantioselectivity is observed (93%, 93:7 er) and diastereoselectivity increased (98:2 vs 93:7 dr). Recent work has noted potential synergistic effects of electronics and ortho substitution on the selectivity of donor/donor carbene chemistry.⁴⁴ Substitution on both phenyl rings was able to achieve excellent yield and good selectivity in **5d** (98% yield, 90:10 dr, 89:11 er), although slightly lower compared to other substitution patterns. The steric and push-pull electronic effects combined improve enantioselectivity compared to symmetrical diazo compounds. These substrates demonstrate that the presence of any ortho-substitution may promote enantioselectivity. Replacing phenyl with a 1-naphthyl group led to decreased diastereoselectivity (**5e**, 85:15 dr) and low enantioselectivity (61:39 er), suggesting other competitive steric effects are present. We sought to explore varied substitution of silanes with prochiral diazo **3a**, given the increase in yield and enantioselectivity compared to using **2a**. Prochiral silanes were tested with diazo **3a** and all demonstrated above 90:10 er for the major diastereomer (Scheme 2, **5f-5j**). Additionally, the reaction performed with 1 gram of **3a** using <1 mol% catalyst affords excellent yield, diastereoselectivity and enantioselectivity (Scheme 2, **5a**). Overall, the data shows that diastereoselectivity is substrate controlled, while enantioselectivity is controlled by the rhodium catalyst. Notably, using a diastereoselective reaction with silane **1c** promotes enantioselectivity with **5f** (94:6 er) compared to **4e** (50:50 er). This result highlights the benefit of using prochiral **3a** to improve enantioselectivity.

A catalytic cycle for the enantioselective Si–H insertion of donor/donor carbenes is proposed (Figure 2A).³⁶ The Rh(II) carboxylate catalyst (**I**) reacts with the diazo compound (**2a** or **3a**) to form complex **II**, which is approached by prochiral silane **1a** to produce the silicon-stereogenic silane and regenerate catalyst. Kinetic isotope effect experiments support the rate-determining insertion step ($k_H/k_D = 1.6$), fitting closely with previous experiments of Si–H insertion with donor/acceptor^{20,41,45} and donor/donor carbenes.²⁹ Off-cycle formation of azine (**6** or **7**) can occur when metal carbene **II** reacts with another diazo reactant. Using *in situ* IR spectroscopy, we determined that the ortho-substituted prochiral diazo **3a** has a significantly reduced rate of azine formation (vs **2a**), which accounts for higher yields of the Si–H insertion products. Relative rates of azine formation ($k_{rel} > 120$) was observed for decomposition of diazo **2a** vs **3a** with Rh₂(*S*-TCPTTL)₄ (in toluene) in the absence of silane (Figure 2A).⁴⁶ Increased yields of Si–H insertion products with ortho-substitution (**5a-i**) are attributed to steric interactions blocking the approach of diazo **3a** to **II**, which reduces off-cycle azine formation. The addition of 4 Å mol sieves reduces off-cycle processes leading to formation of siloxane **8**.⁴² The increase in enantioselectivity observed with prochiral donor/

donor diazo **3** is attributed to a twisting of the ortho substituted aryl ring, which blocks one face of the carbene in **II** to promote selective approach of the silane (Figure 2B).^{44,47,48} There has been a recent report proposing the twisting effect to have electronic contributions similar to that of a donor-acceptor carbene.⁴⁴ However, we hypothesize that the steric effect of an out-of-plane phenyl twist is more significant than the electronic effects, although the analogy is noted.

To demonstrate the utility of enantioenriched silanes, silane **5a** was transformed to silanol, dehydrocoupling, and intramolecular C–H silylation products. Silanes are useful intermediates in stereoselective synthesis, and have versatile reactivity with the remaining Si–H bond.⁴⁹ It is well known that transition metals are capable of oxidative insertion into Si–H bonds with retention of configuration.^{50,51} Pd/C-catalyzed silane hydrolysis affords silanol **9** in 90% yield with 90:10 dr and 93:7 er.^{38,51,52} Under attempted hydrosilylation conditions, an unexpected dehydrocoupling product **10** was isolated in good yield (62%) and 93:7 dr.^{53–55} Exploiting the presence of the ortho-methyl group, diastereoenriched sila-indane **11** was accessed in 90% yield with 90:10 dr using Ir-catalyzed C–H silylation methodology developed by the Hartwig group.^{56,57}

In conclusion, the first example of enantioselective diarylcarbene insertion into Si–H bonds has been accomplished with $\text{Rh}_2(\text{S-TCPTTL})_4$, yielding silicon-stereogenic benzhydryl silanes. While symmetrical diazo compounds demonstrated initial enantioselectivity, using a prochiral diazo reactant dramatically improved the reaction, providing yields up to 98% with 98:2 dr and 95:5 er. A catalytic cycle is proposed and the impact of the off-cycle azine formation is supported based on the structural dependence for relative rates of diazo decomposition. Transformation of the enantioenriched silane affords access to silicon-stereogenic silanol, dehydrocoupling and intramolecular C–H silylation products.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Will Jewell is acknowledged for assistance with acquiring mass spectrometry data. Benjamin Bergstrom, Sarah Dishman and Lucas Souza are acknowledged for helpful discussion.

Funding Sources

We acknowledge the National Science Foundation for support of this research (CHE-1900300, CHE-1363375; AKF) and the dual source diffractometer (CHE-1531193) used in this study. The National Institutes of Health (R01 GM124234; JTS) is also acknowledged for support of this research.

REFERENCES


- (1). Schmidt DR; O'Malle SJ; Leighton JL Catalytic Asymmetric Silane Alcoholysis: Practical Access to Chiral Silanes. *J. Am. Chem. Soc.* 2003, 125 (5), 1190–1191. [PubMed: 12553820] See ref. 2 and 3 for more examples of dehydrocoupling methodology to access stereogenic silanes.
- (2). Li N; Guan B Yttrium–Benzyl Complexes Bearing Chiral Iminophosphonamide Ligands: Synthesis and Application in Catalytic Asymmetric Amine–Silane Dehydrocoupling Reactions. *Adv. Synth. Catal.* 2017, 359 (20), 3526–3531.

- (3). Corriu RJP; Moreau JJE Asymmetric Synthesis at Silicon. *J. Organomet. Chem* 1976, 120 (3), 337–346.
- (4). Koga S; Ueki S; Shimada M; Ishii R; Kurihara Y; Yamanoi Y; Yuasa J; Kawai T; Uchida TA; Iwamura M; Nozaki K; Nishihara H Access to Chiral Silicon Centers for Application to Circularly Polarized Luminescence Materials. *J. Org. Chem* 2017, 82 (12), 6108–6117. [PubMed: 28494580] See ref. 5 for more examples of arylation methodology to access stereogenic silanes.
- (5). Kurihara Y; Nishikawa M; Yamanoi Y; Nishihara H Synthesis of Optically Active Tertiary Silanes via Pd-Catalyzed Enantioselective Arylation of Secondary Silanes. *Chem. Commun* 2012, 48 (94), 11564–11566.
- (6). Igawa K; Yoshihiro D; Ichikawa N; Kokan N; Tomooka K Catalytic Enantioselective Synthesis of Alkenylhydrosilanes. *Angew. Chemie Int. Ed* 2012, 51 (51), 12745–12748. See ref. 7–9 for more examples of hydrosilylation methodology to access stereogenic silanes.
- (7). Hayashi T; Yamamoto K; Kumada M Asymmetric Synthesis of Bifunctional Organosilicon Compounds via Hydrosilylation. *Tetrahedron Lett* 1974, 15 (4), 331–334.
- (8). Ohta T; Ito M; Tsuneto A; Takaya H Asymmetric Synthesis of Silanes with a Stereogenic Centre at Silicon via Hydrosilylation of Symmetric Ketones with Prochiral Diaryl Silanes Catalysed by Binap-Rh Complexes. *J. Chem. Soc. Chem. Commun* 1994, No. 21, 2525–2526.
- (9). Zhao Z-Y; Nie Y-X; Tang R-H; Yin G-W; Cao J; Xu Z; Cui Y-M; Zheng Z-J; Xu L-W Enantioselective Rhodium-Catalyzed Desymmetric Hydrosilylation of Cyclopropenes. *ACS Catal.* 2019, 9 (10), 9110–9116.
- (10). Shintani R Recent Advances in the Transition-Metal-Catalyzed Enantioselective Synthesis of Silicon-Stereogenic Organosilanes. *Asian J. Org. Chem* 2015, 4, 510–514. See ref. 11 for more examples of Si–C activation to access stereogenic silanes.
- (11). Shintani R Recent Progress in Catalytic Enantioselective Desymmetrization of Prochiral Organosilanes for the Synthesis of Silicon-Stereogenic Compounds. *Synlett* 2018, 29 (4), 388–396.
- (12). Bauer JO; Strohmam C Stereocontrol in Nucleophilic Substitution Reactions at Silicon: The Role of Permutation in Generating Silicon-Centered Chirality. *J. Am. Chem. Soc* 2015, 137 (13), 4304–4307. [PubMed: 25803673] See ref. 13 and 14 for more examples of using chiral auxiliaries to access stereogenic silanes.
- (13). Bauer JO; Strohmam C Stereoselective Synthesis of Silicon-Stereogenic Aminomethoxysilanes: Easy Access to Highly Enantiomerically Enriched Siloxanes. *Angew. Chemie Int. Ed* 2014, 53 (3), 720–724.
- (14). Kimiko K; Takayuki K; Masafumi U; Shinji M Asymmetric Synthesis of Organosilicon Compounds Using a C2 Chiral Auxiliary. *Bull. Chem. Soc. Jpn* 1997, 70 (6), 1393–1401.
- (15). Xu L-W; Li L; Lai G-Q; Jiang J-X The Recent Synthesis and Application of Silicon-Stereogenic Silanes: A Renewed and Significant Challenge in Asymmetric Synthesis. *Chem. Soc. Rev* 2011, 40 (3), 1777–1790. [PubMed: 21088772] See ref. 16 and 17 for more examples of transformations using silicon-stereogenic silanes.
- (16). Bauer JO; Strohmam C Recent Progress in Asymmetric Synthesis and Application of Difunctionalized Silicon-Stereogenic Silanes. *Eur. J. Inorg. Chem* 2016, 2016 (18), 2868–2881.
- (17). Shintani R Catalytic Asymmetric Synthesis of Siliconstereogenic Compounds by Enantioselective Desymmetrization of Prochiral Tetraorganosilanes. *Yuki Gosei Kagaku Kyokaiishi/Journal Synth. Org. Chem* 2018, 76 (11), 1163–1169.
- (18). Keipour H; Carreras V; Ollevier T Recent Progress in the Catalytic Carbene Insertion Reactions into the Silicon-Hydrogen Bond. *Org. Biomol. Chem* 2017, 15 (26), 5441–5456. [PubMed: 28639662] See ref 19–25 for work on enantioselective donor/acceptor carbene insertions into Si–H bonds.
- (19). Bagheri V; Doyle MP; Taunton J; Claxton EE A New and General Synthesis of .Alpha.-Silyl Carbonyl Compounds by Silicon-Hydrogen Insertion from Transition Metal-Catalyzed Reactions of Diazo Esters and Diazo Ketones. *J. Org. Chem* 1988, 53 (26), 6158–6160.


- (37). Tran NT; Wilson SO; Franz AK Cooperative Hydrogen-Bonding Effects in Silanediol Catalysis. *Org. Lett* 2012, 14 (1), 186–189. [PubMed: 22148869] See ref. 38 and 39 for our previous work on the synthesis of silanes.
- (38). Diemoz KM; Wilson SO; Franz AK Synthesis of Structurally Varied 1,3-Disiloxanediols and Their Activity as Anion-Binding Catalysts. *Chemistry (Easton)*. 2016, 22 (51), 18349–18353.
- (39). Diemoz KM; Hein JE; Wilson SO; Fettinger JC; Franz AK Reaction Progress Kinetics Analysis of 1,3-Disiloxanediols as Hydrogen-Bonding Catalysts. *J. Org. Chem* 2017, 82 (13), 6738–6747. [PubMed: 28564545]
- (40). Keipour H; Jalba A; Delage-Laurin L; Ollevier T Copper-Catalyzed Carbenoid Insertion Reactions of α -Diazoesters and α -Diazoketones into Si–H and S–H Bonds. *J. Org. Chem* 2017, 82 (6), 3000–3010. [PubMed: 28132507]
- (41). Landais Y; Parra-Rapado L; Planchenault D; Weber V Mechanism of Metal-Carbenoid Insertion into the Si–H Bond. *Tetrahedron Lett.* 1997, 38 (2), 229–232.
- (42). Hydrolysis of Si–H bonds using diRhodium(II) compounds has been previously observed: Doyle MP; High KG; Bagheri V; Pieters R; Lewis PJ; Pearson MM Rhodium(II) Perfluorobutyrate Catalyzed Silane Alcoholysis. A Highly Selective Route to Silyl Ethers. *J. Org. Chem* 1990, 55 (25), 6082–6086.
- (43). Lloret J; Carbó JJ; Bo C; Lledós A; Pérez-Prieto J Influence of the Nature of the Ligand on Dirhodium(II) Carbene Species: A Theoretical Analysis. *Organometallics* 2008, 27 (12), 2873–2876.
- (44). Previous work supports the twisting of the phenyl ring for carbene II caused by ortho substitution, see: Lee M; Ren Z; Musaev DG; Davies HML Rhodium-Stabilized Diarylcarbenes Behaving as Donor/Acceptor Carbenes. *ACS Catal.* 2020, 6240–6247.
- (45). Yasutomi Y; Suematsu H; Katsuki T Iridium(III)-Catalyzed Enantioselective Si–H Bond Insertion and Formation of an Enantioenriched Silicon Center. *J. Am. Chem. Soc* 2010, 132 (13), 4510–4511. [PubMed: 20232868]
- (46). Azines 6 and 7 were formed >90 % yield during kinetic experiments as determined using ¹H NMR spectroscopy with Ph-TMS as an internal standard; therefore the rates of azine formation were assumed to correlate directly with the consumption of diazo compounds 2a and 3a.
- (47). The rationale for the “all up” conformation of ligands is based on: Lindsay VNG; Charette AB Design and Synthesis of Chiral Heteroleptic Rhodium(II) Carboxylate Catalysts: Experimental Investigation of Halogen Bond Rigidification Effects in Asymmetric Cyclopropanation. *ACS Catal.* 2012, 2 (6), 1221–1225.
- (48). The rationale for a twisted phenyl ring in the structure of carbene II is based on X-ray data from: Werlé C; Goddard R; Fürstner A The First Crystal Structure of a Reactive Dirhodium Carbene Complex and a Versatile Method for the Preparation of Gold Carbenes by Rhodium-to-Gold Transmetalation. *Angew. Chemie Int. Ed* 2015, 54 (51), 15452–15456. Refer to ref. 44 and 32 for computational support for this structure.
- (49). Fleming I; Barbero A; Walter D Stereochemical Control in Organic Synthesis Using Silicon-Containing Compounds. *Chem. Rev* 1997, 97 (6), 2063–2192. [PubMed: 11848898]
- (50). The insertion of Rh(I) and Ir(I) is known to occur with retention of stereochemistry: Oestreich M. Chirality Transfer from Silicon to Carbon. *Chem. – A Eur. J* 2006, 12 (1), 30–37. See ref. 51 for more information.
- (51). Corriu RJP; Guérin C; Moreau JJE Stereochemistry at Silicon. *Topics in Stereochemistry*. 1 1, 1984, pp 43–198.
- (52). Hydrolysis of the silane to the silanol using Pd/C/H₂O conditions occurs with inversion of stereochemistry, see ref. 15 and 51 for more information. Hydrolysis of the silane to the silanol under these conditions was performed on all substrates to ensure separation of enantiomers for CSP-HPLC analysis.
- (53). Onopchenko A; Sabourin ET; Beach DL Vinyl- and Allylsilanes from the Rhodium(I)-Catalyzed Hydrosilylation of 1-Alkenes with Trialkylsilanes. *J. Org. Chem* 1984, 49 (18), 3389–3392.
- (54). Kakiuchi F; Nogami K; Chatani N; Seki Y; Murai S Dehydrogenative Silylation of 1,5-Dienes with Hydrosilanes Catalyzed by RhCl(PPh₃)₃. *Organometallics* 1993, 12 (12), 4748–4750.

- (55). Adams C; Riviere P; Riviere-Baudet M; Morales-Verdejo C; Dahrouch M; Morales V; Castel A; Delpech F; Manriquez JM; Chávez I Catalytic Study of Heterobimetallic Rhodium Complexes Derived from Partially Alkylated S-Indacene in Dehydrogenative Silylation of Olefins. *J. Organomet. Chem* 2014, 749, 266–274.
- (56). Cheng C; Hartwig JF Catalytic Silylation of Unactivated C–H Bonds. *Chem. Rev* 2015, 115 (17), 8946–8975. [PubMed: 25714857]
- (57). Karmel C; Chen Z; Hartwig JF Iridium-Catalyzed Silylation of C–H Bonds in Unactivated Arenes: A Sterically Encumbered Phenanthroline Ligand Accelerates Catalysis. *J. Am. Chem. Soc* 2019, 141 (17), 7063–7072. [PubMed: 30971087]

A. Enantioselective donor/acceptor Si–H insertion:



B. Intermolecular donor/donor Si–H insertion:



C. This work: donor/donor Si–H insertion for silicon-stereogenic silanes




Figure 1.
Insertion of carbenes into Si–H bonds.






Figure 2.
A. Proposed catalytic cycle with kinetic isotope effect; B. Diagram of proposed selectivity rationale of donor/donor insertions.


**Scheme 1.**

Scope of enantioselective Si–H insertion with symmetrical diazo compounds^a

^a isolated yields; er determined using ¹H NMR spectroscopy; er Determined using CSP-HPLC analysis of silanol obtained from Pd/C hydrolysis. ^bReaction performed using 1.00 g of **3a** and 0.05 mol % catalyst, at 0.1 M in toluene. ^cdr was determined using ¹⁹F NMR spectroscopy. ^dRelative configuration assigned by X-ray analysis.

**Scheme 2.**Scope of enantioselective Si–H insertion with prochiral diazo reagents^a

^a Isolated yields; dr determined using ¹H NMR spectroscopy; er Determined using CSP-HPLC analysis of silanol obtained from Pd/C hydrolysis. ^bReaction performed using 1.00 g of **3a** and 0.05 mol % catalyst, at 0.1 M in toluene. ^cdr was determined using ¹⁹F NMR spectroscopy. ^dRelative configuration assigned by X-ray analysis.


**Scheme 3.**

Transformations of Si–H insertion products^a



^a Isolated yields; dr determined using ¹H NMR spectroscopy; er determined using CSP-HPLC. ^b Isolated as a 85:15 (major) mixture with the hydrosilylation product. See SI for more information. ^c dr determined using ¹⁹F NMR spectroscopy.

Table 1.

Optimization of donor/donor Si-H insertion



entry	R	Rh_2L_4	% yield ^a	dr ^b	er ^c
1	H	$Rh_2(OAc)_4$	34	-	50:50
2	H	$Rh_2(5R-MEPY)_4$	<5	-	ND
3	H	$Rh_2(S-BTPCP)_4$	<5	-	ND
4	H	$Rh_2(S-DOSP)_4$	65	-	55:45
5	H	$Rh_2(R-PTAD)_4$	67	-	61:39
6	H	$Rh_2(S-PTTL)_4$	62	-	64:36
7	H	$Rh_2(S-BPTTL)_4$	62	-	64:36
8	H	$Rh_2(S-PTV)_4$	67	-	59:41
9	H	$Rh_2(S-TCPTTL)_4$	76	-	76:24
10	H	$Rh_2(S-TCPTTL)_4^d$	78	-	82:18
11	CH ₃	$Rh_2(OAc)_4$	45	55:45	50:50
12	CH ₃	$Rh_2(R-PTAD)_4$	72	60:40	ND
13	CH ₃	$Rh_2(S-DOSP)_4$	75	61:39	ND
14	CH₃	$Rh_2(S-TCPTTL)_4^d$	91	93:7	93:7



entry	R	Rh ₂ L ₄	% yield ^a	dr ^b	er ^c
15 ^e	CH ₃	Rh ₂ (S-TCPTTL) ₄ ^d	81	93:7	93:7

^aNMR yield using Ph-TMS as an internal standard.

^bDetermined using ¹H NMR Spectroscopy.

^cDetermined using CSP-HPLC analysis of silanol obtained from Pd/C hydrolysis; major diastereomer if relevant.

^dToluene used as a solvent.

^eDiazo added via syringe over five minutes.