

# Smoking in pregnancy and risk of cancer among young children: a population-based study

Short title: Smoking in pregnancy and risk of childhood cancer

Julia E Heck <sup>a</sup>, Zuelma A Contreras <sup>a</sup>, Andrew S Park <sup>a</sup>, Tom B. Davidson <sup>b</sup>, Myles Cockburn <sup>c</sup>, Beate Ritz <sup>a</sup>

<sup>a</sup> Department of Epidemiology, Fielding School of Public Health, University of California, Los Angeles, CA, USA;

<sup>b</sup> Department of Pediatrics, Division of Pediatric Hematology/Oncology, David Geffen School of Medicine UCLA, Los Angeles, CA;

<sup>c</sup> Department of Preventive Medicine, USC/Keck School of Medicine, Los Angeles, CA, USA.

## Correspondence:

Julia Heck  
Department of Epidemiology  
650 Charles E. Young Drive, Box 951772  
Los Angeles, CA 90095-1772  
(Tel) 1 310 825 4254  
(Fax) 1 310 206 6039  
Email: jeheck@ucla.edu.

Key words: tobacco, childhood cancer epidemiology, pregnancy, retinoblastoma, brain tumors, leukemia

## Abbreviations:

ALL	Acute Lymphoblastic Leukemia
CBTRUS	Central Brain Tumor Registry of the United States
CI	Confidence Interval
ICCC	International Classification of Childhood Cancer
ICD-O-3	International Classification of Diseases for Oncology
OR	Odds Ratio

Article Category: Short Report

Novelty and Impact: This article is one of the few prospective, population-based studies of maternal pregnancy smoking and childhood cancer, with smoking status collected by medical record review. We report increased risk of retinoblastoma and some childhood brain tumors with maternal smoking during pregnancy.

## **Abstract**

Smoking during pregnancy is a plausible risk factor for childhood cancer, yet previous studies have yielded conflicting results, and few prospective studies have been published.

Data on maternal smoking were obtained from California birth certificates. We linked California birth certificates (births 2007-2011) with California Cancer Registry records for childhood cancer cases (diagnosed January 2007- September 2013) that were ages 5 or younger at diagnosis (N cases=2,021). Controls (N=40,356) were frequency-matched by birth year and randomly selected from birth certificate records. We used unconditional logistic regression to obtain odds ratios (OR) and 95% confidence intervals (CI) to assess the association between smoking during pregnancy and childhood cancer.

We observed positive associations for gliomas (OR=1.8, 95% CI: 1.0-3.4) and retinoblastoma (OR=3.0, 95% CI: 1.4-6.6), particularly bilateral retinoblastoma (OR=9.4, 95% CI 3.6-24.7) with maternal smoking in pregnancy.

Maternal smoking during pregnancy may be a risk factor for retinoblastoma and certain types of childhood brain tumors.

## **Introduction**

Smoking is the leading cause of cancer worldwide, and the constituents of tobacco rapidly cross the placenta and are found in cord blood and fetal tissues.<sup>1</sup> Most of the epidemiologic literature on parental smoking and childhood cancer is comprised of retrospective case-control studies which collected data via parental interview. These studies may be subject to selective participation, recall biases, or a limited willingness to disclose smoking status in pregnancy to research interviewers. There remains a need for prospective population-based studies.

## **Materials and Methods**

In 2007, California began collecting maternal smoking information on the birth certificate. Data on maternal smoking, demographics, and pregnancy and birth-related factors are abstracted by hospital clerks from the medical record, reported by the mother's physician or midwife, or self-reported by mothers.

We linked California birth certificates (births 2007-2011) with California Cancer Registry records for childhood cancer cases (diagnosed January 2007-September 2013) that were ages 5 or younger at diagnosis. Controls were frequency-matched by year of birth. Detailed methods have been previously published.<sup>2</sup> We estimated adjusted odds ratios among cancer types with  $\geq 5$  exposed cases only, and additionally estimated risk for hepatoblastoma because parental smoking was classified as a cause of hepatoblastoma by the International Agency for Research on Cancer.<sup>3</sup> Cancer types were categorized using International Classification of Childhood Cancer (ICCC) codes, however in order to compare results to other studies, we additionally examined risk among all gliomas, high-grade, and low-grade gliomas, defining these subtypes using

International Classification of Diseases for Oncology (ICD-O-3) codes as categorized by the Central Brain Tumor Registry of the United States (CBTRUS).<sup>4</sup> Thus glioma cases overlap with astrocytoma cases. Odds ratios (OR) and 95% confidence intervals were ascertained using unconditional logistic regression, with adjustment for the child's birth year (matching variable), the race/ethnicity of the mother (White non-Hispanic vs. all others), and maternal years of education. Additional adjustment for the number of prenatal care visits and the source of payment for prenatal care (a socioeconomic indicator, as described previously<sup>2</sup>) did not change results and were left out of final models.

Data were collected on smoking status in the three months pre-pregnancy and within each trimester, however smoking status was strongly correlated across trimesters ( $r \sim 0.77-0.91$ ). Thus our analysis examined ever smoking during any point in pregnancy in the main results.

## **Results**

Compared to mothers of cases, a slightly higher proportion of control mothers smoked in the 3 months before pregnancy (3.8% vs. 3.2%). During pregnancy, however, smoking was more common among case mothers (1st trimester, 2.7%; 2<sup>nd</sup> trimester, 2.1%; 3<sup>rd</sup> trimester, 1.8%) than among control mothers (1st trimester, 2.5%; 2<sup>nd</sup> trimester, 1.8%; 3<sup>rd</sup> trimester, 1.7%). Hence, of all the mothers who smoked prior to pregnancy, 16% of case mothers and 34% of control mothers quit smoking during the 1<sup>st</sup> trimester.

We observed positive associations for glial tumors and retinoblastoma with maternal smoking in pregnancy, with a particularly elevated risk estimate for bilateral retinoblastoma (Table 1). With

regards to central nervous system tumor subtypes, gliomas (OR=1.8, 95% confidence interval 1.0- 3.4) were elevated. We also observed an elevated risk estimate for astrocytomas, which made up 123 of the 194 gliomas (adjusted OR=2.0, 95% confidence interval 0.9, 4.2).

We additionally examined the risk from smoking by trimester of pregnancy for gliomas and retinoblastomas. The adjusted estimates for gliomas were fairly similar across trimesters (1<sup>st</sup> trimester, OR=1.7; 2<sup>nd</sup> trimester, OR=1.6; 3<sup>rd</sup> trimester, OR=1.5) whereas adjusted estimates for retinoblastoma appeared to steadily increase with each subsequent trimester (1<sup>st</sup> trimester, OR=3.1; 2<sup>nd</sup> trimester, OR=4.5; 3<sup>rd</sup> trimester, OR=5.0).

The prevalence of smoking was high among mothers of children with both low grade gliomas (4/53 cases, 7.6%) and high-grade gliomas (1/14 cases, 7.1%). Other cancer types were generally null excepting leukemia where point estimates were below unity, with wide confidence intervals.

## **Discussion**

In this records-based prospective study, we observed increased risk for two major types of childhood cancers. Among all cancers, smoking quit rates during the 1<sup>st</sup> trimester were higher among control mothers than among case mothers, and the prevalence of smoking in pregnancy was particularly high for bilateral retinoblastoma (15.4%). Children with bilateral retinoblastoma carry a germline mutation in one allele of a retinal progenitor cell, due to *RBI* inheritance from a parent, *de novo* mutation in parental germline cells, or due to a mutation that occurs during very early embryonic development; these children lose the second allele somatically during the pregnancy. Only a small number of studies have reported on maternal smoking and

retinoblastoma, but several have reported increased effect estimates.<sup>5-7</sup> Of these, one study reported an increased risk for maternal smoking in the first trimester and bilateral retinoblastoma (OR=3.7),<sup>7</sup> and another study an increase with sporadic heritable disease (OR=2.0), which typically presents as bilateral;<sup>5</sup> the third study did not stratify by laterality. Our findings for astrocytoma support those seen in another prospective investigation.<sup>8</sup>

The decreased risk we observed with acute lymphoblastic leukemia (ALL) has been reported in other prospective studies<sup>5,9-11</sup> and may be due to competing risks such as miscarriage or the development of fatal birth defects.<sup>12</sup> If so, our results for all cancer types might be underestimated. Further, this would explain the seemingly counterintuitive results seen in studies which have examined leukemia risk according to cigarettes per day. Several of these studies have reported an higher risk estimate for ALL with light smoking (e.g. <6 or <10 cigarettes per day) but then decreasing point estimates with heavier smoking,<sup>13-18</sup> with reported effect estimates as low as 0.26 for maternal smoking of > 30 cigarettes per day.<sup>16</sup>

Many women attempt to quit smoking during pregnancy, and the smoking patterns exhibited often involve attempts to cut down, quit, followed by relapsing and attempting to quit again. Overall, between 20 and 50% of female smokers attempt to quit or reduce their smoking during pregnancy;<sup>19</sup> of these women, roughly half quit successfully, and the other half change their smoking intensity multiple times.<sup>20</sup> As such, the complex pattern of smoking during pregnancy could not be captured with our study design. Nondifferential misclassification may also be expected due to errors in recall or due to poor willingness to disclose smoking to medical providers, due to the stigma attached to smoking during pregnancy in the United States. Thus

there are concerns for the accuracy of medical-record abstracted smoking status in our data. Validation studies of maternal smoking status on birth certificates suggest that smoking has moderate sensitivity (74-89%) but high specificity (99%),<sup>21, 22</sup> which should be expected to slightly attenuate effect estimates.

California residents smoke less than residents of other US States, with current smoking reported by 17% of adult males and 10% of adult females in 2011.<sup>23</sup> The prevalence of smoking that we observed on birth certificates is lower than that reported among pregnant California women in a 2003 survey (8.7%)<sup>24</sup> although adult smoking rates may dropped since that survey was published, as they have done nationally.<sup>25</sup> In addition, the proportion of births from Hispanic and Asian/Pacific Islander women have risen in California across the last several decades, and those women are less likely to smoke in pregnancy than women of other ethnic groups (3% of Hispanic women and 2% of Asian/Pacific Islander women smoke in pregnancy, as compared to 14% of White non-Hispanics).<sup>26</sup> In the present study, 52% of mothers were Hispanic and 10% were Asian/Pacific Islander.

Interpretation of results and comparison to other studies should be considered in light of the younger age of the children in our study (<6 years at diagnosis). Exposures during pregnancy may be most relevant for childhood cancers which are diagnosed at younger ages. Further, variation in histologic subtype by age for central nervous system and germ cell cancers, among other cancer types, should limit generalizability to older cohorts.<sup>4</sup> In contrast, as the embryonal tumors (neuroblastoma, Wilms tumor, retinoblastoma) occur primarily in younger childhood, most cases diagnosed during the study period would have been included in our analysis.

If a child's cancer is diagnosed during pregnancy or at birth, medical providers might be more vigilant in reporting smoking on the birth certificate. However, this was unlikely to have biased our results because by California law, births must be registered with the State within 10 days of the child's birth. In our data, there were no cases of glioma, astrocytoma, or retinoblastoma diagnosed prior to two weeks of age whose mothers were smokers. A limitation of our study design was that it did not allow exclusion of cases with a family history of disease. We also lacked covariate information on possible confounders such as maternal alcohol consumption, occupational pesticide exposure, and paternal smoking, which could introduce residual confounding.

As we were able to include only 5 years of California births, the sample size of our study was small and data should be viewed as preliminary. Nonetheless our findings corroborate other prospective studies. Further research might ascertain smoking by cotinine assay, and it may be insightful for future meta-analyses on maternal smoking and childhood cancer to compare findings between prospective and retrospective studies.

### **Acknowledgements**

This work was supported by grants from the US National Institutes of Health (grant numbers R21ES018960, R21ES019986, T32CA009142) and the University of California Tobacco-Related Disease Research Program (Award 340657). Dr. Cockburn was supported in part by the National Cancer Institute's Surveillance, Epidemiology and End Results Program under contract

HHSN261201000140C awarded to the Cancer Prevention Institute of California, contract  
HHSN261201000035C awarded to the University of Southern California, and contract  
HHSN261201000034C awarded to the Public Health Institute; and the Centers for Disease  
Control and Prevention's National Program of Cancer Registries, under agreement  
U58DP003862-01 awarded to the California Department of Public Health.

## References

1. Jauniaux E, Burton GJ. Morphological and biological effects of maternal exposure to tobacco smoke on the feto-placental unit. *Early Hum Dev* 2007;83:699-706.
2. Heck JE, Lombardi CA, Cockburn M, Meyers TJ, Wilhelm M, Ritz B. Epidemiology of Rhabdoid Tumors of Early Childhood. *Pediatr Blood Cancer* 2013;60:77-81.
3. IARC (International Agency for Research on Cancer). *A review of human carcinogens, part E: personal habits and indoor combustions*, vol. 100 (E). Lyon, France: WHO Press, 2012, 475p.
4. Ostrom QT, de Blank PM, Kruchko C, Petersen CM, Liao P, Finlay JL, Stearns DS, Wolff JE, Wolinsky Y, Letterio JJ, Barnholtz-Sloan JS. Alex's Lemonade Stand Foundation Infant and Childhood Primary Brain and Central Nervous System Tumors Diagnosed in the United States in 2007-2011. *Neuro-oncology* 2015;16 Suppl 10:x1-x36.
5. Stavrou EP, Baker DF, Bishop JF. Maternal smoking during pregnancy and childhood cancer in New South Wales: a record linkage investigation. *Cancer Causes Control* 2009;20:1551-8.
6. Bunin GR, Meadows AT, Emanuel BS, Buckley JD, Woods WG, Hammond GD. Pre- and postconception factors associated with sporadic heritable and nonheritable retinoblastoma. *Cancer Res* 1989;49:5730-5.
7. Azary S, Ganguly A, Bunin GR, Lombardi C, Park AS, Ritz B, Heck JE. Sporadic retinoblastoma and parental smoking and alcohol consumption before and after conception: a report from the Children's Oncology Group. *PLOS One* 2016;In press.
8. Brooks DR, Mucci LA, Hatch EE, Cnattingius S. Maternal smoking during pregnancy and risk of brain tumors in the offspring. A prospective study of 1.4 million Swedish births. *Cancer Causes Control* 2004;15:997-1005.
9. Cnattingius S, Zack MM, Ekblom A, Gunnarskog J, Kreuger A, Linet M, Adami HO. Prenatal and neonatal risk factors for childhood lymphatic leukemia. *J Natl Cancer Inst* 1995;87:908-14.
10. Podvin D, Kuehn CM, Mueller BA, Williams M. Maternal and birth characteristics in relation to childhood leukaemia. *Paediatr Perinat Epidemiol* 2006;20:312-22.
11. Mucci LA, Granath F, Cnattingius S. Maternal smoking and childhood leukemia and lymphoma risk among 1,440,542 Swedish children. *Cancer Epidemiol Biomarkers Prev* 2004;13:1528-33.
12. Ness RB, Grisso JA, Hirschinger N, Markovic N, Shaw LM, Day NL, Kline J. Cocaine and tobacco use and the risk of spontaneous abortion. *N Engl J Med* 1999;340:333-9.
13. MacArthur AC, McBride ML, Spinelli JJ, Tamaro S, Gallagher RP, Theriault G. Risk of childhood leukemia associated with parental smoking and alcohol consumption prior to conception and during pregnancy: the cross-Canada childhood leukemia study. *Cancer Causes Control* 2008;19:283-95.
14. Shu XO, Ross JA, Pendergrass TW, Reaman GH, Lampkin B, Robison LL. Parental alcohol consumption, cigarette smoking, and risk of infant leukemia: a Childrens Cancer Group study. *J Natl Cancer Inst* 1996;88:24-31.
15. Rudant J, Menegaux F, Leverger G, Baruchel A, Lambilliotte A, Bertrand Y, Patte C, Pacquement H, Verite C, Robert A, Michel G, Margueritte G, Gandemer V, Hemon D, Clavel J. Childhood hematopoietic malignancies and parental use of tobacco and alcohol: the ESCALE study (SFCE). *Cancer Causes Control* 2008;19:1277-90.
16. Sorahan T, McKinney PA, Mann JR, Lancashire RJ, Stiller CA, Birch JM, Dodd HE, Cartwright RA. Childhood cancer and parental use of tobacco: findings from the inter-regional epidemiological study of childhood cancer (IRESCC). *Br J Cancer* 2001;84:141-6.
17. Clavel J, Bellec S, Rebouissou S, Menegaux F, Feunteun J, Bonaiti-Pellie C, Baruchel A, Kebaili K, Lambilliotte A, Leverger G, Sommelet D, Lescoeur B, Beaune P, Hemon D, Lorient MA. Childhood leukaemia, polymorphisms of metabolism enzyme genes, and interactions with maternal tobacco, coffee and alcohol consumption during pregnancy. *Eur J Cancer Prev* 2005;14:531-40.

18. Milne E, Greenop KR, Scott RJ, Bailey HD, Attia J, Dalla-Pozza L, de Klerk NH, Armstrong BK. Parental prenatal smoking and risk of childhood acute lymphoblastic leukemia. *Am J Epidemiol* 2012;175:43-53.
19. Cnattingius S. The epidemiology of smoking during pregnancy: smoking prevalence, maternal characteristics, and pregnancy outcomes. *Nicotine Tob Res* 2004;6 Suppl 2:S125-40.
20. Pickett KE, Wakschlag LS, Dai L, Leventhal BL. Fluctuations of maternal smoking during pregnancy. *Obstet Gynecol* 2003;101:140-7.
21. Roohan PJ, Josberger RE, Acar J, Dabir P, Feder HM, Gagliano PJ. Validation of birth certificate data in New York State. *J Community Health* 2003;28:335-46.
22. Piper JM, Mitchel EF, Jr., Snowden M, Hall C, Adams M, Taylor P. Validation of 1989 Tennessee birth certificates using maternal and newborn hospital records. *Am J Epidemiol* 1993;137:758-68.
23. Centers for Disease Control and Prevention. State Tobacco Activities Tracking and Evaluation system (STATE). Interactive maps: Current smokers - BRFSS: CDC, 2015.
24. California Department of Public Health Tobacco Control Program. Smoking during pregnancy. 2006. Fact sheets and data, available from <http://www.cdph.ca.gov/programs/tobacco/pages/ctcpfactsheets.aspx>.
25. Agaku IT, King BA, Dube SR. Current cigarette smoking among adults - United States, 2005-2012. *MMWR Morb Mortal Wkly Rep* 2014;63:29-34.
26. Tong VT, Dietz PM, Morrow B, D'Angelo DV, Farr SL, Rockhill KM, England LJ. Trends in smoking before, during, and after pregnancy--Pregnancy Risk Assessment Monitoring System, United States, 40 sites, 2000-2010. *MMWR Surveill Summ* 2013;62:1-19.

Table 1. Prevalence of maternal smoking during pregnancy and risk of childhood cancer

Cancer	ICCC or ICD- 0 code	N smoked in pregnancy/total N	% smoked in pregnancy	Crude OR <sup>1</sup>	Adjusted OR (95% CI) <sup>2</sup>
Controls	----	1056/40356	2.6	Referent	Referent
All cancers	011-999	56/2021	2.8	1.1	1.0 (0.8, 1.4)
Leukemias	011-015	8/469	1.7	0.6	0.6 (0.3-1.3)
Acute lymphoblastic leukemia	011	4/282	1.4	0.5	---
Acute myeloid leukemia	012	1/136	0.7	0.3	---
Central nervous system tumors	031-036	13/308	4.2	1.6	1.3 (0.7-2.4)
Gliomas	<sup>3</sup>	12/194	6.2	2.4	1.8 (1.0- 3.4)
Intracranial & intraspinal embryonal tumors	033	2/91	2.2	0.8	---
Neuroblastoma	041	7/238	2.9	1.1	1.2 (0.5-2.5)
Retinoblastoma	050	7/125	5.6	2.2	3.0 (1.4-6.6)
Bilateral	050	6/39	15.4	7.0	9.4 (3.6-24.7)
Wilms tumor	061	6/148	4.1	1.6	1.3 (0.6-3.0)
Hepatoblastoma	071	2/84	2.4	0.9	1.2 (0.3-5.2)
Rhabdomyosarcoma	091	1/58	1.7	0.6	---
Germ cell tumors	101-105	2/75	2.7	1.0	---

<sup>1</sup>Adjusted for the matching variable, birth year.

<sup>2</sup>Adjusted for birth year, maternal race/ethnicity, and maternal years of education.

<sup>3</sup>CBTRUS definition of gliomas, ICD-O-3 codes: 9380-9384, 9391-9460, 9480; low-grade gliomas: 9380, 9382-9384, 9410-9413, 9420, 9424, 9450; high-grade gliomas: 9401, 9440-9442, 9451, 9460, 9480.<sup>4</sup>