

# UC San Diego

## UC San Diego Previously Published Works

### Title

Enzymatic incorporation and utilization of an emissive 6-azauridine

### Permalink

<https://escholarship.org/uc/item/7hn9n05c>

### Journal

Organic & Biomolecular Chemistry, 15(3)

### ISSN

1477-0520

### Authors

Hopkins, Patrycja A

McCoy, Lisa S

Tor, Yitzhak

### Publication Date

2017-01-18

### DOI

10.1039/c6ob02080a

Peer reviewed



Published in final edited form as:

*Org Biomol Chem.* 2017 January 18; 15(3): 684–690. doi:10.1039/c6ob02080a.

## Enzymatic Incorporation and Utilization of an Emissive 6-Aza Uridine


Patrycja A. Hopkins, Lisa S. McCoy, and Yitzhak Tor

<sup>a</sup>Department of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093, USA

### Abstract

To display favorable fluorescent properties, the non-emissive native nucleosides need to be modified. Here we present a motif that relies on conjugating 5-membered aromatic heterocycles (e.g., thiophene) to a 6-azapyrimidine (1,2,4-triazine) core. Synthetic accessibility and desirable photophysical properties make these nucleosides attractive candidates for enzymatic incorporation and biochemical assays. While 6-azauridine triphosphate is known to be poorly tolerated by polymerases in RNA synthesis, we illustrate that conjugating a thiophene ring at position 5 overcomes such limitations, facilitating its T7 RNA polymerase-mediated *in vitro* transcription incorporation into RNA constructs. We further show that the modified transcripts can be ligated to longer oligonucleotides to form singly modified RNAs, as illustrated for an A-site hairpin model RNA construct, which was employed to visualize aminoglycoside antibiotics binding.

### Graphical abstract



### Introduction

The sensitivity and simplicity of fluorescence spectroscopy has made it one of the most invaluable techniques for exploring biomolecules, their structure and dynamics as well as their interactions with diverse ligands.<sup>1, 2</sup> Unlike many proteins, nucleic acids are not intrinsically emissive. The non-emissive pyrimidine and purine nucleobases need to be modified to display usable fluorescent properties.<sup>1</sup> Diverse approaches for emissive nucleosides have indeed been devised over the years,<sup>1, 3–8</sup> but only a small fraction of the analogues made satisfies the criterion of isomorphism.<sup>9–13</sup> Isomorphic nucleobases need to

Correspondence to: Yitzhak Tor.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here].  
See DOI: 10.1039/x0xx00000x

closely resemble their native counterparts with respect to their size, shape and hydrogen bonding pairing abilities. While only few useful purine analogues have been reported,<sup>1, 10, 13–18</sup> the largest number of chemical variations reported are for the pyrimidines, particularly 5-modified uridine analogues.<sup>19</sup> Several groups, including ours, have investigated a variety of isomorphous nucleosides whose design relied on conjugating five-membered rings such as thiophene,<sup>11, 20, 21</sup> furan,<sup>9, 21–27</sup> thiazole<sup>21, 27</sup> and oxazole<sup>9, 24</sup> to the pyrimidine core. The 5-substituted uridine cores extend the chromophore, frequently augmenting the charge separation in the excited state, yielding useful fluorophores (Figure 1a). Interestingly, markedly fewer 6-modified uridines have been utilized in a similar manner.

Modifying the pyrimidine's 5 and 6 positions by judiciously placing conjugated electron poor/electron rich substituents can further enhance the chromophore's "push-pull" interactions, allowing one to potentially tune their photophysical features (Figure 1a). In our primary design, we have replaced the uracil core with the more electron-deficient 6-azauracil, while keeping the electron-rich thiophene ring expecting red-shifted absorption and emission maxima (**1**, Figure 1b). This indeed has been confirmed to be the case.<sup>12</sup> To explore the utility of such a modification within oligonucleotides we have sought to explore the enzymatic incorporation of the corresponding triphosphates. However, enzymatic incorporation of 6-azauridine triphosphate has been reported to be extremely inefficient<sup>28–30</sup> and has been proposed to potentially be due to the nucleoside's favourable *syn* conformation.<sup>31–33</sup>

We have previously observed enhanced T7 RNA polymerase-mediated incorporation of 5-thiopheneUTP (**6**).<sup>11</sup> We therefore hypothesized that conjugating a thiophene ring to 6-azaUTP at position 5 may partially overcome the detrimental effect of the N substituent at the 6 position, potentially providing "polymerase-friendly" substrate (**4**), which will allow the enzymatic fabrication of fluorescent RNAs. Here, we report the synthesis of this triphosphate and its enzymatic incorporation into oligonucleotides (Figure 1b). The impact of the "6-aza modification" on *in vitro* transcription reactions was investigated by comparing the performance of native UTP, 6-azaUTP (**5**) and 5-thiopheneUTP (**6**). We illustrate that the modification at the 5 position indeed significantly improves the enzymatic incorporation with essentially doubled transcription efficiency for 5-thiophene-6azaUTP (**4**) compared to 6-azaUTP (**5**). We further demonstrate that the resulting modified transcript can be ligated to unmodified oligonucleotides to provide longer, fluorescent RNA constructs. As a proof of concept, a singly modified A-site model RNA hairpin is assembled and used to monitor the binding of aminoglycoside antibiotics. The emissive 5-thiophene-6-azauridine can probe antibiotics binding either directly or via FRET-monitored displacement titrations. We conclude that the efficient enzymatic incorporation of this 5-modified 6-azaUTP and its responsiveness to environmental changes make it an attractive reporter nucleoside for exploring RNA and its interactions.

## Results

### Triphosphate synthesis

The parent nucleoside 5-(thiophen-2-yl)-6-azauridine **1** was synthesized as reported.<sup>12</sup> It was then converted to the 5'-triphosphate **4** using POCl<sub>3</sub> and tributylammonium pyrophosphate (Scheme 1), following standard procedures.<sup>11, 34, 35</sup> The analytically pure triphosphate **4** was obtained by ion-exchange chromatography and reverse-phase HPLC. After precipitation with 3% sodium perchlorate in acetone and treatment with Chelex 100, nucleotide **4** was characterized by mass spectrometry and <sup>1</sup>H, <sup>13</sup>C and <sup>31</sup>P NMR spectroscopy (See ESI).

### T7 RNA polymerase-mediated *in vitro* transcription reactions

To evaluate the enzymatic incorporation of the 6-aza-modified nucleosides into short RNA oligonucleotides, transcription reactions with the analytically pure triphosphates **4–6** and T7 RNA polymerase were performed (Figure 2a). A short DNA promoter-template duplex was assembled by annealing the 18-mer consensus promoter with commercial DNA template **A**. All full-length RNA transcripts **7–10** were obtained under standard *in vitro* transcription conditions and purified by denaturing polyacrylamide gel electrophoresis.<sup>11, 14, 23, 36, 37</sup> UV shadowing (254 nm) was used to visualize the transcription reactions. Notably, under long UV illumination (365 nm) transcripts **8** and **10**, which contain fluorescent modifications (lanes 3, 4, 7 and 8), are highly emissive (Figure 3b).

Following extraction and isolation, the incorporation efficiency of triphosphates **4**, **5** and **6** compared to native UTP (lane 1) in the full-length product was determined to be 77±2%, 37±3% and 97±2%, respectively (lanes 3, 6 and 8). Additionally, the aborted transcripts were extracted and characterized by MALDI-TOF MS. Nucleotide **1** produced two main truncated products, where only one is highly fluorescent (lane 3). These were assigned as the emissive 7-mer (5'-pppGCG CCG**1**-3') and the shorter, non-emissive 6-mer (5'-pppGCG CCG-3') (lane 3). The major abortive transcript for **2** was also found to correlate to a 6-mer (lane 6). Interestingly, when only native NTPs were used (lane 1) the truncated construct was characterized as a 5-mer, where the last nucleotide is A instead of C (5'-pppGCG CA-3' instead of 5'-pppGCG CC-3').

Additionally, competitive transcription reactions were performed in the presence of equimolar concentrations of UTP and the modified triphosphates **4** and **5**. The 6-azauridine-containing triphosphates (**4** and **5**) cannot compete with UTP and only the native full length 10-mer is produced. In contrast, 5-thiophene-uridine triphosphate **6**, can favourably compete with UTP.<sup>11</sup> T7 polymerase exhibits approximately 1.5x fold preference for the modified triphosphate (**4**), in which the thiophene ring is conjugated at position 5, over 6-azaUTP (**5**) (lane 7).<sup>‡</sup>

<sup>‡</sup>For full characterization of transcripts **7–10** by MALDI-TOF MS and digestions profiles see supporting information sections S3 and S4.

#### T4 DNA ligase-mediated *in vitro* ligation reaction

A singly modified A-site hairpin was designed for monitoring small molecule binding based on previously designed short RNA constructs (Figure 2b, 5).<sup>23, 38</sup> A singly modified 11-nt RNA construct with **1** at position 7 was transcribed to give transcript **12** using previously described conditions (Figures 2a and S2.2). The modified transcript **12**, serving as an acceptor, was ligated to a 5'-phosphorylated 20-nt donor oligonucleotide containing only native nucleotides (Figure 2b) under standard *in vitro* ligation condition, including T4 DNA ligase, ATP and a bridging DNA splint, and resolved by 20% PAGE (Figure S2.3).<sup>39</sup> The ligation product **14** was extracted and analyzed for its composition by enzymatic digestion in the presence of S1 nuclease and alkaline phosphatase. Integrated reversed-phase HPLC gave the following nucleoside ratio: C/U/A/G/**1** = 10.7/4.0/4.0/12.6/1.1 (Figure 4). Fractions corresponding to each peak were collected and subjected to mass spectrometric analysis, which confirmed the authenticity of the native and modified nucleoside **1**. Thermal melting studies showed that the incorporation of **1** had a small effect on the overall stability of the folded RNA. The unmodified control RNA ligation product **13** displayed a  $T_m$  of 82.8 ( $\pm 0.1$ ) °C, while the  $T_m$  of modified RNA ligation product **14** was 79.3 ( $\pm 0.3$ ) °C (Figure S5).

#### Binding of Aminoglycosides to a Fluorescently Modified A-site

To assess the potential of the 6-aza modified uridine **1** as a reporter nucleoside for small molecule binding assays, we chose the A-site model hairpin **14** as a test system. Note that the modified residue **1** is in close proximity to the binding site but not part of it (Figure 5). A-site hairpin **14** was therefore excited at 320 nm and changes in emission at 420 nm were monitored upon addition of aminoglycosides. Titration curves were generated by plotting fractional fluorescence saturation versus aminoglycoside concentration. Titration of neomycin (**15**) resulted first in slight decrease and then more significant increase of the fluorescence intensity (Figure 6a). The first binding event, corresponding to  $EC_{50}$  value of  $0.9 \pm 0.2 \times 10^{-3}$  mM, in close agreement with literature values, reflects the A-site binding of neomycin.<sup>38</sup> The second binding event, manifested by an increase of emission intensity, corresponding to  $EC_{50}$  value of  $2 \pm 1 \times 10^{-2}$  mM, likely reflects remote neomycin-RNA association. Similarly, when tobramycin (**16**) was titrated into the labelled construct **14**, the fluorescent intensity first slightly increased ( $EC_{50}$   $4 \pm 1 \times 10^{-3}$  mM) and was then followed by more significant emission increase ( $EC_{50}$   $0.24 \pm 0.03$  mM) (Figure S6.2).

#### Displacement of A-site Bound Coumarin-Aminoglycoside with Unlabelled RNA Binders

In our displacement assay format, detection of binding events is dependent on the interactions between two fluorophores, where a labelled RNA serves as a FRET donor and an appropriately labelled aminoglycoside serves as a FRET acceptor.<sup>38</sup> The emissive 5-thiophene-6-azauridine was found to be an excellent FRET donor for 7-diethylaminocoumarin-3-carboxylic acid (Figure 7).

Assembly of the FRET complex was accomplished by adding the coumarin conjugate of 6''-amino-6''-deoxykanamycin **17** ( $0.53 \mu\text{M}$ ) to the modified A-site hairpin **14** ( $1 \mu\text{M}$ ) prior to the addition of an unlabelled aminoglycoside. Titration of unlabelled neomycin (**15**) into a pre-formed FRET complex excited at 320 nm resulted in a slight decrease followed by an

increase of emission intensity of the FRET donor at 420 nm accompanied by a concurrent decrease of the acceptor emission at 470 nm (Figure 6). The corresponding EC<sub>50</sub> values from the displacement assay for each event ( $3.0 \pm 0.1 \times 10^{-3}$  mM and  $3 \pm 1 \times 10^{-2}$  mM, respectively) are similar to the data obtained by the direct binding assay. Note that the first binding event is reflected by changes in the donor's and sensitized acceptor's emission, while the second event is manifested by changes in the donor's emission intensity only, as expected (Figure 6).

## Discussion

Tuning the photophysical properties of emissive nucleoside analogues can be a challenging task. We envision that introducing an electron-withdrawing modification at position 6 can enhance the polarization of 5-modified conjugated biaryl uridines, ideally yielding red-shifted emission maxima and increased fluorescence intensity (Figure 1a). While indeed realized with the 6-azauracil motif, the major challenge remaining was the sluggish recognition of such modified pyrimidines by common RNA polymerases.<sup>28–30</sup> To explore the utility of such a modification within oligonucleotides we have sought to explore the enzymatic incorporation of the corresponding triphosphates, hypothesizing that the conjugated aromatic ring at the 5-position may enhance enzymatic incorporation when compared to 6-azaUTP, a rather poorly incorporating parent nucleotide.

### T7 RNA polymerase-mediated *in vitro* transcription reactions

Although *in vitro* transcription reactions can be very effective, the “pickiness” of RNA polymerase can be challenging. To investigate the “6-aza” effect on *in vitro* transcription we compared the incorporation efficiency of 5-thiophene-6-azauridine (**1**), 6-azauridine (**2**) and 5-thiophene-uridine (**3**).

T7 RNA polymerase accepts the triphosphates **4–6** as substrates and effectively incorporates all three modified nucleosides **1–3**. Triphosphate **6**, which is lacking the nitrogen at position 6 in the uridine core is nearly as effective as the native UTP. 6-substituted uridines are frequently forced into a less favourable *syn* orientation and perform less effectively in transcription reactions.<sup>31–33</sup> Indeed, analyses of the abortive transcripts suggest the enzyme aborts transcription when in a close proximity to a 6-azauridine (**2**) modification, in addition to a lower overall transcription yield. Conjugation of a thiophene ring to the 6-azauridine core, restores “enzyme friendly” behaviour, facilitating the enzymatic generation of full length emissive transcripts, with 2-fold increase of incorporation efficiency compared to 6-azauridine (**2**). Additionally, competitive transcription reactions show, as hypothesized, that conjugating a thiophene ring to 6-azaUTP at position 5 partially overcomes the detrimental effect of the CH/N modification at the 6 position.

### T4 DNA ligase-mediated *in vitro* ligation reaction

A singly modified A-site hairpin was designed for a potential aminoglycoside binding assay (Figure 5). Considering that the construct contains multiple uridine residues and only one modification is desired, its preparation by an enzymatic transcription reaction is not plausible. A potential approach involves transcribing a smaller 11-nt construct with a single

incorporation of **1** (transcript **12**, Figure 2a) and ligating it to a 20-nt oligonucleotide containing native uridines only (donor, Figure 2b).

Using T4 DNA ligase, an ATP-dependent enzyme, which catalyzes the joining of two duplexed oligonucleotides, an efficient ligation reaction between acceptor **12** and native donor yielded the modified A-site construct **14**. Enzymatic digestion followed by reversed-phase HPLC profiling clearly shows that the nucleoside composition matches with the expected 31-nt ligation product **14**'s sequence (Figure 4).

### Binding of Aminoglycosides to a Fluorescently Modified A-site

The bacterial decoding site within the 16S ribosomal RNA, also referred to as the A-site, possesses a small domain comprised of unpaired adenosines, targeted by aminoglycoside antibiotics.<sup>40–42</sup> For biophysical studies, a short RNA construct representing a functional decoding site is commonly used.<sup>43, 44</sup> The dynamics and remodelling of this domain upon aminoglycoside binding can be exploited for fluorescence-based assays as structural changes are likely to translate into changes in the microenvironment of the fluorescent probe.<sup>45–54</sup> Both 2-aminopurine (2-AP) and pyrroloC (pC) have shown promise as such,<sup>55–59</sup> however, both suffer from significantly quenched emission upon incorporation into oligonucleotides.<sup>1, 23</sup> Additionally, the responsiveness of 2-aminopurine is antibiotic-dependent and appears to fail for certain binders, such as neomycin.<sup>57</sup>

We hypothesized that upon titration of aminoglycoside antibiotics (e.g., neomycin, tobramycin) into the labelled RNA construct **14**, the induced conformational and environmental changes would cause changes in emission intensity of the responsive fluorescent nucleoside **1** (Figure 6a), which is sensitive to changes in polarity, viscosity, and pH.<sup>12, 60–62</sup> Indeed, upon titration of neomycin (**15**) and tobramycin (**16**) into the fluorescent A-site hairpin **14** the emission intensity has changed with increasing concentrations of the antibiotic. For neomycin, the fluorescent intensity slightly decreased initially, which was then followed by a significant increase likely reflecting two discrete binding events. Similar behaviour was observed when tobramycin was titrated. The EC<sub>50</sub> values extracted for the first event closely corresponded to previously reported literature values for A-site–aminoglycoside binding.<sup>38</sup> We speculate that the second event potentially reflects remote binding (such as to the hairpin stem or loop present in the RNA construct **14**), which propagates to trigger an environmental change and a photophysical response of the responsive nucleoside.<sup>25, 53, 63, 64</sup>

In a second approach to detecting aminoglycoside–RNA binding, we envisioned an assay where detection is dependent on the interaction between two fluorophores acting as a FRET pair, with 5-thiophene-6-azauridine as a donor and 7-diethylaminocoumarin-3-carboxamide as the acceptor. Following the binding of the singly labelled RNA **14** to a coumarin conjugate of 6''-amino-6''-deoxykanamycin **17** (Figure 5b), which exhibit a high FRET signal, a competitor (e.g., unlabelled neomycin **15**) is titrated in, displacing the fluorescent placeholder (**17**). The intensity of the FRET-acceptor's sensitized emission drops upon titrating neomycin, reflecting its displacement from the A-site. Further titration of neomycin, shows, as before, enhanced emission of the donor azaU nucleotide. This event, likely



reflecting neomycin binding to remote RNA sites (e.g., the tetraloop) nevertheless propagates through the folded structure and alters the probe's immediate environment, yielding enhanced emission.

The A-site and its interactions with aminoglycoside antibiotics have been thoroughly studied.<sup>48–52</sup> Models for specific as well as non-specific binding events have been discussed, including crystal structures showing specifically- and nonspecifically-bound neomycin.<sup>53</sup> In this context, such fluorescence based assays are frequently used for inhibitor discovery and not necessarily for thorough biophysical analyses of RNA–Ligand interactions,<sup>23,38</sup> as the photophysical changes and their physical triggers are rarely fully interpretable, even for established probes.<sup>1</sup>

## Conclusions

The design of emissive nucleosides and their implementation for biochemical and biophysical applications remain an empirical exercise. The relationship between molecular structure and photophysical properties (particularly when it comes to excited state dynamics) is hard to predict and can only be experimentally determined. This issue is particularly challenging for emissive nucleosides since their photophysical features are frequently impacted upon incorporation into oligonucleotides. Here we illustrate that 6-modified uridine analogues represent an intriguing family of emissive nucleosides with potentially valuable applications. Particularly rewarding is the observation that upon modification of the 5-position, such nucleosides can be incorporated into RNA constructs by T7 RNA polymerase-mediated *in vitro* transcription reactions. 5-thiophene-6-azauridine triphosphate (**4**) is a “polymerase-friendly” substrate, and is incorporated twice as efficiently as 6-azauridine triphosphate, the parent nucleotide (77±2% vs 37±3%, respectively), which can then be utilized in RNA–ligand binding assays. We illustrate that this analogue can be used as the sole probe or in conjunction with a FRET acceptor for monitoring RNA binding by aminoglycoside antibiotics, either directly or via displacement assays, respectively. These strategies are not necessarily limited to the A-site and its binders, but can possibly be used for the analysis of other RNA targets with biochemical interest or therapeutic potential.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank the National Institutes of Health for financial support (grant GM 069773), and the Chemistry and Biochemistry MS facility at UCSD.


## Notes and references

1. Sinkeldam RW, Greco NJ, Tor Y. *Chem Rev.* 2010; 110:2579–2619. [PubMed: 20205430]
2. Sameiro M, Goncalves T. *Chem Rev.* 2009; 109:190–212. [PubMed: 19105748]
3. Wilhelmsson LM. *Quarterly reviews of biophysics.* 2010; 43:159–183. [PubMed: 20478079]
4. Tor Y. *Pure Appl Chem.* 2009; 81:263–272.
5. Dodd DW, Hudson RHE. *Mini-Reviews in Organic Chemistry.* 2009; 6:378–391.




6. Wilson JN, Kool ET. *Org Biomol Chem*. 2006; 4:4265–4274. [PubMed: 17102869]
7. Okamoto A, Saito Y, Saito I. *Journal of Photochemistry and Photobiology C*. 2005; 6:108–122.
8. Rist MJ, Marino JP. *Current Organic Chemistry*. 2002; 6:775–793.
9. Greco NJ, Tor Y. *J Am Chem Soc*. 2005; 127:10784–10785. [PubMed: 16076156]
10. Shin D, Sinkeldam RW, Tor Y. *J Am Chem Soc*. 2011; 133:14912–14915. [PubMed: 21866967]
11. Srivatsan SG, Tor Y. *Chem-Asian J*. 2009; 4:419–427. [PubMed: 19072942]
12. Sinkeldam RW, Hopkins PA, Tor Y. *ChemPhysChem*. 2012; 13:3350–3356. [PubMed: 22777983]
13. Rovira AR, Fin A, Tor Y. *J Am Chem Soc*. 2015; 137:14602–14605. [PubMed: 26523462]
14. McCoy LS, Shin D, Tor Y. *J Am Chem Soc*. 2014; 136:15176–15184. [PubMed: 25255464]
15. Liu H, Gao J, Maynard L, Saito YD, Kool ET. *J Am Chem Soc*. 2004; 126:1102–1109. [PubMed: 14746479]
16. Gao J, Liu H, Kool ET. *J Am Chem Soc*. 2004; 126:11826–11831. [PubMed: 15382917]
17. Seela F, Becher G. *Helv Chim Acta*. 2000; 83:928–942.
18. Ward DC, Reich E, Stryer L. *J Biol Chem*. 1969; 244:1228–1237. [PubMed: 5767305]
19. Fin, A., Rovira, AR., Hopkins, PA., Tor, Y. *Modified Nucleic Acids*. Nakatani, K., Tor, Y., editors. Springer International Publishing; Cham: 2016. p. 1-26.
20. Lane RSK, Jones R, Sinkeldam RW, Tor Y, Magennis SW. *ChemPhysChem*. 2014; 15:867–871. [PubMed: 24604669]
21. Wigerinck P, Pannecouque C, Snoeck R, Claes P, De Clercq E, Herdewijn P. *J Med Chem*. 1991; 34:2383–2389. [PubMed: 1652017]
22. Greco NJ, Tor Y. *Tetrahedron*. 2007; 63:3515–3527. [PubMed: 18431439]
23. Srivatsan SG, Tor Y. *J Am Chem Soc*. 2007; 129:2044–2053. [PubMed: 17256858]
24. Srivatsan SG, Tor Y. *Tetrahedron*. 2007; 63:3601–3607. [PubMed: 18431440]
25. Sinkeldam RW, Greco NJ, Tor Y. *ChemBioChem*. 2008; 9:706–709. [PubMed: 18286575]
26. Greco NJ, Sinkeldam RW, Tor Y. *Org Lett*. 2009; 11:1115–1118. [PubMed: 19196162]
27. Gutierrez AJ, Terhorst TJ, Matteucci MD, Froehler BC. *J Am Chem Soc*. 1994; 116:5540–5544.
28. Kahan FM, Hurwitz J. *J Biol Chem*. 1962; 237:3778–3785. [PubMed: 13958267]
29. Shapiro L, August JT. *J Mol Biol*. 1965; 14:214–220. [PubMed: 5883914]
30. Asano S, Kurashin Y, Anraku Y, Mizuno DI. *J Biochem-Tokyo*. 1971; 70:9–20. [PubMed: 4934990]
31. Cody V, Kalman TI. *Nucleosides Nucleotides*. 1985; 4:587–594.
32. Felczak K, Drabikowska AK, Vilpo JA, Kulikowski T, Shugar D. *J Med Chem*. 1996; 39:1720–1728. [PubMed: 8648611]
33. Sochacka E, Czerwinska G, Guenther R, Cain R, Agris PF, Malkiewicz A. *Nucleosides Nucleotides Nucleic Acids*. 2000; 19:515–531. [PubMed: 10843489]
34. Srivatsan SG, Weizman H, Tor Y. *Org Biomol Chem*. 2008; 6:1334–1338. [PubMed: 18385838]
35. Ludwig J. *Acta Biochim Biophys*. 1981; 16:131–133.
36. Milligan JF, Uhlenbeck OC. *Methods Enzymol*. 1989; 180:51–62. [PubMed: 2482430]
37. Milligan JF, Groebe DR, Witherell GW, Uhlenbeck OC. *Nucleic Acids Res*. 1987; 15:8783–8798. [PubMed: 3684574]
38. Xie Y, Dix AV, Tor Y. *J Am Chem Soc*. 2009; 131:17605–17614. [PubMed: 19908830]
39. Lang K, Micura R. *Nat Protoc*. 2008; 3:1457–1466. [PubMed: 18772873]
40. Moazed D, Noller HF. *Nature*. 1987; 327:389–394. [PubMed: 2953976]
41. Tor Y. *Angew Chem Int Ed*. 1999; 38:1579–1582.
42. McCoy LS, Xie Y, Tor Y. *Wiley interdisciplinary reviews RNA*. 2010; 2011(2):209–232.
43. Purohit P, Stern S. *Nature*. 1994; 370:659–662. [PubMed: 8065453]
44. Recht MI, Fourmy D, Blanchard SC, Dahlquist KD, Puglisi JD. *J Mol Biol*. 1996; 262:421–436. [PubMed: 8893854]
45. Zapp ML, Stern S, Green MR. *Cell*. 1993; 74:969–978. [PubMed: 8402886]


46. Mei HY, Mack DP, Galan AA, Halim NS, Heldsinger A, Loo JA, Moreland DW, SannesLowery KA, Sharmeen L, Truong HN, Czarnik AW. *Bioorgan Med Chem*. 1997; 5:1173–1184.
47. Hermann T, Westhof E. *Biopolymers*. 1998; 48:155–165. [PubMed: 10333743]
48. Tor Y, Hermann T, Westhof E. *Chem Biol*. 1998; 5:R277–R283. [PubMed: 9831530]
49. Hermann T, Westhof E. *J Mol Biol*. 1998; 276:903–912. [PubMed: 9566195]
50. Vicens Q, Westhof E. *Biopolymers*. 2003; 70:42–57. [PubMed: 12925992]
51. Vicens Q, Westhof E. *ChemBioChem*. 2003; 4:1018–1023. [PubMed: 14523919]
52. Tor Y. *ChemBioChem*. 2003; 4:998–1007. [PubMed: 14523917]
53. Zhao F, Zhao Q, Blount KF, Han Q, Tor Y, Hermann T. *Angew Chem Int Ed*. 2005; 44:5329–5334.
54. Thomas JR, Hergenrother PJ. *Chem Rev*. 2008; 108:1171–1224. [PubMed: 18361529]
55. Kaul M, Barbieri CM, Pilch DS. *J Am Chem Soc*. 2004; 126:3447–3453. [PubMed: 15025471]
56. Shandrick S, Zhao Q, Han Q, Ayida BK, Takahashi M, Winters GC, Simonsen KB, Vourloumis D, Hermann T. *Angew Chem Int Ed*. 2004; 43:3177–3182.
57. Kaul M, Barbieri CM, Pilch DS. *J Am Chem Soc*. 2006; 128:1261–1271. [PubMed: 16433544]
58. Barbieri CM, Kaul M, Pilch DS. *Tetrahedron*. 2007; 63:3567–3574. [PubMed: 18431442]
59. Marti AA, Jockusch S, Li ZM, Ju JY, Turro NJ. *Nucleic Acids Res*. 2006; 34:e50. [PubMed: 16595796]
60. Pal SK, Zewail AH. *Chem Rev*. 2004; 104:2099–2123. [PubMed: 15080722]
61. Nguyen B, Neidle S, Wilson WD. *Accounts Chem Res*. 2009; 42:11–21.
62. Duboue-Dijon E, Fogarty AC, Hynes JT, Laage D. *J Am Chem Soc*. 2016; 138:7610–7620. [PubMed: 27240107]
63. Cho JH, Rando RR. *Biochemistry-US*. 1999; 38:8548–8554.
64. Blount KF, Zhao F, Hermann T, Tor Y. *J Am Chem Soc*. 2005; 127:9818–9829. [PubMed: 15998086]

**Fig. 1.**


a) Nucleoside design: enhancing the polarization of the conjugated electron poor/electron rich biaryl system. b) Modified uridines: 5-(thiophen-2-yl)-6-azauridine (**1**), 6-azauridine (**2**), 5-(thiophen-2-yl)uridine (**3**) and their corresponding triphosphates **4**, **5**, **6**.




**Fig. 2.** (a) Transcription reaction with T7 promoter and template **A** in the presence of natural NTPs and UTP or modified UTPs **4–6**; resulting in transcripts **7–10** respectively. Transcription reaction with T7 promoter and template **B** in the presence of natural NTPs and UTP or modified **4**, resulting in transcripts **11, 12** respectively. (b) Ligation reaction with T4 DNA ligase, modified acceptor, donor and splint.



**Fig. 3.** Transcription reactions were carried out with T7 promoter and template **A** in the presence of natural NTPs and UTP or modified UTPs **4**, **5**, **6**. Lane 1: control reaction with all natural NTPs; lane 2: competition reaction in the presence of equimolar concentration of UTP and **4**, lane 3: reaction in the presence of **4**, lane 4: reaction in presence of **4** (\*5mM instead of 1mM, as for all other reactions), lane 5: reaction in the presence of equimolar concentration of UTP and **5**, lane 6: reaction in the presence of **5**, lane 7\*\*: reaction in the presence of equimolar concentration of **4** and **5**, lane 8: reaction in the presence of **6**. The reaction was resolved by gel electrophoresis on a denaturing 20% polyacrylamide gel; under UV light at a) 254 nm (on TLC plate); b) 365 nm; c) black and white picture after Stains-All was used to visualize the oligonucleotides.




**Fig. 4.** HPLC profile of enzymatic digestion reactions. (a) A mixture of nucleosides used as a standard and (b) digestion results of ligation product **14**. Digestion of 1–2 nmol of transcript was carried out using S1 nuclease for 2 h at 37 °C and followed by dephosphorylation with alkaline phosphatase for 2 h at 37 °C. The ribonucleoside mixture obtained was analyzed by reverse-phase analytical HPLC with Agilent column eclipse XDB-C18 (5 $\mu$ m, 4.6  $\times$  150mm). Mobile phase: 0–5% acetonitrile (0.1% formic acid) in water (0.1% formic acid) over 10 min and then increase 5–35% acetonitrile (0.1% formic acid) in water (0.1% formic acid) 10–17 min; flow rate 1 mL/min.




**Fig. 5.** (a) Binding of aminoglycosides to a fluorescently modified A-site hairpin **14**. (b) Binding and displacement events of A-site bound coumarin-aminoglycoside with unlabelled RNA binders (where 6-aza modified uridine **1** is a FRET donor and coumarin is a FRET acceptor).






**Fig. 6.** Fluorescence titrations of the A-site construct **14** with neomycin (**15**). (a) Representative emission spectra for titrations of A-site hairpin **14** with neomycin. (b) Curve fit for the titrations of A-site hairpin **14** with neomycin (**15**) with respective values  $EC_{50} 0.9 \pm 0.2 \times 10^{-3}$  mM and  $2 \pm 1 \times 10^{-2}$  mM (fractional saturation vs. antibiotic concentration). (c) Representative emission spectra for displacement studies: titrations of A-site hairpin **14** with neomycin (**15**). (d) Curve fit for the displacement studies: titrations of A-site hairpin **14** with neomycin with respective values  $EC_{50} 3.0 \pm 0.1 \times 10^{-3}$  mM and  $3 \pm 1 \times 10^{-2}$  mM [fractional saturation vs antibiotic concentration; **1**, FRET donor (purple), 6''-amino-6''-deoxykanamycin-coumarin conjugate, FRET acceptor (green)]. Each event was fitted separately (see SI for details).



**Fig. 7.** Normalized absorption (---) and emission (—) spectra of 5-thiophene-6-azauridine in purple (**1**; FRET donor) and coumarin in green (FRET acceptor).

**Scheme 1.**

Synthesis of modified nucleoside triphosphate **4**. Reagents and conditions: i) POCl<sub>3</sub>, (MeO)<sub>3</sub>PO; ii) tributylammonium pyrophosphate, proton sponge, 0–4°C. See Supporting Information for details.